

## Self-adjuvanting polymer-peptide conjugates as therapeutic vaccine candidates against cervical cancer.

Tzu-Yu Liu, Waleed M Hussein, Zhongfan Jia, Zyta M Ziora, Nigel A J McMillan, Michael J. Monteiro, Istvan Toth, and Mariusz Skwarczynski

*Biomacromolecules*, **Just Accepted Manuscript** • DOI: 10.1021/bm400626w • Publication Date (Web): 09 Jul 2013

Downloaded from <http://pubs.acs.org> on July 15, 2013

### Just Accepted

“Just Accepted” manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides “Just Accepted” as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. “Just Accepted” manuscripts appear in full in PDF format accompanied by an HTML abstract. “Just Accepted” manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). “Just Accepted” is an optional service offered to authors. Therefore, the “Just Accepted” Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the “Just Accepted” Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these “Just Accepted” manuscripts.



1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

# Self-adjuvanting polymer-peptide conjugates as therapeutic vaccine candidates against cervical cancer.

*Tzu-Yu Liu,<sup>†</sup> Waleed M. Hussein,<sup>†</sup> Zhongfan Jia,<sup>‡</sup> Zyta M. Ziora,<sup>†</sup> Nigel A. J. McMillan,<sup>□</sup>  
Michael J. Monteiro,<sup>‡</sup> Istvan Toth,<sup>†,§</sup> Mariusz Skwarczynski.<sup>†,\*</sup>*

<sup>†</sup>The University of Queensland, School of Chemistry and Molecular Biosciences, Brisbane, QLD 4072, Australia, <sup>‡</sup>The University of Queensland, Australian Institute for Bioengineering and Nanotechnology, Brisbane, QLD 4072, Australia, <sup>□</sup>Cancer Research Centre, Griffith Health Institute and School of Medical Science, Griffith University, Gold Coast, QLD 4222, Australia, and <sup>§</sup>The University of Queensland, School of Pharmacy, Brisbane, QLD 4072, Australia.

1  
2  
3 **Corresponding Author:**  
4

5  
6 Mariusz Skwarczynski PhD  
7

8  
9 The University of Queensland  
10

11 School of Chemistry and Molecular Biosciences  
12

13  
14 St Lucia, QLD 4072  
15

16  
17 Australia  
18

19  
20 Phone +61 7 33469894  
21

22  
23 Fax: +61 7 33654273  
24

25  
26 e-mail: [m.skwarczynski@uq.edu.au](mailto:m.skwarczynski@uq.edu.au)  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 ABSTRACT  
4  
5

6 Dendrimers are structurally well-defined, synthetic polymers with sizes and physicochemical  
7 properties often resembling those of biomacromolecules (e.g. proteins). As a result they are  
8 promising candidates for peptide-based vaccine delivery platforms. Herein, we established a  
9 synthetic pathway to conjugate a human papillomavirus (HPV) E7 protein-derived peptide  
10 antigen to a star-polymer to create a macromolecular vaccine candidate to treat HPV-related  
11 cancers. These conjugates were able to reduce tumor growth and eradicate E7-expressing TC-  
12 1 tumors in mice after a single immunization, without the help of any external adjuvant.  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30

31 KEYWORDS: peptide subunit vaccine, human papillomavirus, therapeutic anticancer  
32 vaccine, star-polymer, self-adjuvanting, isodipeptide  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## INTRODUCTION


Vaccination is one of the most cost-effective public health interventions, and has been proposed as a promising strategy for the treatment of cancer.<sup>1</sup> Cervical cancer results from the infection of the cervix with human papillomavirus (mainly HPV type 16) which disrupts the cell cycle through interaction between HPV proteins and the host cell.<sup>2</sup> Prophylactic vaccines against HPV have been developed to prevent HPV infection (and by extension, cervical cancer), achieving the best results if administered prior to the commencement of sexual activity.<sup>3</sup> However, a large proportion of the global population is already infected, and many women will continue to become infected with cervical cancer-associated HPV strains until global vaccine coverage is achieved.<sup>4,5</sup> Furthermore, the long-term efficacy of the new HPV vaccines can only be evaluated decades after implementing the existing prophylactic vaccination.<sup>6</sup> Importantly, prophylactic vaccines protect the host against viral infection but are not designed to kill tumor cells. These limitations have prompted research into the development of therapeutic vaccines that treat HPV-related cancer.<sup>2,7-11</sup> Therapeutic vaccine candidates against HPV-related cancers were usually designed to elicit a cytotoxic T-cell response by targeting E7 oncoprotein. This protein is constitutively expressed in cancer cells and is required for the induction and maintenance of tumor growth. T-cells recognize tumor-associated antigen in the form of a defined short peptide sequence (T cell epitope), rapidly and specifically killing target cells that display the antigen.<sup>12</sup> This approach to treating cancer overcomes the many disadvantages of classical chemotherapy, and other therapies like siRNA, which is severely limited by the instability and poor delivery of siRNA molecules into the targeted cells. Therapeutic vaccines use the body's natural fighting system, and have minimal risk of side effects.

1  
2  
3 All reported peptide-based therapeutic HPV vaccine candidates required formulation with  
4 an adjuvant (immunostimulant) to produce the desired immune response.<sup>2</sup> Not only are there  
5 a limited number of adjuvants approved for human use, but many of these adjuvants are  
6 associated with toxicity and low efficacy.<sup>13</sup> Therefore, the discovery and development of  
7 novel adjuvants with potent immunomodulatory activities is of significant importance in the  
8 field of cancer immunotherapy. Specifically, the development of potent and well-defined  
9 adjuvants without adverse toxicity is crucial to the success of therapeutic vaccines.<sup>14</sup>  
10  
11  
12  
13  
14  
15  
16  
17  
18

19 Dendrimers are structurally well-defined, synthetic polymers that can be designed to mimic  
20 the size and physiochemical properties of biomolecules (e.g. proteins).<sup>15</sup> Their hyper-  
21 branched structure allows the attachment and presentation of antigen molecules at the  
22 periphery of the dendrimer, resulting in a highly multifunctional biomacromolecule.<sup>15</sup> It was  
23 demonstrated that a Multiple Antigenic Peptide (MAP) system with a poly-lysine core<sup>16</sup> was  
24 able to generate high antibody titers as a prophylactic vaccine, but only when co-administered  
25 with an adjuvant. Similarly, linear polymers with the peptide epitopes randomly conjugated  
26 along the polymer backbone also generated antibody responses when co-administered with  
27 the toxic complete Freund's adjuvant (CFA).<sup>17</sup> To overcome these disadvantages, well-  
28 defined polymer-based amphiphiles were proposed. Polyacrylate was selected as the most  
29 promising candidate because polyacrylic acid and its esters were easy to synthesize and had  
30 little or no toxicity.<sup>18,19</sup> We recently reported the ability of polyacrylate amphiphilic  
31 dendrimers (decorated with B-cell epitopes on the dendrimer's periphery) to act as a self-  
32 adjuvanting prophylactic vaccine, and induced a protective (humoral) immunity.<sup>20,21</sup>  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
However, while there are a substantial number of self-adjuvanting systems and adjuvants that  
induce humoral immunity, the induction of cellular responses for therapeutic vaccination is  
more challenging. For example, aluminium salts, the only adjuvant that is widely used in

humans, is recognized to essentially only stimulate a humoral immunity.<sup>22,23</sup> Thus, there is an urgent need for alternatives that are able to induce cellular immune responses.

In this work, we hypothesized that a star polymer-based delivery systems might have the adjuvanting potency to generate T-cell mediated immunity against cells that produce the E7 protein, thus could eradicate tumor cells. We designed and synthesized polymer-peptide conjugates with the well-known HPV-16 E7 protein epitope **8Q** (QAEPDRAHYNIVTFCKCD; E7<sub>44-62</sub>) which contains a CTL epitope (CD8<sup>+</sup> cytotoxic T lymphocytes), and T-helper cell (CD4<sup>+</sup>) and B-cell epitopes (Figure 1).<sup>9</sup> This combination makes **8Q** an ideal candidate to stimulate long-term vaccine efficacy to kill cancer cells. We designed a short series of modified epitopes (Figure 1) to avoid uncontrolled aggregation of the cysteine-rich **8Q** epitope. These epitopes were conjugated to the dendritic structure that resembled a 4-arm star-polymer P(<sup>t</sup>BA<sub>37-≡</sub>)<sub>4</sub> (**S4**) and self-assembled into particles under aqueous conditions. All conjugates produced microparticles when formulated in phosphate buffer saline (PBS). A conjugate that contained the **8Q<sub>min</sub>** epitope was able to reduce tumor growth and eradicate E7-expressing TC-1 tumors in mice after a single immunization, without help from any external adjuvant.



**Figure 1.** The amino acid sequence of the original **8Q** epitope and its novel derivatives. The CTL epitope (CD8<sup>+</sup> cytotoxic T lymphocytes), T-helper cell (CD4<sup>+</sup>), and a B-cell epitope are

1  
2  
3 highlighted within the **8Q** sequence. The immunologically redundant pentapeptide sequence  
4  
5 is indicated by italic font.  
6  
7  
8  
9

## 10 11 **EXPERIMENTAL SECTION**

12  
13  
14 **Materials.** Protected L-amino acids were purchased from Novabiochem (Merck Chemicals,  
15 Darmstadt, Germany) and Mimotopes (Melbourne, Australia). pMBHA resin was purchased  
16 from Peptides International (Kentucky, USA). Rink amide MBHA resin, *N,N'*-  
17 dimethylformamide (DMF), dichloromethane (DCM), methanol, *N,N'*-diisopropylethylamine  
18 (DIPEA), piperidine and trifluoroacetic acid (TFA) were obtained from Merck (Hohenbrunn,  
19 Germany). Cu wires were purchased from Aldrich (Steinheim, Germany). *t*-Butyl acrylate  
20 (*t*BA, Aldrich, > 99%) was deinhibited before use by passing through a basic alumina  
21 column, ethyl 2-bromoisobutyrate (EBiB, Aldrich, 98%), tripropargylamine (TPA, Aldrich,  
22 98 %), dimethyl sulfoxide (DMSO, Labscan, AR grade), *N,N,N',N',N''*-  
23 pentamethyldiethylenetri-amine (PMDETA, Aldrich, 99%), copper(I) bromide (Cu(I)Br, MV  
24 Laboratories, INC., 99.999%), copper(II) bromide (CuBr<sub>2</sub>, Aldrich, 99%) were used as  
25 recieved. Propargyl nitroxide was synthesized according the previous procedure.<sup>24</sup> 1-(1*H*-  
26 Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) and 2-(1*H*-7-  
27 azabenzotriazol-1-yl)-1,1,3,3-tetramethyl uronium hexafluorophosphate methanaminium  
28 (HATU) were purchased from Mimotopes (Melbourne, Australia). HPLC grade acetonitrile  
29 was obtained from Labscan (Bangkok, Thailand). All other reagents were obtained at the  
30 highest available purity from Sigma-Aldrich (Castle Hill, NSW, Australia).  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51

52  
53 Microwave assisted Boc-SPPS was carried out using a SPS mode CEM Discovery reactor  
54 (CEM Corporation, Matthews, NC, USA). Anhydrous hydrofluoric acid (HF) was supplied  
55 by BOC gases (Sydney, NSW, Australia). A Kel-F HF apparatus (Peptide Institute, Osaka,  
56  
57  
58  
59  
60



1  
2  
3 Japan) was used for HF cleavage. ESI-MS was performed using a Perkin-Elmer-Sciex  
4 API3000 instrument with Analyst 1.4 software (Applied Biosystems/MDS Sciex, Toronto,  
5 Canada). <sup>1</sup>H nuclear magnetic resonance (<sup>1</sup>H NMR) spectra were recorded with a Bruker  
6 Avance 300 MHz spectrometer (Bruker Biospin, Germany). Analytical RP-HPLC was  
7 performed using an Agilent instrument with a 1 mL/min flow rate and detection at 214 nm.  
8 Separation was achieved using a 0-100% linear gradient of solvent B over 40 min with 0.1%  
9 TFA/H<sub>2</sub>O as solvent A and 90% MeCN/0.1% TFA/H<sub>2</sub>O as solvent B on either a Vydac  
10 analytical C4 column (214TP54; 5 μm, 4.6 mm x 250 mm) or a Vydac analytical C18 column  
11 (218TP54; 5 μm, 4.6 mm x 250 mm). Preparative RP-HPLC was performed on Shimadzu  
12 (Kyoto, Japan) instrumentation (either LC-20AT, SIL-10A, CBM-20A, SPD-20AV, FRC-  
13 10A or LC-20AP x 2, CBM-20A, SPD-20A, FRC-10A) in linear gradient mode using a 10-  
14 20 mL/min flow rate, with detection at 230 nm. Separations were performed with solvent A  
15 and solvent B on either a Vydac preparative C4 column (214TP1022; 10 μm, 22 mm x 250  
16 mm) or a Vydac preparative C18 column (218TP1022; 10 μm, 22 mm x 250 mm). Particle  
17 size was measured by dynamic light scattering (DLS) using a Malvern Zetasizer Nano Series  
18 with DTS software. Sizes were analyzed using a non-invasive backscatter system.  
19 Multiplicate measurements were performed at 25 °C with scattering angle of 173° using  
20 disposable cuvettes and the number-average hydrodynamic particle diameter are reported.  
21 The average particle diameter in PBS was determined with a ZEISS LSM 510 META  
22 confocal microscope. The software used for image acquisition was AIM 4.2 (Carl Zeiss, Ltd)  
23 and Carl Zeiss Zen 2009 was used for image analysis.

24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51 **Synthesis of 8Q Peptide.** 8Q epitope (QAEPDRAHYNIVTFCKCD; E7<sub>44-62</sub>) was  
52 synthesized on *p*MBHA resin (substitution ratio: 0.45 mmol/g, 0.2 mmol scale, 0.44g) using  
53 HBTU/DIPEA Boc-chemistry by Microwave-Assisted Solid-Phase Peptide Synthesis (MW-  
54 SPSS). The temperature was set at 70 °C (at 20 Watt, 10 min) for amino acid coupling except  
55  
56  
57  
58  
59  
60

1  
2  
3 for His, Cys and Asp, which were coupled at 50 °C (at 20 Watt, 15 min). Each amino acid  
4  
5 coupling cycle consisted of Boc-deprotection with neat TFA (2 × 1 min, at rt), a 1 min DMF  
6  
7 flow wash, followed by 10 min coupling with the preactivated amino acid. Amino acid  
8  
9 activation was achieved by dissolving Boc-amino acid (0.84 mmol, 4.2 equiv.), in a 0.5 M  
10  
11 HBTU/DMF solution (1.6 mL, 0.8 mmol, 4.0 equiv.) followed by the addition of DIPEA  
12  
13 (0.22 mL, 1.24 mmol, 6.2 equiv.). Amino acids were preactivated for 1 min prior to their  
14  
15 addition to the resin. For peptides containing His(DNP) residues, the DNP (2,4-dinitrophenyl)  
16  
17 group was cleaved by treating the resin with 20% (v/v) β-mercaptoethanol and 10% (v/v)  
18  
19 DIPEA in DMF for 2 × 1 h treatments prior to peptide cleavage. Upon completion of  
20  
21 synthesis and removal of the dinitrophenyl (DNP) protecting group, the resin was washed  
22  
23 with DMF, DCM, and MeOH, then dried (vacuum desiccator). The peptide was cleaved from  
24  
25 the resin using HF, with *p*-cresol and *p*-thiocresol as scavengers. The cleaved peptide was  
26  
27 precipitated, filtered, and washed thoroughly with ice-cold Et<sub>2</sub>O and dissolved in 50%  
28  
29 MeCN/0.1% TFA/H<sub>2</sub>O. After lyophilization, the crude peptide was obtained as an amorphous  
30  
31 powder. The product was purified by preparative RP-HPLC on a C18 column with a solvent  
32  
33 gradient of 0-50% solvent B over 50 min. HPLC analysis (C18 column):  $t_R = 19.18$  min,  
34  
35 purity > 95%. Yield: 12%, ESI-MS:  $m/z$  2213.2 (calc 2213.5) [M+H]<sup>+</sup>; 1107.7 (calc 1107.2)  
36  
37 [M+2H]<sup>2+</sup>; 738.5 (calc 738.5) [M+3H]<sup>3+</sup>; MW 2212.49.

38  
39  
40  
41  
42  
43 **Synthesis of Azidoacetic Acid (N<sub>3</sub>CH<sub>2</sub>CO<sub>2</sub>H).** Azidoacetic acid was synthesized using a  
44  
45 similar method to published procedure.<sup>25</sup> Sodium azide (6.0 g, 92.3 mmol, 3.0 equiv.) was  
46  
47 dissolved in H<sub>2</sub>O (10 mL) and bromoacetic acid (4.3 g, 30.8 mmol, 1.0 equiv.) was added.  
48  
49 The reaction was stirred continuously in an ice bath for 24 h and subsequently acidified with  
50  
51 32% HCl (10 mL). The product was then extracted with Et<sub>2</sub>O (4 x 50 mL), dried over  
52  
53 anhydrous MgSO<sub>4</sub> and the solvent was evaporated under vacuum. The final product was  
54  
55 obtained as a colorless oil (2.95 g, 95%) after prolonged evaporation under vacuum to  
56  
57  
58  
59  
60

1  
2  
3 remove organic solvent and the last traces of water.  $^1\text{H}$  NMR (300 MHz,  $\text{CDCl}_3$ )  $\delta$  10.60 (br  
4 s, 1H, OH), 3.98 (s, 2H,  $\text{CH}_2$ ).  $^{13}\text{C}$  NMR (100 MHz,  $\text{CDCl}_3$ )  $\delta$  50.0, 173.7.

7  
8 **Synthesis of Boc-Thr(Fmoc-Val)-OH.** The isopeptide unit was synthesized according to  
9 published procedure.<sup>26-28</sup>

11  
12 **Synthesis of 1,3-di(hydroxymethyl)-5-(prop-2-ynoxy)benzene.** This compound was  
13 prepared as previously described.<sup>29,30</sup>  $^1\text{H}$ -NMR (300 MHz,  $\text{CD}_3\text{OD}$ ):  $\delta$  2.52 (t,  $J = 2.4$  Hz,  
14 1H,  $\text{CH}_2\text{C}\equiv\text{CH}$ ), 4.68 (bs, 4H,  $2\times\text{CH}_2\text{OH}$ ), 4.71 (d,  $J = 2.4$  Hz, 2H,  $\text{CH}_2\text{C}\equiv\text{CH}$ ), 6.92 (bs, 2H,  
15 ArH), 7.00-6.98 (m, 1H, ArH).

17  
18  
19  
20  
21 **Synthesis of polyacrylate  $\text{P}(\text{BA}_{37-\equiv})_4$ .** The alkyne-functionalized 4-arm poly(*t*-butyl  
22 acrylate) star (**S4**) was synthesized by atom-transfer radical polymerization. Star-polymer  
23  $\text{P}(\text{BA}_{37-\text{Br}})_4$ <sup>21</sup> ( $M_n=14900$ ,  $\text{PDI}=1.05$ ; 0.700 g,  $3.68 \times 10^{-5}$  mol), propargyl nitroxide (0.078  
24 g,  $3.72 \times 10^{-4}$  mol), *N,N,N',N'',N'''*-pentamethyldiethylenetriamine (PMDETA, 0.043 g,  $2.48 \times$   
25  $10^{-4}$  mol) were dissolved in 6 mL of DMSO/toluene mixture (1:2, vol). The mixture was  
26 purged with argon for 20 min before the addition of CuBr (0.033 g,  $2.32 \times 10^{-4}$  mol). The  
27 reaction was stirred at R.T. for 30 min, then diluted by THF and passed through an activated  
28 basic alumina column. The solution was concentrated and the polymer was recovered by  
29 precipitation into a MeOH/water (50:50 vol) mixture and then filtered. The polymer was  
30 further purified by preparative GPC. GPC:  $M_n=15730$ ,  $\text{PDI}=1.06$ .  $^1\text{H}$  NMR ( $\text{CDCl}_3$ ) :  $\delta$  1.11-  
31 1.21 (m, 15H, methyl group of core and TEMPO ring), 1.45 (b, methyl protons of *t*-BA  
32 repeat units), 1.81, 2.21 (b, methylene and methine protons of polymer backbone), 2.40 (s,  
33 4H,  $(-\text{OCH}_2-\text{C}\equiv\text{CH})_4$ ), 3.76, (b, 4H, methine proton on TEMPO ring), 3.80-4.25, (b, 20H,  
34 (methylene protons of the core, methine proton of *t*-BA unit close to the alkoxyamine  
35 linkage, methylene protons of propargyl group).

36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55 **Synthesis of N-terminus Model Azide.** N-terminus model azide ( $\text{N}_3\text{CH}_2\text{CO-Phe-Cys-NH}_2$ )  
56 was synthesized by manual stepwise SPPS on rink amide MBHA resin (substitution ratio:  
57  
58  
59  
60

0.59 mmol/g, 0.2 mmol scale, 0.34 g) using HBTU/DIPEA Fmoc-chemistry. Amino acid activation was achieved by dissolving Fmoc-amino acid (0.84 mmol, 4.2 equiv.), in 0.5 M HBTU/DMF solution (1.6 mL, 0.8 mmol, 4.0 equiv.) followed by the addition of DIPEA (146  $\mu$ L, 0.84 mmol, 4.2 equiv.). Coupling cycle consisted of Fmoc deprotection with 20% of piperidine in DMF (twice, 10 and 20 min), a 1 min DMF flow-wash, followed by coupling with 4.2 equiv. of preactivated Fmoc-amino acids ( $2 \times 2$  h). The attachment of azidoacetic acid (4.2 equiv.) was achieved using HBTU (3 equiv.)/DIPEA (4.2 equiv.) at room temperature ( $2 \times 2$  h). Upon completion of synthesis, the resin was washed with DMF, DCM, and MeOH, then dried (vacuum desiccator). The cleavage of model azide was carried out by stirring the resin in the solution of TFA (99%)/triisopropylsilane/water (95:2.5:2.5) for 4 h. The cleaved peptide was precipitated, filtered, and washed with ice-cold Et<sub>2</sub>O. After lyophilization the crude peptide was obtained as an amorphous powder. The product was purified by preparative RP-HPLC on C18 column with solvent gradient 0-50% solvent B over 50 min. HPLC analysis (C18 column):  $t_R = 18.4$  min, purity > 95%. Yield: 96% ESI-MS:  $m/z$  702.1 (calc 701.8) [2M+H]<sup>+</sup>; 351.7 (calc 351.4) [M+H]<sup>+</sup>; MW 350.40.

**Synthesis of N-terminus 8Q Azide.** 8Q azide peptide epitope (N<sub>3</sub>CH<sub>2</sub>CO-QAEPDRAHYNIVTFCKCD-NH<sub>2</sub>) was synthesized according to the above procedure. Fmoc deprotection of Thr, Val and Ile were performed with 2% of 1,8-Diazabicycloundec-7-ene (DBU) in DMF (twice, 5 and 10 min) instead of 20% piperidine in DMF. HPLC analysis (C18 column):  $t_R = 17.5$  min, purity > 95%. Yield: 17%. ESI-MS:  $m/z$  1148.8 (calc 1148.8) [M+2H]<sup>2+</sup>; 766.2 (calc 766.2) [M+3H]<sup>3+</sup>; MW 2295.54.

**Synthesis of N-terminus 8Q<sub>min</sub> Azide.** 8Q<sub>min</sub> azide peptide epitope (N<sub>3</sub>CH<sub>2</sub>CO-QAEPDRAHYNIVTF-NH<sub>2</sub>) was synthesized according to the above procedure. Fmoc deprotection of Thr, Val, and Ile were performed with 2% DBU in DMF (twice, 5 and 10 min) instead of 20% piperidine in DMF. HPLC analysis (C18 column):  $t_R = 18.3$  min, purity

1  
2  
3 > 95% (Figure S1). Yield: 42%. ESI-MS:  $m/z$  1743.3 (calc 1743.8)  $[M+H]^+$ ; 872.3 (calc  
4  
5 872.4)  $[M+2H]^{2+}$ ; MW 1742.85.

### 7 **Synthesis of N-terminus 8Q<sub>Ser</sub> Azide.**

9  
10 Method A: 8Q<sub>Ser</sub> azide peptide epitope (N<sub>3</sub>CH<sub>2</sub>CO-QAEPDRAHYNIVTFSSKSD-NH<sub>2</sub>) was  
11 synthesized according to the above procedure. Fmoc deprotection of Thr, Val, and Ile were  
12 performed with 2% DBU in DMF (twice, 5 and 10 min) instead of 20% piperidine in DMF.  
13  
14 HPLC analysis (C18 column):  $t_R$  = 16.66 min, purity > 95%. Yield: 14 %. ESI-MS:  $m/z$   
15  
16 1124.8 (calc 1124.7)  $[M+2H]^{2+}$ ; 750.2 (calc 750.1)  $[M+3H]^{3+}$ ; MW 2247.34.  
17  
18

19  
20 Method B: The isopeptide 8Q<sub>Ser</sub> azide peptide epitope was synthesized according to the  
21 above procedure. Boc-Thr(Fmoc-Val)-OH was used to introduce Thr and Val. Fmoc  
22 deprotection were performed only with 20% piperidine in DMF (twice, 10 and 20 min).  
23  
24 Isopeptide 8Q<sub>Ser</sub> azide was cleaved from resin ( $t_R$  = 15.82 min, C18), lyophilized and  
25 dissolved in 50% aqueous MeCN. Ammonia solution (25%) was added dropwise until the pH  
26 was 8-9. The reaction mixture was stirred for 2 h to form the desired migrated 8Q<sub>Ser</sub> azide.  
27  
28 HPLC analysis (C18 column):  $t_R$  = 16.66 min, purity > 95% (Figure S1). Yield: 55%. ESI-  
29  
30 MS:  $m/z$  1124.8 (calc 1124.7)  $[M+2H]^{2+}$ ; 750.2 (calc 750.1)  $[M+3H]^{3+}$ ; MW 2247.34.  
31  
32  
33

34  
35 **Synthesis of N-terminus 8Q<sub>Lys</sub> Azide.** 8Q<sub>Lys</sub> azide peptide epitope (N<sub>3</sub>CH<sub>2</sub>CO-  
36  
37 QAEPDRAHYNIVTFSSKKK-NH<sub>2</sub>) was synthesized according to the above procedure.  
38  
39 Fmoc deprotection of Thr, Val, and Ile were performed with 2% DBU in DMF (twice, 5 and  
40  
41 10 min) instead of 20% piperidine in DMF. HPLC analysis (C18 column):  $t_R$  = 15.9 min,  
42  
43 purity > 95% (Figure S1). Yield: 18% ESI-MS:  $m/z$  1172.3 (calc 1172.3)  $[M+2H]^{2+}$ ; 781.9  
44  
45 (calc 781.9)  $[M+3H]^{3+}$ ; 586.9 (calc 586.6)  $[M+4H]^{4+}$ ; 469.8 (calc 469.5)  $[M+5H]^{5+}$ ; MW  
46  
47 2342.61.  
48  
49  
50  
51  
52

53  
54 **Copper-Catalyzed Alkyne-Azide Cycloaddition (CuAAC) Reaction - model studies.** To  
55  
56 determine the optimal conditions to avoid disulfide formation during the CuAAC reaction,  
57  
58  
59  
60

1  
2  
3 model azide and 1,3-di(hydroxymethyl)-5-(prop-2-ynyloxy)benzene were employed. The  
4  
5 reactions were conducted by using different copper sources (Cu wires, CuSO<sub>4</sub>, CuI), with or  
6  
7 without adding reducing agent (ascorbic acid, tris(2-carboxyethyl)phosphine), different  
8  
9 solvents (solvent B (90% MeCN/0.1% TFAH<sub>2</sub>O), DMF), and stirring at 50 °C or room  
10  
11 temperature. Briefly, model azide (1 equiv.) and 1,3-di(hydroxymethyl)-5-(prop-2-  
12  
13 ynyloxy)benzene (1 equiv.) were dissolved in DMF (1 mL), and the copper source added.  
14  
15 The air in the reaction mixture was removed either by nitrogen or argon bubbling. The  
16  
17 reaction mixture was covered and protected from light with aluminum foil and stirred at  
18  
19 either 50 °C or room temperature under nitrogen or argon atmosphere. During the reaction,  
20  
21 samples (30 μL) were frequently taken and analyzed by analytical HPLC and ESI-MS to  
22  
23 monitor the progress of reaction.  
24  
25  
26

27  
28 **Synthesis of Vaccine Candidate P(<sup>t</sup>BA<sub>37-≡</sub>)<sub>4</sub>-8Q<sub>min</sub> (S4-8Q<sub>min</sub>).** 8Q<sub>min</sub> azide peptide epitope  
29  
30 (1.8 mg, 1 μmmol, 10 equiv.) and P(<sup>t</sup>BA<sub>37-≡</sub>)<sub>4</sub> (1.9 mg, 0.10 μmol, 1.0 equiv.) were dissolved  
31  
32 in DMF (1 mL). Copper wires (60 mg)—treated with concentrated sulphuric acid (3 min),  
33  
34 subsequently washed with distilled water, methanol and dried under reduced pressure—were  
35  
36 added into the mixture. The mixture was bubbled with nitrogen to remove the air. The  
37  
38 reaction mixtures were covered and protected from light with aluminum foil and stirred at 50  
39  
40 °C in a temperature controlled oil bath under nitrogen atmosphere for 8 h. The wires were  
41  
42 filtered off from the warm solution and washed with 1 mL of DMF. Millipore endotoxin-free  
43  
44 water (7 mL) was slowly added to the solution (0.005 mL/min). Particles formed through the  
45  
46 self-assembly process were exhaustively dialyzed against endotoxin-free water (pH = 6.8)  
47  
48 using presoaked and rinsed dialysis bags (Pierce Snakeskin, MWCO 3K). The resulting  
49  
50 particles were self-assembled into particles with diameters above 1 μm as observed by  
51  
52 dynamic light scattering.  
53  
54  
55

56  
57 **Synthesis of Vaccine Candidate P(<sup>t</sup>BA<sub>37-≡</sub>)<sub>4</sub>-8Q<sub>Ser</sub> (S4-8Q<sub>Ser</sub>).** S4-8Q<sub>Ser</sub> was synthesized in  
58  
59  
60

1  
2  
3 the same manner as above. **8Q<sub>ser</sub>** azide peptide epitope (2.5 mg, 1  $\mu$ mmol, 10 equiv.) and  
4  
5 star-polymer  $p(\text{BA}_{37-\equiv})_4$  (1.9 mg, 0.10  $\mu$ mmol, 1.0 equiv.) were dissolved in DMF (1 mL).  
6  
7 The reaction was terminated after 12 h. The resulting particles were self-assembled into to  
8  
9 yield diameters of around 150 nm and 700 nm as observed by dynamic light scattering.

10  
11 **Synthesis of Vaccine Candidate  $P(\text{BA}_{37-\equiv})_4\text{-8Q}_{\text{Lys}}$  (**S4-8Q<sub>Lys</sub>**).** **S4-8Q<sub>Lys</sub>** was synthesized  
12  
13 in the same manner as above. **8Q<sub>Lys</sub>** azide peptide epitope (2.9 mg, 1  $\mu$ mmol, 10 equiv.) and  
14  
15 star-polymer  $p(\text{BA}_{37-\equiv})_4$  (1.9 mg, 0.10  $\mu$ mmol, 1.0 equiv.) were dissolved in DMF (1 mL).  
16  
17 The reaction was terminated after 12 h. The resulting particles were self-assembled into  
18  
19 particles with a 26 nm diameter as observed by dynamic light scattering.

20  
21 **Mice and cell lines.** Female C57BL/6 (6-8 weeks old) mice were used in this study and  
22  
23 purchased from Animal Resources Centre (Perth, Western Australia). TC-1 cells (murine  
24  
25 C57BL/6 lung epithelial cells transformed with HPV-16 E6/E7 and ras oncogenes) were  
26  
27 obtained from TC Wu.<sup>31</sup> TC-1 cells were cultured and maintained at 37 °C/5% CO<sub>2</sub> in RPMI  
28  
29 1640 medium (Gibco) supplemented with 10% heat inactivated fetal bovine serum (Gibco)  
30  
31 and 1% nonessential amino acid (Sigma-Aldrich). The animal experiments were approved by  
32  
33 the University of Queensland Animal Ethics committee (DI/034/11/NHMRC) in accordance  
34  
35 with National Health and Medical research Council (NHMRC) of Australia guidelines.

36  
37 **In vivo tumor treatment experiments.** To test the efficacy of polymer-peptide conjugates as  
38  
39 a therapeutic vaccine against established tumors, groups of C57BL/6 mice (5 per group) were  
40  
41 first challenged subcutaneously in the right flank with  $1 \times 10^5$ /mouse of TC-1 tumor cells. On  
42  
43 the third day after tumor challenge, the mice were injected subcutaneously at the tail base  
44  
45 with 100  $\mu$ g of polymer-peptide conjugates in a total volume of 100  $\mu$ L of sterile-filtered  
46  
47 phosphate buffered saline (PBS) or control formulations. Polymer-peptide conjugates  
48  
49 consisted of **S4-8Q<sub>min</sub>**, **S4-8Q<sub>ser</sub>** and **S4-8Q<sub>Lys</sub>**. Positive control received 30  $\mu$ g of **8Q**  
50  
51 emulsified in a total volume of 100  $\mu$ L of Montanide ISA51 (Seppic, France)/PBS (1:1, v/v).  
52  
53  
54  
55  
56  
57  
58  
59  
60

1  
2  
3 Three negative control groups were administered PBS, 70  $\mu\text{g}$  of 4-arm star-polymer alone  
4 (S4), or 4-arm star-polymer (70  $\mu\text{g}$ ) + 8Q (30  $\mu\text{g}$ ) physical mixture (S4 + 8Q) in total volume  
5 of 100  $\mu\text{L}$  PBS. The mice were given a single immunization only. The size of the tumor was  
6 measure by palpation and calipers every two days and reported as the average tumor size  
7 across the group of five mice or as tumor size in individual mice.<sup>32,33</sup> Tumor volume was  
8 calculated using the formula  $V (\text{cm}^3) = 3.14 \times [\text{largest diameter} \times (\text{perpendicular}$   
9  $\text{diameter})^2]/6$ .<sup>33</sup> The mice were euthanized when tumor reached 1  $\text{cm}^3$  or started bleeding to  
10 avoid unnecessary suffering.

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21 **Statistical Analysis.** All data were analyzed using GraphPad Prism 5 software. Kaplan-Meier  
22 survival curves for tumor treatment experiments were applied. Differences in survival  
23 treatments were determined using the log-rank (Mantel-Cox) test, with  $p < 0.05$  considered  
24 statistically significant.

## 25 26 27 28 29 30 31 32 **RESULTS AND DISCUSSION**


33  
34  
35  
36 **Synthesis and characterization of the polymer-peptide conjugates.** Copper-  
37 catalyzed alkyne-azide 1,3-dipolar cycloaddition (CuAAC) “click” reaction<sup>34</sup> was used to  
38 synthesize vaccine candidates against HPV-associated cancers. These candidates possessed a  
39 hydrophobic polymeric core and multiple copies of a peptide epitope derived from the HPV-  
40 16 E7 oncoprotein. The 8Q epitope and its new analogues were applied as antigens, while the  
41 polyacrylate 4-arm star-polymer served as a delivery platform. All the peptides contained  
42 CTL, T-helper, and B-cell epitopes.

43  
44  
45  
46  
47  
48  
49  
50  
51 First, the alkyne-functionalized 4-arm poly(*t*-butyl acrylate) star (S4) was synthesized  
52 by atom-transfer radical polymerization (ATRP) and end-group functionalization with a very  
53 narrow molecular-weight distribution (polydispersity index of less than 1.06). 8Q epitope and  
54 its azide ( $\text{N}_3\text{CH}_2\text{CO-QAEPDRAHYNIVTFCKCD-NH}_2$ ) were synthesized using  
55  
56  
57  
58  
59  
60



1  
2  
3 microwave-assisted Boc- and Fmoc-SPPS, respectively. During the initial synthesis, a so-  
4 called “difficult” sequence<sup>35-37</sup> inside **8Q** epitope was identified to be composed of  $\beta$ -  
5 branched amino acids (the IVT tripeptide). It was reported that the presence of these  
6 sequences predisposed the peptide to aggregate on the resin via  $\beta$ -sheet formation and  
7 inhibited completion of Fmoc group deprotection.<sup>38,35</sup> Indeed, a significant amount of amino  
8 acid deletion products were detected during **8Q** synthesis, resulting from incomplete  
9 deprotection of the Fmoc group. Complete deprotection was achieved by replacing piperidine  
10 with DBU for the removal of Fmoc groups after coupling isoleucine, valine and threonine.  
11  
12  
13  
14  
15  
16  
17  
18  
19

20  
21 To examine the 'click' reaction, simple model compounds were designed, in which an  
22 alkyne (**1**) and a dipeptide azide possessing single thiol functionality (**2**) were coupled using  
23 copper-wires as the catalyst (Scheme 1).<sup>21,39-41</sup> Analysis of the reaction showed that instead of  
24 the desired 'click' product **3**, a disulfide bond-mediated dimer of **2** was exclusively produced.  
25 Subsequently, this dimer reacted with alkyne **1** producing disulfide bond-mediated dimer of  
26 compound **3** (Figure S2). Thus, formation of the single product could not be expected during  
27 conjugation of **S4** core with **8Q** peptide which possessed multiple thiol groups. In an effort to  
28 eliminate disulfide bond formation, we investigated alternative CuAAC 'click' reaction  
29 conditions by changing the copper source (copper wire, copper sulfate, copper iodide),  
30 reducing agents (ascorbic acid, tris(2-carboxyethyl)phosphine), solvents (DMF,  
31 acetonitrile/water/trifluoroacetic acid, 90/10/0.1) and temperature (50 °C, RT). Regardless of  
32 the experimental conditions, we were unable to obtain compound **3** (Figure S3).  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




**Scheme 1.** The CuAAC reaction was examined using the model compounds **1** and **2**.


These observations prompted us to modify the **8Q** peptide. Three new epitopes were designed (Figure. 1): **8Q<sub>min</sub>** (QAEPDRAHYNIVTF) which lacked the CCKCD sequence; **8Q<sub>Ser</sub>** (QAEPDRAHYNIVTFSSKSD) where the cysteine moieties were replaced with their structural analog, serine; and **8Q<sub>Lys</sub>** (QAEPDRAHYNIVTFSKKKK) in which the whole hydrophilic pentapeptide CCKCD was replaced with an SKKKK sequence derived from a “solubilizing moiety” incorporated in Pam2Cys/Pam3Cys-based vaccine candidates. Pam2Cys/Pam3Cys are popular adjuvants used to stimulate an immune response in peptide-based vaccines.<sup>15</sup>

**8Q<sub>min</sub>** was conjugated to 4-arm star-polymer to produce **S4-8Q<sub>min</sub>** (Scheme 2). The product, **S4-8Q<sub>min</sub>**, was self-assembled into particles via the solvent replacement method (DMF/water) followed by dialysis against water to remove the organic solvent, excess of unreacted peptide and copper. Formation of the conjugate was confirmed by elemental analysis,<sup>21</sup> which showed a significant increase in the nitrogen/carbon ratio (N/C = 0.085) compared with that of **S4** (N/C = 0.004), due to presence of a nitrogen-rich peptide. A high conjugation efficacy was obtained (3.5 peptide epitopes per polymer, 88% substitution). The substitution ratio was calculated based on comparison of the observed and theoretical N/C ratio. DLS analysis showed that only large particles with diameters over 1  $\mu\text{m}$  were formed

(Figure S4).



**Figure 2.** HPLC chart of crude  $8Q_{Ser}$  (a) synthesized by standard SPPS and (b) O-acyl isopeptide method.



**Scheme 2.** Synthesis of polymer-peptide conjugates.


$8Q_{Ser}$  and  $8Q_{Lys}$  azide derivatives were synthesized using the method developed for

1  
2  
3 synthesis of the **8Q** epitope. Despite the use of DBU for removal of the protective Fmoc  
4 group, **8Q<sub>Ser</sub>** azide was obtained in relatively poor yield after tedious purification. The  
5 isopeptide method developed by Kiso and coworkers was previously applied to the synthesis  
6 of peptides containing a "difficult" sequence.<sup>42-45</sup> Thus, isodipeptide unit (Boc-Thr(Fmoc-  
7 Val)-OH) was synthesized in a similar manner to the reported procedure,<sup>26-28</sup> and applied to  
8 the synthesis of **8Q<sub>Ser</sub>** azide (Scheme S1) to avoid aggregation of the peptide during SPPS.<sup>44</sup>  
9 The final compound was obtained after O-N acyl migration of the O-acyl isoform, with  
10 significantly improved purity and yield (Figure 2). Both epitopes (**8Q<sub>Ser</sub>** and **8Q<sub>Lys</sub>**) were  
11 coupled to **S4** to produce the desired conjugates (Scheme 2). The coupling for **S4-8Q<sub>Ser</sub>** and  
12 **S4-8Q<sub>Lys</sub>** was calculated to be (on average) 3.0 (75%) and 3.0 (74%) epitopes per **S4**,  
13 respectively. Conjugation of peptides to the polymer followed by self-assembly and dialysis  
14 was repeated several times, producing virtually identical products.  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 Both compounds (**S4-8Q<sub>Ser</sub>** and **S4-8Q<sub>Lys</sub>**) were self-assembled in DMF/water and  
30 dialyzed against water. DLS analysis indicated that **S4-8Q<sub>Ser</sub>** formed large nanoparticles (~  
31 530 nm; Figure S4). In contrast, the size distribution of **S4-8Q<sub>Lys</sub>** particles was narrow, and  
32 smaller nanoparticles (26 nm) were detected (Figure S4). To prepare samples for in vivo  
33 experiments, **S4-8Q<sub>min</sub>**, **S4-8Q<sub>Ser</sub>** and **S4-8Q<sub>Lys</sub>** were formulated in PBS. All of the  
34 conjugates formed a milky suspension upon addition of the buffer (pH = 7.4). Confocal  
35 images were used to visualize microparticles and their aggregates (Figure S5). We were  
36 unable to quantitatively determine particles size distribution using DLS, TEM, and analytical  
37 ultracentrifugation.  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48

49  
50 **In vivo tumor treatment.** To evaluate the therapeutic effect of polymer-peptide  
51 conjugates against established HPV tumor using the newly designed epitopes, mice were  
52 immunized on day 3 post-tumor implantation<sup>46,47</sup> with **S4-8Q<sub>min</sub>**, **S4-8Q<sub>Ser</sub>**, **S4-8Q<sub>Lys</sub>** or  
53 control groups (**8Q** emulsified with IFA-like adjuvant (Montanide ISA51), physical mixture  
54  
55  
56  
57  
58  
59  
60

of S4 and 8Q, S4 alone and PBS). Mice were monitored every two days for evidence of tumor growth by palpation and calipers for 60 days.



**Figure 3.** In vivo tumor treatment experiments. C57BL/6 (5 per group) were inoculated subcutaneously in the right flank with  $1 \times 10^5$ /mouse of TC-1 tumor cells (day 0) and vaccinated with different immunogens on day 3 without additional boost. (a) Survival rate monitored over 60 days post implantation and time to death plotted on a Kaplan-Meier survival curve. Mice were euthanized when tumor volume reached  $1 \text{ cm}^3$  or started bleeding. The survival rate of each group was compared to the negative control (PBS) and was analyzed using the log-rank (Mantel-Cox) test (\*\* $p < 0.01$ ). (b) Mean tumor volume (cm<sup>3</sup>) in

1  
2  
3 different groups of mice over 60 days post implantation. Tumor volume ( $\text{cm}^3$ ) in individual  
4  
5 TC-1 tumor-bearing mice treated with (c) **S4-8Q<sub>min</sub>**, (d) **S4 + 8Q** physical mixture, (e) **S4**  
6  
7 alone or (f) PBS shown only up to day 21 of the experiment.  
8  
9

10  
11  
12 The Kaplan-Meier survival curve (Figure 3a) showed that 100% of mice treated with  
13  
14 physical mixture (**S4 + 8Q**), **S4** alone and PBS were moribund (i.e. mice were sacrificed due  
15  
16 to tumor burden) by day 28, 33 and 33, respectively. In contrast, the survival rate in the mice  
17  
18 treated with **S4-8Q<sub>min</sub>**, **S4-8Q<sub>ser</sub>** and **8Q + ISA51** was 60%, 20%, and 40% over two months,  
19  
20 respectively. Among the tested formulations, only **S4-8Q<sub>min</sub>** and positive control showed  
21  
22 significantly better survival compared with tumor-bearing mice treated with negative control  
23  
24 ( $p < 0.0027$ ).  
25  
26

27  
28 As shown in Figure 3b, tumor-bearing mice treated with **S4-8Q<sub>min</sub>** showed slower  
29  
30 tumor growth over time than the other polymer-peptide conjugates and was similar to the  
31  
32 mice treated with positive control. Treatment with **S4-8Q<sub>min</sub>** eradicated tumor by day 21 post-  
33  
34 tumor implantation in three mice, while an initial tumor reduction was observed in remaining  
35  
36 two mice after 14-17 days (Figure 3c). It is noteworthy that an initial reduction in tumor size  
37  
38 was observed in four groups including positive control and all conjugates (see Figure 3c and  
39  
40 Supporting Information, Figure S6a-c). Remission of tumor growth was observed after 19  
41  
42 days, especially in the group immunized with **S4-8Q<sub>Lys</sub>**. Thus, **S4-8Q<sub>min</sub>** showed similar  
43  
44 efficacy to the positive control while the antitumor efficacy of other conjugates was not  
45  
46 statistically significant. No adverse side effects or allergic responses were observed in mice  
47  
48 immunized with the conjugates, while one mouse from the group immunized with an  
49  
50 adjuvant (**8Q + ISA51**) developed an adverse reaction (scratching resulting in patchy hair  
51  
52 loss). Chemical conjugation of epitopes with a polymer core was essential to elicit a  
53  
54 therapeutic effect since **S4 + 8Q** failed to induce an antitumor immune response. The three  
55  
56  
57  
58  
59  
60

1  
2  
3 conjugates (**S4-8Q<sub>min</sub>**, **S4-8Q<sub>ser</sub>** and **S4-8Q<sub>Lys</sub>**) demonstrated different antitumor activity,  
4  
5 which might be associated with the ability of antigen presenting cells to correctly process and  
6  
7 display different epitopes. Correct antigen processing is essential to trigger the desired  
8  
9 cellular response. While **8Q<sub>min</sub>** is a native fragment of the HPV-16 E7 protein, both **8Q<sub>ser</sub>** and  
10  
11 **8Q<sub>Lys</sub>** are chemically altered epitopes where the CTL epitope is flanked with a foreign  
12  
13 peptide on its C-terminal. This alteration might impair antigen processing and consequently  
14  
15 may have reduced the cellular immune response.<sup>48-50</sup> In addition, size can play important role  
16  
17 in immune responses, while some groups reported that immune responses were significantly  
18  
19 stronger for small nanoparticles (less than 100 nm),<sup>51,52</sup> others claimed that microparticles are  
20  
21 required for strong activation of the immune system.<sup>53,54</sup> Small nanoparticles can easily  
22  
23 migrate to lymphatic nodes for antigen presentation, whereas large particles are transported to  
24  
25 lymphatic nodes by dendritic cells and the immunity induced by large particles is often  
26  
27 related to the depot effect.<sup>55</sup> Interestingly, T- cell activation is often facilitated by the  
28  
29 extended presence of antigen through depot-forming adjuvants.<sup>54</sup> The three conjugates form  
30  
31 microparticles and have the tendency to aggregate further. This observation may suggest that  
32  
33 the differences in the conjugates antitumor activity are not related to particle size.  
34  
35  
36  
37

38  
39 Polymer-peptide conjugate **S4-8Q<sub>min</sub>** demonstrated self-adjuvanting activity because  
40  
41 it induced a therapeutic effect against tumor cells without help of any external adjuvant. This  
42  
43 delivery system overcomes the poor immunogenicity of peptide-based vaccines and the  
44  
45 common toxic side effects associated with external adjuvants. Self-adjuvanting properties of  
46  
47 the polymer-based delivery system can be explained by particle-attributed activation of the  
48  
49 immune system.<sup>16,56,57</sup> Particulate vaccines potentially cross-present the antigen, and antigen  
50  
51 cross-presentation is especially important to induce the priming of the CD8<sup>+</sup> T-cell immune  
52  
53 responses that naturally play a crucial role for therapeutic vaccines against cancer. This  
54  
55 delivery system also exploits the depot effect (i.e. retaining the antigen at the injection site),  
56  
57  
58  
59  
60

1  
2  
3 thus increasing the duration of vaccine exposure to the immune cells. In addition, as particles  
4 are covered by multiple copies of the same peptide antigen, they should possess the  
5 advantages of a MAP system. Furthermore, formulation of antigens into particles provides  
6 some protection against enzymatic degradation, which is important for highly susceptible  
7 peptide antigens. Noticeably, in contrast to many previously reported peptide-based vaccine  
8 candidates,<sup>33,46,47,58</sup> **S4-8Q<sub>min</sub>** demonstrated a therapeutic effect after a single immunization.  
9  
10  
11  
12  
13  
14  
15  
16  
17

## 18 **Conclusion**

19  
20 In conclusion, we established a synthetic pathway to produce polymer-peptide  
21 conjugates as macromolecular vaccine candidates against HPV-related cancers. The synthesis  
22 of peptide epitopes was greatly improved by the change of standard SPPS procedure and  
23 application of the isopeptide method. Modification of the immunogenic epitope allowed the  
24 elimination of undesirable disulfide bond-based aggregation/polymerization of the polymer-  
25 peptide conjugates.  
26  
27  
28  
29  
30  
31  
32  
33

34 Chemical alteration of **8Q** epitope allowed us to find the most effective epitope  
35 (**8Q<sub>min</sub>**). Polyacrylate star-polymer conjugated with this epitope produced a robust  
36 therapeutic effect against a tumor without the help of any external adjuvant. This delivery  
37 system overcomes the poor immunogenicity of peptide-based vaccines. In contrast to many  
38 previously reported vaccine candidates, the conjugates demonstrated therapeutic effect after  
39 only a single immunization. Thus, we have developed the first self-adjuvanting delivery  
40 system for a therapeutic peptide-based vaccine against cervical cancer.  
41  
42  
43  
44  
45  
46  
47  
48  
49

## 50 **Acknowledgement**

51  
52 This work was supported by the National Health and Medical Research Council of Australia  
53 (NHMRC 1006454). We thank Thalia Guerin for her critical review of the manuscript. We  
54  
55  
56  
57  
58  
59  
60



1  
2  
3 thank Ai Preston for her assistance in animal studies and Ashwin Kumar Giddam for the  
4  
5 confocal microscope images.  
6  
7

### 8 9 **Supporting Information**

10  
11 Additional information related to model study, isopeptide method (reactions schemes and  
12  
13 HPCL charts), confocal microscope images and animal study. This material is available free  
14  
15 of charge via the Internet at <http://pubs.acs.org>.  
16  
17  
18

### 19 20 **Notes**


21  
22  
23 The authors declare no competing financial interest.  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

## References

1. Oka, Y.; Tsuboi, A.; Fujiki, F.; Shirakata, T.; Nishida, S.; Hosen, N.; Nakajima, H.; Li, Z.; Kawase, I.; Oji, Y.; Sugiyama, H. *Curr. Med. Chem.* **2008**, *15*, 3052-3061.
2. Liu, T. Y.; Hussein, W. M.; Toth, I.; Skwarczynski, M. *Curr. Top. Med. Chem.* **2012**, *12*, 1581-1592.
3. Frazer, I. H. *Nat. Rev. Immunol.* **2004**, *4*, 46-54.
4. Goldie, S. J.; Grima, D.; Kohli, M.; Wright, T. C.; Weinstein, M.; Franco, E. *Int. J. Cancer* **2003**, *106*, 896-904.
5. Schlecht, N. F.; Platt, R. W.; Duarte-Franco, E.; Costa, M. C.; Sobrinho, J. P.; Prado, J. C. M.; Ferenczy, A.; Rohan, T. E.; Villa, L. L.; Franco, E. L. *J. Natl. Cancer Inst.* **2003**, *95*, 1336-1343.
6. Galani, E.; Christodoulou, C. *Clin. Microbiol. Infect.* **2009**, *15*, 977-981.
7. Frazer, I. H.; Quinn, M.; Nicklin, J. L.; Tan, J.; Perrin, L. C.; Ng, P.; O'Connor, V. M.; White, O.; Wendt, N.; Martin, J.; Crowley, J. M.; Edwards, S. J.; McKenzie, A. W.; Mitchell, S. V.; Maher, D. W.; Pearse, M. J.; Bassler, R. L. *Vaccine* **2004**, *23*, 172-181.
8. Trimble, C. L.; Peng, S.; Kos, F.; Gravitt, P.; Viscidi, R.; Sugar, E.; Pardoll, D.; Wu, T. C. *Clin. Cancer Res.* **2009**, *15*, 361-367.
9. Bijker, M. S.; van den Eeden, S. J. F.; Franken, K. L.; Melief, C. J. M.; Offringa, R.; van der Burg, S. H. *J. Immunol.* **2007**, *179*, 5033-5040.
10. Bellone, S.; El-Sahwi, K.; Cocco, E.; Casagrande, F.; Cargnelutti, M.; Palmieri, M.; Bignotti, E.; Romani, C.; Silasi, D. A.; Azodi, M.; Schwartz, P. E.; Rutherford, T. J.; Pecorelli, S.; Santin, A. D. *J. Virol.* **2009**, *83*, 6779-6789.
11. Villa, L. L.; Perez, G.; Kjaer, S. K.; Paavonen, J.; Lehtinen, M.; Munoz, N.; Sigurdsson, K.; Hernandez-Avila, M.; Skjeldestad, F. E.; Thoresen, S.; Garcia, P.; Majewski, S.; Dillner, J.; Olsson, S. E.; Tay, E. H.; Bosch, F. X.; Ault, K. A.; Brown, D. R.; Ferris, D. G.; Koutsky, L. A.; Kurman, R. J.; Myers, E. R.; Barr, E.; Boslego, J.; Bryan, J.; Esser, M. T.; Gause, C. K.; Hesley, T. M.; Lupinacci, L. C.; Sings, H. L.; Taddeo, F. J.; Thornton, A. R.; Boulos, M.; Cox, J. T.; Langmark, F.; Modlin, J.; Munoz, A.; Odlind, V.; Wilkinson, E.; Ferenczy, A.; Kurman, R.; Ronett, B.; Stoler, M.; Andreoni, G.; Bahamondes, L.; Camargos, A.; Costa, R.; De Andrade, R.; Fedrizzi, E.; Ferriani, R.; Goncalves, M.; Laginha, F.; Mendonca, J.; Moreira, E.; Nonnenmacher, B.; Taborda, W.; Zanetta, D.; Ardila, J.; Balcazar, N.; Maldonado, I.; Revollo, F.; Ruiz, A.; Andersen, E. S.; Djursing, H.; Hansen, T.; Jorgensen, J. J.; Nilas, L.; Ottesen, B.; Petersen, L. K.; Thomsen, S. G.; Toftager-Larsen, K.; Apter, D.; Kekki, M.; Kuortti, M.; Lahti, L.; Lindroos, Y.; Lunnas, T.; Palmroth, J.; Lazcano, E.; Zertuche, J.; Dalaker, K.; Eriksen, B.; Erno, L. E.; Fiane, B.; Hesla, K.; Isachsen, M. M.; Iversen, O. E.; Kasin, K.; Kristoffersen, M.; Lunde, T.; Nordmark, P. T.; Nygaard, T.; Onsrud, M.; Riis-Johannessen, G.; Schiøtz, H.; Sundhagen, H.; Sviggum, O.; Trosterud, K. A.; Penny, M.; Vivar, A.; Basta, A.; Czajkowski, K.; Knapp, P.; Spaczynski, M.; Barnes, R.; Elfgren, K.; Hardmeier, E.; Hellsten, C.; Hofte, C.; Jensen, P.; Johannisson, G.; Lowhagen, G. B.; Steinwall, M.; Tamsen, L.; Varnauskas, T.; Wikstrom, A.; Crawford, G.; Allen, B.; Ault, K.; Brown, D.; Edwards, R.; Comerci, J.; Giuliano, A.; Greer, S.; Hatch, K.; Kriesel, J.; Lalezari, J.; Partridge, E.; Sperling, R.; Spruance, S.; Stapleton, J.; Wright, P.; Zedler, P.; McCarroll, K.; Zhang, L.; Zhou, H.; Grp, F. I. S. *N. Engl. J. Med.* **2007**, *356*, 1915-1927.
12. Purcell, A. W.; McCluskey, J.; Rossjohn, J. *Nat. Rev. Drug Discovery* **2007**, *6*, 404-414.

13. Reed, S. G.; Bertholet, S.; Coler, R. N.; Friede, M. *Trends Immunol.* **2009**, *30*, 23-32.
14. Sharma, R. K.; Elpek, K. G.; Yolcu, E. S.; Schabowsky, R. H.; Zhao, H.; Bandura-Morgan, L.; Shirwan, H. *Cancer Res.* **2009**, *69*, 4319-4326.
15. Boas, U.; Heegaard, P. M. H. *Chem. Soc. Rev.* **2004**, *33*, 43-63.
16. Wang, C. Y.; Looney, D. J.; Li, M. L.; Walfield, A. M.; Ye, J.; Hosein, B.; Tam, J. P.; Wongstaal, F. *Science* **1991**, *254*, 285-288.
17. Brandt, E. R.; Sriprakash, K. S.; Hobb, R. I.; Hayman, W. A.; Zeng, W. G.; Batzloff, M. R.; Jackson, D. C.; Good, M. F. *Nat. Med.* **2000**, *6*, 455-459.
18. Hilgers, L. A. T.; Ghenne, L.; Nicolas, I.; Fochesato, M.; Lejeune, G.; Boon, B. *Vaccine* **2000**, *18*, 3319-3325.
19. Hilgers, L. A. T.; Nicolas, I.; Lejeune, G.; Dewil, E.; Strebelle, M.; Boon, B. *Vaccine* **1998**, *16*, 1575-1581.
20. Zaman, M.; Skwarczynski, M.; Malcolm, J. M.; Urbani, C. N.; Jia, Z. F.; Batzloff, M. R.; Good, M. F.; Monteiro, M. J.; Toth, I. *Nanomed.-Nanotechnol. Biol. Med.* **2011**, *7*, 168-173.
21. Skwarczynski, M.; Zaman, M.; Urbani, C. N.; Lin, I. C.; Jia, Z. F.; Batzloff, M. R.; Good, M. F.; Monteiro, M. F.; Toth, I. *Angew. Chem. Int. Ed. Engl.* **2010**, *49*, 5742-5745.
22. Scheerlinck, J. P. Y.; Greenwood, D. L. V. *Drug Discov. Today* **2008**, *13*, 882-887.
23. Peek, L. J.; Middaugh, C. R.; Berkland, C. *Adv. Drug Delivery Rev.* **2008**, *60*, 915-928.
24. Jia, Z.; Bell, C. A.; Monteiro, M. J. *Macromolecules* **2011**, *44*, 1747-1751.
25. Dyke, J. M.; Groves, A. P.; Morris, A.; Ogden, J. S.; Dias, A. A.; Oliveira, A. M. S.; Costa, M. L.; Barros, M. T.; Cabral, M. H.; Moutinho, A. M. C. *J. Am. Chem. Soc.* **1997**, *119*, 6883-6887.
26. Yoshiya, T.; Taniguchi, A.; Sohma, Y.; Fukao, F.; Nakamura, S.; Abe, N.; Ito, N.; Skwarczynski, M.; Kimura, T.; Hayashi, Y.; Kiso, Y. *Org. Biomol. Chem.* **2007**, *5*, 1720-1730.
27. Sohma, Y.; Taniguchi, A.; Skwarczynski, M.; Yoshiya, T.; Fukao, F.; Kimura, T.; Hayashi, Y.; Kiso, Y. *Tetrahedron Lett.* **2006**, *47*, 3013-3017.
28. Hussein, W. M.; Liu, T. Y.; Toth, I.; Skwarczynski, M. *Org. Biomol. Chem.* **2013**, *11*, 2370-2376.
29. Urbani, C. N.; Bell, C. A.; Lonsdale, D.; Whittaker, M. R.; Monteiro, M. J. *Macromolecules* **2008**, *41*, 76-86.
30. Urbani, C. N.; Lonsdale, D. E.; Bell, C. A.; Whittaker, M. R.; Monteiro, M. J. *J. Polym. Sci., Part A: Polym. Chem.* **2008**, *46*, 1533-1547.
31. Lin, K. Y.; Guarnieri, F. G.; StaveleyOcarroll, K. F.; Levitsky, H. I.; August, J. T.; Pardoll, D. M.; Wu, T. C. *Cancer Res.* **1996**, *56*, 21-26.
32. Hung, C. F.; Cheng, W. F.; Chai, C. Y.; Hsu, K. F.; He, L.; Ling, M.; Wu, T. C. *J. Immunol.* **2001**, *166*, 5733-5740.
33. Zeng, Q.; Peng, S.; Monie, A.; Yang, M.; Pang, X.; Hung, C.-F.; Wu, T. C. *Hum. Gene Ther.* **2011**, *22*, 809-819.
34. Kolb, H. C.; Sharpless, K. B. *Drug Discovery Today* **2003**, *8*, 1128-1137.
35. Sohma, Y.; Hayashi, Y.; Skwarczynski, M.; Hamada, Y.; Sasaki, M.; Kimura, T.; Kiso, Y. *Biopolymers* **2004**, *76*, 344-356.
36. Carpino, L. A.; Krause, E.; Sferdean, C. D.; Schumann, M.; Fabian, H.; Bienert, M.; Beyermann, M. *Tetrahedron Lett.* **2004**, *45*, 7519-7523.
37. Sheppard, R. *J. Pept. Sci.* **2003**, *9*, 545-552.


- 1  
2  
3 38. Larsen, B. D.; Holm, A. *Int. J. Pept. Protein Res.* **1994**, *43*, 1-9.  
4 39. Skwarczynski, M.; Parhiz, B. H.; Soltani, F.; Srinivasan, S.; Kamaruzaman, K. A.; Lin,  
5 I.-C.; Toth, I. *Aust. J. Chem.* **2012**, *65*, 35-39.  
6 40. Skwarczynski, M.; Ahmad Fuaad, A. A. H.; Rustanti, L.; Ziora, Z. M.; Aqil, M.; Batzloff,  
7 M. R.; Good, M. F.; Toth, I. *Drug Delivery Lett.* **2011**, *1*, 2-8.  
8 41. Urbani, C. N.; Bell, C. A.; Whittaker, M. R.; Monteiro, M. J. *Macromolecules* **2008**, *41*,  
9 1057-1060.  
10 42. Taniguchi, A.; Skwarczynski, M.; Sohma, Y.; Okada, T.; Ikeda, K.; Prakash, H.; Mukai,  
11 H.; Hayashi, Y.; Kimura, T.; Hirota, S.; Matsuzaki, K.; Kiso, Y. *ChemBioChem* **2008**, *9*,  
12 3055-3065.  
13 43. Skwarczynski, M.; Kiso, Y. *Curr. Med. Chem.* **2007**, *14*, 2813-2823.  
14 44. Sohma, Y.; Taniguchi, A.; Yoshiya, T.; Chiyomori, Y.; Fukao, F.; Nakamura, S.;  
15 Skwarczynski, M.; Okada, T.; Ikeda, K.; Hayashi, Y.; Kimura, T.; Hirota, S.; Matsuzaki,  
16 K.; Kiso, Y. *J. Pept. Sci.* **2006**, *12*, 823-828.  
17 45. Skwarczynski, M.; Sohma, Y.; Noguchi, M.; Kimura, T.; Hayashi, Y.; Kiso, Y. *J. Org.*  
18 *Chem.* **2006**, *71*, 2542-2545.  
19 46. Wu, C. Y.; Monie, A.; Pang, X.; Hung, C. F.; Wu, T. C. *J. Biomed. Sci.* **2010**, *17*, 88.  
20 47. Tang, J.; Yin, R.; Tian, Y.; Huang, Z.; Shi, J.; Fu, X.; Wang, L.; Wu, Y.; Hao, F.; Ni, B.  
21 *Vaccine* **2012**, *30*, 1071-1082.  
22 48. Mellman, I.; Coukos, G.; Dranoff, G. *Nature* **2011**, *480*, 480-489.  
23 49. Timm, J.; Lauer, G. M.; Kavanagh, D. G.; Sheridan, I.; Kim, A. Y.; Lucas, M.; Pillay, T.;  
24 Ouchi, K.; Reyor, L. L.; zur Wiesch, J. S.; Gandhi, R. T.; Chung, R. T.; Bhardwaj, N.;  
25 Klenerman, P.; Walker, B. D.; Allen, T. M. *J. Exp. Med.* **2004**, *200*, 1593-1604.  
26 50. Allen, T. M.; Altfeld, M.; Yu, X. G.; O'Sullivan, K. M.; Lichterfeld, M.; Le Gall, S.;  
27 John, M.; Mothe, B. R.; Lee, P. K.; Kalife, E. T.; Cohen, D. E.; Freedberg, K. A.; Strick,  
28 D. A.; Johnston, M. N.; Sette, A.; Rosenberg, E. S.; Mallal, S. A.; Goulder, P. J. R.;  
29 Brander, C.; Walker, B. D. *J. Virol.* **2004**, *78*, 7069-7078.  
30 51. Skwarczynski, M.; Toth, I. *Curr. Drug Delivery* **2011**, *8*, 282-289.  
31 52. Xiang, S. D.; Scholzen, A.; Minigo, G.; David, C.; Apostolopoulos, V.; Mottram, P. L.;  
32 Plebanski, M. *Methods* **2006**, *40*, 1-9.  
33 53. Oyewumi, M. O.; Kumar, A.; Cui, Z. R. *Expert Rev. Vaccines* **2010**, *9*, 1095-1107.  
34 54. Bachmann, M. F.; Jennings, G. T. *Nat. Rev. Immunol.* **2010**, *10*, 787-796.  
35 55. Manolova, V.; Flace, A.; Bauer, M.; Schwarz, K.; Saudan, P.; Bachmann, M. F. *Eur. J.*  
36 *Immunol.* **2008**, *38*, 1404-1413.  
37 56. Oyewumi, M. O.; Kumar, A.; Cui, Z. *Expert Rev. Vaccines* **2010**, *9*, 1095-1107.  
38 57. Bachmann, M. F.; Jennings, G. T. *Nat. Rev. Immunol.* **2010**, *10*, 787-796.  
39 58. Torrens, I.; Mendoza, O.; Batte, A.; Reyes, O.; Fernandez, L. E.; Mesa, C.; Guillen, G.  
40 *Vaccine* **2005**, *23*, 5768-5774.  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60




26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

The amino acid sequence of the original 8Q epitope and its novel derivatives. CTL epitope (CD8+ cytotoxic T lymphocytes), T-helper cell (CD4+) and a B-cell epitope are highlighted within the 8Q sequence. The immunologically redundant pentapeptide sequence is indicated by italic font.

44x25mm (300 x 300 DPI)




HPLC chart of crude 8QSer (a) synthesized by standard SPPS and (b) O-acyl isopeptide method.  
85x116mm (300 x 300 DPI)




In vivo tumor treatment experiments. C57BL/6 (5 per group) were inoculated subcutaneously in the right flank with  $1 \times 10^5$ /mouse of TC-1 tumor cells (day 0) and vaccinated with different immunogens on day 3 without additional boost. (a) Survival rate monitored over 60 days post implantation and time to death plotted on a Kaplan-Meier survival curve. Mice were euthanized when tumor volume reached 1 cm<sup>3</sup> or started bleeding. The survival rate of each group was compared to the negative control (PBS) and was analyzed using the log-rank (Mantel-Cox) test (\*\* $p < 0.01$ ). (b) Mean tumor volume (cm<sup>3</sup>) in different groups of mice over 60 days post implantation. Tumor volume (cm<sup>3</sup>) in individual TC-1 tumor-bearing mice treated with (c) S4-8Q<sub>min</sub>, (d) S4 + 8Q physical mixture, (e) S4 alone or (f) PBS shown only up to day 21 of the experiment.

243x228mm (300 x 300 DPI)




The CuAAC reaction was examined using the model compounds 1 and 2.  
61x29mm (300 x 300 DPI)





Synthesis of polymer-peptide conjugates.  
85x50mm (300 x 300 DPI)



graphic  
35x14mm (300 x 300 DPI)