

6-1-2013

VEGF and Sympathetic Perivascular Nerves Contribute to Hypoxic Remodeling of Ovine Cranial Arteries

Olayemi Olufikayo Adeoye
Loma Linda University

Follow this and additional works at: <http://scholarsrepository.llu.edu/etd>

 Part of the [Medical Physiology Commons](#)

Recommended Citation

Adeoye, Olayemi Olufikayo, "VEGF and Sympathetic Perivascular Nerves Contribute to Hypoxic Remodeling of Ovine Cranial Arteries" (2013). *Loma Linda University Electronic Theses, Dissertations & Projects*. 110.
<http://scholarsrepository.llu.edu/etd/110>

This Dissertation is brought to you for free and open access by TheScholarsRepository@LLU: Digital Archive of Research, Scholarship & Creative Works. It has been accepted for inclusion in Loma Linda University Electronic Theses, Dissertations & Projects by an authorized administrator of TheScholarsRepository@LLU: Digital Archive of Research, Scholarship & Creative Works. For more information, please contact scholarsrepository@llu.edu.

LOMA LINDA UNIVERSITY
School of Medicine
In conjunction with the
Faculty of Graduate Studies

VEGF and Sympathetic Perivascular Nerves contribute to Hypoxic
Remodeling of Ovine Cranial Arteries

by

Olayemi Olufikayo Adeoye

A Dissertation submitted in partial satisfaction of
the requirements for the degree
Doctor of Philosophy in Physiology

June 2013

© 2013

Olayemi Olufikayo Adeoye
All Rights Reserved

Each person whose signature appears below certifies that this dissertation in his/her opinion is adequate, in scope and quality, as a dissertation for the degree Doctor of Philosophy.

_____, Chairperson
William J Pearce, Professor of Physiology and Associate Director, Center for Perinatal
Biology

Deborah Damon, Professor of Physiology, Eastern Virginia Medical School

Charles Ducsay, Professor of Physiology and Associate Director, Center for Perinatal
Biology

Eugenia Mata-Greenwood, Assistant Professor of Pharmacology

Lubo Zhang, Professor of Pharmacology and Director, Center for Perinatal Biology

ACKNOWLEDGEMENTS

My profound gratitude goes to Dr. William Pearce who provided me with the opportunity and enabling environment to carry out research in quest of answers to my scientific curiosities. He not only taught me how to work professionally in diverse environments but also taught me how to attain a balance between humor, professionalism and hard work. I could not have asked for a better mentor. Thank you Dr. Pearce. Many thanks to my committee members for their patience, advice and availability to me during my study. Thank you for your invaluable contributions and corrections, for your prompt responses to my endless counts of emails. I will forever be grateful. To James Williams and other lab A members, I thank you and appreciate your support with all my projects. Thank you for contributing to a happy work environment I enjoyed for the last 4 and half years. To my loving parents, thank you for always believing in me and giving me the opportunity to have this much education. Indeed I owe everything to you. To my in-laws, thank you for giving me the most beautiful gift of my life – my wife. Thank you also for your prayers and well wishes. To my guardians – Dr and Mrs. Atolagbe, thank you for being my parents here in the United States. To a wonderful and reliable friend, Oludare, I say thank you.

To my team, my rock, my love and my life, my ladies, Kemi and Tami “princess Tata”, and my “crown prince” – Mofeoluwa, thank you for holding on and always being there for me. Kemi, you are a special gift to me from God. Thank you giving me wonderful and beautiful kids and taking care of them while I’m gone to work. Tami, your smile is priceless. Thank you for those late evening calls, and telling me you miss me when I work late into the night.

Finally, to God who is faithful, gracious and marvelously leads us into our destinies, and provides us opportunities beyond our wildest imagination, I owe it all to you. Bless God.

DEDICATION

This doctoral dissertation is dedicated to my beautiful wife – Olukemi Adeoye. You never stopped believing in me and taking care of our jewels – Tami and Mofe while I was gone. Also, this work is dedicated to my uncle, Late (Dr.) Hezekiah Olanrewaju Adesina, who inspired me a lot growing up. Lastly, this work is especially dedicated to my parents Mr. and Dr (Mrs.) Afolabi Adeoye, who denied themselves of many luxuries to get us, the children, educated, I will forever be grateful to you. Love you.

CONTENTS

Approval Page.....	iii
Acknowledgements.....	iv
Dedication.....	vi
Table of Contents.....	vii
List of Figures.....	xi
List of Abbreviations.....	xiii
Dissertation Abstract.....	xiv
Chapter	
1. Introduction.....	1
Hypoxia.....	1
Hypoxic Vascular Remodeling.....	3
Hypoxia Inducible Factor & Vascular Endothelial Growth Factor.....	5
Vascular Endothelial Growth Factor (VEGF).....	7
VEGF signaling pathway.....	12
Neurogenic regulation of cerebral blood flow.....	13
VEGF and the neuron.....	13
VEGF, Transmural Gradients and Phenotypic Transformation.....	13
References.....	16
2. Contribution of Increased VEGF Receptors to Hypoxic Changes in Fetal Ovine Carotid Artery Contractile Proteins.....	25
Abstract.....	26
Keywords.....	27
Introduction.....	28
Materials and Methods.....	32
Tissue Harvest and Preparation.....	32

Contractility Studies.....	33
Fluorescent Immunohistochemistry.....	34
Confocal Microscopy.....	35
Western Blotting.....	36
Organ Culture.....	38
Data Analysis and Statistics.....	39
Results.....	40
Effects of chronic hypoxia on carotid artery structure and stress-strain relations.....	40
Effects of chronic hypoxia on contractile protein abundance and colocalization.....	42
Effects of long-term hypoxia on abundances of VEGF and VEGF receptors.....	45
VEGF receptors mediate VEGF-induced changes in contractile protein organization.....	48
Effects of VEGF on contractile protein abundance and colocalization.....	50
Discussion.....	53
Effects of hypoxia on structure and contractility.....	53
Effects of hypoxia on contractile protein abundances.....	54
Effects of hypoxia on contractile protein organization.....	56
Effects of Chronic Hypoxia on VEGF and VEGF Receptors.....	57
Effects of VEGF on contractile protein abundance and organization.....	58
Overview.....	60
Acknowledgements.....	62
References.....	63
3. VEGF receptors mediate hypoxic remodeling of adult ovine carotid arteries.....	69
Abstract.....	70
Keywords.....	71
Introduction.....	72
Materials and Methods.....	75
Tissue Harvest and Preparation.....	75
Contractility Studies.....	76
Fluorescent Immunohistochemistry.....	77
Confocal Microscopy.....	77
Western Blotting.....	78

Organ Culture.....	80
Data Analysis and Statistics.....	81
Results.....	82
Chronic hypoxia alters arterial structure and contractility.....	82
Chronic hypoxia alters smooth muscle contractile protein expression	84
Effects of chronic hypoxia on contractile protein colocalization	86
Effects of chronic hypoxia on VEGF and VEGF receptor expression	88
VEGF-induced changes in contractile protein interactions are maintained through VEGF receptor upregulation.....	91
Effect of VEGF in culture on contractile protein abundances.....	93
Effect of VEGF on contractile protein interactions	95
Discussion.....	97
Long-term hypoxia alters arterial structure and contractility	97
Chronic hypoxia differentially modulates contractile protein abundances.....	98
Chronic hypoxia differentially modulates contractile protein organization.....	100
Effects of long-term hypoxia on VEGF and VEGF receptor levels	101
VEGF receptors mediate arterial remodeling	102
VEGF differentially modulates contractile protein abundances and colocalization	103
Overview.....	105
References.....	109
4. Sympathetic perivascular nerves contribute to long-term hypoxic transformation of smooth muscle phenotype in ovine cerebral arteries	117
Abstract.....	118
Keywords	118
Introduction.....	119
Materials and Methods.....	121
Tissue Harvest and Preparation	121
Electrical Transmural Stimulation	122
Western Immunoblotting	123
Fluorescent Immunohistochemistry.....	124
Wall thickness and stiffness.....	125
Confocal Microscopy.....	125

Data analysis and Statistic	126
Results.....	127
Effects of long-term hypoxia on arterial reactivity to nerve stimulation.....	127
Effects of long-term hypoxia on expression of dopamine- β -hydroxylase.....	129
Effects of sympathetic perivascular nerves on arterial wall thickness and stiffness.....	131
Effect of sympathetic perivascular nerves on contractile protein organization.....	133
Sympathetic perivascular nerves mediate hypoxic remodeling through NE and it's α -1 adrenoceptors.....	138
Discussion.....	140
Long-term hypoxia enhances arterial reactivity to electrical stimulation.....	140
NE from sympathetic perivascular nerves mediates response to transmural stimulation in hypoxic arteries.....	142
Sympathetic perivascular nerves mediate hypoxic changes in arterial structure	142
Sympathetic perivascular nerves mediate hypoxia-induced changes in arterial function through alteration of contractile protein colocalization	143
NE mediates hypoxic reorganization of contractile proteins via it's α -1 adrenoceptors	145
Overview.....	146
Acknowledgment	147
References.....	148
5. Conclusions and Future Directions.....	154

FIGURES

Figures	Page
Chapter 1	
1. Molecular mechanisms of oxygen homeostasis.....	6
2. Ribbon representation of the receptor-binding domain of VEGF	9
3. VEGF signaling pathway	12
Chapter 2	
1. Schematic for hypothesis	31
2. Effects of chronic hypoxia on carotid artery structure and stress-strain relations.....	41
3. Effects of chronic hypoxia on contractile protein abundance.....	43
4. Effects of chronic hypoxia on contractile protein colocalization	44
5. Effects of long-term hypoxia on abundances of VEGF	46
6. Effects of long-term hypoxia on abundances of VEGF receptors	47
7. VEGF Receptors Mediate VEGF-induced Changes in Contractile Protein Organization	49
8. Effects of VEGF on contractile protein abundance	51
9. Effects of VEGF on contractile protein colocalization.....	52
Chapter 3	
1. Chronic hypoxia alters arterial structure and contractility in adult carotid arteries.....	83
2. Chronic hypoxia alters smooth muscle contractile protein expression.....	85
3. Effects of chronic hypoxia on contractile protein colocalization	87
4. Effects of chronic hypoxia on VEGF expression	89
5. Effects of chronic hypoxia on VEGF receptor expression	90

6. VEGF-induced changes in contractile protein interactions are maintained through VEGF receptor upregulation	92
7. Effect of VEGF in culture on contractile protein abundances	94
8. Effect of VEGF on contractile protein interactions	96
9. Schematic linking hypoxia to VEGF, angiogenesis and remodeling in adult ovine carotid arteries	107
 Chapter 4	
1. Effects of long-term hypoxia on arterial reactivity to nerve stimulation.....	128
2. Effects of long-term hypoxia on expression of dopamine- β -hydroxylase.....	130
3. Effects of sympathetic perivascular nerves on arterial wall thickness and stiffness	132
4. Effect of sympathetic perivascular nerves on contractile protein organization (SM _{emb} -SM α A)	135
5. Effect of sympathetic perivascular nerves on contractile protein organization (SM1-SM α A)	136
6. Effect of sympathetic perivascular nerves on contractile protein organization (SM2-SM α A)	137
7. Sympathetic perivascular nerves mediate hypoxic remodeling through NE and it's α -1 adrenoceptors.....	139

ABBREVIATIONS

VEGF	Vascular Endothelial Growth Factor
SM	Smooth Muscle
SM α A	Smooth Muscle Alpha Actin
MLCK	Myosin Light Chain Kinase
MLC ₂₀	Myosin light chain 20 / Regulatory light chain 20
HIF	Hypoxia Inducible Factor
PIGF	Placental Growth Factor
PDGF	Platelet Derived Growth Factor
Flt-1	Fms-related tyrosine kinase
Flk-1	Fetal Liver Kinase 1
KDR	Kinase Insert Domain Receptor
D β H	Dopa beta hydroxylase

ABSTRACT OF THE DISSERTATION

VEGF and Sympathetic Perivascular Nerves Contribute to Hypoxic Remodeling of Ovine Cranial Arteries

By

Olayemi Olufikayo Adeoye

Doctor of Philosophy, Graduate Program in Physiology
Loma Linda University, June 2013
Dr. William J. Pearce, Chairperson

Chronic hypoxia complicates many pregnancies and can result in postnatal pathologies that include compromised fetal cardiovascular structure and function. Mechanisms involved remain unclear. Because hypoxia increases production of VEGF, known to modulate smooth muscle (SM) phenotype, this thesis explored the hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through direct effects on SM cells and indirectly through perivascular nerves. Using a chronic hypoxia sheep model, this work demonstrated that: 1) hypoxia potently upregulates VEGF receptor expression but not endogenous VEGF level in fetal ovine carotid arteries; 2) both chronic hypoxia and VEGF exert similar effects on smooth muscle contractile proteins; 3) both chronic hypoxia and VEGF exert similar effects on contractile protein colocalizations; and lastly, sympathetic autonomic nerves contribute to hypoxic reorganization of structure and function of vascular contractile proteins. Together, these findings advance understanding of how hypoxia precipitates fetal vascular remodeling and offer an essential first step toward finding new treatments for infants that survive in-utero hypoxia.

CHAPTER ONE

INTRODUCTION

Hypoxia

Situations ranging from physiological increases in physical activity leading to transient decreases in vascular oxygen concentration to partial or complete obliteration of oxygen supply to body tissues result in hypoxia. In many cases, organisms are able to mount adaptive mechanisms to compensate for these decreases in oxygen tension, however in other cases, they succumb to the deleterious effects of prolonged periods of hypoxia. In more general terms, hypoxia results from conditions that create a mismatch between cellular oxygen demand and hematological supply of oxygen. Everyone gets exposed to varying levels of hypoxia such as ascending to high altitude, deep sea diving, inhalation of mixture of gases with very low oxygen content such as kitchen smoke, cigarette smoke, smog from vehicles etc. Factors that determine outcome of exposure to hypoxia include but not limited to: levels of exposure, duration of exposure, age, and time to intervention. Individual genetic makeup also has a role to play in prognosis of hypoxia induced pathologies. Under many circumstances, pulse oximetry readings of between 95 and 100 are taken as sufficient indicator of adequate tissue oxygenation; levels below 90 are generally low. Whereas a lot of studies have focused on responses of adults to hypoxic conditions, not much is known about fetal responses to hypoxia.

Normal pregnancy involves a chronic state of mild glucose intolerance as well as blood volume expansion associated with considerable cardiovascular and metabolic

adaptations which when compromised result in maldevelopment of fetus (31). Whereas hypoxia is commonly referred to as a pathological phenomenon, the developing mammalian embryo which develops in a low-oxygen tension environment benefits from regulators of hypoxia which target specific genes to promote anaerobic metabolism hence reducing oxygen consumption. At the same time, other regulators of hypoxia alleviate adverse effects of hypoxia by expanding the surrounding vasculature (21).

Chronic prenatal fetal hypoxia occurs during many obstetric pathologies, including those associated with impaired placental perfusion (51) , pregnancy associated hypertension / eclampsia /preeclampsia (31, 49) and maternal diabetes (61). A 2006 study, which identified maternal influences on stillbirths and neonatal deaths in Australia, showed that, intrauterine hypoxia and asphyxia were the most common etiologies of stillbirth and were initiated during the perinatal period. Factors such as prematurity, low birth weight, smoking during pregnancy and poor prenatal care complicate further the deleterious effects of hypoxia on the developing fetus (49). Both of these modifiable and intrinsic/genetic factors culminate in conditions that result in angiopathies, which alter utero-placental perfusion and consequently intrauterine fetal hypoxia.

Although sustained or prolonged severe hypoxia is lethal to the fetus, multiple homeostatic or compensatory mechanisms enable the fetus to survive more moderate sub-lethal hypoxia. Such mechanisms typically involve increases in fetal erythropoietin levels and hematocrit. In many clinical scenarios, erythropoietin levels in fetal plasma and amniotic fluid have been used as an indicator of chronic intrauterine hypoxia, since it does not cross the placenta and is not stored (72). Chronic hypoxia also orchestrates a redistribution of cardiac output by the fetal cardiovascular system to favor the developing

brain and heart, which have very low tolerance to low oxygen tension (70). In addition to upregulation of hypoxia inducible molecular regulators, chronic fetal hypoxia also stimulates a broad variety of structural and functional changes in the developing cardiovascular system through a process collectively known as *hypoxic vascular remodeling* (70).

Hypoxic Vascular Remodeling

Since the coining of the term hypoxic vascular remodeling (63), several studies have attempted to characterize mechanisms as well as changes initiated and maintained by decreased oxygen tension. Hypoxia has been shown extensively to bring about changes in structure, biochemical composition as well as functional changes in the diverse cell types that make up the fetal vasculature (70). These changes are effected differently in different age groups, species, sites of hypoxic exposures, duration of exposure, developmental stage of exposure amongst others (70). The whole concept of hypoxic site and time specific alterations could then be reasonably suggested to be responsible for the diverse cellular and molecular alterations in different cells, tissues and organs in the body (75). A vast majority of studies done to elucidate processes involved with hypoxic vascular remodeling have shown that this process is particularly prominent in the pulmonary circulation where dramatic increases in arterial wall thicknesses and decreased contractility occur (33, 48). These structural and functional changes occur differently in large and small arteries (70), between different artery types (74), between cells of different phenotypes (75) and between mature and immature arteries (48).

Hypoxic vascular remodeling is a common and continuous process in many vascular beds that matches vascular structure and function to metabolic demand in both

the pulmonary and systemic circulations (20, 55). It is particularly important during periods of rapid growth such as those typical of intrauterine growth during pregnancy and postnatal maturation (14, 47). It is also characteristic of other pathological processes including changes due to ischemia (50) and cancer (11, 59). Because the vasculature is made up of a heterogeneous mix of cells, responses to chronic hypoxia are distinctly regulated in each of these cell types (15). Cellular responses to hypoxia involve gradients of a myriad of growth factors released from both the endothelium and the adventitial stromal cells (10, 70) as well as recruitment of various cell types e.g. bone marrow derived cells (35). Chronic exposure of fetuses to hypoxia result in deficits in transitioning from in-utero life to newborn and adult life (58). Hypoxia induced structural changes result in functional reactivity to various trophic factors. Among the many functional consequences is a marked decrease in vascular tone (2, 56). Just as the hypoxia orchestrated vascular changes are diverse, the molecular mechanisms that mediate them are equally diverse. Numerous influences including the mitogenic effects of endothelium expressed molecules such as endothelin (43), nitric oxide and soluble guanylate cyclase (66) as well as serotonin and rho-kinase (23) amongst others have been suggested as potential mediators of hypoxic vascular remodeling.

Hypoxia Inducible Factor (HIF) and Vascular Endothelial Growth Factor (VEGF)

Hypoxia Inducible Factor-1 (HIF-1) has been suggested as the global mediator of hypoxia (76). HIF-1 is a basic helix-loop-helix-PAS domain transcription factor composed of an alpha and beta subunits. Whereas under normoxic conditions, HIF-1 α is synthesized but rapidly ubiquitinated and targeted for degradation by the 26S proteasome (67), under hypoxic conditions, it stabilizes, accumulates and dimerizes with the β subunit, recruits coactivators such as Creb Binding Protein (CBP) and Protein 300 (p300) subsequently activating transcription of hundreds of genes encoding angiogenic cytokines in a cell type-specific manner (68). Hypoxic increases in HIF-1 results in transcriptional upregulation of glycolytic enzymes, erythropoietin (EPO), Placental Growth Factor (PLGF) and Vascular Endothelial Growth Factor (VEGF) amongst others.




Figure 1. Molecular mechanisms of oxygen homeostasis. Left, under normoxic conditions, HIF-1 α is synthesized, but is rapidly subjected to prolyl hydroxylation by the PHD2-OS9 complex and asparaginyl hydroxylation by FIH-1 (not shown). Prolyl hydroxylated HIF-1 α is bound by VHL, which together with SSAT2 recruits Elongin C, which in turn recruits a ubiquitin ligase complex containing Elongin B (B), ring box protein 1 (RBX1), cullin 2 (CUL2), and an E2 ubiquitin conjugating enzyme. Ubiquitination of HIF-1 α targets the protein for degradation by the 26S proteasome. Right, under hypoxic conditions, HIF-1 α accumulates, dimerizes with HIF-1 β , recruits the coactivators CBP/p300, and activates the transcription of genes encoding angiogenic cytokines including PLGF, VEGF, and SDF-1. The protein products of these genes are secreted and bind to their cognate receptors (VEGFR1, VEGFR1/VEGFR2, and CXCR4, respectively), which are located on the plasma membrane of vascular endothelial cells and circulating angiogenic cells. The diagram is simplified: cells may express one, two, or three of the receptors shown, as well as receptors for other cytokines, which are not shown. (68)

Vascular Endothelial Growth Factor

The path to discovering this very important molecule started in the late 1800s when some German pathologists observed that tumors were highly vascularized (24). This was a very fascinating finding that stirred up scientists' curiosity about possible mediators of tumor vascularity, prompting various researches to discover potent inhibitors of such mediators of extensive vascular supply of tumors. One of the pioneers of this field was Judah Folkman who made the first attempt to isolate a specific proangiogenic signal he named "tumor angiogenic factor-TAF" in 1971 (28) and suggested that application of potent inhibitors of this isolated molecule had the potential to arrest growth and differentiation of solid tumors at very early stages. In addition to the well-established establishment of extensive vascular network in tumors, Senger and Dvorak in 1983 showed that a partially purified protein later named "Vascular Permeability Factor" was able to induce leakages in blood vessels (69). In 1989, Ferrara in Genentech, isolated and cloned an endothelial cell mitogen named Vascular Endothelial Growth Factor and was later confirmed to be same as the earlier discovered Vascular Permeability Factor (26).

VEGF, also known as Vasculotropin is a 34 to 42 KDa homodimeric, heparin-binding glycoprotein. It is one of the most important growth factors with proangiogenic properties as well as serving to potentiate microvascular hyperpermeability, which precedes or occurs simultaneously as angiogenesis. The VEGF family is made up of seven members: VEGF-A, VEGF-B, VEGF-C, VEGF-D, VEGF-E, VEGF-F and PlGF. Each one of these VEGF members has a common core VEGF homology domain composed of a cysteine knot motif with eight invariant cysteine residues involved in

inter- and intramolecular disulfide bonds at one end of a conserved central four-stranded beta sheet within each monomer, which dimerize in an antiparallel, side by side orientation (36, 52).




Figure 2. Ribbon representation of the receptor-binding domain of VEGF showing a monomer in a and a dimer in b. The two-disulfide bonds are shown as black lines. Helix $\alpha 1$ consists of residues 16 to 24, and the central four-stranded β -sheet is formed by $\beta 1$ (residues 27–34), $\beta 3$ (51–58), $\beta 5$ (73–83), and $\beta 6$ (89–99), with the characteristic cysteine knot motif at one end [strands $\beta 4$ (67–69) and $\beta 7$ (103–105)], and a short three-stranded β -sheet [strands $\beta 2$ (46–49), $\beta 5$, and $\beta 6$] at the other end. VEGF monomers dimerize in an antiparallel way.

Members of the VEGF family are grouped based on the length of their tails and their ability to bind to the extracellular matrix. VEGF-A is the most important and abundant of all 7 VEGF types. In normal tissues, the highest amounts of VEGF-A are found in adult lung, kidney, heart and adrenal gland. There are at least seven homodimeric isoforms of VEGF-A containing: 121, 145, 148, 165, 183, 189 or 206 amino acids. The primary VEGF-A transcript derives from a single VEGF-A gene is derived from a single VEGF-A gene, coding for eight exons. Amino acids encoded by exons 1 to 5 and 8 are conserved in all isoforms except VEGF-148. Variable alternative splicing occurs in exons 6 and 7, which encode two distinct heparin-binding domains. These domains are important determinants of solubility and receptor binding propensities. VEGF A-121 which lacks heparin binding domains encoded by both exons 6 and 7 is highly diffusible, VEGF A-165 which has only one heparin-binding region encoded by exon 7 is moderately diffusible and VEGF A-145, 189 and 206 which contain heparin-binding domains encoded by exon 6 are tightly bound to cell surface heparin-containing proteoglycans in the extracellular matrix (54). Hypoxia results in both upregulation and stabilization of VEGF mRNA. Several studies have shown that an RNA-binding protein known as HuR mediates stabilization of VEGF mRNA. HuR does this through either direct inhibition of specific endonucleases involved with VEGF mRNA degradation or alteration of structure of the adenylate-rich elements (AREs), thus rendering those sites inaccessible to putative endonucleases (42).

VEGF has been shown to be a key regulator of physiologic – embryogenesis, reproductive functions and skeletal growth as well as pathological angiogenesis – tumors, diabetes and intraocular neovascular diseases (25). Some of the valuable physiologic

roles of VEGF shown both in-vitro and in-vivo include stimulation of surfactant production by alveolar type II cells (16), prevention of apoptosis induced by serum starvation (30), promotion of monocyte chemotaxis (13) resulting in colony formation by mature subsets of progenitor cells in the bone marrow. Its ability to induce vascular leakages (69) (22) has been implicated in inflammatory and other pathological processes. Biological effects of VEGF are mediated by three tyrosine kinase VEGF receptors: The fms-like tyrosine kinase Flt-1 (VEGFR-1/Flt-1), the kinase domain region, also known as fetal liver kinase (VEGFR-2/KDR/Flk-1) and Flt-4 (VEGFR-3) (36). Most previous work suggests that VEGF acts mainly on the vascular endothelium (44) (27); some actually say it's a specific mitogen for the endothelium (36) (46). In recent times however, there has been a growing evidence that VEGF has potent trophic effects on non-endothelial cells including smooth muscle cells (8) (12) (57) (34) (4), placenta (37) (3) (62) and neurons (7) (53). VEGF primarily acts through VEGF tyrosine receptor kinases resident on these cell types. VEGF has also been shown to exert potent chemoattractant effect on various cell types, trafficking them into blood vessel walls, these cells include pericytes (5), progenitor cells (19, 35) (18) etc.




Figure 3. Pathway showing VEGF signaling through its VEGF receptor 2 (Flk-1 in sheep and KDR in humans). VEGF mediates processes of cell survival through Src / PI3K / AKT pathway, migration through p38 / MAPK pathway and proliferation through PLC γ / PKC pathway.

Neurogenic Regulation of Cerebral Blood Flow

VEGF and The Neuron

Of much clinical relevance is the discovery that VEGF can promote the growth and differentiation of the sympathetic perivascular innervation (45). This exciting finding spurred research that has shown neuroprotective potential for VEGF and hence its promising therapeutic advantages (65). This non-angiogenic activity of VEGF on neurons stems from the finding that VEGF receptors are resident on neurons and astrocytes in the central nervous system (64). Together, these studies provide evidence for a potential role of VEGF in hypoxic vascular remodeling, because the sympathetic perivascular innervation has been shown to exert a robust trophic influence on vascular smooth muscle (6) (9) (17). One of the main neurotransmitter vasotrophic factor released from sympathetic nerves is Norepinephrine (71), which exerts its trophic influence through activation of alpha-adrenergic-1A receptors and subsequent activation of ERK (77). Other potent nerve released factors include neuropeptide-Y, which has been shown to colocalize with Norepinephrine in synaptic vesicles (73). NPY stimulates smooth muscle contraction through activation of Y1 receptors, which also mediate activation of calmodulin dependent kinase and ERK thereby influencing transcription (73) (60). Sympathetic nerves can also release ATP which together with its breakdown product adenosine, can exert additional trophic effects on smooth muscle (14) (9) (73) (1).

VEGF, Transmural Gradients and Phenotype Transformation

Blood vessels are composed of a highly heterogeneous group of cells that are able to adopt various phenotypes depending on the prevailing intrinsic and extrinsic stimuli (29) (32) (38). A huge variety of vasotrophic factors have been suggested to exert effects

resulting in phenotypic changes in these cells. Vascular endothelium, which is closest in proximity to the circulating blood and affected by factors such as perfusion and shear stress as well as non-vascular parenchyma have both been suggested as potential sources of various vasotrophic factors (40) (41). Some studies have suggested that vasotrophic factor gradients are set up between vascular endothelium and parenchyma (29) (32). One of such factors with great propensity to set up gradients in the vascular bed is VEGF. With a potent ability to mediate vascular permeability, promote endothelial cell transformation and promote angiogenesis, VEGF remains a logical candidate to be considered in vascular remodeling. Our research group tested the hypothesis that smooth muscle cells are organized into lamina of similar phenotype with characteristics that depend on the relative position between the lumen and the adventitia and involve the direct effects of VEGF on arterial smooth muscle independent of the vascular endothelium (10). We showed that VEGF directly alters arterial structure and contractility through endothelium-independent effects on cells that occupy the vascular intima (10). Various experimental data also showed that despite the fact that both VEGF receptors 1 (flt-1) and 2 (flk-1) were present in both fetal and adult arteries, VEGF effects mediated contractile effects only in the adult arteries. Previous data also show that the expression and colocalization between contractile proteins vary with vasotrophic gradients across the artery wall. VEGF has also been shown to alter phenotype of medial wall cells as evidenced by a transition in expression and colocalization between heavy chain myosin isoforms and their contractile thin actin filaments. In our previous studies, whereas non-muscle myosin heavy chain isoform (NM-MHC) increased with hypoxia in both fetal and adult arteries, total smooth muscle myosin heavy chain (SM-MHC)

decreased only in the fetal with a significant increase in adult arteries (39). Low dose VEGF in culture produced similar patterns of change as hypoxia on NM-MHC abundance in fetal but not adult arteries. VEGF had no effect on SM-MHC abundance in both fetal and adult arteries. Changes in NM-MHC correlate with contractile changes observed in both age groups. Since SM-MHC abundance increases in adult arteries could not explain decreased myogenic tone observed, colocalization studies were done. Colocalization data shows that chronic hypoxia and VEGF decrease colocalization of SM-MHC with smooth muscle alpha actin, which explains why contractile tone reduces with hypoxia (39).

Together these initial work suggests that long-term hypoxia significantly alters tone, contractile protein abundance and colocalization and that VEGF contributes to this process through alteration of contractile protein abundance and colocalization as well as setting up of gradients to which different groups of cells react differently (10). This dissertation explored the hypothesis that VEGF and sympathetic perivascular nerves contribute to hypoxic remodeling of ovine cranial arteries.

References

1. **Adair TH.** Growth regulation of the vascular system: an emerging role for adenosine. *Am J Physiol Regul Integr Comp Physiol* 289: R283-R296, 2005: PMID. 16014444.
2. **Adeoye OO, Butler SM, Hubbell MC, Semotiuk AJ, Williams JM, and Pearce WJ.** Contribution of Increased VEGF Receptors to Hypoxic Changes in Fetal Ovine Carotid Artery Contractile Proteins. *Am J Physiol Cell Physiol*, 2013: PMID. 23325408.
3. **Ahmed A, Dunk C, Ahmad S, and Khaliq A.** Regulation of placental vascular endothelial growth factor (VEGF) and placenta growth factor (PIGF) and soluble Flt-1 by oxygen--a review. *Placenta* 21 Suppl A: S16-24, 2000: PMID. 10831117.
4. **Banerjee S, Mehta S, Haque I, Sengupta K, Dhar K, Kambhampati S, Van Veldhuizen PJ, and Banerjee SK.** VEGF-A165 induces human aortic smooth muscle cell migration by activating neuropilin-1-VEGFR1-PI3K axis. *Biochemistry* 47: 3345-3351, 2008: PMID. 18284215.
5. **Benjamin LE, Hemo I, and Keshet E.** A plasticity window for blood vessel remodelling is defined by pericyte coverage of the preformed endothelial network and is regulated by PDGF-B and VEGF. *Development* 125: 1591-1598, 1998: PMID. 9521897.
6. **Bevan RD, Tsuru H, and Bevan JA.** Cerebral artery mass in the rabbit is reduced by chronic sympathetic denervation. *Stroke* 14: 393-396, 1983: PMID. 6362090.
7. **Bogaert E, Van Damme P, Poesen K, Dhondt J, Hersmus N, Kiraly D, Scheveneels W, Robberecht W, and Van Den Bosch L.** VEGF protects motor neurons against excitotoxicity by upregulation of GluR2. *Neurobiol Aging* 31: 2185-2191, 2010: PMID. 19185395.
8. **Brogi E, Schatteman G, Wu T, Kim EA, Varticovski L, Keyt B, and Isner JM.** Hypoxia-induced paracrine regulation of vascular endothelial growth factor receptor expression. *J Clin Invest* 97: 469-476, 1996: PMID. 8567969.

9. **Burnstock G.** Dual control of vascular tone and remodelling by ATP released from nerves and endothelial cells. *Pharmacol Rep* 60: 12-20, 2008: PMID. 18276981.
10. **Butler SM, Abrassart JM, Hubbell MC, Adeoye O, Semotiuk A, Williams JM, Mata-Greenwood E, Khorram O, and Pearce WJ.** Contributions of VEGF to age-dependent transmural gradients in contractile protein expression in ovine carotid arteries. *Am J Physiol Cell Physiol* 301: C653-666, 2011: PMID. 21653901.
11. **Carmeliet P.** Angiogenesis in health and disease. *Nat Med* 9: 653-660, 2003: PMID. 12778163.
12. **Christou H, Yoshida A, Arthur V, Morita T, and Kourembanas S.** Increased vascular endothelial growth factor production in the lungs of rats with hypoxia-induced pulmonary hypertension. *Am J Respir Cell Mol Biol* 18: 768-776, 1998: PMID. 9618381.
13. **Clauss M, Gerlach M, Gerlach H, Brett J, Wang F, Familletti PC, Pan YC, Olander JV, Connolly DT, and Stern D.** Vascular permeability factor: a tumor-derived polypeptide that induces endothelial cell and monocyte procoagulant activity, and promotes monocyte migration. *J Exp Med* 172: 1535-1545, 1990: PMID. 2258694.
14. **Clyman RI, Seidner SR, Kajino H, Roman C, Koch CJ, Ferrara N, Waleh N, Mauray F, Chen YQ, Perkett EA, and Quinn T.** VEGF regulates remodeling during permanent anatomic closure of the ductus arteriosus. *Am J Physiol Regul Integr Comp Physiol* 282: R199-206, 2002: PMID. 11742839.
15. **Cogo A, Napolitano G, Michoud MC, Barbon DR, Ward M, and Martin JG.** Effects of hypoxia on rat airway smooth muscle cell proliferation. *J Appl Physiol* 94: 1403-1409, 2003: PMID. 12626471.
16. **Compernelle V, Brusselmans K, Acker T, Hoet P, Tjwa M, Beck H, Plaisance S, Dor Y, Keshet E, Lupu F, Nemery B, Dewerchin M, Van Veldhoven P, Plate K, Moons L, Collen D, and Carmeliet P.** Loss of HIF-2alpha and inhibition of VEGF impair fetal lung maturation, whereas treatment with VEGF prevents fatal

- respiratory distress in premature mice. *Nat Med* 8: 702-710, 2002: PMID. 12053176.
17. **Damon DH.** Sympathetic innervation promotes vascular smooth muscle differentiation. *Am J Physiol Heart Circ Physiol* 288: H2785-2791, 2005: PMID. 15665063.
 18. **Daniel JM and Sedding DG.** Circulating smooth muscle progenitor cells in arterial remodeling. *J Mol Cell Cardiol* 50: 273-279, 2011: PMID. 21047514.
 19. **Davie NJ, Crossno JT, Jr., Frid MG, Hofmeister SE, Reeves JT, Hyde DM, Carpenter TC, Brunetti JA, McNiece IK, and Stenmark KR.** Hypoxia-induced pulmonary artery adventitial remodeling and neovascularization: contribution of progenitor cells. *Am J Physiol Lung Cell Mol Physiol* 286: L668-678, 2004: PMID. 12754186.
 20. **Distler JH, Hirth A, Kurowska-Stolarska M, Gay RE, Gay S, and Distler O.** Angiogenic and angiostatic factors in the molecular control of angiogenesis. *Q J Nucl Med* 47: 149-161, 2003: PMID. 12897707.
 21. **Dunwoodie SL.** The role of hypoxia in development of the Mammalian embryo. *Dev Cell* 17: 755-773, 2009: PMID. 20059947.
 22. **Dvorak HF, Brown LF, Detmar M, and Dvorak AM.** Vascular permeability factor/vascular endothelial growth factor, microvascular hyperpermeability, and angiogenesis. *Am J Pathol* 146: 1029-1039, 1995: PMID. 7538264.
 23. **Fagan KA, Oka M, Bauer NR, Gebb SA, Ivy DD, Morris KG, and McMurtry IF.** Attenuation of acute hypoxic pulmonary vasoconstriction and hypoxic pulmonary hypertension in mice by inhibition of Rho-kinase. *Am J Physiol Lung Cell Mol Physiol* 287: L656-664, 2004: PMID. 14977625.
 24. **Ferrara N.** VEGF and the quest for tumour angiogenesis factors. *Nat Rev Cancer* 2: 795-803, 2002: PMID. 12360282.
 25. **Ferrara N, Gerber HP, and LeCouter J.** The biology of VEGF and its receptors. *Nat Med* 9: 669-676, 2003: PMID. 12778165.

26. **Ferrara N and Henzel WJ.** Pituitary follicular cells secrete a novel heparin-binding growth factor specific for vascular endothelial cells. *Biochem Biophys Res Commun* 161: 851-858, 1989: PMID. 2735925.
27. **Fischer S, Clauss M, Wiesnet M, Renz D, Schaper W, and Karliczek GF.** Hypoxia induces permeability in brain microvessel endothelial cells via VEGF and NO. *Am J Physiol* 276: C812-820, 1999: PMID. 10199811.
28. **Folkman J, Merler E, Abernathy C, and Williams G.** Isolation of a tumor factor responsible for angiogenesis. *J Exp Med* 133: 275-288, 1971: PMID. 4332371.
29. **Frid MG, Moiseeva EP, and Stenmark KR.** Multiple phenotypically distinct smooth muscle cell populations exist in the adult and developing bovine pulmonary arterial media in vivo. *Circ Res* 75: 669-681, 1994: PMID. 7923613.
30. **Gerber HP, McMurtrey A, Kowalski J, Yan M, Keyt BA, Dixit V, and Ferrara N.** Vascular endothelial growth factor regulates endothelial cell survival through the phosphatidylinositol 3'-kinase/Akt signal transduction pathway. Requirement for Flk-1/KDR activation. *J Biol Chem* 273: 30336-30343, 1998: PMID. 9804796.
31. **Gilbert JS, Nijland MJ, and Knoblich P.** Placental ischemia and cardiovascular dysfunction in preeclampsia and beyond: making the connections. *Expert Rev Cardiovasc Ther* 6: 1367-1377, 2008: PMID. 19018690.
32. **Giuriato L, Scatena M, Chiavegato A, Tonello M, Scannapieco G, Pauletto P, and Sartore S.** Non-muscle myosin isoforms and cell heterogeneity in developing rabbit vascular smooth muscle. *Journal of cell science* 101 (Pt 1): 233-246, 1992: PMID. 1533224.
33. **Griffith SL, Rhoades RA, and Packer CS.** Pulmonary arterial smooth muscle contractility in hypoxia-induced pulmonary hypertension. *J Appl Physiol* 77: 406-414, 1994: PMID. 7961262.
34. **Grosskreutz CL, Anand-Apte B, Duplaa C, Quinn TP, Terman BI, Zetter B, and D'Amore PA.** Vascular endothelial growth factor-induced migration of vascular smooth muscle cells in vitro. *Microvasc Res* 58: 128-136, 1999: PMID. 10458928.

35. **Hayashida K, Fujita J, Miyake Y, Kawada H, Ando K, Ogawa S, and Fukuda K.** Bone marrow-derived cells contribute to pulmonary vascular remodeling in hypoxia-induced pulmonary hypertension. *Chest* 127: 1793-1798, 2005: PMID. 15888860.
36. **Hoeben A, Landuyt B, Highley MS, Wildiers H, Van Oosterom AT, and De Bruijn EA.** Vascular endothelial growth factor and angiogenesis. *Pharmacol Rev* 56: 549-580, 2004: PMID. 15602010.
37. **Hoffmann P, Feige JJ, and Alfaidy N.** Placental expression of EG-VEGF and its receptors PKR1 (prokineticin receptor-1) and PKR2 throughout mouse gestation. *Placenta* 28: 1049-1058, 2007: PMID. 17531315.
38. **Holifield B, Helgason T, Jemelka S, Taylor A, Navran S, Allen J, and Seidel C.** Differentiated vascular myocytes: are they involved in neointimal formation? *J Clin Invest* 97: 814-825, 1996: PMID. 8609239.
39. **Hubbell MC, Semotiuk AJ, Thorpe RB, Adeoye OO, Butler SM, Williams JM, Khorram O, and Pearce WJ.** Chronic hypoxia and VEGF differentially modulate abundance and organization of myosin heavy chain isoforms in fetal and adult ovine arteries. *Am J Physiol Cell Physiol* 303: C1090-1103, 2012: PMID. 22992677.
40. **Kermani P and Hempstead B.** Brain-derived neurotrophic factor: a newly described mediator of angiogenesis. *Trends Cardiovasc Med* 17: 140-143, 2007: PMID. 17482097.
41. **Leung DW, Cachianes G, Kuang WJ, Goeddel DV, and Ferrara N.** Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 246: 1306-1309, 1989: PMID. 2479986.
42. **Levy AP.** Hypoxic regulation of VEGF mRNA stability by RNA-binding proteins. *Trends Cardiovasc Med* 8: 246-250, 1998: PMID. 14987559.
43. **Li H, Chen SJ, Chen YF, Meng QC, Durand J, Oparil S, and Elton TS.** Enhanced endothelin-1 and endothelin receptor gene expression in chronic hypoxia. *J Appl Physiol* 77: 1451-1459, 1994: PMID. 7836152.

44. **Liu Y, Cox SR, Morita T, and Kourembanas S.** Hypoxia regulates vascular endothelial growth factor gene expression in endothelial cells. Identification of a 5' enhancer. *Circ Res* 77: 638-643, 1995: PMID. 7641334.
45. **Marko SB and Damon DH.** VEGF promotes vascular sympathetic innervation. *Am J Physiol Heart Circ Physiol* 294: H2646-2652, 2008: PMID. 18408130.
46. **Mata-Greenwood E, Meyrick B, Soifer SJ, Fineman JR, and Black SM.** Expression of VEGF and its receptors Flt-1 and Flk-1/KDR is altered in lambs with increased pulmonary blood flow and pulmonary hypertension. *Am J Physiol Lung Cell Mol Physiol* 285: L222-231, 2003: PMID. 12665467.
47. **Ment LR, Stewart WB, Fronc R, Seashore C, Mahooti S, Scaramuzzino D, and Madri JA.** Vascular endothelial growth factor mediates reactive angiogenesis in the postnatal developing brain. *Brain Res Dev Brain Res* 100: 52-61, 1997: PMID. 9174246.
48. **Meyrick B and Reid L.** Normal postnatal development of the media of the rat hilar pulmonary artery and its remodeling by chronic hypoxia. *Lab Invest* 46: 505-514, 1982: PMID. 7078094.
49. **Mohsin M, Bauman AE, and Jalaludin B.** The influence of antenatal and maternal factors on stillbirths and neonatal deaths in New South Wales, Australia. *J Biosoc Sci* 38: 643-657, 2006: PMID. 16867210.
50. **Mu D, Jiang X, Sheldon RA, Fox CK, Hamrick SE, Vexler ZS, and Ferriero DM.** Regulation of hypoxia-inducible factor 1alpha and induction of vascular endothelial growth factor in a rat neonatal stroke model. *Neurobiol Dis* 14: 524-534, 2003: PMID. 14678768.
51. **Neerhof MG and Thaete LG.** The fetal response to chronic placental insufficiency. *Semin Perinatol* 32: 201-205, 2008: PMID. 18482622.
52. **Neufeld G, Cohen T, Gengrinovitch S, and Poltorak Z.** Vascular endothelial growth factor (VEGF) and its receptors. *FASEB J* 13: 9-22, 1999: PMID. 9872925.

53. **Ogunshola OO, Stewart WB, Mihalcik V, Solli T, Madri JA, and Ment LR.** Neuronal VEGF expression correlates with angiogenesis in postnatal developing rat brain. *Brain Res Dev Brain Res* 119: 139-153, 2000: PMID. 10648880.
54. **Ortega N, Hutchings H, and Plouet J.** Signal relays in the VEGF system. *Front Biosci* 4: D141-152, 1999: PMID. 9924144.
55. **Owens GK, Kumar MS, and Wamhoff BR.** Molecular regulation of vascular smooth muscle cell differentiation in development and disease. *Physiol Rev* 84: 767-801, 2004: PMID. 15269336.
56. **Packer CS, Roepke JE, Oberlies NH, and Rhoades RA.** Myosin isoform shifts and decreased reactivity in hypoxia-induced hypertensive pulmonary arterial muscle. *Am J Physiol* 274: L775-785, 1998: PMID. 9612293.
57. **Partovian C, Adnot S, Eddahibi S, Teiger E, Levame M, Dreyfus P, Raffestin B, and Frelin C.** Heart and lung VEGF mRNA expression in rats with monocrotaline- or hypoxia-induced pulmonary hypertension. *Am J Physiol* 275: H1948-1956, 1998: PMID. 9843792.
58. **Pearce WJ, Hull AD, Long DM, and Longo LD.** Developmental changes in ovine cerebral artery composition and reactivity. *Am J Physiol* 261: R458-465, 1991: PMID. 1877701.
59. **Plate KH, Breier G, Millauer B, Ullrich A, and Risau W.** Up-regulation of vascular endothelial growth factor and its cognate receptors in a rat glioma model of tumor angiogenesis. *Cancer Res* 53: 5822-5827, 1993: PMID. 7694795.
60. **Pons J, Kitlinska J, Jacques D, Perreault C, Nader M, Everhart L, Zhang Y, and Zukowska Z.** Interactions of multiple signaling pathways in neuropeptide Y-mediated bimodal vascular smooth muscle cell growth. *Can J Physiol Pharmacol* 86: 438-448, 2008: PMID. 18641693.
61. **Rackham O, Paize F, and Weindling AM.** Cause of death in infants of women with pregestational diabetes mellitus and the relationship with glycemic control. *Postgrad Med* 121: 26-32, 2009: PMID. 19641265.

62. **Ratajczak MZ, Ratajczak J, Machalinski B, Majka M, Marlicz W, Carter A, Pietrkowski Z, and Gewirtz AM.** Role of vascular endothelial growth factor (VEGF) and placenta-derived growth factor (PIGF) in regulating human haemopoietic cell growth. *Br J Haematol* 103: 969-979, 1998: PMID. 9886308.
63. **Reid LM.** The pulmonary circulation: remodeling in growth and disease. The 1978 J. Burns Amberson lecture. *Am Rev Respir Dis* 119: 531-546, 1979: PMID. 443626.
64. **Rosenstein JM and Krum JM.** New roles for VEGF in nervous tissue--beyond blood vessels. *Exp Neurol* 187: 246-253, 2004: PMID. 15144851.
65. **Ruiz de Almodovar C, Lambrechts D, Mazzone M, and Carmeliet P.** Role and therapeutic potential of VEGF in the nervous system. *Physiol Rev* 89: 607-648, 2009: PMID. 19342615.
66. **Schermuly RT, Stasch JP, Pullamsetti SS, Middendorff R, Muller D, Schluter KD, Dingendorf A, Hackemack S, Kolosionek E, Kaulen C, Dumitrascu R, Weissmann N, Mittendorf J, Klepetko W, Seeger W, Ghofrani HA, and Grimminger F.** Expression and function of soluble guanylate cyclase in pulmonary arterial hypertension. *Eur Respir J* 32: 881-891, 2008: PMID. 18550612.
67. **Semenza GL.** Hypoxia-inducible factor 1 (HIF-1) pathway. *Sci STKE* 2007: cm8, 2007: PMID. 17925579.
68. **Semenza GL.** Vasculogenesis, angiogenesis, and arteriogenesis: mechanisms of blood vessel formation and remodeling. *J Cell Biochem* 102: 840-847, 2007: PMID. 17891779.
69. **Senger DR, Galli SJ, Dvorak AM, Perruzzi CA, Harvey VS, and Dvorak HF.** Tumor cells secrete a vascular permeability factor that promotes accumulation of ascites fluid. *Science* 219: 983-985, 1983: PMID. 6823562.
70. **Stenmark KR, Fagan KA, and Frid MG.** Hypoxia-induced pulmonary vascular remodeling: cellular and molecular mechanisms. *Circ Res* 99: 675-691, 2006: PMID. 17008597.

71. **Teeters JC, Erami C, Zhang H, and Faber JE.** Systemic alpha 1A-adrenoceptor antagonist inhibits neointimal growth after balloon injury of rat carotid artery. *Am J Physiol Heart Circ Physiol* 284: H385-392, 2003: PMID. 12388268.
72. **Teramo KA and Widness JA.** Increased fetal plasma and amniotic fluid erythropoietin concentrations: markers of intrauterine hypoxia. *Neonatology* 95: 105-116, 2009: PMID. 18776724.
73. **Wier WG, Zang WJ, Lamont C, and Raina H.** Sympathetic neurogenic Ca²⁺ signalling in rat arteries: ATP, noradrenaline and neuropeptide Y. *Exp Physiol* 94: 31-37, 2009: PMID. 18931047.
74. **Williams JM and Pearce WJ.** Age-dependent modulation of endothelium-dependent vasodilatation by chronic hypoxia in ovine cranial arteries. *J Appl Physiol* 100: 225-232, 2006: PMID. 16179402.
75. **Wohrley JD, Frid MG, Moiseeva EP, Orton EC, Belknap JK, and Stenmark KR.** Hypoxia selectively induces proliferation in a specific subpopulation of smooth muscle cells in the bovine neonatal pulmonary arterial media. *J Clin Invest* 96: 273-281, 1995: PMID. 7615796.
76. **Yamakawa M, Liu LX, Date T, Belanger AJ, Vincent KA, Akita GY, Kuriyama T, Cheng SH, Gregory RJ, and Jiang C.** Hypoxia-inducible factor-1 mediates activation of cultured vascular endothelial cells by inducing multiple angiogenic factors. *Circ Res* 93: 664-673, 2003: PMID. 12958144.
77. **Zhang H, Chalothorn D, Jackson LF, Lee DC, and Faber JE.** Transactivation of epidermal growth factor receptor mediates catecholamine-induced growth of vascular smooth muscle. *Circ Res* 95: 989-997, 2004: PMID. 15486316.

CHAPTER TWO
CONTRIBUTION OF INCREASED VEGF RECEPTORS TO
HYPOXIC CHANGES IN FETAL OVINE CAROTID ARTERY
CONTRACTILE PROTEINS

**Olayemi Adeoye, Stacy M. Butler, Margaret C. Hubbell, Andrew Semotiuk,
James M. Williams and William J. Pearce**

Division of Physiology, Pharmacology and Biochemistry
Center for Perinatal Biology
Loma Linda University School of Medicine
Loma Linda, CA 92350

Abstract

Recent studies suggest that vascular endothelial growth factor (VEGF) can modulate smooth muscle phenotype and consequently the composition and function of arteries upstream from the microcirculation where angiogenesis occurs. Given that hypoxia potently induces VEGF, the present study explores the hypothesis that in fetal arteries VEGF contributes to hypoxic vascular remodeling through changes in the abundance, organization and function of contractile proteins. Pregnant ewes were acclimatized at sea level or at altitude (3820 m) for the final 110 days of gestation. Carotid arteries collected from the fetuses at term were denuded of endothelium and used fresh or after organ culture for 24 h in a physiological concentration (3 ng/ml) of VEGF. After 110 days, hypoxia had no effect on VEGF abundance, but markedly increased abundances of both the Flk-1 (171%) and Flt-1 (786%) VEGF receptors. Hypoxia had no effect on smooth muscle α -actin (SM α A), decreased myosin light chain kinase (MLCK), and increased 20 kDa regulatory myosin light chain (MLC₂₀) abundances. Hypoxia also increased colocalization of MLCK with SM α A, MLC₂₀ with SM α A, and MLCK with MLC₂₀. Compared to hypoxia, organ culture with VEGF produced the very same pattern of changes in contractile protein abundances and colocalization. Effects of VEGF on colocalization were blocked by the VEGF receptor antagonists vatalanib (240 nM) and dasatinib (6.3 nM). Thus, through increases in VEGF receptor density, hypoxia can recruit VEGF to help mediate remodeling of fetal arteries upstream from the microcirculation. The results support the hypothesis that VEGF contributes to hypoxic vascular remodeling through changes in the abundance, organization and function of contractile proteins.

Key Words

Myosin Light Chain Kinase, Organ Culture, Regulatory Myosin Light Chain,
Smooth Muscle α -Actin, VEGF Receptors

Introduction

In-utero hypoxia secondary to maternal diabetes (49), smoking (3), pulmonary insufficiency or placental malformation (36) is a frequent etiologic factor in many complicated pregnancies (25). These hypoxic insults result in numerous perinatal and postnatal morbidities (30) that often include altered fetal cardiovascular function secondary to atypical patterns of vascular structure and contractility collectively known as remodeling (16, 32). Whereas the exact molecular mechanisms governing hypoxic fetal vascular remodeling remain poorly understood, considerable evidence in adult vasculature demonstrates that chronic hypoxia orchestrates arterial wall thickening (17, 39) together with changes in contractile protein abundance (48) and organization (55). The functional consequences of these changes include altered arterial contractility (8, 51) due at least in part to modified reactivity of both thick and thin filament contractile proteins (35).

Efforts to understand the mechanisms that drive hypoxic vascular remodeling have focused largely on the vasotrophic factors released by hypoxia, the most prominent of which is hypoxia inducible factor (HIF) (46, 56). Upregulation, stabilization and dimerization of HIF mediate transcription of multiple angiogenic genes including VEGF (Figure 1), which in turn is known traditionally to mediate capillary angiogenesis (18). In further support of angiogenesis, some studies have also suggested that hypoxia can modulate levels of the VEGF receptors 1 and 2 (Flt-1 and Flk-1) in the vascular endothelium as well as in certain tumors of rodent brain (43, 50) (Figure 1). Whereas abundant previous work suggests that VEGF acts mainly on the vascular endothelium (18), recent findings further suggest that VEGF can exert potent trophic effects on

multiple non-endothelial cell types (24) suggesting possible expression of VEGF receptors on such cell types. Of particular relevance to vascular remodeling is growing evidence that VEGF has potent trophic effects on smooth muscle cells (12) that result in alteration of contractile protein expression and organization (7).

The current study explores the hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through changes in the abundance, organization and function of contractile proteins. Through increases in VEGF and/or its receptors in smooth muscle, we propose that hypoxia could recruit VEGF to help mediate changes in ovine fetal vascular structure and function that are characteristic of chronic hypoxia. Given the central importance of the rate-limiting enzyme myosin light chain kinase (MLCK) (41), its substrate, regulatory myosin light chain (MLC₂₀) (20), and their contractile partner smooth muscle α -Actin (SM α A) (15), our experimental design focused on mechanisms involved in hypoxia-induced changes in these contractile proteins. Comparisons between fetal arteries harvested from sheep maintained at sea level and those maintained at high altitude (3820m) for 110 days served to define effects of chronic hypoxia, as previously described in detail (32). To assess the role of VEGF and its receptors in hypoxic remodeling (Figure 1), we used organ cultures of whole carotid arteries, which preserved the spatial organization of all cell types in the arterial wall (24). To explain changes in smooth muscle phenotype orchestrated by hypoxia, which ultimately dictate contractile protein abundance and organization, we assessed contractile protein abundances via Western blots and protein organization via a novel confocal colocalization imaging technique (9). Stress-strain measurements of myogenic reactivity normalized to arterial wall cross-sectional area (34) identified the functional consequences of arterial

remodeling induced by hypoxia. Together, these studies provided a unique perspective of the direct contribution of VEGF and its receptors to hypoxic vascular remodeling in fetal ovine carotid arteries.




Figure 1: This schematic summarizes our approach to test the hypothesis that VEGF contributes to hypoxic vascular remodeling through changes in the abundance, organization and function of contractile proteins in fetal arteries. First, we propose that hypoxia induces short-term increases in VEGF (**Arrow 1**) through upregulation of the transcription factor hypoxia inducible factor. We further propose that these increases in VEGF act on VEGF receptors expressed by smooth muscle cells (**Arrow 2**). In addition, we propose that chronic hypoxia increases expression of smooth muscle VEGF receptors (**Arrow 3**). Finally, we propose that activation of smooth muscle VEGF receptors leads to changes in contractile protein abundance and organization that result in changes in arterial structure and function (**Arrow 4**). In this manner, we propose that hypoxic increases in VEGF mediate not only microcirculatory angiogenesis, but also arterial remodeling. Separate experiments were performed to test each of the numbered arrows in fetal arteries.

Materials and Methods

All techniques, protocols and procedures used in these studies received approval from the Animal Research Committee of Loma Linda University and complied with all policies and codes of practice outlined in the National Institutes of Health Guide for the Care and Use of Laboratory Animals. Procedures related to tissue harvesting, tissue preparation and animal surgery have been previously described in detail (54).

Tissue Harvest and Preparation

All experiments used common carotid arteries harvested from term fetal lambs (139 – 142 days gestation) delivered by Caesarian using strict sterile techniques. Pregnant ewes were anesthetized with 30 mg/kg pentobarbital, intubated and maintained on 1.5-2.0% halothane. The fetus was subsequently exteriorized via a midline vertical incision and killed by exsanguination via rapid removal of the heart. Normoxic arteries (FN) were harvested from animals maintained at sea level whereas hypoxic arteries (FH) were harvested from animals acclimatized at altitude (3820m above sea level) for the final 110 days of gestation, as previously described in detail (26, 32). This chronic hypoxia model yields arterial oxygen tensions (PaO_2) of 19 ± 1 Torr in fetal sheep (26). Corresponding normoxic arterial oxygen tensions average 23 ± 1 Torr. Harvested common carotid arteries were kept in sterile HEPES buffer solution composed of (in mM) 122.1 NaCl, 25 HEPES, 5.16 KCl, 2.4 MgSO_4 , 11.1 dextrose, 1.6 CaCl_2 and 50 μM EDTA. After a thorough and careful removal of the loose connective tissues and blood, the arteries were denuded of endothelium via mechanical abrasion. Next, 3 mm lengths of artery were cut and distributed to the various experimental protocols. Medial thicknesses and exact arterial segment length measurements were completed using an Olympus U-PMTVC

Optical microscope mounted with a Scion Visicapture Twain 1394 Camera for image capture with ImagePro software (v6.0, Media Cybernetics).

Contractility Studies

The 3-mm arterial segments were mounted on tungsten wires between isometric force transducers and micrometers used for precise positioning. Each artery was equilibrated for one hour in calcium replete Na^+ -Krebs buffer (pH 7.4) containing (in mM): 122 NaCl, 25.6 NaHCO_3 , 5.17 KCl, 2.56 dextrose, 2.49 MgSO_4 , 1.60 CaCl_2 , 0.114 ascorbic acid and 0.027 EGTA. Artery viability was preserved by diffusional oxygen supply enabled by continuous bubbling with 95% O_2 with 5% CO_2 at normal ovine core temperature (38 °C) (40). Unstressed artery diameters were measured immediately following equilibration at a passive tension of 0.03 g. Relative to unstressed diameters (D_0), working diameters (D) required to attain arterial strain ratios (D/D_0) of 1.5, 1.8, 2.1, 2.3, 2.7, 3.0 and 3.3 were calculated. Under resting conditions and after addition of high potassium buffer solutions, contractile stresses (dynes/cm^2) were measured at each of the strain ratios to determine spontaneous myogenic tone and potassium-induced active tone respectively. The constituents of the high potassium Krebs solution were (in mM) 122 KCl, 11.1 dextrose, 5.16 NaCl, 2.50 MgSO_4 , 2.15 NaHCO_3 , 1.60 CaCl_2 , 0.114 ascorbic acid and 0.027 EDTA. Contractile responses to high potassium were recorded until stabilized after which the arteries were returned to basal conditions using Na^+ -Krebs buffer and then were equilibrated at the next stretch ratio. After responses were recorded at the highest strain ratios, the arterial segments were frozen in liquid nitrogen to eliminate any active component of stress (7, 40). The segments were subsequently

incubated in calcium depleted Na^+ -Krebs containing 3 mM EGTA. Passive stresses produced at each strain ratio from the highest to lowest were then recorded.

Differences between spontaneous tone measured before and after freezing with liquid nitrogen and EGTA at each level of strain were taken as spontaneous myogenic tone. Stiffness coefficients were determined by assessing the relations between strain ratios and passive stresses using non-linear regression fits to a monotonic exponential model (Young's modulus) **(13)**.

Fluorescent Immunohistochemistry

Segments of common carotid arteries were fixed overnight in 4% neutral buffered EM-grade formaldehyde (Electron Microscopy Sciences, Hatfield, #15713S), paraffin embedded and sectioned at 5 μm . Slides were deparaffinized in HistoClear solution (National Diagnostic, Atlanta, #HS-200) and rehydrated in descending concentrations of alcohol. The samples were microwaved in citrate buffer (pH 6.03) to recover antigenicity after which the sections were permeabilized in 0.1% Triton X-100 (Sigma Aldrich, St Louis, #T-8787) and then incubated in 1% Bovine Serum Albumin (Santa Cruz Biotechnology, Santa Cruz, #SC-2323) to block non-specific binding. Incubations with primary antibodies were carried out overnight at 4 °C. Antibody selectivity was confirmed with Western blots and the titers used for IHC were the lowest that optimized signal to noise ratio. The primary antibodies used included monoclonal anti-SM α A (Sigma Aldrich, St Louis, A5228) @1:200, polyclonal anti-SM α A (abcam, ab5694) @1:200, polyclonal anti-MLCK (Santa Cruz Biotechnology, Santa Cruz, SC-25428) @ 1:50 and monoclonal anti-MLC₂₀ (Sigma-Aldrich, St Louis, #M4401) @ 1:300. Slides

were washed in PBS for two 10-minute cycles on the morning of the next day before the secondary antibody (DyLight 488 Conjugated – Pierce Chemical, Rockford, #35502) was applied to the tissues on the slides for two hours at room temperature. Keeping the slides in the dark post-incubation preserved photosensitivity. The slides were subsequently covered and washed for two 10-minute cycles in PBS. SlowFade Gold anti-fade reagent with DAPI was then added and the slides were coverslipped and stored in the dark until imaged. Tissue slides were imaged using a Zeiss Imager A1 on an AX10 Fluorescence microscope with Spot software (Diagnostic Instruments, Inc. Ver. 4.6.4.5).

Confocal Microscopy

Artery sections of 5 μm thicknesses were double-stained with antibodies against MLCK and SM α A as described in the “Fluorescent Immunohistochemistry” section. Following primary antibody incubation, the sections were washed in PBS and equilibrated in a dark room with two secondary antibodies labeled with Dylight – 488 (SM α A) and 649 (MLCK) for two hours at room temperature. The sections were imaged with an Olympus FV1000 at an optical section thickness of 700 nm, a lateral resolution of 200 nm and a numerical aperture of 1.8. The extent of colocalization between the two markers was determined using a custom-written non-parametric quadrant analysis that calculated the fraction of total pixels in the upper half of the intensity range for both markers relative to the number of pixels in the upper half of the intensity range for SM α A. We refer to this index as the percentage of pixels in the upper right quadrant (%UR). This method of quadrant analysis was adapted from flow-cytometry theory (1, 2) and has been described previously (9, 21). Double staining was also done on 5 μm

sections on slides for MLC₂₀ with SM α A and MLCK with MLC₂₀. Primary antibodies used were monoclonal anti-MLC₂₀ (Sigma-Aldrich, St Louis, #M4401) @ 1:300 for MLC₂₀, monoclonal anti- SM α A (Sigma Aldrich, St Louis, A5228) @ 1:200 for SM α A and polyclonal MYLK (Santa Cruz Biotechnology, Santa Cruz, SC -25428) @ 1:50 for MLCK. Secondary antibodies for MLC₂₀ colocalization with SM α A (DyLight 488 - SM α A and 633 -MLC₂₀) and for MLCK colocalization with MLC₂₀ (Dylight 488 -MLC₂₀ and 633 -MLCK) (Conjugated – Pierce Chemical, Rockford, #35502) were applied to the tissues after which the sections were imaged as earlier described under “Fluorescent Immunohistochemistry”.

Western Blotting

Known weights of frozen arterial segments were homogenized via glass-on-glass in 8M Urea containing (in mM) 500 NaCl, 20 Tris, 23 Glycine, 10 EGTA and 10% Glycerol at pH 8.6 with protease inhibitor cocktail at 5 μ l/ml of buffer (Sigma-Aldrich, Saint Louis, #M1745). Centrifugation of the homogenates at 5,000G for 20 min yielded supernatants whose protein concentrations were determined by the Bio-Rad Bradford assay. Protein homogenates were separated via SDS-PAGE together with increasing concentrations of standards used to calibrate target protein abundance. The tissues for the standards were harvested from pooled adult ovine common carotid arteries. SDS-PAGE separated proteins were transferred onto nitrocellulose membranes at 200 mA for 90 minutes in Towbin’s buffer (25 mM Tris, 192 mM Glycine, 10% and 20% Methanol) on ice. Using 5% milk in Tris buffered saline at pH 7.5 (M-TBS); the membranes were blocked for 1h at room temperature using continuous gentle shaking. After the initial

blocking, subsequent washes and incubations were done in M-TBS containing 0.1% Tween-20. Incubations with primary antibody were performed for 3 hours using the following dilutions for SM α A (1:3000), MLCK (1:10,000) and MLC₂₀ (1:200) and VEGF-A₁₆₅ (1:750). All antibodies were obtained from same sources as described under the immunohistochemistry section except for anti-VEGF antibody that was purchased from Abcam (Cambridge, #AB119). Visualization was performed after a 90-minute secondary incubation using a secondary antibody conjugated to Dylight 800 (Pierce Chemical, Rockford, #46422). Imaging was completed on a LI-COR Bioscience Odyssey system.

For the VEGF receptor Western blot assay, artery segments were homogenized using a glass pestle and mortar containing buffer with (in mM) 500 NaCl, 50 Tris, and 5 EDTA at pH 7.4 with protease inhibitors including (in μ M) 500 AEBSF, 400 pepstatin-A, 20 bestatin, 10 E-64, 7.5 leupeptin, and 7 aprotinin (Sigma-Aldrich, Saint Louis, MO) at 1:50 tissue buffer ratio. Homogenates were centrifuged at 100,000 XG for 1hr at 4 °C. Pellets were resuspended at a 1:10 tissue buffer ratio with (in mM) 150 NaCl, 50 Tris, 10 DTT and 1% Triton X-100, 0.5% Sodium Deoxycholate, 0.2% SDS and 10% Glycerol with protease inhibitor cocktail as above for 1hr with gentle shaking at room temperature. Samples were ultra-sonicated 6 x 5 sec at 20% amplitude to shear DNA, centrifuged at 10,000 XG for 15 min after which the supernatants were collected. Protein concentrations were determined using Bradford's protein assay. Protein homogenates were separated on 5% SDS-PAGE with addition of 35 mM β -mercaptoethanol (BME) in the upper buffer reservoir, along with a pooled reference to normalize samples. Separated proteins were transferred onto 0.2 μ m nitrocellulose (Whatman, BA83) using

350 mA for 90 min in Towbin's buffer containing (in mM) 25 Tris, 35 BME, 192 Glycine, 0.01% SDS and 20% Methanol. Following transfer, the membranes were blocked in 5% milk with Tris-buffered saline at pH 7.5 (M-TBS) for 1 hour at room temperature using continuous shaking. All subsequent washes and incubations were performed in M-TBS with 0.1% Tween-20 included. Primary antibodies were incubated for 3 hours with the following dilutions for Fms-like tyrosine kinase receptor (Flt-1) at 1:200 (SC-316) and Fetal Liver Kinase 1 (Flk-1) at 1:200 (SC-504) from Santa Cruz. Membranes were washed 6 X 5min followed by a 90 min incubation of secondary antibody conjugated to DyLight 800 (Pierce Chemical, Rockford, IL). Membranes were washed 6 X 5min then washed in TBS only. Membranes were imaged on LI-COR Bioscience's Odyssey system.

Organ Culture

As previously described (7), serial segments of individual arteries from both experimental groups (FN and FH) were maintained in Dulbecco's Modified Eagle Media (DMEM) fortified with 3.7 g/l of Na₂HCO₃, 0.5% amino acid solution (Sigma Aldrich, St. Louis, #M5550), 1% non-essential amino acid solution (Sigma Aldrich, St. Louis, #M7145), 4 mM glutamine (Sigma Aldrich, St. Louis, #G7513), 2% antibiotic-antimycotic solution (Gibco, Carlsbad, #15240-096), and 70 µg/ml of Gentamycin (Gibco, Carlsbad, #15750-060) in an untreated 12-well plate, and subsequently maintained in a humidified incubator with 5% CO₂ in room air at 37 °C. Matched sets of artery segments left in DMEM media for 24 hrs served as control arteries while those treated with low dose (3 ng/ml) of VEGF were used to assess the effects of VEGF. Segments treated with vatalanib (240 nM) and dasatinib (6.3 nM) (42) were used to

assess the effects of VEGF tyrosine kinase receptors; the concentrations used were identified as the lowest effective concentrations in preliminary dose-finding experiments. In all organ culture experiments, the artery segments were first maintained in media without growth factors (DMEM) for 24hrs. Control arteries remained in DMEM for another 24 h, and treated arteries were exposed to 3 ng/ml VEGF for 24 hours. This low dose of VEGF was used to minimize non-specific binding of VEGF to other non-VEGF receptors and represented physiological serum levels measured in gravid sheep (52).

Data Analysis and Statistics

Contractile stresses generated from applied graded strains were calculated as ratios of force per cross-sectional area where force equals tension measured in grams multiplied by acceleration due to gravity, and cross sectional area equals working wall thickness (μm) multiplied by segment length. Each animal used contributed to the fresh, control, VEGF and VEGF+vatalanib+dasatinib groups. Contractile stress measurements, thicknesses and stiffness values were compared using analysis of variance (ANOVA). All data sets were confirmed to be normally distributed using the D'Agostino-Pearson analysis, and homogeneity of variance within ANOVA was verified using a Bartlett's-Cochran test as previously described (9). Statistical power was routinely at least 0.8.

All Western immune blot measurements were calibrated against a standard curve generated from a pooled reference of common carotid tissues. Regional abundance values were compared using a 2-way analysis of variance (ANOVA) with age and treatment as variables. Duncan's Multiple Range analysis was used to make post-hoc comparisons between means within the ANOVA.

Results

The study is based on a total of 124 segments from 17 normoxic fetal lambs and 93 segments from 12 hypoxic fetal lambs. In all cases, “n” indicates the number of animals used. Statistical significance implies $P < 0.05$. All values are presented as mean \pm SEM.

Effects of Chronic Hypoxia on Carotid Artery Structure and Stress-Strain Relations

Medial thicknesses in fresh normoxic arteries ($341 \pm 20 \mu\text{m}$) were significantly less than in fresh hypoxic arteries ($400 \pm 28 \mu\text{m}$) (**Figure 2**); chronic hypoxia significantly increased medial thicknesses by 17%. Similarly, arterial stiffness values in fresh normoxic arteries (8.02 ± 0.42) were significantly less than stiffness values in fresh hypoxic arteries (9.20 ± 0.69) (**Figure 2**); hypoxic increases in stiffness averaged to 15%. Peak myogenic tone averaged $102.5 \pm 21.8 \text{ dynes/cm}^2$ in normoxic fetal arteries but only $25.9 \pm 6.3 \text{ dynes/cm}^2$ in hypoxic arteries. On average, chronic hypoxia significantly decreased peak myogenic tone by 75%.




Figure 2: Hypoxia remodels ovine carotid artery structure and function. When compared to normoxic arteries, arteries from chronically hypoxic animals exhibited an increased thickness of the medial layer (left panel) and also increased stiffness (middle panel). Determination of active stress-strain relations revealed that chronic hypoxia also significantly depressed myogenic tone but did not alter the strain values at which contractile force was maximal. Results are presented as mean \pm SEM for arteries from normoxic (n =17) and hypoxic (n=12) fetuses. Significant differences denoted by asterisks (*) imply $P < 0.05$.

Effects of Chronic Hypoxia on Contractile Protein Abundance and Colocalization

The effects of chronic hypoxia on contractile protein abundances were highly protein specific (**Figure 3**). For SM α A, Western blot quantification revealed similar abundances (relative to standard) in both normoxic (0.62 ± 0.07) and hypoxic arteries (0.68 ± 0.08). For MLCK, abundances were 90% less in hypoxic than in normoxic arteries. In contrast, hypoxic abundances of MLC₂₀ were 61% greater than in normoxic arteries.

Chronic hypoxia significantly increased colocalization among all three pairs of contractile proteins examined (**Figure 4**). For MLCK- SM α A, chronic hypoxia increased the coefficient of colocalization 42% relative to the normoxic group. Corresponding increases for MLC₂₀- SM α A and MLCK-MLC₂₀ colocalization averaged 123% and 237%, respectively.




Figure 3: Chronic hypoxia alters smooth muscle contractile protein abundances. Western blot quantification of SM α A abundance yielded similar values in normoxic and hypoxic fetal carotid arteries (Left Panel). For MLCK abundance, the values were markedly less in hypoxic than in normoxic arteries. In contrast, MLC₂₀ abundances were significantly greater in hypoxic than in normoxic arteries. These results demonstrate that the effects of chronic hypoxia on smooth muscle contractile protein abundances are highly protein specific. Results are presented as mean \pm SEM for $n \geq 5$ in all experimental groups. Significant differences denoted by asterisks (*) imply $P < 0.05$ via ANOVA.




Figure 4: Chronic hypoxia increases colocalization among smooth muscle contractile proteins. As revealed by confocal microscopy, colocalization of MLCK with SM α A was 42% greater in hypoxic than normoxic fetal arteries. Similarly, colocalization of MLC₂₀ with SM α A was 42% greater, and colocalization of MLCK with MLC₂₀ was 123% greater, in hypoxic than in normoxic fetal arteries. All these differences were statistically significant and suggest that the contractile proteins were becoming more compact and highly organized in response to chronic hypoxia. Results are presented as mean \pm SEM for $n \geq 5$ in all experimental groups. Significant differences denoted by asterisks (*) imply $P < 0.05$ via ANOVA.

Effects of Long-term Hypoxia on Abundances of VEGF and VEGF Receptors

Western blot quantification of endogenous VEGF levels in FN (0.29 ± 0.07) and FH arteries (0.25 ± 0.06) revealed similar VEGF abundances (**Figure 5**), suggesting that hypoxia had little effect on VEGF levels after 110 days of hypoxic acclimatization. In contrast, the abundances of the endogenous VEGF receptors 1 (flt-1) and 2 (flk-1) were 786% and 171% greater in chronically hypoxic fetal arteries than in normoxic fetal arteries (**Figure 6**).




Figure 5: Effects of long-term hypoxia on VEGF abundance. Endogenous VEGF levels quantified via western blot analysis exhibited similar abundances in normoxic and hypoxic fetal sheep. These results suggest that the well-documented increases in VEGF induced upon exposure to hypoxia are transient and disappear after 110 days of hypoxic acclimatization. Results are presented as mean \pm SEM for arteries from normoxic (n=7) and hypoxic (n=5) fetal arteries.




Figure 6: Effects of long-term hypoxia on VEGF receptor expression. In homogenized endothelium-denuded fetal arteries, abundances of VEGF receptor 1 (Flt-1) and VEGF receptor 2 (Flk-1) were significantly increased by chronic hypoxia. The multiple bands indicated on the western blots (lower panel) indicate the different glycosylation states of the receptors. The results shown in the bar charts (upper panel) represent the total of all glycosylation states for Flk-1. Due to the absence of the 230 kD form of Flt-1 in hypoxic arteries (lower panel), the bar charts compare abundances of the 250 kD form. The standards shown on the western blots were prepared from mixed samples of normoxic adult arteries. Results are presented as mean \pm SEM for $n=6$ in all experimental groups. Significant differences denoted by asterisks (*) imply $P<0.05$ via ANOVA.

VEGF Receptors Mediate VEGF-induced Changes in Contractile Protein Organization

To confirm that VEGF altered contractile protein organization through a VEGF receptor-dependent mechanism, MLC₂₀-SM α A colocalization was used as a reporter assay (**Figure 7**). In organ culture, a low physiological concentration (3 ng/ml) of VEGF increased the colocalization of MLC₂₀ with SM α A by 237%. This effect of VEGF was blocked completely in the presence of the VEGF receptor blockers vatalanib (240 nM) and dasatanib (6.3 nM).




Figure 7: VEGF activation of its tyrosine kinase receptors mediates increased contractile protein colocalization. Organ culture of endothelium-denuded normoxic fetal arteries with VEGF (3ng/ml) significantly increased colocalization of MLC₂₀ with SM α A. Addition of the VEGF receptor antagonists vatalanib (240nM) and dasatinib (6.3nM) completely blocked this effect of VEGF. Results are presented as mean \pm SEM for n = 5 in all experimental groups. Significant differences denoted by (*) imply P < 0.05 via ANOVA.

Effects of VEGF on Contractile Protein Abundance and Colocalization

In organ culture, treatment with 3 ng/ml VEGF had no significant effect on SM α A abundances in either normoxic (0.59 ± 0.05 vs. 0.64 ± 0.07) or hypoxic (0.66 ± 0.07 vs. 0.64 ± 0.09) fetal arteries (**Figure 8**). For MLCK however, VEGF significantly increased MLCK abundances in normoxic arteries by 100% but decreased them in hypoxic arteries by 33%; following VEGF treatment MLCK abundances were significantly greater in normoxic than hypoxic arteries. For MLC₂₀, VEGF had no individually significant effect in normoxic (0.23 ± 0.04 vs. 0.27 ± 0.05) or hypoxic arteries (0.38 ± 0.03 vs. 0.44 ± 0.08), but after treatment with VEGF, MLC₂₀ abundances were significantly less in normoxic than hypoxic arteries. As indicated by comparison of Figures 3 and 8, chronic hypoxia and VEGF had highly similar effects on the abundances of all three contractile proteins.

The effects of VEGF and chronic hypoxia on patterns of contractile protein colocalization were also highly similar. In parallel with the effects of chronic hypoxia (**Figure 4**), organ culture with 3 ng/ml VEGF increased colocalization of MLCK and SM α A by 55%, increased colocalization of MLC₂₀ with SM α A by 237%, and increased colocalization of MLCK with MLC₂₀ by 75% (**Figure 9**). All of these effects of VEGF on contractile protein colocalization were statistically significant.




Figure 8: Effects of VEGF on contractile protein abundances are highly protein specific. As revealed by western blot quantification, organ culture with 3 ng/ml VEGF had no significant effect on SM α A abundance in endothelium-denuded arteries from either normoxic or hypoxic fetuses. In contrast, organ culture with 3 ng/ml VEGF significantly increased MLCK abundance in endothelium-denuded arteries from normoxic fetuses, but decreased it in arteries from hypoxic fetuses. Conversely, organ culture with 3 ng/ml VEGF had no significant effect on MLC₂₀ abundance in arteries from normoxic fetuses, but significantly increased MLC₂₀ in arteries from hypoxic fetuses. This pattern of effects emphasizes that the effects of VEGF on contractile protein abundance are highly protein specific, are markedly influenced by hypoxic acclimatization, and are closely similar to the effects of chronic hypoxia (**Figure 3**). Results are presented as mean \pm SEM for n = 5 in all experimental groups. Significant differences denoted by asterisks (*) imply P<0.05 via ANOVA.




Figure 9: VEGF increases colocalization among smooth muscle contractile proteins. In endothelium-denuded arteries from normoxic fetuses, organ culture of with 3 ng/ml VEGF increased colocalization of MLCK with SM α A by 55%, MLC₂₀ with SM α A by 237%, and MLCK with MLC₂₀ by 75%, compared to corresponding untreated controls. These results demonstrate that VEGF can act directly on arterial smooth muscle to enhance contractile protein colocalization. This pattern of effects was also closely similar to the effects of chronic hypoxia on contractile protein colocalization (Figure 4). Results are presented as mean \pm SEM for $n \geq 5$ in all experimental groups. Significant differences denoted by (*) imply $P < 0.05$ via ANOVA.

Discussion

This study offers three original findings suggesting a non-angiogenic role of VEGF in hypoxic remodeling of fetal ovine carotid arteries. First, in homogenates of endothelium-denuded fetal carotids, 110 days of hypoxic acclimatization had no effect on VEGF levels but significantly increased abundances of both main VEGF receptors (Flt-1 and Flk-1), relative to normoxic controls. Second, both chronic hypoxia and organ culture with VEGF were without effect on SM α A, decreased MLCK, and increased MLC₂₀ abundances in endothelium-denuded fetal carotids. Third, both chronic hypoxia and organ culture with VEGF increased colocalization of MLCK with SM α A, MLC₂₀ with SM α A, and MLCK with MLC₂₀. Together, these findings support the general hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through changes in the abundance, organization and function of contractile proteins.

Effects of Hypoxia on Structure and Contractility

Previous studies of hypoxic vascular remodeling have reported hypoxic increases in total arterial wall thickness involving both the medial and adventitial layers (17, 37, 47). In the present study, chronic hypoxia increased medial thicknesses (Figure 2), reinforcing the view that hypoxia promotes expansion of arterial smooth muscle (11, 47), an important determinant of overall arterial stiffness. Correspondingly, hypoxia also increased arterial stiffness, suggesting altered smooth muscle composition and structure (31) and possibly increased collagen cross-linking and collagen-to-elastin ratios (53). The associated hypoxic decreases in contractile stresses, measured in units of dynes/cm² (Figure 2), suggest that hypoxia increased the ratio of non-contractile to contractile

proteins in the artery wall in these fetal ovine carotid arteries (7, 17). Such changes were probably mediated by a unique but unknown combination of numerous possible mechanisms (31).

Consistent with previous studies (17), hypoxic structural changes in the fetal arteries were associated with decreased myogenic reactivity (Figure 2). Hypoxic depression of myogenic tone must involve changes in either regulation of cytosolic calcium or myofilament calcium sensitivity (34, 41). Whereas effects of chronic hypoxia on calcium signaling in non-pulmonary smooth muscle remain largely unreported, evidence from our group suggests that chronic hypoxia significantly alters myofilament calcium sensitivity in ovine cranial arteries (35). Correspondingly, the present study explored the hypothesis that hypoxia may alter myofilament calcium sensitivity and contractility through changes in contractile protein abundance and organization.

Effects of Hypoxia on Contractile Protein Abundances

The most abundant contractile protein in smooth muscle is SM α A, a \approx 42 kDa monomer that polymerizes to form the thin filaments essential for contraction (38). Expression of SM α A did not differ significantly from normoxic basal levels after hypoxic acclimatization for 110 days (Figure 3, Left) suggesting that the signaling pathways that govern expression of this contractile protein were not altered by hypoxia.

Another contractile protein whose expression varies with smooth muscle phenotype is MLCK (4, 7, 22), the rate-limiting enzyme that phosphorylates and activates regulatory MLC₂₀ (19). Hypoxia potently decreased MLCK (Figure 2, Middle), due possibly to phenotypic transformation or an effect at the level of transcription, translation, or

turnover. For transcription, hypoxia might act through HIF-1 α to activate an HRE that decreases MLCK gene transcription. No evidence for a repressor element in the MLCK gene has yet been published, although Qi (44) has reported that HIF can upregulate MLCK expression in cultured pulmonary vein endothelial cells suggesting the possible presence of an HRE in the MLCK promoter. Alternatively, hypoxia might enhance MLCK degradation; MLCK abundance dropped precipitously in arteries organ cultured under serum-starved conditions (7). This drop in MLCK would require rapid degradation, which might be stimulated by hypoxia (28). Hypoxia might also drive phenotypic transformation toward a less contractile, more synthetic phenotype that expresses less MLCK (19). Hypoxia's ability to promote such transformation has been demonstrated in cultured rat pulmonary artery smooth muscle (23, 57), but not previously in whole arteries adapted to chronic hypoxia.

In sharp contrast to the pattern of change observed for MLCK, MLC₂₀ abundances significantly increased in hypoxic compared to normoxic arteries (Figure 3, Right). Although hypoxic increases in MLC₂₀ abundance could result from increased transcription due to an HRE in the promoter for MLC₂₀, literature searches yielded no evidence of an HRE in the promoters for MLC₂₀ or myosin heavy chain with which MLC₂₀ should be co-expressed (14). Hypoxic increases in MLC₂₀ abundance also might result from increases in translation efficiency, but available evidence suggests only inhibition of translation by hypoxia (33). Hypoxic increases in MLC₂₀ might be explained by phenotypic transformation toward a synthetic phenotype, which is characteristic of hypoxia (21). Such a transformation would increase MLC₂₀ as a component of increased Non-Muscle MHC expression, which is induced by chronic

hypoxia in fetal carotid arteries (21). In light of evidence that hypoxia can increase protein degradation (45), this mechanism might also contribute to hypoxic increases in MLC₂₀ abundance. Equally important, the diverse effects of hypoxia on SM α A, MLCK, and MLC₂₀ abundances emphasize that the effects of hypoxia on contractile protein expression are unique and highly specific for each protein.

Effects of Hypoxia on Contractile Protein Organization

Vascular contractility is determined not only by contractile protein abundances, but also by protein organization within smooth muscle cells (6). Chronic hypoxia increased colocalization of MLCK with SM α A (**Figure 4, Left**) suggesting that hypoxia depressed release of an inhibitory factor, or enhanced release of a stimulatory factor, from either the endothelium or adventitia to enhance MLCK-SM α A colocalization. Although the molecular mechanisms responsible remain unknown, this result is consistent with evidence that MLCK binds strongly with SM α A (29) through mechanisms that are physiologically regulated (6).

As for colocalization of MLCK with SM α A, hypoxia also enhanced colocalization of MLC₂₀ with SM α A (**Figure 4, Middle**) and colocalization of MLCK with MLC₂₀ (**Figure 4, Right**), demonstrating that hypoxia orchestrated processes that enhanced interaction among multiple contractile proteins. Importantly, this pattern of increased colocalization occurred concurrently with markedly decreased MLCK abundance (**Figure 3**), which raises the possibility that increased colocalization may help preserve hypoxic contractility (**Figure 2**). Together, these results emphasize that hypoxia exerts

coordinated but separate influences on contractile protein abundance, organization, and function through mechanisms that remain largely unidentified.

Effects of Chronic Hypoxia on VEGF and VEGF Receptors

To better understand the mechanisms that mediate the effects of hypoxia on contractile protein abundance, organization, and function we studied established mediators of hypoxic effects. The most prominent of these is hypoxia inducible factor (HIF) (46, 56), which is well known to increase capillary angiogenesis through induction of VEGF (18). In light of recent evidence from our laboratory that VEGF can act directly on arterial smooth muscle to alter phenotype, contractile protein expression and contractility (7, 21), we formulated the hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through changes in the abundance, organization and function of contractile proteins (**Figure 1**). Our first test of this hypothesis (**Figure 1, Arrow 1**) was to examine the effects of 110 days of hypoxia on VEGF levels in endothelium-denuded fetal carotid homogenates. VEGF levels were unchanged by chronic hypoxia (**Figure 5**), in agreement with previous reports that VEGF rises then returns to baseline within 21 days of chronic hypoxia (10). In addition, this finding also suggested that elevated VEGF could not explain the sustained changes in contractile protein abundance and colocalization observed in hypoxic arteries.

In light of evidence that hypoxia can modulate levels of VEGF receptors in the vascular endothelium and certain tumors of rodent brain (43, 50), we also tested the possibility that chronic hypoxia increased VEGF receptor levels in fetal arterial smooth muscle (**Figure 1, Arrow 3**). These measurements yielded the novel finding that both main VEGF receptors (Flk-1 and Flt-1) were expressed in fetal ovine carotid arteries, as suggested by previous measurements of VEGF receptor mRNA (7). More importantly,

these measurements revealed that chronic hypoxia significantly enhanced VEGF receptor abundances (**Figure 6**). Together with our VEGF measurements (**Figure 5**), these findings support the novel view that hypoxia promotes VEGF-mediated effects in the short term through increases in VEGF abundance (**10**), but in the long term through increases in the abundances of VEGF receptors.

Effects of VEGF on Contractile Protein Abundance and Organization

To confirm that our previously reported effects of VEGF on contractile proteins in fetal lamb carotids (7, 21) were mediated by action on smooth muscle VEGF receptors (**Figure 1, Arrow 2**), we examined the effects of VEGF on colocalization of MLC₂₀ with SM α A in the presence and absence of vatalanib and dasatinib, two well-characterized VEGF receptor antagonists (27, 42). Organ culture of endothelium-denuded fetal carotid arteries with a low physiological concentration (3 ng/ml) of VEGF (7, 52) significantly increased colocalization of MLC₂₀ with SM α A (**Figure 7**). At concentrations found to be optimal in previous dose-finding experiments (21), vatalanib at 240 nM with dasatinib at 6.3 nM in coculture with VEGF completely blocked the effects of VEGF on colocalization of MLC₂₀ with SM α A. These results thus verified that VEGF can act through VEGF receptors to alter contractile protein organization in fetal carotid arteries.

As a further test of the hypothesis that VEGF contributes to hypoxic vascular remodeling, our next series of experiments examined the effects of VEGF on contractile protein abundances (**Figure 1, Arrow 4**). Organ culture with a physiological concentration (3 ng/ml) of VEGF (**52**) had no effect on SM α A abundance, significantly depressed MLCK abundance and significantly increased MLC₂₀ abundance, in hypoxic

relative to normoxic arteries (**Figure 8**). Most importantly, this pattern of effects was closely similar to the effects of chronic hypoxia on the same contractile proteins in fresh arteries. When we examined the effects of organ culture with VEGF on contractile protein colocalization, VEGF increased colocalization of MLCK with SM α A, of MLC₂₀ with SM α A, and of MLCK with MLC₂₀. Again this pattern of effects was quite similar to the effects of chronic hypoxia on contractile protein colocalization. Together, these abundance and colocalization measurements support the hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through changes in the abundance, organization and function of contractile proteins.

To confirm that our previously reported effects of VEGF on contractile proteins in fetal lamb carotids (7, 21) were mediated by action on smooth muscle VEGF receptors (**Figure 1, Arrow 2**), we examined the effects of VEGF on colocalization of MLC₂₀ with SM α A in the presence and absence of vatalanib and dasatinib, two well-characterized VEGF receptor antagonists (27, 42). Organ culture of endothelium-denuded fetal carotid arteries with a low physiological concentration (3 ng/ml) of VEGF (7, 52) significantly increased colocalization of MLC₂₀ with SM α A (**Figure 7**). At concentrations found to be optimal in previous dose-finding experiments (21), vatalanib at 240 nM with dasatinib at 6.3 nM in coculture with VEGF completely blocked the effects of VEGF on colocalization of MLC₂₀ with SM α A. These results thus verified that VEGF can act through VEGF receptors to alter contractile protein organization in fetal carotid arteries.

As a further test of the hypothesis that VEGF contributes to hypoxic vascular remodeling, our next series of experiments examined the effects of VEGF on contractile protein abundances (**Figure 1, Arrow 4**). Organ culture with a physiological

concentration (3 ng/ml) of VEGF (**52**) had no effect on SM α A abundance, significantly depressed MLCK abundance and significantly increased MLC₂₀ abundance, in hypoxic relative to normoxic arteries (**Figure 8**). Most importantly, this pattern of effects was closely similar to the effects of chronic hypoxia on the same contractile proteins in fresh arteries. When we examined the effects of organ culture with VEGF on contractile protein colocalization, VEGF increased colocalization of MLCK with SM α A, of MLC₂₀ with SM α A, and of MLCK with MLC₂₀. Again this pattern of effects was quite similar to the effects of chronic hypoxia on contractile protein colocalization. Together, these abundance and colocalization measurements support the hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through changes in the abundance, organization and function of contractile proteins.

Overview

The present study explores the mechanisms that couple chronic hypoxia to changes in the structure and function of large systemic fetal arteries (**Figure 1**). Consistent with numerous previous studies (21, 32), our findings demonstrate that chronic hypoxia alters fetal artery structure and contractility (**Figure 2**). A diverse and abundant literature also supports the idea that hypoxia potently increases VEGF in the short-term (10, 46). The present study expands this concept by demonstrating that in the long term, chronic hypoxia has little effect on VEGF (**Figure 5**) but potently increases the VEGF receptor abundances (**Figure 6**). Our results also augment previous findings that VEGF can modulate the expression, organization and function of smooth muscle contractile proteins (7, 21), by demonstrating that these effects are dependent on activation of smooth muscle

VEGF receptors (**Figure 7**). Finally, the present study demonstrates that the qualitative effects of chronic hypoxia on the abundance (**Figure 3**) and colocalization (**Figure 4**) of SM α A, MLCK, and MLC₂₀ are closely similar to the corresponding effects of VEGF (**Figures 8 and 9**). Together with the other evidence obtained, the findings as a whole are highly consistent with the general hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through changes in the abundance, organization and function of contractile proteins. Without doubt, VEGF is not the only factor contributing to hypoxic vascular remodeling (5), and future in vivo studies that involve selective interruption of VEGF signaling in large systemic arteries will be required to quantitatively assess the role of VEGF in hypoxic vascular remodeling. The present findings suggest that such future studies are warranted, and advance the idea that hypoxic increases in VEGF act simultaneously in the microcirculation to increase capillary density, while also acting in the large upstream arteries to alter the abundance, organization and function of contractile proteins.

Acknowledgments

The work reported in this manuscript was supported by USPHS Grants HL54120, HD31266, HL64867, NS076945 and the Loma Linda University School of Medicine.

References

1. Agnati LF, Fuxe K, Torvinen M, Genedani S, Franco R, Watson S, Nussdorfer GG, Leo G, and Guidolin D. New methods to evaluate colocalization of fluorophores in immunocytochemical preparations as exemplified by a study on A2A and D2 receptors in Chinese hamster ovary cells. *J Histochem Cytochem* 53: 941-953, 2005. PMID: 16055748
2. Bagwell CB, Hudson JL, and Irvin GL, 3rd. Nonparametric flow cytometry analysis. *J Histochem Cytochem* 27: 293-296, 1979. PMID: 374589
3. Bakker H and Jaddoe VW. Cardiovascular and metabolic influences of fetal smoke exposure. *Eur J Epidemiol* 26: 763-770, 2011. PMID: 21994150
4. Belik J, Kerc E, and Pato MD. Rat pulmonary arterial smooth muscle myosin light chain kinase and phosphatase activities decrease with age. *Am J Physiol Lung Cell Mol Physiol* 290: L509-516, 2006. PMID: 16214816
5. Bir SC, Kolluru GK, Fang K, and Kevil CG. Redox balance dynamically regulates vascular growth and remodeling. *Semin Cell Dev Biol* 23: 745-757, 2012. PMID: 22634069
6. Blue EK, Goeckeler ZM, Jin Y, Hou L, Dixon SA, Herring BP, Wysolmerski RB, and Gallagher PJ. 220- and 130-kDa MLCKs have distinct tissue distributions and intracellular localization patterns. *Am J Physiol Cell Physiol* 282: C451-460, 2002. PMID: 11832329
7. Butler SM, Abrassart JM, Hubbell MC, Adeoye O, Semotiuk A, Williams JM, Mata-Greenwood E, Khorram O, and Pearce WJ. Contributions of VEGF to age-dependent transmural gradients in contractile protein expression in ovine carotid arteries. *Am J Physiol Cell Physiol* 301: C653-666, 2011. PMID: 21653901
8. Chakrabarty K and Fahim M. Modulation of the contractile responses of guinea pig isolated tracheal rings after chronic intermittent hypobaric hypoxia with and without cold exposure. *J Appl Physiol* 99: 1006-1011, 2005. PMID: 16103517
9. Charles SM, Zhang L, Cipolla MJ, Buchholz JN, and Pearce WJ. Roles of cytosolic Ca²⁺ concentration and myofilament Ca²⁺ sensitization in age-dependent cerebrovascular myogenic tone. *Am J Physiol Heart Circ Physiol* 299: H1034-1044, 2010. PMID: 20639216
10. Chavez JC, Agani F, Pichiule P, and LaManna JC. Expression of hypoxia-inducible factor-1alpha in the brain of rats during chronic hypoxia. *J Appl Physiol* 89: 1937-1942, 2000. PMID: 11053346

11. Christou H, Yoshida A, Arthur V, Morita T, and Kourembanas S. Increased vascular endothelial growth factor production in the lungs of rats with hypoxia-induced pulmonary hypertension. *Am J Respir Cell Mol Biol* 18: 768-776, 1998. PMID: 9618381
12. Clyman RI, Seidner SR, Kajino H, Roman C, Koch CJ, Ferrara N, Waleh N, Mauray F, Chen YQ, Perkett EA, and Quinn T. VEGF regulates remodeling during permanent anatomic closure of the ductus arteriosus. *Am J Physiol Regul Integr Comp Physiol* 282: R199-206, 2002. PMID: 11742839
13. Cox RH. Three-dimensional mechanics of arterial segments in vitro: methods. *J Appl Physiol* 36: 381-384, 1974. PMID: 4814310
14. Eddinger TJ and Meer DP. Myosin II isoforms in smooth muscle: heterogeneity and function. *Am J Physiol Cell Physiol* 293: C493-508, 2007. PMID: 17475667
15. Frid MG, Moiseeva EP, and Stenmark KR. Multiple phenotypically distinct smooth muscle cell populations exist in the adult and developing bovine pulmonary arterial media in vivo. *Circ Res* 75: 669-681, 1994. PMID: 7923613
16. Giussani DA, Camm EJ, Niu Y, Richter HG, Blanco CE, Gottschalk R, Blake EZ, Horder KA, Thakor AS, Hansell JA, Kane AD, Wooding FB, Cross CM, and Herrera EA. Developmental programming of cardiovascular dysfunction by prenatal hypoxia and oxidative stress. *PLoS One* 7: e31017, 2012. PMID: 22348036
17. Griffith SL, Rhoades RA, and Packer CS. Pulmonary arterial smooth muscle contractility in hypoxia-induced pulmonary hypertension. *J Appl Physiol* 77: 406-414, 1994. PMID: 7961262
18. Hoeben A, Landuyt B, Highley MS, Wildiers H, Van Oosterom AT, and De Bruijn EA. Vascular endothelial growth factor and angiogenesis. *Pharmacol Rev* 56: 549-580, 2004. PMID: 15602010
19. Hong F, Haldeman BD, Jackson D, Carter M, Baker JE, and Cremo CR. Biochemistry of smooth muscle myosin light chain kinase. *Arch Biochem Biophys*, 2011. PMID: 21565153
20. Horowitz A, Menice CB, Laporte R, and Morgan KG. Mechanisms of smooth muscle contraction. *Physiol Rev* 76: 967-1003, 1996. PMID: 8874491
21. Hubbell MC, Semotiuk AJ, Thorpe RB, Adeoye OO, Butler SM, Williams JM, Khorram O, and Pearce WJ. Chronic hypoxia and VEGF differentially modulate abundance and organization of myosin heavy chain isoforms in fetal and adult ovine arteries. *Am J Physiol Cell Physiol* 303: C1090-1103, 2012. PMID: 22992677

22. Injeti ER, Sandoval RJ, Williams JM, Smolensky AV, Ford LE, and Pearce WJ. Maximal stimulation-induced in situ myosin light chain kinase activity is upregulated in fetal compared with adult ovine carotid arteries. *Am J Physiol Heart Circ Physiol* 295: H2289-2298, 2008. PMID: 18835918
23. Jie W, Guo J, Shen Z, Wang X, Zheng S, Wang G, and Ao Q. Contribution of myocardin in the hypoxia-induced phenotypic switching of rat pulmonary arterial smooth muscle cells. *Exp Mol Pathol* 89: 301-306, 2010. PMID: 20621093
24. Jin KL, Mao XO, and Greenberg DA. Vascular endothelial growth factor: direct neuroprotective effect in in vitro ischemia. *Proc Natl Acad Sci U S A* 97: 10242-10247, 2000. PMID: 10963684
25. Julian CG. High altitude during pregnancy. *Clin Chest Med* 32: 21-31, vii, 2011. PMID: 21277446
26. Kamitomo M, Longo LD, and Gilbert RD. Right and left ventricular function in fetal sheep exposed to long-term high-altitude hypoxemia. *Am J Physiol* 262: H399-405, 1992. PMID: 1539699
27. Karaman MW, Herrgard S, Treiber DK, Gallant P, Atteridge CE, Campbell BT, Chan KW, Ciceri P, Davis MI, Edeen PT, Faraoni R, Floyd M, Hunt JP, Lockhart DJ, Milanov ZV, Morrison MJ, Pallares G, Patel HK, Pritchard S, Wodicka LM, and Zarrinkar PP. A quantitative analysis of kinase inhibitor selectivity. *Nat Biotechnol* 26: 127-132, 2008. PMID: 18183025
28. Kim HA, Lim S, Moon HH, Kim SW, Hwang KC, Lee M, Kim SH, and Choi D. Hypoxia-inducible vascular endothelial growth factor gene therapy using the oxygen-dependent degradation domain in myocardial ischemia. *Pharm Res* 27: 2075-2084, 2010. PMID: 20607367
29. Kohama K, Ye LH, Hayakawa K, and Okagaki T. Myosin light chain kinase: an actin-binding protein that regulates an ATP-dependent interaction with myosin. *Trends Pharmacol Sci* 17: 284-287, 1996 Aug. PMID: 8810874
30. Kurinczuk JJ, White-Koning M, and Badawi N. Epidemiology of neonatal encephalopathy and hypoxic-ischaemic encephalopathy. *Early Hum Dev* 86: 329-338, 2010. PMID: 20554402
31. Lacolley P, Challande P, Osborne-Pellegrin M, and Regnault V. Genetics and pathophysiology of arterial stiffness. *Cardiovascular research* 81: 637-648, 2009. PMID: 19098299
32. Longo LD, Hull AD, Long DM, and Pearce WJ. Cerebrovascular adaptations to high-altitude hypoxemia in fetal and adult sheep. *Am J Physiol* 264: R65-72, 1993. PMID: 8430888

33. Magagnin MG, van den Beucken T, Sergeant K, Lambin P, Koritzinsky M, Devreese B, and Wouters BG. The mTOR target 4E-BP1 contributes to differential protein expression during normoxia and hypoxia through changes in mRNA translation efficiency. *Proteomics* 8: 1019-1028, 2008. PMID: 18219697
34. Meininger GA and Davis MJ. Cellular mechanisms involved in the vascular myogenic response. *Am J Physiol* 263: H647-659, 1992. PMID: 1415587
35. Nauli SM, Williams JM, Gerthoffer WT, and Pearce WJ. Chronic hypoxia modulates relations among calcium, myosin light chain phosphorylation, and force differently in fetal and adult ovine basilar arteries. *J Appl Physiol* 99: 120-127, 2005. PMID: 16036903
36. Neerhof MG and Thaete LG. The fetal response to chronic placental insufficiency. *Semin Perinatol* 32: 201-205, 2008. PMID: 18482622
37. Ooi CY, Wang Z, Tabima DM, Eickhoff JC, and Chesler NC. The role of collagen in extralobar pulmonary artery stiffening in response to hypoxia-induced pulmonary hypertension. *Am J Physiol Heart Circ Physiol* 299: H1823-1831, 2010. PMID: 20852040
38. Owens GK, Loeb A, Gordon D, and Thompson MM. Expression of smooth muscle-specific alpha-isoactin in cultured vascular smooth muscle cells: relationship between growth and cytodifferentiation. *J Cell Biol* 102: 343-352, 1986. PMID: 3944187
39. Packer CS, Roepke JE, Oberlies NH, and Rhoades RA. Myosin isoform shifts and decreased reactivity in hypoxia-induced hypertensive pulmonary arterial muscle. *Am J Physiol* 274: L775-785, 1998. PMID: 9612293
40. Pearce WJ, Hull AD, Long DM, and Longo LD. Developmental changes in ovine cerebral artery composition and reactivity. *Am J Physiol* 261(2 Pt 2): R458-R465, 1991. PMID: 1877701
41. Pfitzer G. Invited review: regulation of myosin phosphorylation in smooth muscle. *J Appl Physiol* 91: 497-503, 2001 Jul. PMID: 11408468
42. Pichot CS, Hartig SM, Xia L, Arvanitis C, Monisvais D, Lee FY, Frost JA, and Corey SJ. Dasatinib synergizes with doxorubicin to block growth, migration, and invasion of breast cancer cells. *Br J Cancer* 101: 38-47, 2009. PMID: 19513066
43. Plate KH, Breier G, Millauer B, Ullrich A, and Risau W. Up-regulation of vascular endothelial growth factor and its cognate receptors in a rat glioma model of tumor angiogenesis. *Cancer Res* 53: 5822-5827, 1993. PMID: 7694795

44. Qi H, Wang P, Liu C, Li M, Wang S, Huang Y, and Wang F. Involvement of HIF-1alpha in MLCK-dependent endothelial barrier dysfunction in hypoxia. *Cell Physiol Biochem* 27: 251-262, 2011. PMID: 21471714
45. Ramchandran R, Pilipenko E, Bach L, Raghavan A, Reddy SP, and Raj JU. Hypoxic regulation of pulmonary vascular smooth muscle cyclic guanosine monophosphate-dependent kinase by the ubiquitin conjugating system. *Am J Respir Cell Mol Biol* 46: 323-330, 2012. PMID: 21997485
46. Semenza GL. Expression of hypoxia-inducible factor 1: mechanisms and consequences. *Biochem Pharmacol* 59: 47-53, 2000. PMID: 10605934
47. Stenmark KR, Fagan KA, and Frid MG. Hypoxia-induced pulmonary vascular remodeling: cellular and molecular mechanisms. *Circ Res* 99: 675-691, 2006. PMID: 17008597
48. Taggart M and Wray S. Hypoxia and smooth muscle function: key regulatory events during metabolic stress. *J Physiol (Lond)* 509 (Pt 2): 315-325, 1998. PMID: 9575282
49. Teramo KA. Obstetric problems in diabetic pregnancy - The role of fetal hypoxia. *Best Pract Res Clin Endocrinol Metab* 24: 663-671, 2010. PMID: 20832744
50. Tipoe GL and Fung ML. Expression of HIF-1alpha, VEGF and VEGF receptors in the carotid body of chronically hypoxic rat. *Respir Physiol Neurobiol* 138: 143-154, 2003. PMID: 14609506
51. Villamor E, Kessels CG, Ruijtenbeek K, van Suylen RJ, Belik J, de Mey JG, and Blanco CE. Chronic in ovo hypoxia decreases pulmonary arterial contractile reactivity and induces biventricular cardiac enlargement in the chicken embryo. *Am J Physiol Regul Integr Comp Physiol* 287: R642-651, 2004. PMID: 15117730
52. Vonnahme KA, Wilson ME, Li Y, Rupnow HL, Phernetton TM, Ford SP, and Magness RR. Circulating levels of nitric oxide and vascular endothelial growth factor throughout ovine pregnancy. *J Physiol* 565: 101-109, 2005. PMID: 15774525
53. Wang Z and Chesler NC. Role of collagen content and cross-linking in large pulmonary arterial stiffening after chronic hypoxia. *Biomech Model Mechanobiol* 11: 279-289, 2012. PMID: 21538012
54. Williams JM, White CR, Chang MM, Injeti ER, Zhang L, and Pearce WJ. Chronic hypoxic decreases in soluble guanylate cyclase protein and enzyme activity are age dependent in fetal and adult ovine carotid arteries. *J Appl Physiol* 100: 1857-1866, 2006. PMID: 16469937

55. Yablonka-Reuveni Z, Christ B, and Benson JM. Transitions in cell organization and in expression of contractile and extracellular matrix proteins during development of chicken aortic smooth muscle: evidence for a complex spatial and temporal differentiation program. *Anat Embryol (Berl)* 197: 421-437, 1998. PMID: 9682974
56. Yamakawa M, Liu LX, Date T, Belanger AJ, Vincent KA, Akita GY, Kuriyama T, Cheng SH, Gregory RJ, and Jiang C. Hypoxia-inducible factor-1 mediates activation of cultured vascular endothelial cells by inducing multiple angiogenic factors. *Circ Res* 93: 664-673, 2003. PMID: 12958144
57. Yin H, Li Q, Qian G, Wang Y, Li Y, Wu G, and Wang G. Rab1 GTPase regulates phenotypic modulation of pulmonary artery smooth muscle cells by mediating the transport of angiotensin II type 1 receptor under hypoxia. *Int J Biochem Cell Biol* 43: 401-408, 2011. PMID: 2109523

CHAPTER THREE
VEGF RECEPTORS MEDIATE HYPOXIC REMODELING OF
ADULT OVINE CAROTID ARTERIES

Olayemi Adeoye, Vincent Bouthors, James M. Williams, and William J. Pearce

Divisions of Physiology, Pharmacology, and Biochemistry
Center for Perinatal Biology
Loma Linda University School of Medicine
Loma Linda, CA 92350

Running Title: VEGF receptors mediate hypoxic remodeling

Address for Correspondence:
William J. Pearce, Ph.D.
Center for Perinatal Biology
Loma Linda University School of Medicine
Loma Linda, CA 92350
Phone: 909-558-4325
FAX: 909-558-4029
E-Mail: wpearce@som.llu.edu

Abstract

In response to hypoxia, Hypoxia inducible factor (HIF-1) in living cells induces transcription of angiogenic genes; one of which is VEGF. Though principally known for its regulation of angiogenesis and vasculogenesis, recent studies have suggested VEGF involvement with hypoxic vascular remodeling. Many regard VEGF as a specific mitogen for endothelial cells, which exert paracrine effects on other cell types but recent studies provide evidence that VEGF exerts direct trophic effects on cells in the intima layer of blood vessels. This current study explores the hypothesis that chronic hypoxia produces a sustained increase in the density of vascular VEGF receptors, which in turn helps mediate the long-term effects of hypoxia on adult ovine carotid arteries. Carotid arteries from animals maintained at altitude for 110 days or at sea level were harvested and denuded of endothelium. Fresh normoxic and hypoxic as well as organ cultured segments in 3 ng/ml VEGF were used to test the effects of hypoxia and VEGF respectively. Whereas VEGF levels were similar to basal levels after 110 days of hypoxia, VEGF receptor levels were increased by 107% (flt-1) and 156% (flk-1). Also VEGF replicated effects of hypoxia on SM α A and MLC₂₀ abundances as well as MLC₂₀ - SM α A colocalization. VEGF did not replicate hypoxic effects on MLCK abundance, MLCK- SM α A colocalization and MLCK- MLC₂₀ colocalization, suggesting that postnatal development alters MLCK gene and its consequent reactivity to hypoxia and VEGF. VEGF effects were potently inhibited by VEGF receptor blockers – vatalinib (240 nM) and dasatinib (6.3 nM). Taken together, the current study supports the hypothesis that long-term upregulation of VEGF receptors mediate hypoxic vascular

remodeling through differential alteration of contractile protein abundance and colocalization.

Key Words

Myosin Light Chain Kinase, Organ Culture, Regulatory Myosin Light Chain, Smooth Muscle, VEGF

Introduction

Blood vessels regularly undergo structural changes in response to changing patterns of endogenous and exogenous stimuli (4, 42, 50, 51). These continuous changes in blood vessel cellular and extracellular structure are collectively known as remodeling and enable functional adaptation and specialization (25, 46). Vascular remodeling serves both normal physiologic adaptation, as with uterine artery remodeling during pregnancy (4), and also with pathological remodeling, as occurs during atherosclerosis (48) and tumorigenesis (8, 10, 35). Whereas vascular remodeling is clearly a prominent and dynamic process in all vascular beds, the factors involved with its initiation and maintenance remain largely unknown.

One of the most widely studied initiators of vascular remodeling is hypoxia, which can promote both physiological (13) and pathological (19) changes in vessel structure and function. Reduced oxygen availability can act through multiple pathways to alter the phenotypic characteristics and contractile function of vascular smooth muscle, and thereby optimize tissue perfusion and oxygenation (17, 24, 34, 57). Whereas a plethora of evidence supports endothelial (33) and adventitial (14) expansion as major components of vascular remodeling, more recent findings suggest that hypoxic remodeling also involves the medial layer of the artery wall (24, 26). In addition, vascular remodeling appears to involve not only myocytes in the medial layer, but also smooth muscle progenitor cells that originate in the bone marrow, migrate to and reside in the adventitial layer of the vessel wall, and then differentiate into myofibroblasts in the medial layer (32). Another key feature of medial smooth muscle cells is that they retain remarkable plasticity and can dedifferentiate from contractile cells into more synthetic, proliferative,

and migratory phenotypes in response to many types of environmental stimuli (54). Not surprisingly, smooth muscle phenotypic plasticity is particularly pronounced in term fetuses and neonates and facilitates the rapid changes in vascular structure and function that are characteristic of postnatal development (43). How hypoxia influences smooth muscle plasticity and differentiation remains poorly understood, particularly in immature blood vessels.

Efforts to better understand the mechanisms of hypoxic vascular remodeling have strongly implicated the transcription factor HIF as a primary initiator of multiple gene transcription events that coordinate responses to hypoxia and culminate in increased production of erythropoietin, glycolytic enzymes and angiogenic factors (5, 39, 47). The most important angiogenic factor induced by HIF is Vascular Endothelial Growth Factor (VEGF), which directly stimulates angiogenesis and neovascularization through activation of broadly distributed specific tyrosine kinase receptors (22). Interestingly, hypoxia-induced increases in VEGF typically attain peak values within 48 hours and then return back to basal levels within three weeks of hypoxic exposure (28) (28). Despite the transient nature of the transient responses of VEGF to hypoxia, the remodeling effects of hypoxia can persist for many weeks and months (21, 29, 36). This pattern of vascular responses to hypoxia raises multiple questions about the role of VEGF in long-term hypoxic vascular remodeling. One possible explanation is that hypoxia promotes short-term increases in VEGF, but long-term increases in VEGF receptors, as suggested to occur in fetal ovine carotid arteries from ewes acclimatized at altitude for 110 days (1). Consistent with this observation, other studies have also shown that hypoxia can increase the density of VEGF receptors (flt-1 and flk-1) (3, 20, 53).

In light of the published evidence, the current study explores the hypothesis that chronic hypoxia produces a sustained increase in the density of vascular VEGF receptors, which in turn helps mediate the long-term effects of hypoxia on vascular remodeling. This hypothesis builds on our previous work demonstrating that VEGF can contribute to hypoxic vascular remodeling of fetal arteries through alteration of contractile protein abundances and colocalization (1, 9, 24). To test this hypothesis, the experimental approach employed a well-established model of chronic hypoxia in which adult sheep were rendered hypoxic by housing them at an altitude of 3820m for 110 days; normoxic controls were maintained at sea level (30). Assessments of hypoxic changes in composition and interactions among contractile protein focused on: 1) SM α A, which is the most abundant contractile protein in smooth muscle cells (55); 2) regulatory Myosin Light Chain (MLC₂₀), which is a key regulator of contraction; and 3) Myosin Light Chain Kinase (MLCK), which is the dedicated kinase that phosphorylates and activates MLC₂₀ (16, 49). To assess the direct effects of VEGF on arterial wall remodeling, the experimental approach included organ-culture with VEGF, as previously described in detail (1, 9). Together, these experiments provided a unique perspective of the role of VEGF and VEGF receptors in long-term hypoxic vascular remodeling.

Materials and Methods

Protocols, techniques and procedures used in all animal experiments were approved by the Animal Research Committee of Loma Linda University and also complied with policies and codes of practice recommended by the National Institutes of Health in the NIH-Guide for Care and Use of Laboratory Animals.

Tissue Harvest and Preparation

Common carotid arteries harvested from adult non-pregnant sheep were used in all experiments. These sheep were sacrificed while fully anesthetized with 100 mg/kg pentobarbital administered intravenously. Control animals (sheep maintained at sea level) yielded normoxic arteries while sheep subjected to our chronic hypoxia model generated hypoxic arteries. This model involves acclimatization of non-gravid sheep at altitude of 3280m above sea level thereby yielding arterial oxygen tension of 64 ± 2 Torr in the animals. Correspondingly normoxic arterial oxygen tensions equal 102 ± 2 Torr. Harvested arteries were kept in sterile HEPES buffer solution composed of (in mM) 122.1 NaCl, 25 HEPES, 5.16 KCl, 2.4 MgSO₄, 11.1 dextrose, 1.6 CaCl₂ and 50 μ M EDTA. Loose parenchymal and adventitial tissue as well as blood was removed from the arteries after which endothelial denudation was achieved via luminal mechanical abrasion. All experimental protocols made use of segments 3mm length arterial segments. Precise arterial length and medial thicknesses were determined using an Olympus U-PMTVC Optical microscope mounted with a Scion Visicapture Twain 1394 Camera for image capture. ImagePro software (v6.0, Media Cybernetics) was used to measure the variables from the images captured by the camera.

Contractility Studies

For contractility measurements, artery segments measuring 3-mm in length were mounted on paired tungsten wires between a low-compliance isometric force transducer and a rod attached to a micrometer used for precise variation of resting tension. Each segment was equilibrated at 38.5 °C (normal ovine core temperature) for one hour in calcium replete Na bicarbonate Krebs solution (pH7.4) composed of (in mM) 122 NaCl, 25.6 NaHCO₃, 5.17 KCl, 2.56 dextrose, 2.49 MgSO₄, 1.60 CaCl₂, 0.114 ascorbic acid and 0.027 EGTA. Artery segments were kept viable by continuous bubbling of the incubation buffer with 95% oxygen with 5% carbon IV oxide. Working diameters denoted as (D) required to attain strain ratios of 1.5, 1.8, 2.1, 2.3, 2.7, 3.0 and 3.3 were calculated relative to unstressed artery diameter, D₀ (measured at a passive tension of 0.03 g. Contractile responses at each stretch ratio to isotonic potassium Krebs composed of (in mM) 122 KCl, 11.1 dextrose, 5.16 NaCl, 2.50 MgSO₄, 2.15 NaHCO₃, 1.60 CaCl₂, 0.114 ascorbic acid and 0.027 EDTA, were recorded after equilibration in Na Krebs. After each maximum contraction, arteries were washed and left to stabilize in Na Krebs before the next stretch ratio was achieved. Once contractions in response to the highest stretch ratio were recorded, the active component of the artery was eliminated by freezing in liquid nitrogen and left to equilibrate in 3 mM EGTA solution. Passive arterial stresses were subsequently measured at each strain ratio starting with the highest stretch ratio to the lowest. Spontaneous myogenic tone was defined as the difference in tone generated before and after freezing the artery segment while arterial stiffness was calculated from the passive stresses using the Young's modulus model as previously described.

Fluorescent Immunohistochemistry

Freshly harvested as well as organ-cultured arteries were fixed overnight in EM-grade 4% paraformaldehyde (Electron Microscopy Sciences, Hatfield, #15713S) then embedded in paraffin before sectioned at 5 μ m. Immunohistochemical processing of the tissues commenced with initial deparaffinization in HistoClear solution (National Diagnostic, Atlanta, #HS-200) and rehydration in graded concentrations of alcohol in descending order. The tissue samples were subsequently microwaved in a 6.03-pH citrate buffer in order to recover antigenicity. Permeabilization of the sections in 0.1% Triton X-100 (Sigma Aldrich, St Louis, #T-8787) was done prior to incubation in 1% Bovine Serum Albumin (Santa Cruz Biotechnology, Santa Cruz, #SC-2323) to ensure blockade of non-specific interactions.

Confocal Microscopy

Using the 5 μ m arterial sections generated as previously described under Fluorescent immunohistochemistry section, double staining of the sections was done using primary antibodies reactive with 3 different pairs of contractile proteins (SM α A - MLCK, SM α A -MLC₂₀ and MLCK-MLC₂₀). Following an overnight primary incubation, the sections were washed in phosphate buffer solution (PBS) and left to equilibrate with species matched secondary antibodies labeled with Dylight – 488 (SM α A) and 633 (MLCK) for SM α A -MLCK colocalization, 488 (SM α A) and 633 (MLC₂₀) for SM α A -MLC₂₀ colocalization, and 488(MLC₂₀) and 633 (MLCK) for MLCK-MLC₂₀ colocalization. Secondary incubation was done for two hours at room temperature in a dark room to minimize photo bleaching. Imaging was done with an Olympus FV1000 at a lateral resolution of 200 nm, optical section width of 700 nm and a numerical aperture

of 18. Colocalization indices were completed using a non-parametric quadrant analysis as previously described. Sources and titer ratios of the primary antibodies include: Sigma-Aldrich, St Louis for monoclonal anti-MLC₂₀ (#M4401) @ 1:300, polyclonal anti-SM α A (abcam, ab5694) @1:200, monoclonal anti- SM α A (A5228) at 1:200 and Santa Cruz Biotechnology for polyclonal MYLK (SC-25428) @ 1:50.

Western Blotting

Carotid artery segments of known weights were homogenized using a glass-on-glass method in a high urea (8M) extraction buffer containing (in mM) 500 NaCl, 23 Glycine, 20 Tris, 10 EGTA and 10% Glycerol at pH 8.6. Protease inhibitor cocktail (Sigma-Aldrich, Saint Louis, #M1745) at 5 μ l/ml of buffer was also added to the extraction buffer. After the tissues were grinded in the extraction buffer, the homogenate samples were centrifuged at 5,000G for 20 min. Protein concentrations were determined using the Bio-Rad Bradford assay. Post centrifugation, homogenates along with progressively increasing concentrations of standards used to calibrate target protein abundance was separated with an SDS-PAGE set-up. Separated proteins were electrophoretically transferred from the gel matrix onto nitrocellulose membranes at 200 mA current for 1.5 hr in Towbin's buffer containing 192 mM Glycine, 25 mM Tris, 10% and 20% Methanol. The transfer process was conducted with ice packs on both sides of the transfer tank to mitigate the heat generated via electrophoresis. Following transfer onto nitrocellulose, the membranes were blocked for 1 hr with Tris buffered saline containing 5% nonfat dry milk (M-TBS) while applying gentle shaking. Afterwards, membranes were washed in a detergent (0.1% Tween-20) containing MTBS and then

incubated with primary antibodies for 3 hr using 1:3000 for SM α A, 1:10,000 for MLCK, 1:200 for MLC₂₀ (all three from Sigma Aldrich as stated under “Confocal Microscopy section” and 1:750 for VEGF-A165 (from Abcam, Cambridge, #AB119). Following primary incubation, secondary antibody conjugated to Dylight 800 sourced from Pierce Chemical, Rockford, #46422 was applied to the membranes for 90 min before imaging was completed via a LI-COR Bioscience Odyssey system.

Tissue homogenization for quantifying VEGF receptors was initiated via a glass pestle and mortar using an extraction buffer containing (in mM) 500 NaCl, 50 Tris and 5 EDTA at pH 7.4. Six different protease inhibitors including (in μ M) 500 AEBSF, 400 Pepstatin-A, 20 Bestatin, 10 E-64, 7.5 Leupeptin and 7 Aprotinin all purchased from Sigma-Aldrich, Saint Louis, MO were also added to the buffer. A 1:50 tissue extraction buffer ratio was used. Next, centrifugation of the homogenate was done at 100,000 XG for 1 hr at 4°C after which the pellet was resuspended into the buffer at a 1:10 ratio with the addition of (in mM) 150 NaCl, 50 Tris, 10 DTT, 1% Triton X-100, 0.5% Na Deoxycholate, 0.2% SDS and 10% Glycerol with same concentrations of protease inhibitors listed above. Following this, the homogenate was ultra-sonicated at 20% amplitude to shear DNA for 6 times @ 5sec each, centrifuged again at 10,000 XG for 15 min. Collected supernatants were assayed for total protein concentrations via Bradford’s protein assay. Separation of the proteins was completed using a 5% SDS-PAGE with the addition of b-mercaptoethanol (BME- 35 mM) in the upper part of the tank with the buffer. As stated in the above paragraph, standard pooled reference samples were included in the separation gel lanes. Proteins from the electrophoresis gel were transferred onto a nitrocellulose membrane using Towbin’s buffer at 350mA for 1.5 hr.

Towbin's buffer used here is same as the one described in the previous paragraph but with the addition of 35 mM of BME, 0.01% SDS and 20% Methanol only. With the proteins now successfully transferred on membranes, blocking was done with MTBS for 60 min at room temperature while applying gentle shaking. Membranes were subsequently washed in 0.1% Tween containing MTBS. Incubation with primary antibodies was done for 3 hrs using dilutions for Flk-1 (Fetal Liver Kinase 1) (SC-504) and for Flt-1 (Fms-like tyrosine kinase receptors) (SC-316) both purchased from Santa Cruz using titers of 1:200. Following primary incubation, membranes were washed for 5 mins X 6 before secondary incubation commenced for 90 min. The same secondary antibody as the one described in the previous paragraph was used. Next, membranes were washed in TBS only for 5 mins X 6 before the membranes were imaged with a LI-COR Bioscience's Odyssey system.

Organ Culture

To determine the effects of VEGF and its receptor antagonists compared to those of hypoxia, arterial segments from both normoxic and hypoxic sheep were maintained in Media (Dulbecco's Modified Eagle Media) containing 70 µg/ml of Gentamycin (Gibco, Carlsbad, #15750-060), 4 mM glutamine (Sigma Aldrich, St. Louis, #G7513), 3.7 g/l of Na₂HCO₃, 2% antibiotic-antimycotic solution (Gibco, CARLSBAD, #15240-096), 1% non-essential amino acid solution (Sigma Aldrich, St Louis, #M7145), 0.5% amino acid solution (Sigma Aldrich, St. Louis, #M5550) in a 12-well untreated plate. Incubation was done in a humidified incubator with 5% CO₂ at 37°C. Control arteries were defined as those kept in Media for an additional 24 hrs after the initial 24hrs while those incubated in media containing 3 ng/ml of VEGF, 240 nM of Vatalanib and 6.3 nM of Dasatinib

served to determine the effect of VEGF and VEGF tyrosine kinase receptors. Treatment groups were defined as those that were kept in media for an initial 24 hr and then an additional 24 hr in media containing the various drugs.

Data Analysis and Statistics

Each animal sacrificed contributed to segments used in fresh, control, VEGF and VEGF receptor antagonist treatment groups. Normal distribution of data sets was tested using the D'Agostino-Pearson analysis whereas contractile stress measurements, wall thicknesses and stiffness were compared using analysis of variance (ANOVA). Within ANOVA, homogeneity was verified using a Bartlett's-Cochran test and statistical power was at least 0.8. In calculating contractile stresses, a robust computation involving force generated (g) by graded strains per unit area (length X working wall thickness - μm) was used.

For western blotting, unknown concentrations of protein of interest were measured against a standard curve generated from a pooled reference of ovine adult carotid tissues. Abundances from normoxic and hypoxic groups were compared using 2-way analysis of variance (ANOVA).

Results

A total of 10 normoxic and 11 hypoxic sheep were used in this study. The normoxic sheep contributed 108 arterial segments and the hypoxic ones contributed 91 segments. In all cases, “n” denotes number of sheep used in each protocol and $p < 0.05$ represents comparative results of statistical significance. Data here was presented as mean \pm SEM.

Chronic Hypoxia Alters Arterial Structure and Contractility in Adult Carotid Arteries

Medial thicknesses averaged $685 \pm 62 \mu\text{m}$ and $846 \pm 51 \mu\text{m}$ in normoxic and hypoxic arteries respectively (**Figure 1, left panel**); thus chronic hypoxia significantly increased medial thickness by 24%. Likewise, stiffness coefficients averaged 5.99 ± 0.40 and 7.36 ± 0.41 in fresh normoxic and hypoxic arteries respectively (**Figure 1, middle panel**); chronic hypoxia significantly increased arterial stiffness by 23%. Stretch ratio at peak myogenic tone averaged 2.42 ± 0.12 and 2.05 ± 0.11 for in normoxic and hypoxic arteries respectively (**Figure 1, right panel**); thus chronic hypoxia significantly decreased stretch ratio to attain peak myogenic tone by 15%. Similarly, peak myogenic tone averaged 71.4 ± 14.6 and 40.8 ± 14.3 in normoxic and hypoxic arteries respectively (**Figure 1, right panel**); chronic hypoxia thus decreased peak myogenic tone by 43%.




Figure 1: Arterial structure and contractility are altered by chronic hypoxia. Arteries from sheep subjected to hypoxic acclimatization exhibited increased medial thickness (left panel) and stiffness (middle panel) compared to the normoxic control group. In contrast, stress strain experiments showed a decreased maximum stretch ratio as well as peak myogenic tone in arteries from hypoxic sheep compared to normoxic control sheep. Results are presented as mean \pm SEM for $n=8$ for normoxic arteries and $n=11$ for hypoxic arteries. $P<0.05$ denotes comparative significant differences illustrated by an asterisk (*).

Chronic Hypoxia Alters Smooth Muscle Contractile Protein Expression

Chronic hypoxia exerted unique and protein specific changes in abundance of smooth muscle contractile proteins. For SM α A, quantification via Western immunoblotting showed a significant difference in abundance in normoxic (1.42 ± 0.08) compared to hypoxic (2.19 ± 0.2) arteries. Hypoxia resulted in a 54% increase in SM α A abundance. For MLCK, abundances in normoxic and hypoxic arteries averaged 1.34 ± 0.14 and 0.71 ± 0.31 respectively; hypoxia produced a 47% decrease in MLCK abundance. For MLC₂₀, abundances averaged 0.65 ± 0.09 and 1.10 ± 0.18 in normoxic and hypoxic arteries respectively; hypoxia resulted in a 69% increase in MLC₂₀ abundance (**Figure 2**).




Figure 2: Chronic Hypoxia differentially alters smooth muscle contractile protein expression. Quantification of smooth muscle contractile proteins via Western immunoblotting assays revealed a significantly higher level of SM α A and MLC₂₀ in hypoxic arteries compared to normoxic control arteries. In sharp contrast, MLCK abundance decreased with chronic hypoxia compared to normoxic values. These results suggest a highly protein specific alteration effect of hypoxia in these ovine carotid arteries. Results here are presented as mean \pm SEM for n=7 for all normoxic groups and n=6 for all hypoxic groups. Comparative significant differences denoted by asterisk (*) denotes p<0.05.

Effects of Chronic Hypoxia on Contractile Protein Colocalization

Coefficients of colocalization between MLCK and SM α A were similar in both normoxic (2.43 ± 0.78) and hypoxic (2.08 ± 0.84) (**Figure 3, left panel**) arteries. For MLC₂₀- SM α A, colocalization coefficients averaged 3.09 ± 1.63 and 24.83 ± 6.66 for normoxic and hypoxic arteries respectively (**Figure 3, middle panel**); hypoxia produced a 704% increase in colocalization between MLC₂₀ and SM α A. Similarly, for MLCK-MLC₂₀ colocalization, coefficients averaged 7.71 ± 4.36 and 33.56 ± 13.44 for normoxic and hypoxic arteries respectively (**Figure 3, right panel**); hypoxia produced a 335% increase in colocalization between MLCK and MLC₂₀.




Figure 3: Effects of chronic hypoxia on contractile protein colocalization. Interactions between smooth muscle contractile proteins assayed through confocal colocalization microscopy revealed a 704% increase in MLC₂₀-SM α A colocalization and a 335% increase in MLCK- MLC₂₀ colocalization in hypoxic compared to normoxic arteries. For MLCK-SM α A colocalization however, coefficients were similar in both normoxic and hypoxic arteries. Results are presented as mean \pm SEM for $n \geq 5$ for all experimental groups. Comparative significant differences denoted by an asterisk (*) represents $p < 0.05$.

Effects of Chronic Hypoxia on VEGF and VEGF Receptor Expression

Following 110 days of hypoxic acclimatization, expression of VEGF (1.44 ± 0.23) was not significantly different from normoxic control value (1.39 ± 0.17), suggesting that chronic hypoxia had no appreciable effect on VEGF abundance (**Figure 4**). In contrast, VEGF receptors (flk-1 andflt-1) expression potentially increased with chronic hypoxia. For flk-1, abundances increased from 5.87 ± 0.99 in normoxic arteries to 15.01 ± 1.87 in hypoxic arteries. Also for flt-1, abundances increased from 0.87 ± 0.17 in normoxic arteries to 1.80 ± 0.38 in hypoxic arteries (**Figure 5**).




Figure 4: Chronic hypoxia does not significantly increase VEGF expression. Western blot quantification of endogenous VEGF levels in arteries harvested from sheep acclimatized to chronic hypoxia for 110 days yielded values similar to those obtained in arteries harvested from normoxic control sheep. This result suggests that initial increases in VEGF levels (well documented) were not sustained chronically. Results are presented as mean \pm SEM for $n=7$ and 6 for normoxic and hypoxic arteries respectively.




Figure 5: Chronic hypoxia upregulates VEGF receptors. Whereas Western blot quantification of VEGF levels were not altered by chronic hypoxia, abundances of VEGF tyrosine kinase receptors were significantly increased by chronic hypoxia. VEGF receptor 1 – Flt-1 level was increased by 107% and VEGF receptor 2- Flk-1 was increased by 156% in hypoxic arteries. Results are presented as mean \pm SEM for n=6 for all experimental groups. Comparative significant differences denoted by an asterisk (*) imply $p < 0.05$ via ANOVA.

VEGF-induced Changes in Contractile Protein Interactions are Maintained Through VEGF Receptor Upregulation

To test the upregulation of VEGF tyrosine kinase receptors as a possible mechanism for VEGF alteration of contractile protein interactions, MLC₂₀- SM α A colocalization was used as a reporter assay. Whereas VEGF (3 ng/ml) in culture produced a 377% increase in MLC₂₀- SM α A colocalization, cocultures with VEGF receptor antagonists' vatalinib (240 nM) and dasatinib (6.3 nM) produced a complete blockade of this effect (**Figure 6**).




Figure 6: Chronic hypoxia alters contractile protein colocalization through upregulation of VEGF receptors. VEGF in culture produced significant increases (377%) in colocalization between MLC₂₀ and SM α A. Co-cultures with VEGF receptor antagonists, Vatalinib (240 nM) and Dasatinib (6.3 nM), completely blocked the potent effect of VEGF on these contractile proteins. These results suggest that VEGF induced changes in interactions between the contractile proteins are mediated through the receptors. Results are presented as mean \pm SEM for n = 5 in all experimental groups. Comparative significant differences denoted by an asterisk (*) imply p<0.05 via ANOVA.

Effect of VEGF in Culture on Contractile Protein Abundances

Low dose (3 ng/ml) VEGF in culture had no significant effect on $SM\alpha A$ (1.38 ± 0.09) MLCK (0.09 ± 0.04) or MLC_{20} abundances (0.42 ± 0.08) in normoxic arteries. However in hypoxic arteries, VEGF produced 51%, 83% and 161% increases in $SM\alpha A$, MLCK and MLC_{20} abundances; thus replicating effects of hypoxia on $SM\alpha A$ and MLC_{20} but not MLCK in culture.




Figure 7: VEGF replicates hypoxic effects on SM α A and MLC₂₀ expression. Organ culture of endothelium-denuded common carotid arteries with low dose VEGF (3 ng/ml) resulted in significant increases in SM α A (left panel) and MLC₂₀ (right panel) abundance. For MLCK, levels in VEGF treated arteries were similar to levels in normoxic control arteries (middle panel). VEGF thus replicates effects of chronic hypoxia in a highly protein specific manner, suggesting that VEGF might be mediating hypoxic changes in contractile protein expression in ovine carotid smooth muscle cells. Data are presented as mean \pm SEM for n=7 and 6 for normoxic and hypoxic arteries respectively. Comparative significant differences denoted by an asterisk (*) imply p<0.05 via ANOVA.

Effect of VEGF on Contractile Protein Interactions

Again, VEGF produced distinct effects on contractile protein colocalization. For MLCK- SM α A colocalization, coefficients increased from 3.85 ± 0.93 in control arteries to 7.99 ± 0.86 in VEGF treated arteries; VEGF produced a 108% increase in colocalization between the two contractile proteins. Similarly, for MLC₂₀- SM α A colocalization, coefficients increased from 1.00 ± 0.45 in control arteries to 4.77 ± 3.01 in VEGF treated arteries; VEGF produced a 377% increase in colocalization between the two contractile proteins. For MLCK-MLC₂₀ colocalization, coefficients were similar for control and VEGF treated arteries; VEGF had no effect on colocalization between the two contractile proteins. VEGF produced similar effects as hypoxia only for MLC₂₀- SM α A colocalization.




Figure 8: VEGF replicates hypoxic effects on MLCK-SM α A and MLC₂₀-SM α A colocalization. Organ culture of normoxic arteries with 3 ng/ml of VEGF increased colocalization of MLCK with SM α A and MLC₂₀ with SM α A. For MLCK-MLC₂₀ colocalization, VEGF had no significant effect on their colocalization coefficient. These results show that VEGF replicates hypoxic effects on MLC₂₀-SM α A colocalization, suggesting that VEGF mediates hypoxic alteration of contractile protein interaction in a highly protein specific manner. Data here are presented as mean \pm SEM for $n \geq 5$ for all experimental groups. Comparative significant differences denoted by an asterisk (*) imply $p < 0.05$ via ANOVA.

Discussion

This study offers four main original findings. First, long-term hypoxia (110 days at altitude of 3820 m) increased medial wall thickness and overall stiffness but decreased spontaneous myogenic tone. Second, long-term hypoxia had no effect on tissue VEGF levels but increased abundances of both the R1 and R2 subtypes of VEGF receptors. Third, both long-term hypoxia and organ culture with VEGF increased abundances of SM α A and MLC₂₀, but had opposite effects on MLCK abundance. Fourth, VEGF in organ culture replicated effects of long-term hypoxia on contractile protein colocalization of MLC₂₀ with SM α A, but not for MLCK with SM α A nor for MLCK with MLC₂₀, suggesting that factors other than VEGF contribute to hypoxic vascular remodeling. Overall, the current study demonstrates that hypoxic vascular remodeling involves a long-term up-regulation of VEGF receptors that can explain some, but not all, of the effects of chronic hypoxia on contractile protein abundance, organization, and function.

Long-term Hypoxia Alters Arterial Structure and Contractility

Chronic exposure to hypoxia typically promotes increased wall thickness in most arteries (30, 46) due largely to expansion of the adventitial layer (14). Some studies have also found, however, that resident cells in the medial layer can undergo hypertrophy (41, 45) and hyperplasia (7) upon chronic exposure to hypoxia. This medial hyperplasia, in turn, can result from hypoxia induced de-differentiation of fully contractile smooth muscle cells into synthetic cells with potent abilities to proliferate, migrate, and secrete extracellular matrix proteins (6). Recent findings further suggest that adventitial fibroblasts and smooth muscle progenitor cells can invade the medial layer and differentiate into myofibroblasts (32), adding further to hypoxic increases in medial

thickness. The current findings (**Figure 1**) reinforce previous reports that chronic hypoxia can increase medial thickness, and further demonstrate that this response includes an increase in passive arterial stiffness. This increase in stiffness, in turn, suggests either an increase in the collagen-to-elastin ratio, or an increase in collagen cross-linking (17, 58). Accompanying these increases in thickness and stiffness, the data also revealed that chronic hypoxia depressed myogenic reactivity (**Figure 1**). Because stiffness was increased, these results together suggest that hypoxia decreased either the efficiency of mechanotransduction, the strength of its coupling to the myogenic response, or overall contractile capacity (2, 15, 31). In light of previous results suggesting that hypoxia can promote the transformation of smooth muscle phenotype from a contractile to a more synthetic phenotype (6) and thereby decrease contractility (24), our next series of experiments examined the effects of hypoxia on the abundance and organization of three key contractile proteins.

Chronic Hypoxia Differentially Modulates Contractile Protein Abundances

Smooth muscle α -actin (SM α A) is the most abundant contractile protein in vascular smooth muscle and is the main component of the thin filament cytoskeleton essential for force production via interactions with activated myosin cross-bridges (27, 61). Immunoblot quantification demonstrated a significant increase in SM α A abundance in response to chronic hypoxia (**Figure 2**). This effect could be explained by the presence of a hypoxia responsive element (HRE) in the SM α A promoter, but evidence supporting this hypothesis has yet to be reported. Another possibility is that the translation efficiency for SM α A could be increased by hypoxia. Although a variety of interesting evidence

suggests that miRNAs can significantly modulate actin abundance and assembly (63), evidence that chronic hypoxia can stimulate the actions of any of these miRNAs is not available in the literature. Given that SM α A is one of the first contractile proteins expressed in newly differentiated smooth muscle (44), hypoxic increases in SM α A might also reflect increased smooth muscle cell proliferation (11, 58). Aside from the mechanism through which chronic hypoxia increased SM α A abundance, it is clear that the direction of this change cannot explain the parallel hypoxic inhibition of myogenic contractility (**Figure 1**).

Regulatory myosin light chain, also known as 20 kDa myosin light chain (MLC₂₀), is another critically important contractile protein in smooth muscle; phosphorylation of its serine 19 is prerequisite for contraction (60). As observed for SM α A, chronic exposure to hypoxia significantly increased the abundance of MLC₂₀, again suggesting possible hypoxic enhancement of the transcription or translation of MLC₂₀ mRNA. Evidence supporting any influence of hypoxia on either the transcription or translation of mRNA for MLC₂₀, however, has not been reported. Alternatively, hypoxic increases in smooth muscle proliferation (11, 58) could also explain increased abundance of MLC₂₀. As for hypoxic increases in SM α A abundance, these increase in MLC₂₀ abundance appear to be generally inconsistent with the observed hypoxic depression of myogenic contractility (**Figure 1**), unless these increases are occurring more in non-contractile synthetic smooth muscle than in fully differentiated contractile smooth muscle (52).

Myosin Light Chain Kinase (MLCK) is the dedicated kinase that phosphorylates serine 19 of MLC₂₀, and thus initiates smooth muscle contraction (23). In contrast to the patterns observed for SM α A and MLC₂₀, acclimatization to chronic hypoxia decreased

MLCK abundance by 47% (**Figure 2**). In turn, this result suggests a possible hypoxic inhibition of either transcription or translation of mRNA for MLCK, but as for SM α A and MLC₂₀, the published literature provides no support for such effects of hypoxia in smooth muscle. Another possibility is that hypoxia promotes the degradation and turnover of MLCK, but here too evidence of rapid turnover of MLCK has been offered by in vitro organ culture studies (9), but not by studies of vascular effects of long-term hypoxia. As for the effects of chronic hypoxia on SM α A and MLC₂₀, hypoxia-induced shifts in smooth muscle phenotype may also help explain the marked decrease in MLCK abundance observed in hypoxic arteries, particularly if MLCK is one of the last contractile proteins to be expressed during contractile differentiation (56). Although the mechanisms that mediate the protein-specific effects of chronic hypoxia on abundances of SM α A, MLC₂₀, and MLCK remain unclear, the present results strongly suggest that the effects of chronic hypoxia on myogenic contractility can be explained, at least in part, by decreased abundance of MLCK.

Chronic Hypoxia Differentially Modulates Contractile Protein Organization

Smooth muscle contraction involves rapid and efficient interactions among multiple contractile proteins whose distribution and organization, as well as their abundances, powerfully influence contractile force (60). Correspondingly, under some conditions phosphorylated MLC₂₀ can be uncoupled from force generation, even though all essential contractile proteins are present at appropriate abundances (37, 38). Such observations support the hypothesis that the organization of contractile proteins within smooth muscle is an important determinant of contractility that is physiologically regulated, particularly

during differentiation and functional maturation (64). To explore the hypothesis that chronic hypoxia could modulate contractility through changes in contractile protein organization, the experimental approach focused on the colocalization of several pairs of contractile proteins.

Consistent with the effects of chronic hypoxia on the abundances of both $SM\alpha A$ and MLC_{20} , chronic hypoxia increased the fraction of MLC_{20} colocalized with $SM\alpha A$ (**Figure 3**). In contrast, chronic hypoxia had no effect on the fraction of MLCK colocalized with $SM\alpha A$ despite the finding that hypoxia dramatically reduced MLCK abundance (**Figure 2**). More importantly, the fraction of MLCK bound to MLC_{20} was increased by adaptation to chronic hypoxia. These latter findings suggest that the MLCK lost during adaptation to chronic hypoxia came primarily from the fraction not colocalized with either $SM\alpha A$ or MLC_{20} . This interpretation is consistent with the hypothesis that MLCK serves important non-kinase and possible structural functions within smooth muscle (40, 62) that can be sacrificed during adaptation to chronic hypoxia. Conversely, the data further suggest that contractility is diminished but preserved in hypoxic arteries through increased colocalization of the remaining MLCK with MLC_{20} . Taken together, these results emphasize that the organization of contractile proteins may be more important than their abundance for generating contractile force.

Effects of Long-term Hypoxia on VEGF and VEGF Receptor Levels

To explore the mechanisms mediating hypoxic changes in vascular structure and function, our experimental approach examined the involvement of VEGF, a well-documented mediator of hypoxic angiogenesis (22, 59) that can influence the abundances

and organization of contractile proteins in both normoxic (9) and hypoxic (24) arteries. Consistent with previous reports that chronic hypoxia transiently increases VEGF (1, 12), chronic hypoxia of 110 days duration did not significantly change tissue VEGF levels (**Figure 4**).

Conversely, chronic hypoxia did significantly elevate the abundances of both the Flk-1 and Flt-1 subtypes of VEGF receptors in arterial tissue (**Figure 5**). Together, these findings support the hypothesis that hypoxic remodeling of large adult arteries is mediated in the short term (<21 days) through increases in VEGF levels, and in the long term (>28 days) through increases in VEGF receptor levels, as previously suggested in fetal arteries (1).

VEGF Receptors Mediate Arterial Remodeling

To confirm that VEGF receptors can mediate VEGF-induced modulation of contractile protein organization, a separate series of organ culture experiments quantified the effects of exposure to VEGF, in the presence and absence of VEGF receptor antagonists, on contractile protein colocalization. Given that colocalization between MLC₂₀ and SM α A was increased by chronic hypoxia (**Figure 3**), we used this pair of proteins to monitor the effects of a low, physiological concentration of VEGF (3 ng/ml), as previously reported (1). Consistent with our findings in fetal carotid arteries (1), cocultures of VEGF with optimal doses of vatalinib (240 nm) alone or with dasatinib (6.3 nm) completely inhibited VEGF effects on MLC₂₀ – SM α A colocalization (**Figure 6**). These results support the hypothesis that contributions of VEGF to hypoxic vascular remodeling occur via VEGF receptors 1 and 2 in ovine adult carotid arteries. To further

elucidate mechanisms through which VEGF contributes to hypoxic vascular remodeling, we conducted a series of western immunoblotting and colocalization studies on VEGF treated arterial segments.

VEGF Differentially Modulates Contractile Protein Abundances and Colocalization

Quantification of the SM α A via immunoblot analysis showed that similar to hypoxia, VEGF in culture produced a significant increase in SM α A abundance (**Figure 7**). This result suggests that VEGF might be contributing to the process of hypoxic vascular remodeling through modulation of smooth muscle cellular proliferation that results in SM α A upregulation (11, 58). Comparison of this result to our earlier work which showed that VEGF in culture upregulated SM α A abundance in fetal arteries suggests that post natal maturation does not alter smooth muscle cell reactivity to VEGF in terms of SM α A abundance. This might be related to the fact that SM α A is the most abundance marker contractile protein in smooth muscle cells regardless of age group (27, 52).

Again for MLC₂₀, similar to hypoxic effect on fresh arteries, VEGF in organ culture produced a significant increase in MLC₂₀ abundance (**Figure 7**). This result once again suggests that VEGF contributes to hypoxic vascular remodeling through upregulation of MLC₂₀, which though is inconsistent with decreased myogenic contractility observed, but nonetheless is crucial for the induction and maintenance of contractility (60).

For MLCK, in contrast to the pattern observed with hypoxic effect on fresh arteries, VEGF in culture had no significant on MLCK abundance (**Figure 7**). This suggests that contribution of VEGF to hypoxic vascular remodeling as it relates to contractile protein

abundance is unique and highly protein specific (9). Also this result suggests that some fundamental factors, yet to be elucidated, promote postnatal maturation alteration of cellular reactivity to VEGF. Our previous finding in fetal sheep (1), which showed that VEGF replicated hypoxic effects on MLCK abundance, supports this hypothesis.

Having shown the differential effects of VEGF on contractile protein abundances, we tested the hypothesis that VEGF contributes to hypoxic vascular remodeling through alteration of contractile protein interactions. Whereas hypoxia had no effect on MLCK – SM α A colocalization, the first set of experiments showed that VEGF produced significant increases in colocalization between the two proteins (**Figure 8**).

Also for MLCK – MLC₂₀ colocalization, whereas hypoxia produced significant increases in colocalization between them, VEGF in culture had no effect on their colocalization (**Figure 8**). Taken together, VEGF neither replicated hypoxic effects on MLCK abundance nor colocalization of other contractile proteins with MLCK. These results support our earlier hypothesis that certain fundamental factors alter reactivity of smooth muscle cells to VEGF postnatally with consequent alteration of MLCK production. Perhaps these same factors are the ones responsible for the precipitous drop in MLCK abundance secondary to long-term hypoxic acclimatization (1, 9). A striking decrease in MLCK abundance which has been shown to trigger apoptosis (18) and observed here with hypoxic acclimatization might prefer leads to those factors which might be involved with altered MLCK signaling postnatally.

For MLC₂₀ – SM α A colocalization, VEGF in culture replicated hypoxic effects on fresh arteries (**Figure 8**). This might be related to increased abundances in both MLC₂₀ and SM α A abundances observed with both hypoxia and VEGF in culture. Together, this

result suggests that at least in part, VEGF contributes to hypoxic vascular remodeling through differential and protein specific alterations in contractile protein abundance and interactions.

Overview

This study focuses on elucidating mechanisms involved with hypoxic vascular remodeling through a direct effect on arterial smooth muscle cells. This process is a dynamic one necessary for living cells to adapt and contend with prevailing internal and external stimuli (25, 46). In this study, we explored a well-established mediator of angiogenesis (VEGF), which has recently been suggested as a pivotal factor in vascular remodeling. Hypoxic increases in HIF-1 α result in modulated angiogenic transcription of VEGF as early as 24 hr of hypoxic exposure but falls back to basal levels by day 7 – 21 (28). Here we tested the role of endogenous VEGF in vascular remodeling after 110 days of hypoxic acclimatization (**Figure 9, arrow 1**). VEGF binds to its tyrosine kinase receptors 1 (Flt-1) and 2 (Flk-1) present on smooth muscle cells (**Figure 9, arrow 2**), ultimately leading to processes of angiogenesis (22), arterial structural changes and altered contractility (**Figure 9, arrow 4**). Parallel studies were done to quantify abundances of VEGF receptors after long-term hypoxic acclimatization (**Figure 9, arrow 3**) owing to their important role in VEGF signaling.

Consistent with literature, hypoxia resulted in increased arterial medial thickness and stiffness but decreased spontaneous myogenic reactivity (**Figure 1**). Also, hypoxic acclimatization produced significant increases in SM α A and MLC₂₀ but markedly decreased MLCK abundances (**Figure 2**). MLCK decreases correlated the most with

decreased contractile reactivity observed. Despite this decrease in MLCK, colocalization studies showed that its colocalization with MLC_{20} was significantly increased after hypoxic acclimatization. Similarly, MLC_{20} – $SM\alpha A$ colocalization increased with hypoxia but no changes occurred in $MLCK$ – $SM\alpha A$ colocalization (**Figure 3**). To test the hypothesis that VEGF contributes to these hypoxic changes in structure and function we quantified endogenous VEGF and VEGF receptor levels. Whereas after 110 days of hypoxia, VEGF levels were not different from basal levels (**Figure 4**), both VEGF receptors significantly increased (**Figure 5**), suggesting that remodeling effects of hypoxia are maintained through long-term upregulation of VEGF receptors (**Figure 6**). Also VEGF was shown to replicate hypoxic increases in $SM\alpha A$ and MLC_{20} in culture but not $MLCK$ abundances (**Figure 7**). Similarly, $MLCK$ colocalizations with other contractile proteins in VEGF treated segments were dissimilar to those observed with hypoxia. In contrast, VEGF replicated hypoxic increases in MLC_{20} – $SM\alpha A$ colocalization (**Figure 8**). Together these results suggest that trends of VEGF involvement in hypoxic vascular remodeling change with postnatal maturation. Also $MLCK$ decreases observed with hypoxia and shown to orchestrate apoptotic processes and factors probably alter cellular synthesis and turnover of $MLCK$, hence affects its colocalization with other contractile proteins. Finally, these results largely support the hypothesis that VEGF contributes to hypoxic vascular remodeling through differential and protein specific changes in contractile protein abundances and interactions.




Figure 9. This schematic illustrates the steps we took in testing the hypothesis that VEGF contributes to hypoxic vascular remodeling in adult ovine carotid arteries through alteration of contractile protein abundance, organization and function. First, we tested the propensity of long-time time hypoxia to directly increase endogenous VEGF (**Arrow 1**) and also VEGF receptors 1 (Flt-1) and 2 (Flk-1) (**Arrow 3**). Also we tested the possibility of VEGF signaling through its receptors (**Arrow 2**) to effect changes in arterial structure (arterial wall thickness and stiffness, contractile protein abundances) and contractility (contractile protein interaction and myogenic tone) (**Arrow 4**). Using this experimental layout, we proposed to explore involvement of VEGF in the non-angiogenic role of remodeling.

Acknowledgment

The work reported in this manuscript was supported by USPHS Grants HL54120, HD31266, HL64867, NS076945 and the Loma Linda University School of Medicine.

References

1. **Adeoye OO, Butler SM, Hubbell MC, Semotiuk AJ, Williams JM, and Pearce WJ.** Contribution of Increased VEGF Receptors to Hypoxic Changes in Fetal Ovine Carotid Artery Contractile Proteins. *Am J Physiol Cell Physiol*, 2013: PMID. 23325408.
2. **Ainslie KM, Garanich JS, Dull RO, and Tarbell JM.** Vascular smooth muscle cell glycocalyx influences shear stress-mediated contractile response. *Journal of applied physiology* 98: 242-249, 2005: PMID. 15322072.
3. **Ara J, Shukla P, and Frank M.** Enhanced expression of the Flt-1 and Flk-1 receptor tyrosine kinases in a newborn piglet model of ischemic tolerance. *J Neurochem* 124: 735-746, 2013: PMID. 23199256.
4. **Ashton SV, Whitley GS, Dash PR, Wareing M, Crocker IP, Baker PN, and Cartwright JE.** Uterine spiral artery remodeling involves endothelial apoptosis induced by extravillous trophoblasts through Fas/FasL interactions. *Arterioscler Thromb Vasc Biol* 25: 102-108, 2005: PMID. 15499040.
5. **Bao B, Ahmad A, Kong D, Ali S, Azmi AS, Li Y, Banerjee S, Padhye S, and Sarkar FH.** Hypoxia induced aggressiveness of prostate cancer cells is linked with deregulated expression of VEGF, IL-6 and miRNAs that are attenuated by CDF. *PLoS One* 7: e43726, 2012: PMID. 22952749.
6. **Beamish JA, He P, Kottke-Marchant K, and Marchant RE.** Molecular regulation of contractile smooth muscle cell phenotype: implications for vascular tissue engineering. *Tissue Eng Part B Rev* 16: 467-491, 2010: PMID. 20334504.
7. **Bevan RD, van Marthens E, and Bevan JA.** Hyperplasia of vascular smooth muscle in experimental hypertension in the rabbit. *Circ Res* 38: 58-62, 1976: PMID. 1269112.
8. **Briet M and Burns KD.** Chronic kidney disease and vascular remodelling: molecular mechanisms and clinical implications. *Clin Sci (Lond)* 123: 399-416, 2012: PMID. 22671427.

9. **Butler SM, Abrassart JM, Hubbell MC, Adeoye O, Semotiuk A, Williams JM, Mata-Greenwood E, Khorram O, and Pearce WJ.** Contributions of VEGF to age-dependent transmural gradients in contractile protein expression in ovine carotid arteries. *Am J Physiol Cell Physiol* 301: C653-666, 2011: PMID. 21653901.
10. **Carmeliet P.** Angiogenesis in health and disease. *Nat Med* 9: 653-660, 2003: PMID. 12778163.
11. **Chanakira A, Dutta R, Charboneau R, Barke R, Santilli SM, and Roy S.** Hypoxia differentially regulates arterial and venous smooth muscle cell proliferation via PDGFR-beta and VEGFR-2 expression. *American journal of physiology Heart and circulatory physiology* 302: H1173-1184, 2012: PMID. 22159994.
12. **Chavez JC, Agani F, Pichiule P, and LaManna JC.** Expression of hypoxia-inducible factor-1alpha in the brain of rats during chronic hypoxia. *J Appl Physiol* 89: 1937-1942, 2000 Nov: PMID.
13. **Clyman RI, Seidner SR, Kajino H, Roman C, Koch CJ, Ferrara N, Waleh N, Mauray F, Chen YQ, Perkett EA, and Quinn T.** VEGF regulates remodeling during permanent anatomic closure of the ductus arteriosus. *Am J Physiol Regul Integr Comp Physiol* 282: R199-206, 2002: PMID. 11742839.
14. **Davie NJ, Crossno JTJ, Frid MG, Hofmeister SE, Reeves JT, Hyde DM, Carpenter TC, Brunetti JA, McNiece IK, and Stenmark KR.** Hypoxia-induced pulmonary artery adventitial remodeling and neovascularization: contribution of progenitor cells. *Am J Physiol Lung Cell Mol Physiol* 286: L668-678, 2004 PMID. 12754186.
15. **Davis MJ, Wu X, Nurkiewicz TR, Kawasaki J, Davis GE, Hill MA, and Meininger GA.** Integrins and mechanotransduction of the vascular myogenic response. *American journal of physiology Heart and circulatory physiology* 280: H1427-1433, 2001: PMID. 11247750.

16. **de Lanerolle P and Paul RJ.** Myosin phosphorylation/dephosphorylation and regulation of airway smooth muscle contractility. *The American journal of physiology* 261: L1-14, 1991: PMID. 1651660.
17. **Dodson RB, Rozance PJ, Fleenor BS, Petrash CC, Shoemaker LG, Hunter KS, and Ferguson VL.** Increased arterial stiffness and extracellular matrix reorganization in intrauterine growth-restricted fetal sheep. *Pediatr Res* 73: 147-154, 2013: PMID. 23154756.
18. **Fazal F, Gu L, Ihnatovych I, Han Y, Hu W, Antic N, Carreira F, Blomquist JF, Hope TJ, Ucker DS, and de Lanerolle P.** Inhibiting myosin light chain kinase induces apoptosis in vitro and in vivo. *Mol Cell Biol* 25: 6259-6266, 2005: PMID. 15988034.
19. **Galis ZS and Khatri JJ.** Matrix metalloproteinases in vascular remodeling and atherogenesis: the good, the bad, and the ugly. *Circ Res* 90: 251-262, 2002: PMID. 11861412.
20. **Gerber HP, Condorelli F, Park J, and Ferrara N.** Differential transcriptional regulation of the two vascular endothelial growth factor receptor genes. Flt-1, but not Flk-1/KDR, is up-regulated by hypoxia. *J Biol Chem* 272: 23659-23667, 1997: PMID. 9295307.
21. **Hampel V, Bibova J, Ostadalova I, Povysilova V, and Herget J.** Gender differences in the long-term effects of perinatal hypoxia on pulmonary circulation in rats. *Am J Physiol Lung Cell Mol Physiol* 285: L386-392, 2003: PMID. 12691957.
22. **Hoeben A, Landuyt B, Highley MS, Wildiers H, Van Oosterom AT, and De Bruijn EA.** Vascular endothelial growth factor and angiogenesis. *Pharmacol Rev* 56: 549-580, 2004: PMID. 15602010.
23. **Hong F, Haldeman BD, Jackson D, Carter M, Baker JE, and Cremo CR.** Biochemistry of smooth muscle myosin light chain kinase. *Arch Biochem Biophys*, 2011: PMID. 21565153.
24. **Hubbell MC, Semotiuk AJ, Thorpe RB, Adeoye OO, Butler SM, Williams JM, Khorram O, and Pearce WJ.** Chronic hypoxia and VEGF differentially modulate

- abundance and organization of myosin heavy chain isoforms in fetal and adult ovine arteries. *Am J Physiol Cell Physiol* 303: C1090-1103, 2012: PMID. 22992677.
25. **Intengan HD and Schiffrin EL.** Vascular remodeling in hypertension: roles of apoptosis, inflammation, and fibrosis. *Hypertension* 38: 581-587, 2001: PMID. 11566935.
 26. **Ishida A, Murray J, Saito Y, Kanthou C, Benzakour O, Shibuya M, and Wijelath ES.** Expression of vascular endothelial growth factor receptors in smooth muscle cells. *J Cell Physiol* 188: 359-368, 2001: PMID. 11473363.
 27. **Kim HR, Appel S, Vetterkind S, Gangopadhyay SS, and Morgan KG.** Smooth muscle signalling pathways in health and disease. *J Cell Mol Med* 12: 2165-2180, 2008: PMID. 19120701.
 28. **Kuo NT, Benhayon D, Przybylski RJ, Martin RJ, and LaManna JC.** Prolonged hypoxia increases vascular endothelial growth factor mRNA and protein in adult mouse brain. *Journal of applied physiology* 86: 260-264, 1999: PMID. 9887138.
 29. **Lewis AM, Mathieu-Costello O, McMillan PJ, and Gilbert RD.** Effects of long-term, high-altitude hypoxia on the capillarity of the ovine fetal heart. *Am J Physiol* 277: H756-762, 1999: PMID. 10444503.
 30. **Longo LD, Hull AD, Long DM, and Pearce WJ.** Cerebrovascular adaptations to high-altitude hypoxemia in fetal and adult sheep. *Am J Physiol* 264: R65-72, 1993: PMID. 8430888.
 31. **Martinez-Lemus LA, Wu X, Wilson E, Hill MA, Davis GE, Davis MJ, and Meininger GA.** Integrins as unique receptors for vascular control. *J Vasc Res* 40: 211-233, 2003: PMID. 12902635.
 32. **Marx SO, Totary-Jain H, and Marks AR.** Vascular smooth muscle cell proliferation in restenosis. *Circ Cardiovasc Interv* 4: 104-111, 2011: PMID. 21325199.
 33. **Mata-Greenwood E, Meyrick B, Soifer SJ, Fineman JR, and Black SM.** Expression of VEGF and its receptors Flt-1 and Flk-1/KDR is altered in lambs with

- increased pulmonary blood flow and pulmonary hypertension. *American journal of physiology Lung cellular and molecular physiology* 285: L222-231, 2003: PMID. 12665467.
34. **McQuillan LP, Leung GK, Marsden PA, Kostyk SK, and Kourembanas S.** Hypoxia inhibits expression of eNOS via transcriptional and posttranscriptional mechanisms [published errata appear in *Am J Physiol* 1995 Feb;268(2 Pt 2):section H following table of contents and 1995 Jun;268(6 Pt 3):section H following table of contents]. *Am J Physiol* 267: H1921-1927, 1994: PMID. 7526714.
 35. **Ment LR, Stewart WB, Fronc R, Seashore C, Mahooti S, Scaramuzzino D, and Madri JA.** Vascular endothelial growth factor mediates reactive angiogenesis in the postnatal developing brain. *Brain Res Dev Brain Res* 100: 52-61, 1997 May 20: PMID. 9174246.
 36. **Mlynarczyk M, Imamura T, Umezaki H, Kaushal KM, Zhang L, and Ducsay CA.** Long-term hypoxia changes myometrial responsiveness and oxytocin receptors in the pregnant ewe: differential effects on longitudinal versus circular smooth muscle. *Biol Reprod* 69: 1500-1505, 2003: PMID. 12855597.
 37. **Morgado M, Cairrao E, Santos-Silva AJ, and Verde I.** Cyclic nucleotide-dependent relaxation pathways in vascular smooth muscle. *Cell Mol Life Sci* 69: 247-266, 2012: PMID. 21947498.
 38. **Morgan KG and Gangopadhyay SS.** Invited Review: Cross-bridge regulation by thin filament-associated proteins. *J Appl Physiol* 91: 953-962, 2001 Aug: PMID.
 39. **Mu D, Jiang X, Sheldon RA, Fox CK, Hamrick SE, Vexler ZS, and Ferriero DM.** Regulation of hypoxia-inducible factor 1alpha and induction of vascular endothelial growth factor in a rat neonatal stroke model. *Neurobiol Dis* 14: 524-534, 2003: PMID. 14678768.
 40. **Nakamura A, Xie C, Zhang Y, Gao Y, Wang HH, Ye LH, Kishi H, Okagaki T, Yoshiyama S, Hayakawa K, Ishikawa R, and Kohama K.** Role of non-kinase activity of myosin light-chain kinase in regulating smooth muscle contraction, a

- review dedicated to Dr. Setsuro Ebashi. *Biochem Biophys Res Commun* 369: 135-143, 2008: PMID. 18053800.
41. **Olivetti G, Anversa P, Melissari M, and Loud AV.** Morphometry of medial hypertrophy in the rat thoracic aorta. *Lab Invest* 42: 559-565, 1980: PMID. 7382430.
 42. **Osol G and Mandala M.** Maternal uterine vascular remodeling during pregnancy. *Physiology (Bethesda)* 24: 58-71, 2009: PMID. 19196652.
 43. **Owens GK.** Regulation of differentiation of vascular smooth muscle cells. *Physiological reviews* 75: 487-517, 1995: PMID. 7624392.
 44. **Owens GK, Loeb A, Gordon D, and Thompson MM.** Expression of smooth muscle-specific alpha-isoactin in cultured vascular smooth muscle cells: relationship between growth and cytodifferentiation. *J Cell Biol* 102: 343-352, 1986: PMID. 3944187.
 45. **Owens GK and Schwartz SM.** Alterations in vascular smooth muscle mass in the spontaneously hypertensive rat. Role of cellular hypertrophy, hyperploidy, and hyperplasia. *Circ Res* 51: 280-289, 1982: PMID. 7116579.
 46. **Packer CS, Roepke JE, Oberlies NH, and Rhoades RA.** Myosin isoform shifts and decreased reactivity in hypoxia-induced hypertensive pulmonary arterial muscle. *Am J Physiol* 274: L775-785, 1998: PMID. 9612293.
 47. **Pages G and Pouyssegur J.** Transcriptional regulation of the Vascular Endothelial Growth Factor gene--a concert of activating factors. *Cardiovasc Res* 65: 564-573, 2005: PMID. 15664382.
 48. **Pasterkamp G.** Imaging of de novo atherosclerotic arterial remodeling: clinical sense or research sensibility? *J Am Coll Cardiol* 53: 1716-1717, 2009: PMID. 19406348.
 49. **Pfitzer G.** Invited review: regulation of myosin phosphorylation in smooth muscle. *Journal of applied physiology* 91: 497-503, 2001: PMID. 11408468.

50. **Pries AR, Reglin B, and Secomb TW.** Remodeling of blood vessels: responses of diameter and wall thickness to hemodynamic and metabolic stimuli. *Hypertension* 46: 725-731, 2005: PMID. 16172421.
51. **Reglin B, Secomb TW, and Pries AR.** Structural adaptation of microvessel diameters in response to metabolic stimuli: where are the oxygen sensors? *Am J Physiol Heart Circ Physiol* 297: H2206-2219, 2009: PMID. 19783778.
52. **Rensen SS, Doevendans PA, and van Eys GJ.** Regulation and characteristics of vascular smooth muscle cell phenotypic diversity. *Neth Heart J* 15: 100-108, 2007: PMID. 17612668.
53. **Sandner P, Wolf K, Bergmaier U, Gess B, and Kurtz A.** Hypoxia and cobalt stimulate vascular endothelial growth factor receptor gene expression in rats. *Pflugers Arch* 433: 803-808, 1997: PMID. 9049173.
54. **Schwartz SM, Campbell GR, and Campbell JH.** Replication of smooth muscle cells in vascular disease. *Circ Res* 58: 427-444, 1986: PMID. 3516443.
55. **Skalli O, Pelte MF, Pecllet MC, Gabbiani G, Gugliotta P, Bussolati G, Ravazzola M, and Orci L.** Alpha-smooth muscle actin, a differentiation marker of smooth muscle cells, is present in microfilamentous bundles of pericytes. *J Histochem Cytochem* 37: 315-321, 1989: PMID. 2918221.
56. **Steinbach SK, El-Mounayri O, DaCosta RS, Frontini MJ, Nong Z, Maeda A, Pickering JG, Miller FD, and Husain M.** Directed differentiation of skin-derived precursors into functional vascular smooth muscle cells. *Arteriosclerosis, thrombosis, and vascular biology* 31: 2938-2948, 2011: PMID. 21852558.
57. **Stenmark KR, Fagan KA, and Frid MG.** Hypoxia-induced pulmonary vascular remodeling: cellular and molecular mechanisms. *Circ Res* 99: 675-691, 2006: PMID. 17008597.
58. **Thompson JA, Richardson BS, Gagnon R, and Regnault TR.** Chronic intrauterine hypoxia interferes with aortic development in the late gestation ovine fetus. *J Physiol* 589: 3319-3332, 2011: PMID. 21540340.

59. **Tipoe GL, Lau TY, Nanji AA, and Fung ML.** Expression and functions of vasoactive substances regulated by hypoxia-inducible factor-1 in chronic hypoxemia. *Cardiovasc Hematol Agents Med Chem* 4: 199-218, 2006: PMID. 16842206.
60. **Walsh MP.** Vascular smooth muscle myosin light chain diphosphorylation: mechanism, function, and pathological implications. *IUBMB Life* 63: 987-1000, 2011: PMID. 21990256.
61. **Walsh MP and Cole WC.** The role of actin filament dynamics in the myogenic response of cerebral resistance arteries. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism* 33: 1-12, 2013: PMID. 23072746.
62. **Wang HH, Nakamura A, Matsumoto A, Yoshiyama S, Qin X, Ye LH, Xie C, Zhang Y, Gao Y, Ishikawa R, and Kohama K.** Nonkinase activity of MLCK in elongated filopodia formation and chemotaxis of vascular smooth muscle cells toward sphingosylphosphorylcholine. *Am J Physiol Heart Circ Physiol* 296: H1683-1693, 2009: PMID. 19234090.
63. **Xin M, Small EM, Sutherland LB, Qi X, McAnally J, Plato CF, Richardson JA, Bassel-Duby R, and Olson EN.** MicroRNAs miR-143 and miR-145 modulate cytoskeletal dynamics and responsiveness of smooth muscle cells to injury. *Genes Dev* 23: 2166-2178, 2009: PMID. 19720868.
64. **Yablonka-Reuveni Z, Christ B, and Benson JM.** Transitions in cell organization and in expression of contractile and extracellular matrix proteins during development of chicken aortic smooth muscle: evidence for a complex spatial and temporal differentiation program. *Anat Embryol (Berl)* 197: 421-437, 1998 Jun: PMID.

CHAPTER FOUR

SYMPATHETIC PERIVASCULAR NERVES CONTRIBUTE TO LONG-TERM
HYPOXIC TRANSFORMATION OF SMOOTH MUSCLE PHENOTYPE IN OVINE
CEREBRAL ARTERIES

**Olayemi O. Adeoye, Jinjutha Silpanisong, Vanessa Hlebowski, Vincent Bouthors,
James M. Williams, and William J. Pearce**

Divisions of Physiology, Pharmacology, and Biochemistry
Center for Perinatal Biology
Loma Linda University School of Medicine
Loma Linda, CA 92350

Running Title: Perivascular Nerves mediate Hypoxic Vascular Remodeling

Address for Correspondence:
William J. Pearce, Ph.D.
Center for Perinatal Biology
Loma Linda University School of Medicine
Loma Linda, CA 92350
Phone: 909-558-4325
FAX: 909-558-4029
E-Mail: wpearce@som.llu.edu

Abstract

In addition to its role in angiogenesis, VEGF may also contribute to arterial remodeling, possibly through activation of VEGF receptors on sympathetic nerves leading to altered trophic input to the arterial wall. To test this hypothesis normoxic and chronically hypoxic fetal sheep (110 days at 3820m) underwent unilateral superior cervical sympathectomy (SANX) at 124d gestation, and then were harvested at 138d (term). In middle cerebral arteries (MCA), hypoxia increased dopamine beta hydroxylase 368% and SANX ablated this increase. Hypoxia also decreased by 34% colocalization between Non-Muscle Myosin Heavy Chain (SMemb) and smooth muscle α -actin (AA) indicating increased contractile differentiation; SANX increased this colocalization 106%. Conversely, hypoxia increased by 125% SM2-MHC colocalization with AA, again indicating increased differentiation; SANX decreased this colocalization by 81%. Organ culture of MCA with NE increased colocalization of SM2-MHC with AA by 150% in SANX arteries; the effects of NE were blocked by prazosin. These results suggest that hypoxic increases in VEGF alter sympathetic release of NE and/or other factors that transform the contractile phenotype of cerebrovascular smooth muscle. This transformation appears to be a central feature of hypoxic cerebrovascular remodeling in the fetus.

Key Words

Hypoxia, Sympathetic, Perivascular nerve, Myosin Light Chain Kinase, Regulatory Myosin Light Chain, Smooth Muscle, VEGF

Introduction

Animal species are endowed with the intrinsic ability to maintain tissue perfusion and oxygenation, which is critical to their survival (31). During periods of reduced oxygen tension i.e. hypoxia, HIF-1 α via its hypoxia response element results in transcription of angiogenic genes that produce cytokines which aid angiogenesis (40). One of such cytokines is Vascular Endothelial Growth Factor (VEGF), which has been extensively shown to mediate angiogenesis and vasculogenesis via its tyrosine kinase receptors (13, 19).

As the name suggests, VEGF from endothelial cells has been shown to produce autocrine effects on endothelial cells (27, 30, 44) as well as paracrine remodeling effects on the medial and adventitial layers (39). Of recent, some studies have suggested a direct remodeling effect of VEGF on the vascular medial layer via VEGF receptors present on the resident smooth muscle cells (1, 25, 47, 51).

Both autocrine and paracrine effects of VEGF have been suggested to be signaled through VEGF receptors 1 and 2 present on the vessel wall (37). Downstream, VEGF mediates genomic changes producing alterations in contractile protein abundances and organization (1, 6). Some studies have also suggested a phenotypic switch in myosin isoforms as a possible mechanism for VEGF mediated hypoxic vascular remodeling in ovine carotid arteries (21). A recent line of studies has shown the presence of VEGF receptors on sympathetic nerves and also the involvement of VEGF in differentiation and growth of nerve fibers (29, 41). In like manner, sympathetic nerves have been shown to modulate differentiation, growth and functioning of the intimal layer of blood vessels

(10). This important finding suggests the possibility of paracrine effects of VEGF on the vessel wall via VEGF receptors expressed on resident nerve cells.

With these background findings in mind, this current study explores the hypothesis that hypoxic increases in VEGF contributes to arterial remodeling in ovine cerebral arteries via sympathetic perivascular nerve alteration of smooth muscle cell phenotype. Using a well established long-term hypoxia model in sheep, we compared the effects of animals acclimatized to altitude and their normoxic controls kept at sea level for patterns of arterial remodeling (23, 28). To test the hypothesis that sympathetic perivascular nerves mediate remodeling effects of hypoxia, we conducted unilateral superior cervical sympathectomy (2) on time-dated normoxic and hypoxic fetuses at 124 day gestation and harvesting both ipsilateral and contralateral middle cerebral arteries at term for various biochemical analyses. To determine sympathetic perivascular nerve alteration of smooth muscle cell phenotype, we quantified abundances of known markers for both synthetic and contractile smooth muscle cell phenotype i.e. non-muscle isoform of heavy chain myosin (SMemb), heavy chain myosin-1 (SM-1) and heavy chain myosin-2 (SM2).

Taken together, this study provides some mechanistic explanations for involvement of sympathetic perivascular nerve in hypoxic vascular remodeling and thus expands the current knowledge on reactivity of the vasculature to hypoxia, which is involved in a great percentage of clinical pathologies.

Materials and Methods

Techniques, protocols and procedures employed in these studies received approval from the Animal Research Committee of Loma Linda University and were in line with the policies and code of practices outlined in the National Institutes of Health Guide for the Care and Use of Laboratory Animals. Procedures related to tissue harvesting, preparation and animal surgery have been previously described in detail (1, 6).

Tissue Harvest and Preparation

Experiments conducted in this study made use of middle cerebral arteries harvested from term fetal (139 – 142 days gestation) sheep harvested and prepared using very strict sterile procedures. Contractility, electrical nerve stimulation and immunohistochemical experiments involved use of arteries from two main groups of animals in terms of percentage oxygen they were exposed to; first, normoxic arteries generated from animals maintained at sea level and second, hypoxic arteries generated from animals chronically acclimatized at altitude of 3820m above sea level for the final 110 days of gestation. The chronic hypoxia model used in this study exposes fetal sheep to arterial oxygen tensions (PaO_2) of 19 ± 1 within the hypoxic period (23). Experiments involving use of denervated arteries begins with performing a unilateral superior cervical ganglionectomy on fetuses at 125 – 128 days gestation through a head exteriorization via a midline incision while still attached to the pregnant ewe. After removal of the ganglion, the surgical wound is repaired and fetus is reduced back into the uterine cavity and left for another two weeks before the fetus is harvested.

The process of harvesting the fetus begins with anesthetization of pregnant ewes with 30mg/kg pentobarbital, intubated and maintained on 1.5 – 2.0% halothane gas. With the use of midline vertical incision, fetuses are subsequently exteriorized and rapidly exsanguinated by removal of the heart. The pregnant ewes from which the fetuses are harvested are subsequently sacrificed using 100mg/kg eutosol administered IV. Harvested arteries are placed in sterile HEPES buffer solution made up of (in mM) 122.1 NaCl, 25 HEPES, 5.16 KCl, 2.4 MgSO₄, 11.1 dextrose, 1.6 CaCl₂ and 50 μM EDTA. Arteries are carefully cleaned of loose connective tissues, blood and subsequently cut into 3mm lengths of coronal slices then distributed into various experimental groups. Precise arterial segment length measurements were done using an Olympus U-PMTVC Optical microscope mounted with a Scion Visicapture Twain 1394 Camera for image capture with ImagePro Software (v6.0, Media Cybernetics) for analysis.

Electrical Transmural Stimulation

5mm segments of cleaned MCA from both normoxic and hypoxic animals were mounted on tungsten wires between force transducers and micrometers that provide precise diameter increases, while immersed in Na⁺ Krebs solution (pH7.4) containing (in mM): 122 NaCl, 25.6 NaHCO₃, 5.17 KCl, 2.56 dextrose, 2.49 MgSO₄, 1.60 CaCl₂, 0.114 ascorbic acid and 0.027 EGTA. The solution was continuously bubbled with 95% O₂ and 5% CO₂ at 38°C. Successive increases in stretch were applied to the tissues until the optimum resting tension was obtained for each segment. Following this, the segments were left to equilibrate for 30 minutes before maximum K⁺ induced active tone was measured by adding K⁺ Krebs buffer which consists of (in mM) 122 KCl, 11.1 dextrose,

5.16 NaCl, 2.50 MgSO₄, 2.15 NaHCO₃, 1.60 CaCl₂, 0.114 ascorbic acid, and 0.027 EDTA. Arterial responses to electrical stimulation frequency were characterized by delivering supramaximal voltage and constant current to the tissues through two platinum field electrodes placed parallel and on either side of the arterial segment. These trains of square-wave pulses were of 0.3 ms durations. Each segment was stimulated at each of five frequencies: 8, 1, 2, 4, 8 and 16 Hz for 2.5 minutes, with a 5 minute rest period between successive stimulations.

Western Immunoblotting

Known weights of frozen arterial tissues were lysed via glass-on-glass in 8M Urea, (in mM) 10 EGTA, 20 Tris, 23 Glycine, 500 NaCl and 10% Glycerol at pH 8.6 with addition of a protease inhibitor cocktail at 5 µl/ml of buffer (Sigma-Aldrich, Saint Louis, #M1745). Homogenates were centrifuged at 5,000G for 20 minutes after which protein concentration of the supernatants were determined using the Bio-Rad Bradford assay. Total cell lysate was analyzed by SDS-PAGE together with increasing concentrations of standards used to calibrate target abundance. Tissues for the standards were harvested from pooled fetal ovine middle cerebral arteries. Separated proteins from SDS-PAGE were transferred onto nitrocellulose membranes at 200 mA for 90 minutes in Towbin's buffer (in mM) 25 Tris, 192 Glycine and 10% Methanol) on ice. Membranes were blocked with 5% milk in Tris buffered saline at pH 7.5 (M-TBS) for 1 hr at room temperature using continuous gentle shaking. After the initial blocking, subsequent washes and incubations were done in M-TBS containing 0.1% Tween-20. Incubations with primary antibody were performed for 3 hours using anti-Dopa Beta Hydroxylase

antibody (1:500) purchased from Abcam (Cambridge, #AB 43868). Visualization was done after 90 minutes of secondary incubation with a secondary antibody (1:5000) conjugated to Dylight 800 (Pierce Chemical, Rockford, #46422). Imaging was done on a LI-COR Bioscience Odyssey system.

Fluorescent Immunohistochemistry

Posterior communicating artery segments were embedded in Tissue-Tek OCT media (#14-373-65, Fisher Scientific) on dry ice then sectioned using a Leica CM3050S cryostat at 10 μ m thick. Sections were dried onto Superfrost Plus Gold slides (#22-035-813, Fisher Scientific) at 37°C for 2hr followed by fixation using -20°C acetone for 10minutes. After Phosphate buffered saline pH 7.4 (PBS) rinses, sections were incubated with 1% bovine serum albumin (BSA), 5% normal goat serum (NGS) and 0.2% Triton X-100 in PBS using CoverWell incubation chambers (#247001, Research Products International) for 1 hr. All incubations were performed with gentle vibrations achieved on a stirrer plate (Jacobsen-K.X and Staines-WA Journal of Neuroscience Methods 137 (2004) 71-77). Sections were incubated with a rabbit polyclonal antibody against Dopamine β -Hydroxylase (ab43868, Abcam) at 1:1000 in 1% BSA, 2% NGS and 0.1% Triton X-100 in PBS overnight at 4oC. Sections were washed in PBS + 0.1% Tween-20 then incubated 2hrs with a goat anti-rabbit IgG conjugated with Dylight 488 (#35553, Pierce Chemical) at room temperature. Following washing, sections were mounted with SlowFade Gold antifade mounting media (S36937, Invitrogen) and coverslipped. Images were obtained using a Zeiss Axio Imager A1 with Spot software (Ver. 4.6.4.5, Diagnostic Instruments).

Wall thickness and Stiffness

Using ImagePro software, immunohistochemically stain slides of tissue coronal segments were used in determining medial wall thicknesses of arteries from both normoxic and hypoxic animals. Stiffness coefficients were calculated from contractility experiments that made use of 3mm cut arterial segments mounted on tungsten wires between isometric force transducers and micrometers used for precise positioning. Initial tissue equilibration was conducted for one hour in calcium replete Na⁺ Krebs buffer (pH7.4). The equilibration solution was continuously bubbled with 95% O₂ with 5% CO₂ at normal ovine core temperature of 38°C. Following equilibration, unstressed artery diameters (D₀) were measured at a passive tension of 0.03 g. Working diameters (D) needed to attain stresses of (in g) 0.3, 0.6, 0.9, 1.2, 1.5 were also calculated. Stiffness coefficient were determined after elimination of the active component of tone by freezing and equilibrating the tissues in EGTA solution. Stiffness calculations were completed by normalizing stress values obtained to gravity, arterial length and thickness.

Confocal Microscopy

Segments of MCA were fixed overnight in 4% neutral buffered EM-grade formaldehyde (Electron Microscopy Sciences, Hatfield, #15713S), paraffin embedded and sectioned at 5µm. Following deparaffinization in histoclear solution (National Diagnostic, Atlanta, #HS-200), rehydration in successive dilute concentrations of alcohol, antigen retrieval in citrate buffer (pH 6.03) and permeabilization in 0.1% Triton X-100 (Sigma Aldrich, St Louis, #T-8787), the sections were incubated in 1% bovine serum albumin (Santa Cruz Biotechnology, Santa Cruz, #SC-2323) to block non-specific

binding. Next, the artery sections were double-stained with monoclonal anti- SM α A (Sigma-Aldrich, St Louis, #M4401) @ 1:100 with polyclonal anti-human SMEMB (Santa Cruz Biotechnology, Santa Cruz, SC-25428) @ 1:500 or monoclonal anti-SM1 @ 1:3000 or monoclonal anti-SM2 @1:400 overnight @ 4°C. The following day, the sections were washed in PBS and equilibrated in a dark room with secondary antibodies labeled with Dylight- 488 (SM α A) and 633 (SMemb, SM1 or SM2) for two hours at room temperature. Imaging of the sections was done using an Olympus FV1000 at an optical section thickness of 1 μ m, a lateral resolution of 2 nm and a numerical aperture of 18. Extent of colocalization between the two markers was determined using a non-parametric quadrant analysis, which calculates the fraction of total pixels in the upper half of the intensity range for both markers relative to the number of pixels in the upper half of the intensity range for either SMEMB, SM1 or SM2. Adeoye et al (1) described details of this method.

Data Analysis and Statistics

Contractile stresses generated from raw grams of strain were calculated as ratios of force per cross-sectional area where force equals tension (g) X acceleration due to gravity and cross-sectional area equals wall thickness (μ m) X artery segment length. Comparisons of contractile stresses were made using analysis of variance (ANOVA). All data sets were confirmed to be normally distributed using the D'Agostino-Pearson analysis and statistical power was at least 0.8.

One-way ANOVA was used to test the hypothesis in each experiment. Duncan's Multiple Range analysis was used to make post-hoc comparisons between means within the ANOVA.

Concentration-response curves were analyzed by computer-assisted excel sigmoid regression fits. Statistical significance was considered a P value <0.05 and n value represents the number of arteries / animals used in each experimental group.

Results

Effects of Long-term Hypoxia on Arterial Reactivity to Nerve Stimulation

Maximum responses to transmural electrical stimulation of the adrenergic nerve terminals occurred at 16 Hz in both normoxic (0.009 ± 0.007) and hypoxic (0.137 ± 0.14) cerebral artery segments (**Figure 1**). Hypoxic acclimatization produced a 1422% increase in frequency-dependent constriction compared to responses in normoxic arteries. This frequency-dependent constriction was sensitive to blockade by either 10^{-7} M tetrodotoxin or 10^{-6} M phentolamine.




Figure 1: Hypoxia upregulates sympathetic perivascular activity in ovine cerebral arteries. In fetal ovine middle cerebral arteries, contractile responses to electrical transmural stimulation of sympathetic perivascular adrenergic nerves were enhanced following 110 days of hypoxic acclimatization. Results are presented as mean \pm SEM for $n = 9$ for fetal normoxic arteries and $n = 6$ for fetal hypoxic arteries. Significant differences imply $P < 0.05$.

Effects of Long-term Hypoxia on Expression of Dopamine- β -Hydroxylase

In fetal ovine cerebral arteries, western blot analysis of the enzyme, dopamine- β -hydroxylase, required for the rate limiting reaction for the synthesis of norepinephrine, yielded (per μg standard protein) 0.25 ± 0.05 and 1.17 ± 0.23 in normoxic and hypoxic arteries respectively (**Figure 2**). Hypoxic acclimatization thus produced ~ 4.5 fold increase in DbH expression compared to normoxic arteries. Similarly, fluorescent immunohistochemistry analysis showed greater fluorescence for D β H in hypoxic arteries compared to normoxic arteries.




Figure 2: Hypoxia upregulates expression of dopa-β-hydroxylase in ovine cerebral arteries. The schematic shows the steps leading to the synthesis of Norepinephrine (Figure 2, upper panel). Dopamine-β-hydroxylase is the rate-limiting step in the synthesis of NE. In fetal ovine middle cerebral arteries, western blot analysis showed that hypoxic acclimatization resulted in about 290% increase in dopa-β-hydroxylase expression (Figure 2 - left panel). Immunohistochemical analysis also showed that hypoxic acclimatization upregulates expression of dopa-β-hydroxylase (Figure 2 – right panel). Sympathectomy downregulates dopa-β-hydroxylase expression in both normoxic and hypoxic arteries (Figure 2 – right panel).

Effects of Sympathetic Perivascular Nerves on Arterial Wall Thickness and Stiffness

In nerve intact arteries, arterial medial wall thicknesses averaged (in μm) 36.36 ± 3.3 and 48.81 ± 3.14 in normoxic and hypoxic arteries respectively. Hypoxic acclimatization thus produced a 34% increase in arterial wall thickness compared to normoxic arteries (**Figure 3**). Medial wall thicknesses of arteries from denervated animals averaged (in μm) 31.99 ± 2.17 and 41.56 ± 1.32 in normoxic and hypoxic groups respectively (**Figure 3**). Hypoxic acclimatization thus produced a 30% increase in medial wall thickness compared to values obtained in normoxic arteries. Whereas denervation had no significant effect on normoxic arteries, it produced a 15% decrease in medial wall thickness in hypoxic arteries.

Also in nerve intact arteries, stiffness coefficients averaged 30.70 ± 18.52 for normoxic arteries and 60.53 ± 20.05 for hypoxic arteries at maximum strain. Hypoxic acclimatization thus produced a 97% increase in stiffness (**Figure 3, Right panel**). In denervated arteries, stiffness coefficients averaged 65.11 ± 21.93 for normoxic arteries and 174.82 ± 61.68 for hypoxic arteries at maximum strain. Hypoxia acclimatization thus produced a 168% increase in stiffness in denervated arteries. Denervation resulted in a 112% increase in stiffness in normoxic arteries and a 189% increase in hypoxic arteries.




Figure 3: Sympathetic perivascular nerves mediate hypoxic effects on cerebral artery structure. Hypoxic acclimatization significantly increased arterial wall thickness by 34% and 30% in nerve intact and SANX arteries respectively. Whereas sympathectomy had no effect on normoxic arteries, it produced a 15% decrease in medial wall thickness in hypoxic arteries (Figure 3, left panel). Data is presented as mean \pm SEM for $n = 5$ for all experimental groups. Also hypoxic acclimatization produced a 97% and 168% increase in stiffness in nerve intact and SANX arteries respectively. Sympathectomy resulted in a 112% and 189% increase in stiffness in normoxic and hypoxic arteries respectively. Data here represents mean \pm SEM for $n = 5$ for both normoxic groups and hypoxic SANX group and $n = 4$ for hypoxic intact group. Statistical significance denoted by (*) represents $P < 0.05$.

Effect of Sympathetic Perivascular Nerves on Contractile Protein Organization

For SMemb - SM α A, colocalization coefficients averaged 0.008 ± 0.001 and 0.0056 ± 0.0013 in nerve intact normoxic and hypoxic arteries respectively. Hypoxic acclimatization thus resulted in a 30% decrease in SMemb - SM α A colocalization compared to normoxic arteries. In denervated arteries, colocalization coefficients averaged 0.012 ± 0.002 and 0.0116 ± 0.0019 in normoxic and hypoxic arteries respectively. Hypoxic acclimatization thus had no effect on SMemb - SM α A in denervated arteries. Denervation produced a 50% and 107% increase in SMemb - SM α A colocalization in normoxic and hypoxic arteries respectively (**Figure 4**).

For SM1- SM α A, colocalization coefficients averaged 0.023 ± 0.003 and 0.005 ± 0.002 in nerve intact normoxic and hypoxic arteries respectively. Hypoxic acclimatization thus resulted in a 78% decrease in SM1- SM α A colocalization compared to normoxic arteries. In denervated arteries, colocalization coefficients averaged 0.015 ± 0.002 and 0.013 ± 0.003 in normoxic and hypoxic arteries respectively. Hypoxic acclimatization thus had no effect on SM1- SM α A colocalization in denervated arteries. Whereas denervation produced a 35% decrease in SM1- SM α A, colocalization in normoxic arteries, it produced a 160% increase in SM1- SM α A, colocalization in hypoxic arteries (**Figure 5**).

For SM2 - SM α A, colocalization coefficients averaged 0.0069 ± 0.0016 and 0.0155 ± 0.00250 in nerve intact normoxic and hypoxic arteries. Hypoxic acclimatization thus resulted in a 125% increase in SM2 - SM α A colocalization in nerve intact arteries. In denervated arteries, colocalization coefficients averaged 0.0030 ± 0.00029 and $0.003 \pm$

0.00068 in normoxic and hypoxic arteries respectively. Hypoxic acclimatization hence had no effect on SM2 - SM α A colocalization in denervated arteries. Denervation significantly decreased SM2 - SM α A colocalization in both normoxic (57%) and hypoxic arteries (81%) (**Figure 6**).




Figure 4: Sympathetic perivascular nerves alter interactions between SMemb and SM α A. Hypoxic acclimatization resulted in a 30% decrease in SMemb - SM α A colocalization in nerve intact arteries but had no effect in SANX arteries. In contrast, sympathectomy resulted in 50% and 107% increase in colocalization between the two proteins in normoxic and hypoxic arteries respectively. Data here represent mean \pm SEM for n = 5 for experimental groups. Statistical significance denoted by (*) represents P < 0.05




Figure 5: Sympathetic perivascular nerves alter interactions between SM1 and SM α A. Hypoxic acclimatization resulted in a 78% decrease in SM1 - SM α A colocalization in nerve intact arteries but had no effect on colocalization between the two contractile proteins in SANX arteries. Similarly, sympathectomy resulted in 35% decrease in SM1 - SM α A colocalization in normoxic arteries but produced a 160% increase in SM1 - SM α A colocalization in hypoxic arteries. Data here represent mean \pm SEM for n = 5 for all experimental groups except FN SANX in which n = 6. Statistical significance denoted by (*) represents P < 0.05




Figure 6: Sympathetic perivascular nerves alter interactions between SM2 and SM α A. Hypoxic acclimatization resulted in a 125% increase in SM2 - SM α A colocalization in nerve intact arteries but had no effect on colocalization between the two contractile proteins in SANX arteries. In sharp contrast, sympathectomy resulted in 57% and 81% decrease in SM2 - SM α A colocalization in normoxic and hypoxic arteries respectively. Data here represent mean \pm SEM for n = 5 for all experimental groups. Statistical significance denoted by (*) represents p < 0.05.

Sympathetic Perivascular Nerves Mediate Hypoxic Remodeling Through NE and its α -1 Adrenoceptors

In hypoxic denervated arteries, SM2 - SM α A colocalization coefficients averaged 0.002 ± 0.001 , 0.002 ± 0.0004 , 0.005 ± 0.001 and 0.002 ± 0.001 in fresh, starved, NE treated and NE+prazosin tissues respectively. Starved segments were incubated in Dulbecco's modified medium (DMEM) with basic essential and non-essential amino acids for 48 hr; NE and NE+Prazosin treated segments were starved for 24 hrs and then treated with the respective drugs for 24hrs. 48 hr starvation had no statistically significant effect on SM2 - SM α A colocalization, whereas NE increased SM2 - SM α A colocalization by 150% and addition of prazosin reversed this increase by 60% (**Figure 7**).




Figure 7: NE acts through its α -1 adrenoceptors in mediating trophic effects of sympathetic perivascular nerves on middle cerebral arteries. In hypoxic denervated arteries, 48 hr starvation treatment i.e. incubation in media with basic essential and non-essential amino acids, produced no significant effect on SM2 - SM α A colocalization. NE treatment produced a 150% increase in SM2 - SM α A colocalization and addition of prazosin to NE treated arteries decreased SM2 - SM α A by 60%. Data here represent mean \pm SEM for n = 6 for all experimental groups except fresh arteries in which n = 4. Statistical significance denoted by (*) represents P < 0.05

Discussion

This study offers 5 novel findings in endothelium denuded ovine cerebral arteries. First, reactivity to electrical nerve stimulation was significantly increased in fetal hypoxic compared to normoxic control arteries. Second, dopa beta hydroxylase (D β H), the rate limiting enzyme in the synthesis of NE is significantly upregulated in fetal hypoxic compared to normoxic control arteries; and denervation ablates DBH. Third, medial wall thicknesses were significantly increased in nerve intact fetal hypoxic arteries compared to normoxic arteries and denervation produced opposite effects. In like manner, hypoxia also significantly increased arterial stiffness and this effect was modulated by sympathectomy in both normoxic and hypoxic arteries. Fourth, hypoxia significantly decreased colocalization of non-contractile (SMemb) and intermediate (SM1) myosin isoforms with SM α A, but increased colocalization of myosin contractile isoform (SM2) with SM α A. In contrast, sympathectomy significantly increased colocalization of SMemb with SM α A, but decreased colocalization of SM1 and SM2 with SM α A. Fifth, perivascular nerve induced NE mediates hypoxic vascular remodeling of ovine cerebral arteries through α -1 adrenoceptors.

Long-term Hypoxia Enhances Arterial Reactivity to Electrical Stimulation

Hypoxia has been shown to induce structural changes (34) in blood vessels that consequently alter their function (34, 43). In response to low oxygen levels, hypoxia inducible factor (HIF1- α) binds to hypoxia response elements present on promoter regions of angiogenic and glycolytic genes (24). VEGF is widely known as the most prominent of the angiogenic genes and has been overwhelmingly shown to mediate

processes of angiogenesis and vasculogenesis in response to hypoxia (19). Of recent, in addition to mediating angiogenesis, VEGF has been implicated in vascular remodeling which is a dynamic process involving structural changes secondary to different endogenous and exogenous stimuli (13). VEGF signals through its receptor-2/ fetal liver kinase-1 (Flk-1) to enhance angiogenesis, mitogenesis and cell survival and through its receptor-1/fms-like tyrosine kinase 1 (Flt-1) to enhance ligand binding during angiogenesis (1, 15). Though many studies have suggested VEGF to be a highly specific mitogen for endothelial cells (14), more recent findings have shown its effects on non-endothelial cells such as smooth muscle cells (18), tumor cells (8), chondrocytes (7) and neurons (35, 38). In testing the hypothesis that hypoxic remodeling of arteries occurs indirectly through the perivascular sympathetic nerves, we conducted electrical nerve stimulation of ovine fetal middle cerebral arteries. Results here showed a significant increase in response to transmural nerve stimulation in hypoxic arteries compared to their normoxic controls. Consistent with previous studies conducted in non-ovine species, a possible mechanistic explanation for this observation could be an upregulated sympathetic innervation of the arterial system during periods of hypoxia {Ruijtenbeek, 2000, 11104750}. Damon in her 2005 publication also suggested that sympathetic innervation modulates arterial medial cell differentiation (10). Sympathetic hyperinnervation and enhanced smooth muscle cell differentiation might ultimately result in an increase in cell populations of a potent contractile phenotype during periods of hypoxia. To test the hypothesis that upregulated perivascular nerve activity in hypoxic arteries mediates enhanced reactivity to nerve stimulation, we quantified D β H, the rate limiting enzyme in NE synthesis.

NE From Sympathetic Perivascular Nerves Mediates Response to Transmural Stimulation in Hypoxic Arteries

To determine the possible involvement of NE, the main sympathetic postganglionic neurohormone in the upregulated reactivity of hypoxic arteries to transmural stimulation, we quantified DbH, the rate-limiting enzyme in the synthesis of NE. To minimize influence of the endothelium involving nitric oxide (NO) co-release of NE (4) and also to test the indirect remodeling effect of VEGF via sympathetic perivascular nerves, we made use of endothelium-denuded arteries from both normoxic and hypoxic arteries. Western blot quantification of D β H showed a 354% increase in hypoxic arteries compared to their normoxic controls, suggesting that sympathetic perivascular nerve activity was significantly upregulated during hypoxic acclimatization. Similarly, immunohistochemical analysis revealed a greater D β H fluorescence in hypoxic compared to normoxic arteries and denervation ablates this response. This result is consistent with non-ovine studies that suggest a hypoxia induced increase in sympathetic inputs from perivascular nerves (16) {Ruijtenbeek, 2000, 11104750} (26). Other possible mechanisms may involve a hypoxia-induced attenuation of presynaptic inhibition of current-induced release of NE (5, 28) or an enhancement of maturation of fetal arteries through shortening of the synaptic cleft width (28) both of which remain to be morphometrically tested. Together these findings suggest that hypoxic acclimatization results in potent upregulation of sympathetic perivascular innervation and/or activity.

Sympathetic Perivascular Nerves Mediate Hypoxic Changes in Arterial Structure

Long-term acclimatization to hypoxia has been shown to result in adaptive changes in arterial structure that consequently alter function (34, 43). To test the hypothesis that

hypoxic upregulation of sympathetic innervations alters arterial structure we measured arterial thicknesses and stiffness coefficients in arteries harvested from intact and denervated normoxic and hypoxic sheep. Consistent with literature, our results showed that hypoxia significantly increased medial wall thickness (42, 50) and denervation produced the opposite effect in hypoxic arteries with no effect in normoxic arteries. These results suggest that hypoxia might be inducing cell hyperplasia (3) or hypertrophy (32, 33, 36) in the wall intima and that perivascular nerves support this process possibly through enhanced cellular differentiation and proliferation (10). Similarly, consistent with literature, hypoxia significantly increased stiffness of the cerebral arteries (11, 45), however denervation increased stiffness further and more so in hypoxic arteries. Increases in stiffness observed in hypoxic arteries could be possibly due to hypoxia induced increases in cellular collagen content or increased ratio of collagen to elastin fibers in smooth muscle cells (11, 45). Sympathectomy enhanced hypoxia induced arterial stiffness suggesting that perivascular nerves attenuate arterial stiffness through mechanisms that might involve release certain vasotrophic substances such as NO (9) that enhance vasodilatation and consequently reduce arterial stiffness (17, 22).

Sympathetic Perivascular Nerves Mediate Hypoxia-induced Changes in Arterial Function Through Alteration of Contractile Protein Colocalization

Coupling between smooth muscle thick filaments (Myosin chains) and thin filaments (SM α A) in the presence of ATP initiates contractile force, which is propagated to the cell membrane by cytoskeletal proteins (20, 48). Recent works from our lab have shown potent changes in contractile protein abundances and interactions (6) secondary to

hypoxic acclimatization (1, 21). To test the hypothesis that sympathetic perivascular nerves mediate hypoxic changes in smooth muscle cell interactions between the three myosin heavy chain isoforms (SMEMB, SM1 and SM2) and SM α A, we conducted immunohistochemistry and confocal colocalization imaging. Results showed that SMemb, a smooth muscle myosin heavy chain isoform with characteristic synthetic phenotype, colocalization with SM α A was shown to be significantly downregulated by hypoxia, and denervation produced the opposite effect. This result suggests a.) That long-term hypoxia promotes cellular reorganization which downregulates colocalization of non-contractile and synthetic myosin with SM α A b.) That sympathetic perivascular nerves mediate hypoxic dedifferentiation of certain subsets of the smooth muscle cells with consequent decreases in SMemb - SM α A colocalization. Similarly, SM1 (smooth muscle myosin heavy chain intermediate contractile isoform) colocalization with SM α A decreased with hypoxia suggesting that hypoxia promotes factors that inhibit coupling between both non-contractile and intermediate contractile isoforms of myosin with SM α A. Whereas denervation decreased SM1- SM α A colocalization in normoxic arteries, denervation increased their colocalization coefficient in hypoxic arteries, suggesting that sympathetic perivascular nerves preferentially support a downregulation of colocalization between synthetic myosin phenotypes and SM α A in hypoxic arteries. In sharp contrast to SMemb and SM1, long-term hypoxia significantly increased SM2 (fully contractile myosin isoform) colocalization with SM α A and denervation potentially produced opposite effects. This result shows that whereas hypoxia downregulated colocalization of SMemb and SM1 with SM α A, it potentially enhanced SM2- SM α A colocalization, which supports contractile force, and perivascular sympathetic nerves seem to mediate this process. The

exact mechanisms by which sympathetic perivascular nerves alter the phenotype of smooth muscle cells and their marker proteins remain exciting topics of future. Some studies in non-ovine species have suggested that sensory nerves have the ability to limit vasoconstriction of arteries during periods of perivascular nerve stimulation (49). It is possible that this protective effect is compromised during periods of hypoxia.

NE Mediates Hypoxic Reorganization of Contractile Proteins via its α -1 Adrenoceptors

Sympathetic nerves have been overwhelming shown to exert a protective effect on vital organs during periods of stress through the release of catecholamines (5); of which norepinephrine is the most common. Also perivascular nerve stimulation has been shown to result in the release of norepinephrine from the nerve terminals in the end organs (12). NE signals predominantly through α -1 adrenoceptors present on blood vessels (46). Using SM2- SM α A colocalization as a reporter assay, we tested the hypothesis that NE mediates hypoxic changes in contractile protein interactions via its α -1 adrenoceptors. In arteries harvested from sympathectomized hypoxic fetal sheep, 48 hrs of starvation in organ culture produced no change; NE produced a significant increase in SM2- SM α A colocalization and addition of prazosin (an α -1 adrenoceptor blocker) produced a marked decrease in SM2- SM α A colocalization. This result suggests that remodeling effects of hypoxia characterized by a preferential increase in colocalization between myosin contractile isoform, SM2 and SM α A occur through NE signaling via its α -1 adrenoceptors present on ovine cerebral arteries.

Overview

VEGF has been suggested to mediate hypoxic vascular remodeling in addition to its well-established role in angiogenesis and vasculogenesis through its receptor tyrosine kinases. Here we have explored the possibility of VEGF contributing to hypoxic vascular remodeling of fetal ovine cerebral arteries through VEGF receptors present on sympathetic perivascular nerves. First set of experiments showed that response to electrical transmural stimulation was significantly increased in hypoxic fetal (FH) arteries compared with their normoxic controls (FN) (**Figure 1**). This was partially explained by a 3-fold increase in dopa- β -hydroxylase in FH group compared to FN, which is the rate-limiting enzyme for NE synthesis (Figure 2). In both groups denervation ablated DBH increases (**Figure 2**). Consistent with existing literature, long term hypoxia produced significant increases in medial arterial wall thickness and stiffness (**Figure 3**). Whereas denervation decreased wall thickness, it further modulated stiffening of the arteries, suggesting that sympathetic perivascular nerves support hypoxic increases in arterial wall thickness but attenuate stiffening of arteries possibly through nerve induced increases in NO. Long-term hypoxia resulted in significant decreases in SMemb – SM α A colocalization (**Figure 4**) as well as SM1 – SM α A colocalization (**Figure 5**) but increased SM2 – SM α A colocalization (**Figure 6**). Denervation produced opposite effects on colocalization of these pairs of contractile proteins suggesting that sympathetic perivascular nerves mediate hypoxic shifts in phenotype from a synthetic to a more contractile one resulting in increased reactivity to transmural current stimuli. Organ culture denervated hypoxic arterial segments with NE produced significant increases in SM2 – SM α A colocalization (**Figure 7**) and co-cultures with prazosin ablated these

increases. Together these results support our hypothesis that perivascular nerves mediate hypoxic vascular remodeling through phenotypic shifts in smooth muscle cell phenotype.

Acknowledgments

The work reported in this manuscript was supported by USPHS Grants HL54120, HD31266, HL64867 and the Loma Linda University School of Medicine.

References

1. Adeoye OO, Butler SM, Hubbell MC, Semotiuk AJ, Williams JM, and Pearce WJ. Contribution of Increased VEGF Receptors to Hypoxic Changes in Fetal Ovine Carotid Artery Contractile Proteins. *Am J Physiol Cell Physiol*, 2013: PMID. 23325408.
2. Appleton AB and Waites GM. A surgical approach to the superior cervical ganglion and related structures in the sheep. *J Physiol* 135: 52-57, 1957: PMID. 13398964.
3. Bevan RD, van Marthens E, and Bevan JA. Hyperplasia of vascular smooth muscle in experimental hypertension in the rabbit. *Circ Res* 38: 58-62, 1976: PMID. 1269112.
4. Buchholz J and Duckles SP. Chronic hypoxia alters prejunctional alpha(2)-receptor function in vascular adrenergic nerves of adult and fetal sheep. *Am J Physiol Regul Integr Comp Physiol* 281: R926-934, 2001: PMID. 11507010.
5. Buchholz J, Edwards-Teunissen K, and Duckles SP. Impact of development and chronic hypoxia on NE release from adrenergic nerves in sheep arteries. *Am J Physiol* 276: R799-808, 1999: PMID. 10070141.
6. Butler SM, Abrassart JM, Hubbell MC, Adeoye O, Semotiuk A, Williams JM, Mata-Greenwood E, Khorram O, and Pearce WJ. Contributions of VEGF to age-dependent transmural gradients in contractile protein expression in ovine carotid arteries. *Am J Physiol Cell Physiol* 301: C653-666, 2011: PMID. 21653901.
7. Carlevaro MF, Cermelli S, Cancedda R, and Descalzi Cancedda F. Vascular endothelial growth factor (VEGF) in cartilage neovascularization and chondrocyte differentiation: auto-paracrine role during endochondral bone formation. *J Cell Sci* 113 (Pt 1): 59-69, 2000: PMID. 10591625.
8. Carmeliet P. Angiogenesis in health and disease. *Nature medicine* 9: 653-660, 2003: PMID. 12778163.
9. Chen FY and Lee TJ. Arginine synthesis from citrulline in perivascular nerves of cerebral artery. *J Pharmacol Exp Ther* 273: 895-901, 1995: PMID. 7752095.

10. Damon DH. Sympathetic innervation promotes vascular smooth muscle differentiation. *Am J Physiol Heart Circ Physiol* 288: H2785-2791, 2005: PMID. 15665063.
11. Dodson RB, Rozance PJ, Fleenor BS, Petrash CC, Shoemaker LG, Hunter KS, and Ferguson VL. Increased arterial stiffness and extracellular matrix reorganization in intrauterine growth-restricted fetal sheep. *Pediatr Res* 73: 147-154, 2013: PMID. 23154756.
12. Elenkov IJ, Wilder RL, Chrousos GP, and Vizi ES. The sympathetic nerve--an integrative interface between two supersystems: the brain and the immune system. *Pharmacol Rev* 52: 595-638, 2000: PMID. 11121511.
13. Ferrara N and Davis-Smyth T. The biology of vascular endothelial growth factor. *Endocr Rev* 18: 4-25, 1997: PMID. 9034784.
14. Ferrara N, Houck K, Jakeman L, and Leung DW. Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocr Rev* 13: 18-32, 1992: PMID. 1372863.
15. Gerber HP, McMurtrey A, Kowalski J, Yan M, Keyt BA, Dixit V, and Ferrara N. Vascular endothelial growth factor regulates endothelial cell survival through the phosphatidylinositol 3'-kinase/Akt signal transduction pathway. Requirement for Flk-1/KDR activation. *J Biol Chem* 273: 30336-30343, 1998: PMID. 9804796.
16. Giussani DA, Riquelme RA, Sanhueza EM, Hanson MA, Blanco CE, and Llanos AJ. Adrenergic and vasopressinergic contributions to the cardiovascular response to acute hypoxaemia in the llama fetus. *J Physiol* 515 (Pt 1): 233-241, 1999: PMID. 9925892.
17. Grignola JC, Domingo E, Aguilar R, Vazquez M, Lopez-Messeguer M, Bravo C, and Roman A. Acute absolute vasodilatation is associated with a lower vascular wall stiffness in pulmonary arterial hypertension. *Int J Cardiol* 164: 227-231, 2013: PMID. 21788089.

18. Grosskreutz CL, Anand-Apte B, Duplaa C, Quinn TP, Terman BI, Zetter B, and D'Amore PA. Vascular endothelial growth factor-induced migration of vascular smooth muscle cells in vitro. *Microvasc Res* 58: 128-136, 1999: PMID. 10458928.
19. Hoeben A, Landuyt B, Highley MS, Wildiers H, Van Oosterom AT, and De Bruijn EA. Vascular endothelial growth factor and angiogenesis. *Pharmacol Rev* 56: 549-580, 2004: PMID. 15602010.
20. Horowitz A, Menice CB, Laporte R, and Morgan KG. Mechanisms of smooth muscle contraction. *Physiol Rev* 76: 967-1003, 1996: PMID. 8874491.
21. Hubbell MC, Semotiuk AJ, Thorpe RB, Adeoye OO, Butler SM, Williams JM, Khorram O, and Pearce WJ. Chronic hypoxia and VEGF differentially modulate abundance and organization of myosin heavy chain isoforms in fetal and adult ovine arteries. *Am J Physiol Cell Physiol* 303: C1090-1103, 2012: PMID. 22992677.
22. Joannides R, Richard V, Moore N, Godin M, and Thuillez C. Influence of sympathetic tone on mechanical properties of muscular arteries in humans. *Am J Physiol* 268: H794-801, 1995: PMID. 7864207.
23. Kamitomo M, Longo LD, and Gilbert RD. Right and left ventricular function in fetal sheep exposed to long-term high-altitude hypoxemia. *Am J Physiol* 262: H399-405, 1992: PMID. 1539699.
24. Ke Q and Costa M. Hypoxia-inducible factor-1 (HIF-1). *Mol Pharmacol* 70: 1469-1480, 2006: PMID. 16887934.
25. Kranenburg AR, de Boer WI, Alagappan VK, Sterk PJ, and Sharma HS. Enhanced bronchial expression of vascular endothelial growth factor and receptors (Flk-1 and Flt-1) in patients with chronic obstructive pulmonary disease. *Thorax* 60: 106-113, 2005: PMID. 15681497.
26. Kummer W. Pulmonary vascular innervation and its role in responses to hypoxia: size matters! *Proc Am Thorac Soc* 8: 471-476, 2011: PMID. 22052922.

27. Langer I, Vertongen P, Perret J, Fontaine J, Atassi G, and Robberecht P. Expression of vascular endothelial growth factor (VEGF) and VEGF receptors in human neuroblastomas. *Med Pediatr Oncol* 34: 386-393, 2000: PMID. 10842244.
28. Longo LD and Pearce WJ. Fetal cerebrovascular acclimatization responses to high-altitude, long-term hypoxia: a model for prenatal programming of adult disease? *Am J Physiol Regul Integr Comp Physiol* 288: R16-24, 2005: PMID. 15590993.
29. Marko SB and Damon DH. VEGF promotes vascular sympathetic innervation. *Am J Physiol Heart Circ Physiol* 294: H2646-2652, 2008: PMID. 18408130.
30. Masood R, Cai J, Zheng T, Smith DL, Hinton DR, and Gill PS. Vascular endothelial growth factor (VEGF) is an autocrine growth factor for VEGF receptor-positive human tumors. *Blood* 98: 1904-1913, 2001: PMID. 11535528.
31. Michiels C. Physiological and pathological responses to hypoxia. *Am J Pathol* 164: 1875-1882, 2004: PMID. 15161623.
32. Olivetti G, Anversa P, Melissari M, and Loud AV. Morphometry of medial hypertrophy in the rat thoracic aorta. *Lab Invest* 42: 559-565, 1980: PMID. 7382430.
33. Owens GK and Schwartz SM. Alterations in vascular smooth muscle mass in the spontaneously hypertensive rat. Role of cellular hypertrophy, hyperploidy, and hyperplasia. *Circ Res* 51: 280-289, 1982: PMID. 7116579.
34. Pearce WJ and Ashwal S. Developmental changes in thickness, contractility, and hypoxic sensitivity of newborn lamb cerebral arteries. *Pediatr Res* 22: 192-196, 1987: PMID. 3658545.
35. Piltonen M, Planken A, Leskela O, Myohanen TT, Hanninen AL, Auvinen P, Alitalo K, Andressoo JO, Saarma M, and Mannisto PT. Vascular endothelial growth factor C acts as a neurotrophic factor for dopamine neurons in vitro and in vivo. *Neuroscience* 192: 550-563, 2011: PMID. 21767614.
36. Plate KH, Breier G, Millauer B, Ullrich A, and Risau W. Up-regulation of vascular endothelial growth factor and its cognate receptors in a rat glioma model of tumor angiogenesis. *Cancer Res* 53: 5822-5827, 1993: PMID. 7694795.

37. Roberts DM, Kearney JB, Johnson JH, Rosenberg MP, Kumar R, and Bautch VL. The vascular endothelial growth factor (VEGF) receptor Flt-1 (VEGFR-1) modulates Flk-1 (VEGFR-2) signaling during blood vessel formation. *Am J Pathol* 164: 1531-1535, 2004: PMID. 15111299.
38. Rosenstein JM, Krum JM, and Ruhrberg C. VEGF in the nervous system. *Organogenesis* 6: 107-114, 2010: PMID. 20885857.
39. Santhanam AV, Smith LA, He T, Nath KA, and Katusic ZS. Endothelial progenitor cells stimulate cerebrovascular production of prostacyclin by paracrine activation of cyclooxygenase-2. *Circ Res* 100: 1379-1388, 2007: PMID. 17395873.
40. Semenza GL. Expression of hypoxia-inducible factor 1: mechanisms and consequences. *Biochem Pharmacol* 59: 47-53, 2000: PMID. 10605934.
41. Sondell M, Sundler F, and Kanje M. Vascular endothelial growth factor is a neurotrophic factor which stimulates axonal outgrowth through the flk-1 receptor. *Eur J Neurosci* 12: 4243-4254, 2000: PMID. 11122336.
42. Stenmark KR, Fagan KA, and Frid MG. Hypoxia-induced pulmonary vascular remodeling: cellular and molecular mechanisms. *Circ Res* 99: 675-691, 2006: PMID. 17008597.
43. Stenmark KR and McMurtry IF. Vascular remodeling versus vasoconstriction in chronic hypoxic pulmonary hypertension: a time for reappraisal? *Circ Res* 97: 95-98, 2005: PMID. 16037575.
44. Strizzi L, Catalano A, Vianale G, Orecchia S, Casalini A, Tassi G, Puntoni R, Mutti L, and Procopio A. Vascular endothelial growth factor is an autocrine growth factor in human malignant mesothelioma. *J Pathol* 193: 468-475, 2001: PMID. 11276005.
45. Thompson JA, Richardson BS, Gagnon R, and Regnault TR. Chronic intrauterine hypoxia interferes with aortic development in the late gestation ovine fetus. *J Physiol* 589: 3319-3332, 2011: PMID. 21540340.
46. van Brummelen P, Jie K, and van Zwieten PA. Alpha-adrenergic receptors in human blood vessels. *Br J Clin Pharmacol* 21 Suppl 1: 33S-39S, 1986: PMID. 2871855.

47. Vera Janavel GL, De Lorenzi A, Cortes C, Olea FD, Cabeza Meckert P, Bercovich A, Criscuolo M, Laguens R, and Crottogini A. Effect of vascular endothelial growth factor gene transfer on infarct size, left ventricular function and myocardial perfusion in sheep after 2 months of coronary artery occlusion. *J Gene Med* 14: 279-287, 2012: PMID. 21954009.
48. Webb RC. Smooth muscle contraction and relaxation. *Adv Physiol Educ* 27: 201-206, 2003: PMID. 14627618.
49. Westcott EB and Segal SS. Ageing alters perivascular nerve function of mouse mesenteric arteries in vivo. *J Physiol* 591: 1251-1263, 2013: PMID. 23247111.
50. Xu XF, Lv Y, Gu WZ, Tang LL, Wei JK, Zhang LY, and Du LZ. Epigenetics of hypoxic pulmonary arterial hypertension following intrauterine growth retardation rat: epigenetics in PAH following IUGR. *Respir Res* 14: 20, 2013: PMID. 23406533.
51. Zhang J, Silva T, Yarovinsky T, Manes TD, Tavakoli S, Nie L, Tellides G, Pober JS, Bender JR, and Sadeghi MM. VEGF blockade inhibits lymphocyte recruitment and ameliorates immune-mediated vascular remodeling. *Circ Res* 107: 408-417, 2010: PMID. 20538685.

CHAPTER FIVE

CONCLUSIONS & FUTURE DIRECTIONS

A broad spectrum of maternal conditions ranging from smoking, diabetes, chronic hypertension or pregnancy induced hypertensive disorders create a hypoxic environment to the developing fetus. Hypoxia remains one of the most common etiologic and/or contributory factors to increased fetal morbidity and mortality both in the USA and the world at large. Whereas severe acute episodes are lethal, moderate to mild chronic episodes though compatible with life result in altered cardiovascular functioning. Grounded knowledge on mechanisms of these altered cardiovascular patterns becomes extremely important to developing therapeutics to ensure survival of the developing fetus as well as treating future complications.

HIF-1 has been regarded as the global mediator of hypoxia since its discovery almost three decades ago. The activity, stability and post translational modifications of its oxygen sensing and regulated subunit (α subunit) determines regulate the induction of transcription of other hypoxia responsive genes including : glycolytic enzyme genes, erythropoietin and VEGF amongst others. VEGF is largely known as the mediator of processes of angiogenesis and vasculogenesis in normal embryonic development, collagen constriction and neovascularization during wound healing and tumorigenesis involving expansion of vascular network supplying the cancer cells. VEGF overwhelmingly contributes to processes of proliferation and cell survival which is desirable during physiologic developmental processes but adverse in pathologic

conditions such as tumorigenesis. Although VEGF is famously known for its role in angiogenesis and vasculogenesis, recent studies have implicated this transcription factor in vascular remodeling, which is a dynamic process involving changes in vessel structure resulting in altered function. Because of the important role of VEGF in angiogenesis and vascular remodeling in physiologic and pathologic processes, it becomes important and of high clinical relevance to elucidate mechanisms through which this vasotrophic factor signals.

The first chapter of this dissertation introduces HIF-1 α , its induction of transcription of VEGF gene and translational synthesis of the cytokine. The overall hypothesis “VEGF mediates hypoxic vascular remodeling through a direct effect on the vascular smooth muscle and indirectly through the sympathetic perivascular nerves” for this thesis was introduced. Whereas VEGF is generally known as a specific mitogen for endothelial cells, recent findings have suggested its direct effect on non-endothelial cells such as chondrocytes, neural cells and smooth muscle cells resident in the intimal layer of arteries. Current knowledge on VEGF and its principal tyrosine receptor kinases flt-1 (VEGF receptor 1) and flk-1 / kdr (VEGF receptor 2) were highlighted in this chapter.

Chapter 2 tested and provided data that supports the hypothesis that smooth muscle cells are organized into lamina of similar phenotype with characteristics that depend on the relative position between the lumen and the adventitia and involve the direct effects of vasotrophic factors such as VEGF, which could potentially act independently of the vascular endothelium in an age-dependent manner.

Chapter 3 highlights our novel study which tested and provided data to support the hypothesis that hypoxic increases in VEGF not only promotes angiogenesis but

contributes to remodeling of artery wall structure and contractility through phenotypic transformation of smooth muscle. In this study low dose VEGF (3 ng/ml) replicated hypoxic decreases in colocalization of SMemb with SM α A in fetal arteries and decreases in colocalization of SM2 with SM α A in adult arteries.

With most of the current literature known on arterial remodeling focused on pathologies in adult vasculature, it becomes important to conduct research on mechanisms involved with hypoxic vascular remodeling in fetal vasculature. Chapter 4 of this dissertation tested and provided data to support the hypothesis that VEGF contributes to hypoxic fetal vascular remodeling through changes in the abundance, organization and function of contractile proteins, with emphasis on SM α A, MLCK and MLC₂₀. VEGF in organ culture replicated hypoxic increases in SM α A and MLC₂₀ abundances as well as precipitous decreases in MLCK. Similarly, VEGF in organ culture mimicked hypoxic increases in colocalization of MLCK with SM α A, MLC₂₀ with SM α A and MLCK with MLC₂₀ despite marked decreases in MLCK abundance, thus suggesting that hypoxia and VEGF exert distinct effects on each contractile protein and that contractile protein abundance and organization are differentially regulated. Of remarkable importance in this study is the short term increase in VEGF and long term increases in VEGF receptor tyrosine kinases (flt-1 and flk-1), suggesting that hypoxic vascular remodeling initiated by VEGF are maintained long-term by potent increases in VEGF receptors.

Having tested mechanisms involved with hypoxic vascular remodeling in fetal ovine carotid arteries, we conducted a parallel study to test the hypothesis that chronic hypoxia produces a sustained increase in the density of vascular VEGF receptors, which in turn helps mediate the long-term effects of hypoxia on vascular remodeling in adult

ovine carotid arteries. Chapter five shows that whereas VEGF levels were similar to basal levels VEGF receptors 1 and 2 (flt-1 and flk-1) were significantly increased after 110 days of hypoxic acclimatization in ovine adult carotid arteries, suggesting that the mechanism through which remodeling effects of hypoxia are maintained long term is through upregulation of VEGF receptor densities. Also this study showed that VEGF in culture replicated hypoxic increases in SM α A and MLC₂₀ but produced no significant effect on MLCK suggesting that post-natal maturation alters reactivity to MLCK. Also, VEGF in culture replicated hypoxic effect only on MLC₂₀ colocalization with SM α A, further supporting the hypothesis that reactivity of arteries to hypoxia and VEGF are fundamentally different in fetal and adult ovine species.

Chapter 6 explored the hypothesis that hypoxic increases in VEGF contributes to arterial remodeling possibly through activation of VEGF receptors on sympathetic perivascular nerve leading to altered trophic input to the arterial wall. This chapter shows that increased response to transmural electrical stimulation in hypoxic arteries compared to normoxic controls might be related to upregulation of sympathetic inputs to the artery. Also long term hypoxia was shown to modulate contractile differentiation as evidenced by decreased colocalization of non-muscle myosin heavy chain (SMemb) with SM α A as well as simultaneous increase in colocalization between fully contractile smooth muscle heavy myosin chain (SM2) with SM α A. Sympathectomy ablated the effects of hypoxia on phenotypic shifts of the contractile proteins. Also NE in organ culture potently increased colocalization between SM2 and SM α A and prazosin (α -1 adrenoceptor blocker) attenuated this increase.

Future relevant work geared towards advancement of the VEGF remodeling hypothesis would involve testing involvement of other vasotrophic factors in hypoxic remodeling since there is overwhelming evidence that VEGF does not work in isolation. Platelet derived growth factor (PDGF) and fibroblast growth factor (FGF) in particular have been closely associated with the role VEGF plays in hypoxic remodeling. With VEGF able to bind with PDGF receptors, this vasotrophic factor occupies the front line in future studies on involvement of other cytokines in hypoxic vascular remodeling. This dissertation focused on the role of VEGF and its receptors in long-term hypoxic vascular remodeling. One of the novel findings was upregulation of VEGF receptors during long-term hypoxic acclimatization; clearly it will be logical to assess levels of VEGF receptors during acute phases of hypoxia. Relevant experiments here will involve sheep made acutely hypoxic via blowing of nitrogen (N₂) through a tracheostomy for a week and then assessing VEGF receptor density via western blotting on the carotid arteries and middle cerebral arteries. Since our work primarily focused on role of growth factors and their receptors on hypoxic vascular remodeling with emphasis on VEGF, the next logical line of study will be to assess intracellular kinases that couple these agonist-receptor signaling to the nucleus of the cell. Assessment of potential intracellular kinases that have been suggested in VEGF signaling would involve – quantification of ERK 1 and 2 and P-I-3 Kinase and their targets (Akt, phospho-Akt and phospho-ERK), use of their antagonists in culture and then assessing their effect on contractile protein expression, organization and contractility of the arteries. ERK 1 and 2 antagonists that have been suggested in the literature include U0126 and PD98059 while those of PI3 kinase are Wortmannin and LY294002. These studies will assess potential changes of these intracellular kinases with

hypoxia , VEGF or with ganglionectomy. The concept of short-term hypoxic increases in VEGF but long-term return to baseline creates a research question of testing what mechanisms mediate this process. Studies involving assessment of possible VEGF mRNA post-transcriptional regulation via micro RNAs, alternative splicing or riboswitches or VEGF post translational regulation via methylation, acetylation or ubiquitination become exciting topics for future research.

Other than HIF and VEGF that have known hypoxia response elements on their genes, studies have only hinted at a possibility of presence of HREs on contractile protein genes and other molecules that respond to hypoxia. In our studies, MLCK has been overwhelmingly shown to be very sensitive hypoxia necessitating the importance of future studies to elucidating potential presence of HREs on the contractile protein genes. In this study, we have hypothesized and tested the involvement of sympathetic perivascular nerves with hypoxic vascular remodeling via release of NE; many studies have however suggested that NPY is co-released with NE. Future studies would be necessary to test the involvement of peptidergic vasotrophic factors such as NPY as well as ATP with hypoxic vascular remodeling. Quantification of these agonists and their receptors in nerve intact and denervated sheep as well as organ culture with them will provide novel information about their possible involvement with hypoxic vascular remodeling.

VEGF is known to mediate both physiologic and pathological processes involving angiogenesis and vasculogenesis. Tumorigenesis is one of the pathologic processes that VEGF mediates. Over the years researchers have discovered small molecule inhibitors of VEGF necessitating testing via clinical trials at various levels. Our organ culture model

would serve a clinical purpose of testing the efficacy and safety of many of these small molecule inhibitors that could serve future uses in combating the scourge of cancer malignancy and recurrence.

Overall, from a clinical perspective, many fetal as well as maternal conditions result in varying levels of hypoxia that adversely affect the developing fetus. These conditions result in long hospital stay in the NICU, increased morbidity and mortality as well as chronic altered functioning of various organs in the fetal body. Gaining a better understanding of processes involved with altered functioning of the vasculature is a valuable first step in improving and optimizing strategies geared towards better clinical outcomes. Manipulation of vasotrophic factors such as VEGF, clearly involved with remodeling, may provide valuable opportunities for therapy in neonates that have survived hypoxia in-utero. The purpose of this thesis was to elucidate mechanisms through which VEGF mediates hypoxic remodeling and by inference first steps towards reduction of hypoxia related morbidities and mortalities.