



ISSN 1600-5368

Received 20 September 2014

Accepted 28 October 2014

Edited by H. Stoeckli-Evans, University of Neuchâtel, Switzerland

Keywords: crystal structure; azobenzene; benzoic acid; liquid crystal; nematic phase**CCDC reference:** 1031374**Supporting information:** this article has supporting information at journals.iucr.org/e



Crystal structure of (*E*)-4-[2-[4-(allyloxy)phenyl]-diazenyl]benzoic acid

Md. Lutfor Rahman,^{a*} Mashitah Mohd. Yusoff,^a Jamil Ismail,^a Huey Chong Kwong^b and Ching Kheng Quah^c^aUniversity Malaysia Pahang, Faculty of Industrial Sciences and Technology, 26300 Gambang, Kuantan, Pahang, Malaysia, ^bSchool of Chemical Sciences, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia, and ^cX-ray Crystallography Unit, School of Physics, Universiti Sains Malaysia, 11800 USM, Penang, Malaysia. *Correspondence e-mail: lutfor73@gmail.com

The title compound, C₁₆H₁₄N₂O₃, has an *E* conformation about the azobenzene [$-\text{N}=\text{N}- = 1.2481(16) \text{ \AA}$] linkage. The benzene rings are almost coplanar [dihedral angle = $1.36(7)^\circ$]. The O atoms of the carboxylic acid group are disordered over two sets of sites and were refined with an occupancy ratio of 0.5:0.5. The two disordered components of the carboxylic acid group make dihedral angles of $1.5(14)$ and $3.8(12)^\circ$ with the benzene ring to which they are attached. In the crystal, molecules are linked *via* pairs of O—H...O hydrogen bonds, forming inversion dimers. The dimers are connected *via* C—H...O hydrogen bonds, forming ribbons lying parallel to [120]. These ribbons are linked *via* C—H... π interactions, forming slabs parallel to (001).

1. Chemical context


It is interesting to note that the title compound shows a nematic phase (Cr 190 N 218 I). Hence, liquid crystallinity may be induced by the formation of hydrogen-bonded dimers. A number of liquid crystal (LC) systems containing hydrogen bonds that function between identical molecules have been reported (Kang & Samulski, 2000; Rahman *et al.*, 2012). Much attention has been paid to hydrogen-bonded supramolecular LCs, including LC dimers based on hydrogen-bonding interactions and several supramolecular LC trimers based on hydrogen-bonding interactions (Lee *et al.*, 2001; Paleos & Tsiourvas, 2001; Takahashi *et al.*, 2003; Bai *et al.*, 2007). A particular aspect of photonics, in which the molecular geometry can be controlled by light, is being proposed as a future technology for optical storage devices (Ikeda & Tsutsumi, 1995; Jayalaxmi *et al.*, 2009). The heart of the phenomenon in such systems is the reversible photo-induced shape transformation of the molecules containing the photochromic azobenzene groups. The title compound contains an azo ($-\text{N}=\text{N}-$) linkage, it was easy to synthesize and hence cost-effective for the possibility of photochromism and photoisomerization usage (Lutfor *et al.*, 2013a,b). We report herein on its synthesis and crystal structure.



2. Structural commentary

The molecular structure of the title molecule is illustrated in Fig. 1. The oxygen atoms forming the carboxylic acid group




Figure 1

The molecular structure of the title compound, showing the atom labelling. Displacement ellipsoids are drawn at the 30% probability level. Only one component of the disordered carboxylic acid group is shown.

Table 1

Hydrogen-bond geometry (Å, °).

Cg1 is the centroid of the C4–C9 ring.

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$O3-H3\cdots O2^i$	0.82	1.90	2.71 (3)	166
$C6-H6A\cdots O2^{ii}$	0.93	2.59	3.367 (15)	145
$C3-H3A\cdots Cg1^{iii}$	0.97	2.66	3.504 (2)	145

Symmetry codes: (i) $-x-1, -y+2, -z$; (ii) $-x+1, -y+1, -z$; (iii) $x+1, y, z$.

are each disordered over two positions and were refined with half occupancy. The carboxylic acid group (C16/O2/O3) is almost coplanar with the attached benzene ring (C10–C15), making dihedral angles of 3.44 (9) and 3.65 (8)° for the two disorder components. The title compound has an *E* conformation about the azobenzene ($-N=N-$) linkage, the length of the N1–N2 bond is 1.2481 (16) Å and the torsion angle for the azo unit (C7–N1=N2–C10) is 179.99 (10)°, which is comparable with the values of $ca \pm 180^\circ$ observed in 4,4-


azinodibenzoic acid (Yu & Liu, 2009) and (*E*)-ethyl-4-[[4-(decanoxloxy)phenyl]diazenyl] benzoate (Lai *et al.*, 2002). The benzene rings (C4–C9) and (C10–C15) are almost coplanar, making a dihedral angle of 1.38 (7)°, compared with 6.79 (9)° in the previously reported compound 4-[(*E*)-2-[4-(but-3-en-1-yloxy)phenyl]-diazene-1-yl]benzoic acid, (Rahman *et al.*, 2012).

3. Supramolecular features

In the crystal, molecules are linked *via* pairs of O–H \cdots O hydrogen bonds, forming inversion dimers (Table 1 and Fig. 2). The dimers are connected *via* C–H \cdots O hydrogen bonds, forming two-molecule-thick ribbons lying parallel to [120]; see Table 1 and Fig. 3. Adjacent ribbons are linked *via* C–H \cdots π interactions, forming slabs parallel to (001), as shown in Fig. 3 (Table 1).

4. Synthesis and crystallization

The title compound was synthesized by a literature procedure (Rahman *et al.*, 2012). The diazonium salt was prepared with sodium nitrite and subsequent coupling with phenol to afforded the ethyl 4-[(4-hydroxyphenyl)diazenyl]benzoate, which was purified by crystallization and recrystallization from methanol. The azobenzene compound was alkylated with allyl bromide to give ethyl 4-[[4-(allyloxy)phenyl]diazenyl]benzoate, which was purified by crystallization from methanol/chloroform. The terminal double bonds-containing azobenzene compound was hydrolysed under basic conditions to yield the title compound. Red plate-like crystals were obtained by crystallization from an ethanol–ethyl acetate


Figure 2

A partial view along the *a*-axis of the crystal packing of the title compound, with hydrogen bonds shown as dashed lines (see Table 1 for details).




Figure 3

A partial view of the crystal packing of the title compound. Blue dashed lines represent the intermolecular hydrogen bonds within two-molecule-thick chains and the green dashed lines represent the weak intermolecular C–H... π interactions (see Table 1 for details).

mixture (1:1); m.p. 494 K. $^1\text{H NMR}$ (CDCl_3): δ 8.18 (*d*, 2H, J = 8.2 Hz), 7.94 (*d*, 2H, J = 7.1 Hz), 7.93 (*d*, 2H, J = 6.7 Hz), 7.05 (*d*, 2H, J = 8.9 Hz), 6.04 (*m*, 1H, CH=), 5.45 (*d*, 1H, J = 16.6 Hz, =CH₂), 5.31 (*d*, 1H, J = 10.2 Hz, =CH₂), 4.60 (*d*, 2H, J = 4.1 Hz, OCH₂).

Table 2
Experimental details.

Crystal data	
Chemical formula	$\text{C}_{16}\text{H}_{14}\text{N}_2\text{O}_3$
M_r	282.29
Crystal system, space group	Triclinic, $P\bar{1}$
Temperature (K)	294
a , b , c (Å)	5.0279 (4), 8.9678 (7), 15.9913 (13)
α , β , γ (°)	80.571 (2), 83.874 (2), 88.371 (2)
V (Å ³)	707.19 (10)
Z	2
Radiation type	Mo $K\alpha$
μ (mm ⁻¹)	0.09
Crystal size (mm)	0.78 × 0.22 × 0.09
Data collection	
Diffractionmeter	Bruker APEX DUO CCD area detector
Absorption correction	Multi-scan (SADABS; Bruker, 2009)
T_{\min} , T_{\max}	0.931, 0.992
No. of measured, independent and observed [$I > 2\sigma(I)$] reflections	12171, 3276, 2344
R_{int}	0.023
$(\sin \theta/\lambda)_{\text{max}}$ (Å ⁻¹)	0.651
Refinement	
$R[F^2 > 2\sigma(F^2)]$, $wR(F^2)$, S	0.043, 0.138, 1.04
No. of reflections	3276
No. of parameters	211
H-atom treatment	H-atom parameters constrained
$\Delta\rho_{\text{max}}$, $\Delta\rho_{\text{min}}$ (e Å ⁻³)	0.24, -0.16

Computer programs: APEX2 and SAINT (Bruker, 2009), SHELXS97, SHELXL97 and SHELXTL (Sheldrick, 2008), Mercury (Macrae *et al.*, 2008), and PLATON (Spek, 2009).

5. Refinement

Crystal data, data collection and structure refinement details are summarized in Table 2. Atoms O2 and O3 of the carboxylic acid group are each disordered over two positions and were refined with half occupancy each. The position of the O-bound H atom was located in a difference Fourier map and refined as a riding atom: O–H = 0.82 Å with $U_{\text{iso}}(\text{H}) = 1.5 U_{\text{eq}}(\text{O})$. The C-bound H atoms were positioned geometrically and refined using a riding model: C–H = 0.93–0.97 Å with $U_{\text{iso}}(\text{H}) = 1.2 U_{\text{eq}}(\text{C})$. Two outlier reflections, 341 and 309, were omitted from the refinement.

Acknowledgements

This research was supported by a PRGS Research Grant (No. RDU 130803).

References

- Bai, B. L., Wang, H., Xin, H., Long, B. & Li, M. (2007). *Liq. Cryst.* **34**, 659–665.
- Bruker (2009). APEX2, SAINT and SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.
- Ikeda, T. & Tsutsumi, O. (1995). *Science*, **268**, 1873–1875.
- Jayalakshmi, V., Hegde, G., Nair, G. & Prasad, S. K. (2009). *Phys. Chem. Chem. Phys.* **11**, 6450–6454.
- Kang, S. K. & Samulski, E. T. (2000). *Liq. Cryst.* **27**, 371–376.
- Lai, L.-L., Su, F.-Y., Lin, Y.-J., Ho, C.-H., Wang, E., Hung, C.-H., Liu, Y.-H. & Wang, Y. (2002). *Helv. Chim. Acta*, **85**, 1517–1522.
- Lee, J. W., Jin, J. I., Achard, M. F. & Hardouin, F. (2001). *Liq. Cryst.* **28**, 663–671.
- Lutfor, M. R., Hegde, G., Pour, M. A., Yusoff, M. M. & Kumar, S. (2013a). *J. Fluorine Chem.* **156**, 230–235.
- Lutfor, M. R., Yusoff, M. M., Srinivasa, H. T., Samah, N. A., Malek, N. F. M. A. & Kumar, S. (2013b). *New J. Chem.* **37**, 2460–2467.
- Macrae, C. F., Bruno, I. J., Chisholm, J. A., Edgington, P. R., McCabe, P., Pidcock, E., Rodriguez-Monge, L., Taylor, R., van de Streek, J. & Wood, P. A. (2008). *J. Appl. Cryst.* **41**, 466–470.

- Paleos, C. M. & Tsiourvas, D. (2001). *Liq. Cryst.* **28**, 1127–1161.
- Rahman, M. L., Kwong, H. C., Mohd. Yusoff, M., Hegde, G., Mohamed Tahir, M. I. & Rahman, M. Z. A. (2012). *Acta Cryst.* **E68**, o2958.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.
- Takahashi, A., Mallia, V. A. & Tamaoki, N. (2003). *J. Mater. Chem.* **13**, 1582–1587.
- Yu, Q.-D. & Liu, Y.-Y. (2009). *Acta Cryst.* **E65**, o2326.

supporting information

Acta Cryst. (2014). E70, 499-502 [doi:10.1107/S1600536814023745]

Crystal structure of (*E*)-4-{2-[4-(allyloxy)phenyl]diazenyl}benzoic acid

Md. Lutfor Rahman, Mashitah Mohd. Yusoff, Jamil Ismail, Huey Chong Kwong and Ching Kheng Quah

Computing details

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINTE* (Bruker, 2009); data reduction: *SAINTE* (Bruker, 2009); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *Mercury* (Macrae *et al.*, 2008); software used to prepare material for publication: *SHELXTL* (Sheldrick, 2008) and *PLATON* (Spek, 2009).

(*E*)-4-{2-[4-(Allyloxy)phenyl]diazenyl}benzoic acid

Crystal data

C₁₆H₁₄N₂O₃

M_r = 282.29

Triclinic, *P* $\bar{1}$

Hall symbol: -P 1

a = 5.0279 (4) Å

b = 8.9678 (7) Å

c = 15.9913 (13) Å

α = 80.571 (2)°

β = 83.874 (2)°

γ = 88.371 (2)°

V = 707.19 (10) Å³

Z = 2

F(000) = 296

D_x = 1.326 Mg m⁻³

Melting point: 494 K

Mo *K* α radiation, λ = 0.71073 Å

Cell parameters from 3848 reflections

θ = 2.5–27.4°

μ = 0.09 mm⁻¹

T = 294 K

Plate, red

0.78 × 0.22 × 0.09 mm

Data collection

Bruker APEX DUO CCD area-detector diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

φ and ω scans

Absorption correction: multi-scan

(*SADABS*; Bruker, 2009)

T_{min} = 0.931, *T_{max}* = 0.992

12171 measured reflections

3276 independent reflections

2344 reflections with *I* > 2 σ (*I*)

R_{int} = 0.023

θ_{\max} = 27.6°, θ_{\min} = 1.3°

h = -6→6

k = -11→11

l = -20→20

Refinement

Refinement on *F*²

Least-squares matrix: full

R [*F*² > 2 σ (*F*²)] = 0.043

wR(*F*²) = 0.138

S = 1.04

3276 reflections

211 parameters

0 restraints

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$$w = 1/[\sigma^2(F_o^2) + (0.0696P)^2 + 0.0848P]$$

where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} < 0.001$

$$\Delta\rho_{\max} = 0.24 \text{ e } \text{\AA}^{-3}$$

$$\Delta\rho_{\min} = -0.16 \text{ e } \text{\AA}^{-3}$$

Special details

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
O1	1.26119 (18)	-0.00858 (10)	0.39744 (5)	0.0464 (3)	
O2	-0.234 (5)	0.8701 (17)	-0.0212 (8)	0.068 (3)	0.50
O3	-0.424 (5)	0.940 (2)	0.1072 (12)	0.0525 (17)	0.50
H3	-0.5052	1.0045	0.0769	0.079*	0.50
O2X	-0.392 (5)	0.943 (2)	0.0945 (13)	0.062 (3)	0.50
O3X	-0.267 (5)	0.8749 (16)	-0.0231 (8)	0.065 (3)	0.50
H3X	-0.3393	0.9554	-0.0404	0.098*	0.50
N1	0.6027 (2)	0.37818 (12)	0.20184 (7)	0.0442 (3)	
N2	0.4446 (2)	0.45802 (12)	0.24161 (7)	0.0444 (3)	
C1	1.6508 (4)	-0.33243 (19)	0.43299 (12)	0.0807 (6)	
H1A	1.5953	-0.3863	0.3931	0.097*	
H1B	1.7412	-0.3818	0.4775	0.097*	
C2	1.6019 (3)	-0.18848 (16)	0.42651 (9)	0.0538 (4)	
H2A	1.6599	-0.1378	0.4674	0.065*	
C3	1.4597 (3)	-0.09989 (14)	0.35815 (8)	0.0437 (3)	
H3A	1.5844	-0.0364	0.3182	0.052*	
H3B	1.3758	-0.1671	0.3273	0.052*	
C4	1.1092 (2)	0.08615 (13)	0.34590 (7)	0.0373 (3)	
C5	1.1259 (3)	0.09639 (14)	0.25801 (7)	0.0421 (3)	
H5A	1.2504	0.0384	0.2297	0.051*	
C6	0.9539 (3)	0.19454 (14)	0.21297 (8)	0.0438 (3)	
H6A	0.9626	0.2013	0.1541	0.053*	
C7	0.7698 (3)	0.28242 (13)	0.25406 (8)	0.0395 (3)	
C8	0.7558 (3)	0.27159 (14)	0.34268 (8)	0.0434 (3)	
H8A	0.6323	0.3302	0.3709	0.052*	
C9	0.9240 (3)	0.17465 (14)	0.38793 (8)	0.0436 (3)	
H9A	0.9147	0.1678	0.4468	0.052*	
C10	0.2754 (3)	0.55478 (13)	0.18998 (8)	0.0412 (3)	
C11	0.2795 (3)	0.56604 (17)	0.10240 (9)	0.0572 (4)	
H11A	0.3976	0.5065	0.0729	0.069*	
C12	0.1078 (3)	0.66589 (17)	0.05898 (9)	0.0579 (4)	
H12A	0.1103	0.6733	0.0002	0.069*	

C13	−0.0685 (3)	0.75530 (14)	0.10282 (8)	0.0427 (3)
C14	−0.0730 (3)	0.74195 (15)	0.19032 (8)	0.0463 (3)
H14A	−0.1914	0.8010	0.2201	0.056*
C15	0.0973 (3)	0.64144 (15)	0.23369 (8)	0.0455 (3)
H15A	0.0919	0.6322	0.2926	0.055*
C16	−0.2538 (3)	0.86292 (15)	0.05719 (8)	0.0461 (3)

Atomic displacement parameters (Å²)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0458 (5)	0.0547 (5)	0.0370 (4)	0.0192 (4)	−0.0066 (4)	−0.0037 (4)
O2	0.062 (4)	0.097 (5)	0.039 (3)	0.042 (3)	−0.008 (2)	0.009 (3)
O3	0.052 (3)	0.062 (3)	0.041 (5)	0.029 (2)	0.000 (3)	−0.009 (3)
O2X	0.067 (7)	0.076 (3)	0.041 (4)	0.039 (4)	−0.006 (4)	−0.009 (3)
O3X	0.071 (6)	0.079 (4)	0.045 (3)	0.045 (3)	−0.012 (2)	−0.012 (3)
N1	0.0475 (6)	0.0424 (6)	0.0412 (6)	0.0075 (5)	−0.0088 (5)	−0.0013 (4)
N2	0.0459 (6)	0.0442 (6)	0.0423 (6)	0.0092 (5)	−0.0089 (5)	−0.0032 (4)
C1	0.1015 (15)	0.0604 (10)	0.0731 (11)	0.0305 (10)	−0.0072 (10)	0.0038 (8)
C2	0.0522 (8)	0.0555 (8)	0.0526 (8)	0.0157 (7)	−0.0097 (6)	−0.0056 (6)
C3	0.0425 (7)	0.0438 (7)	0.0446 (6)	0.0098 (5)	−0.0055 (5)	−0.0081 (5)
C4	0.0359 (6)	0.0389 (6)	0.0359 (6)	0.0044 (5)	−0.0062 (5)	−0.0020 (5)
C5	0.0418 (7)	0.0462 (7)	0.0372 (6)	0.0100 (5)	−0.0021 (5)	−0.0067 (5)
C6	0.0482 (8)	0.0497 (7)	0.0324 (6)	0.0071 (6)	−0.0062 (5)	−0.0031 (5)
C7	0.0411 (7)	0.0372 (6)	0.0391 (6)	0.0038 (5)	−0.0081 (5)	−0.0011 (5)
C8	0.0464 (7)	0.0426 (6)	0.0410 (6)	0.0123 (5)	−0.0040 (5)	−0.0088 (5)
C9	0.0472 (7)	0.0497 (7)	0.0339 (6)	0.0105 (6)	−0.0060 (5)	−0.0072 (5)
C10	0.0414 (7)	0.0389 (6)	0.0425 (6)	0.0045 (5)	−0.0097 (5)	−0.0014 (5)
C11	0.0620 (9)	0.0640 (9)	0.0432 (7)	0.0305 (7)	−0.0043 (6)	−0.0079 (6)
C12	0.0668 (10)	0.0678 (9)	0.0360 (6)	0.0298 (8)	−0.0064 (6)	−0.0039 (6)
C13	0.0418 (7)	0.0426 (7)	0.0421 (6)	0.0094 (5)	−0.0067 (5)	−0.0028 (5)
C14	0.0465 (8)	0.0483 (7)	0.0448 (7)	0.0132 (6)	−0.0061 (6)	−0.0110 (5)
C15	0.0492 (8)	0.0498 (7)	0.0383 (6)	0.0073 (6)	−0.0093 (5)	−0.0076 (5)
C16	0.0440 (8)	0.0490 (7)	0.0436 (7)	0.0149 (6)	−0.0046 (6)	−0.0048 (6)

Geometric parameters (Å, °)

O1—C4	1.3613 (14)	C5—C6	1.3868 (17)
O1—C3	1.4317 (15)	C5—H5A	0.9300
O2—C16	1.238 (13)	C6—C7	1.3812 (18)
O3—C16	1.36 (2)	C6—H6A	0.9300
O3—H3	0.8200	C7—C8	1.3989 (17)
O2X—C16	1.18 (2)	C8—C9	1.3700 (18)
O3X—C16	1.280 (14)	C8—H8A	0.9300
O3X—H3X	0.8200	C9—H9A	0.9300
N1—N2	1.2481 (16)	C10—C15	1.3781 (18)
N1—C7	1.4195 (16)	C10—C11	1.3855 (19)
N2—C10	1.4254 (16)	C11—C12	1.3817 (19)
C1—C2	1.297 (2)	C11—H11A	0.9300

C1—H1A	0.9300	C12—C13	1.3899 (18)
C1—H1B	0.9300	C12—H12A	0.9300
C2—C3	1.4790 (19)	C13—C14	1.3825 (18)
C2—H2A	0.9300	C13—C16	1.4813 (18)
C3—H3A	0.9700	C14—C15	1.3801 (18)
C3—H3B	0.9700	C14—H14A	0.9300
C4—C5	1.3870 (16)	C15—H15A	0.9300
C4—C9	1.3956 (17)		
C4—O1—C3	117.90 (9)	C8—C9—C4	120.20 (11)
C16—O3—H3	109.5	C8—C9—H9A	119.9
C16—O3X—H3X	109.5	C4—C9—H9A	119.9
N2—N1—C7	114.00 (10)	C15—C10—C11	119.92 (12)
N1—N2—C10	114.63 (10)	C15—C10—N2	114.92 (11)
C2—C1—H1A	120.0	C11—C10—N2	125.15 (12)
C2—C1—H1B	120.0	C12—C11—C10	119.87 (12)
H1A—C1—H1B	120.0	C12—C11—H11A	120.1
C1—C2—C3	124.08 (15)	C10—C11—H11A	120.1
C1—C2—H2A	118.0	C11—C12—C13	120.24 (12)
C3—C2—H2A	118.0	C11—C12—H12A	119.9
O1—C3—C2	107.64 (10)	C13—C12—H12A	119.9
O1—C3—H3A	110.2	C14—C13—C12	119.40 (12)
C2—C3—H3A	110.2	C14—C13—C16	119.70 (11)
O1—C3—H3B	110.2	C12—C13—C16	120.89 (11)
C2—C3—H3B	110.2	C15—C14—C13	120.30 (12)
H3A—C3—H3B	108.5	C15—C14—H14A	119.9
O1—C4—C5	124.57 (11)	C13—C14—H14A	119.9
O1—C4—C9	115.09 (10)	C10—C15—C14	120.24 (12)
C5—C4—C9	120.33 (11)	C10—C15—H15A	119.9
C6—C5—C4	118.92 (11)	C14—C15—H15A	119.9
C6—C5—H5A	120.5	O2X—C16—O2	123.9 (14)
C4—C5—H5A	120.5	O2X—C16—O3X	117.6 (15)
C7—C6—C5	121.13 (11)	O2—C16—O3X	8 (2)
C7—C6—H6A	119.4	O2X—C16—O3	7.0 (19)
C5—C6—H6A	119.4	O2—C16—O3	128.6 (14)
C6—C7—C8	119.40 (11)	O3X—C16—O3	121.9 (13)
C6—C7—N1	116.43 (11)	O2X—C16—C13	120.1 (11)
C8—C7—N1	124.16 (11)	O2—C16—C13	115.8 (10)
C9—C8—C7	120.02 (11)	O3X—C16—C13	122.3 (9)
C9—C8—H8A	120.0	O3—C16—C13	115.6 (9)
C7—C8—H8A	120.0		
C7—N1—N2—C10	-179.99 (10)	C15—C10—C11—C12	1.0 (2)
C4—O1—C3—C2	178.33 (11)	N2—C10—C11—C12	-179.04 (13)
C1—C2—C3—O1	132.68 (17)	C10—C11—C12—C13	0.1 (3)
C3—O1—C4—C5	2.60 (19)	C11—C12—C13—C14	-0.9 (2)
C3—O1—C4—C9	-178.51 (11)	C11—C12—C13—C16	179.98 (14)
O1—C4—C5—C6	178.13 (11)	C12—C13—C14—C15	0.4 (2)

C9—C4—C5—C6	-0.70 (19)	C16—C13—C14—C15	179.59 (12)
C4—C5—C6—C7	0.6 (2)	C11—C10—C15—C14	-1.5 (2)
C5—C6—C7—C8	-0.3 (2)	N2—C10—C15—C14	178.60 (11)
C5—C6—C7—N1	-179.63 (11)	C13—C14—C15—C10	0.7 (2)
N2—N1—C7—C6	-178.28 (11)	C14—C13—C16—O2X	4.5 (14)
N2—N1—C7—C8	2.47 (19)	C12—C13—C16—O2X	-176.3 (14)
C6—C7—C8—C9	0.1 (2)	C14—C13—C16—O2	178.8 (10)
N1—C7—C8—C9	179.33 (12)	C12—C13—C16—O2	-2.0 (11)
C7—C8—C9—C4	-0.2 (2)	C14—C13—C16—O3X	-176.7 (11)
O1—C4—C9—C8	-178.46 (11)	C12—C13—C16—O3X	2.5 (11)
C5—C4—C9—C8	0.5 (2)	C14—C13—C16—O3	-1.5 (10)
N1—N2—C10—C15	178.98 (11)	C12—C13—C16—O3	177.7 (10)
N1—N2—C10—C11	-1.0 (2)		

Hydrogen-bond geometry (Å, °)

Cg1 is the centroid of the C4—C9 ring.

<i>D</i> —H... <i>A</i>	<i>D</i> —H	H... <i>A</i>	<i>D</i> ... <i>A</i>	<i>D</i> —H... <i>A</i>
O3—H3...O2 ⁱ	0.82	1.90	2.71 (3)	166
C6—H6 <i>A</i> ...O2 ⁱⁱ	0.93	2.59	3.367 (15)	145
C3—H3 <i>A</i> ...Cg1 ⁱⁱⁱ	0.97	2.66	3.504 (2)	145

Symmetry codes: (i) $-x-1, -y+2, -z$; (ii) $-x+1, -y+1, -z$; (iii) $x+1, y, z$.