

**THE INFLUENCE OF PHYSICAL BARRIERS TO  
FISH ASSEMBLAGE ALONG SUNGAI ENAM IN  
TEMENGOR RESERVOIR, PERAK**

**NURUL FAZLINDA BINTI MOHD FADZIL**

**UNIVERSITI SAINS MALAYSIA**

**2016**

**THE INFLUENCE OF PHYSICAL BARRIERS TO  
FISH ASSEMBLAGE ALONG SUNGAI ENAM IN  
TEMENGOR RESERVOIR, PERAK**

by

**NURUL FAZLINDA BINTI MOHD FADZIL**

**Thesis submitted in fulfillment of the requirements  
for the degree of  
Master of Science**

**December 2016**

## ACKNOWLEDGEMENT

In the name of Allah, the Most Gracious and the Most Merciful

Alhamdulillah, praise to Allah, the Almighty, for the strengths and guidance in completing this study. I wish to express my gratitude to my supervisor, Dr. Zarul Hazrin Hashim, for his guidance and constant support. I have been extremely lucky to have a supervisor who cared so much about my work. His invaluable help of constructive comments and suggestion throughout the study helped me to complete this thesis writing successfully.

My sincere thanks goes to these people, Amirul, Abg Mijan, Abg Syaiful, Muneera, Kak Muzza, Kak Syikin, Kak Iza, Farah, Abg Shafiq, Athira and others who had helped me directly or indirectly, for their kindness and moral support during my study.

Not to forget for funding and financial support under research grants (RUI: 1001/PBIOLOGI/811248, FRGS: 203/PBIOLOGI/6711358, USM Short-Term: 304/PBIOLOGI/6312132) throughout the study.

Last but not least, my deepest gratitude goes to my beloved parents, Mohd Fadzil Kassim and Roshadah Said for allowing me to realize my own potential. Thank you both for giving me strength to chase my dreams. All the support they have provided me over the years was the greatest gift anyone has ever given me. To my siblings, thank you for your support, prayers and endless love. Sweet thank is also for my nephews for their puerile support.

*'Remember, a dead fish can float downstream, but it takes a life one to swim upstream'.*

W. C. Fields

## TABLE OF CONTENTS

<b>ACKNOWLEDGEMENT</b>	ii
<b>TABLE OF CONTENTS</b>	iii
<b>LIST OF TABLES</b>	vii
<b>LIST OF FIGURES</b>	ix
<b>LIST OF PLATES</b>	xi
<b>LIST OF APPENDICES</b>	xii
<b>LIST OF SYMBOLS AND ABBREVIATIONS</b>	xiii
<b>ABSTRAK</b>	xiv
<b>ABSTRACT</b>	xvi
<b>CHAPTER 1: INTRODUCTION</b>	
1.1 Background	1
1.2 Problem statement	3
1.3 Significance of study	4
1.4 Limitations of study	5
1.5 Objectives	6
<b>CHAPTER 2 : LITERATURE REVIEW</b>	
2.1 Running water	7
2.1.1 River and stream	7
2.1.2 River Continuum Concept	9
2.1.3 Malaysian river system with special emphasis on Sungai Enam	10
2.2 Assessment of freshwater fishes in streams	10
2.2.1 Fish assemblages in streams	11
2.2.2 Previous studies of fish assemblages in the Temengor Reservoir	11
2.2.3 Electrofishing	12
2.2.4 Altitude as a key factor limiting fish distribution in streams	14
2.3 Assessment of the impacts of barriers on fish species	15
2.3.1 The influence of man-made barriers to fish species	15
2.3.2 The influence of natural barriers to fish species	17

2.4	Assessment of fish condition on the basis of length-weight relationships and relative condition factor	18
2.4.1	Length-weight relationships (LWR)	18
2.4.2	Relative condition factor ( $K_n$ )	19
2.5	Fish movement over barriers	20
2.5.1	Importance of fish movement	20
2.5.2	Previous studies of fish movement against barriers	21
2.5.3	Tagging or marking methods in detecting fish movement	24
2.5.4	Fins regeneration	30
2.5.5	Ability of fish to survive in extreme environment	32
2.6	Environmental variables that affect fish assemblages in streams	33
2.6.1	<i>In-situ</i> parameters	34
2.6.2	Nutrients	35

### **CHAPTER 3: MATERIALS AND METHODS**

3.1	Study area	39
3.1.1	Sungai Enam	39
3.1.2	Sungai Kooi	39
3.2	General features of Sungai Enam	41
3.3	Sampling design	41
3.3.1	Fish collection and identification	48
3.3.2	Fish measurements and assessment of fish condition	48
3.3.3	Fish marking	50
3.3.4	Short experimental justification on fish marking with fin clipping	52
3.3.5	<i>In-situ</i> parameters	52
3.3.6	Nutrients	52
3.4	Data and statistical analysis	53
3.5	An intensive study of Sungai Kooi	60
3.5.1	General features of Sungai Kooi	60
3.5.2	Sampling design	60
3.5.3	Fish collection and identification	63
3.5.4	Data analysis	63

## CHAPTER 4: RESULTS

4.1	To determine fish assemblages and their similarities among reaches in Sungai Enam, primarily between above and below the barriers	64
4.1.1	Fish checklist and distribution by reaches	64
4.1.2	Fish distribution based on elevation	71
4.1.3	Families composition of fish species	73
4.1.4	Cluster analysis	75
4.2	To determine and to compare the relative condition factor ( $K_n$ ) of selected fish species among reaches in Sungai Enam, primarily between above and below the barriers	77
4.2.1	Length-weight relationship	77
4.2.2	Relative condition factor ( $K_n$ )	77
4.3	To determine fish movement along reaches in Sungai Enam, primarily between above and below the barriers	79
4.3.1	Fish movements along reaches by sampling excursions	79
4.3.2	Percentage of fish recovery	88
4.3.3	Fins regeneration experiment	90
4.4	To correlate environmental variables with fish parameters in Sungai Enam	92
4.5	Intensive study of Sungai Kooi	95
4.5.1	Fish checklist and distribution by reaches	95
4.5.2	Fish distribution based on elevation	95
4.5.3	Families composition of fish species	98

## CHAPTER 5: DISCUSSIONS

5.1	Fish assemblages in Sungai Enam	100
5.1.1	Longitudinal distribution of fish in altitudinal variation stream	105
5.1.2	The influence of physical barriers to fish distribution	107
5.2	Fish condition based on length-weight relationship and relative condition factor of <i>P. smedleyi</i>	110
5.2.1	Length-weight relationship (LWR)	110
5.2.2	Relative condition factor ( $K_n$ )	112
5.3	Fish movements along reaches in Sungai Enam	114

5.3.1	Longitudinal movements along Sungai Enam	114
5.3.2	Fin clipping as a selected marking method	119
5.4	Correlation of environmental variables to fish parameters in Sungai Enam	121
5.4.1	Correlation between <i>in-situ</i> parameters and fish parameters	122
5.4.2	Correlation between nutrients and fish parameters	125
<b>CHAPTER 6: CONCLUSIONS AND RECOMMENDATIONS</b>		
6.1	General conclusion	129
6.2	Recommendations	130
<b>REFERENCES</b>		131
<b>APPENDICES</b>		
<b>LIST OF PUBLICATIONS</b>		

## LIST OF TABLES

	<b>Page</b>
Table 2.1 Advantages and disadvantages of electrofishing	13
Table 2.2 Physical and biological impacts of dams	16
Table 2.3 List of fish which are reported able to move upstream and present above barriers	22
Table 2.4 Short description of fish tags or marks including their benefits and drawbacks	26
Table 2.5 Positive and negative effects of environmental variables on freshwater fishes	36
Table 3.1 General characteristics of the nine reaches in Sungai Enam with GPS coordinates	47
Table 3.2 Types of fin that being clipped during this study	51
Table 3.3 Strength of correlation coefficient	59
Table 3.4 General characteristics of reaches with coordinates in Sungai Kooi	62
Table 4.1 Checklist and distribution of fish species along Sungai Enam based on data collected from October 2014 to June 2015 using a backpack of electroshocker	65
Table 4.2 Homogenous subset of the number of fish species among the nine reaches.	68
Table 4.3 Homogenous subset of the number of fish species between above and below the waterfalls in Sungai Enam.	68
Table 4.4 Species number, Shannon-Wiener Diversity Index and Evenness Index of fish in three areas of Sungai Enam based on data collected from October 2014 to June 2015 using a backpack electroshocker.	70
Table 4.5 Number of recaptured fish, marked fish, downstream and upstream movements, and marked fish that was recorded in the same reach by sampling excursions along Sungai Enam from October 2014 to June 2015 using a backpack electroshocker.	80



Table 4.6	Percentage of fish recovery (%) calculation based on total marked fish in each sampling excursions along Sungai Enam based on data collected from October 2014 to June 2015 using a backpack electroshocker.	89
Table 4.7	Correlation between environmental variables and fish parameters along Sungai Enam based on data collected from October 2014 to June 2015 using a backpack electroshocker and YSI 556 MPS.	93
Table 4.8	Checklist and distribution of fish species along Sungai Kooi based on data collected on one-day intensive sampling excursion (February 2016) using a backpack electroshocker.	96

## LIST OF FIGURES

		<b>Page</b>
Figure 2.1	Chronology on the studies of river and stream ecosystem	8
Figure 2.2	Considerations that have to be noted before the applying the tagging / marking methods	29
Figure 2.3	Caudal fin regeneration of zebrafish	31
Figure 3.1	Location of Sungai Enam and Sungai Kooi in Temengor Reservoir, Perak	40
Figure 3.2	Sketch diagrams of the nine reaches in Sungai Enam	43
Figure 3.3	Equipment and steps in measuring fish length	49
Figure 3.4	Measurement of fish length	49
Figure 3.5	Fin clipping method	51
Figure 3.6	Sketch diagrams of the two reaches (above and below the waterfall) in Sungai Kooi	61
Figure 4.1	Number of fish species recorded among the nine reaches along Sungai Enam based on data collected from October 2014 to June 2015 using a backpack electroshocker	66
Figure 4.2	The presence of fish species, according to elevation in the nine reaches along Sungai Enam based on data collected from October 2014 to June 2015 using a backpack electroshocker	72
Figure 4.3	Percentage of fish species by family based on data collected in Sungai Enam from October 2014 to June 2015 using a backpack electroshocker	74
Figure 4.4	Dendrogram of Jaccard's Similarity Coefficient (JSC) based on fish assemblages among the nine reaches along Sungai Enam from October 2014 to June 2015 using a backpack electroshocker	76
Figure 4.5	The power function of LWR of <i>P. smedleyi</i> based on data collected in the nine reaches along Sungai Enam from October 2014 to June 2015 using a backpack electroshocker	78

Figure 4.6	Average $\pm$ standard error of relative condition factor ( <i>Kn</i> ) of <i>P. smedleyi</i> by reaches based on data collected in the nine reaches along Sungai Enam from October 2014 to June 2015 using a backpack electroshocker	78
Figure 4.7	Number of individuals that had moved downstream recorded along Sungai Enam from October 2014 to June 2015 using a backpack electroshocker	81
Figure 4.8	Number of individuals that had moved upstream recorded along Sungai Enam from October 2014 to June 2015 using a backpack electroshocker	81
Figure 4.9	Fish movement along the reaches in Sungai Enam by sampling excursions based on data collected in the nine reaches along Sungai Enam from October 2014 to June 2015 using a backpack electroshocker	83
Figure 4.10	Percentage of fish recovery (%) based on total marked fish in each sampling excursions along Sungai Enam based on data collected from October 2014 to June 2015 using a backpack electroshocker	89
Figure 4.11	The length of regenerated fin of <i>P. smedleyi</i> in a control tank (Tank A) and two treatment tanks (Tank B and Tank C) that reared for one month in captivity	91
Figure 4.12	The presence of fish species, according to the elevation of Sungai Kooi based on one-day sampling excursion (February 2016) using a backpack electroshocker	97
Figure 4.13	Percentage of fish species by family captured in Sungai Kooi based on one-day intensive sampling excursion (February 2016) using a backpack electroshocker	99

## LIST OF PLATES

	<b>Page</b>
Plate 3.1 Physical barriers in Sungai Enam	42
Plate 3.2 The nine reaches that were selected in Sungai Enam	44
Plate 3.3 Physical barrier in Sungai Kooi	61

## LIST OF APPENDICES

- Appendix A Family, scientific and local name of fish species captured in Sungai Enam during study period
- Appendix B Fish checklist of Sungai Enam, Temengor Reservoir, Perak, Malaysia from 1994 to 2014
- Appendix C Missing and new recorded species observed for Sungai Enam in 2014 compared to previous studies from 1994 to 2013
- Appendix D Abandoned chalets in Sungai Enam
- Appendix E Normality test of the number of fish species in Sungai Enam
- Appendix F One-way ANOVA with permutation of the number of fish species among reaches in Sungai Enam
- Appendix G One-way ANOVA with permutation of the number of fish species between above and below the waterfalls of Enam
- Appendix H One-way ANOVA with permutation of the number of fish species between above and below the dams of Sungai Enam
- Appendix I Normality test of relative condition factor of *P. smedleyi*
- Appendix J One-way ANOVA with permutation of relative condition factor ( $Kn$ ) of *P. smedleyi* among reaches in Sungai Enam
- Appendix K Normality test of environmental variables in Sungai Enam
- Appendix L One-way ANOVA with permutation of *in-situ* parameters and nutrients among reaches in Sungai Enam
- Appendix M Correlation between environmental variables and the number of fish species
- Appendix N Correlation between environmental variables and average of relative condition factor ( $Kn$ ) of *P. smedleyi*
- Appendix O Correlation between environmental variables and fish movements
- Appendix P Family, scientific and local name of fish species captured in Sungai Kooi during study period
- Appendix Q Equipments used during the study period

## LIST OF SYMBOLS AND ABBREVIATION

ANOVA	Analysis Of Variance
cm	Centimeter
DO	Dissolved Oxygen
g	Gram
GIS	Geographical Information System
GPS	Global Positioning System
$H'$	Shannon-Wiener Diversity Index
$J'$	Pielou's Evenness Index
$Kn$	Relative Condition Factor
L	Litre
LWR	Length-Weight Relationship
mg	Miligram
MVSP	Multivariate Statistical Package
NH <sub>3</sub> -N	Ammoniacal-Nitrogen
NO <sub>2</sub> -N	Nitrite-Nitrogen
NO <sub>3</sub> -N	Nitrate-Nitrogen
PO <sub>4</sub> -P	Ortho-Phosphate
$r$	Pearson Correlation Coefficient
$R^2$	Regression Coefficient
RCC	River Continuum Concept
s	Second
SE	Standard Error
SL	Standard Length
SPSS	Statistical Package For Social Science
TDS	Total Dissolved Solids
TL	Total Length
UPGMA	Unweighted Pair Group Method
W	Weight
$\mu$ S	Micro Siemens
$b$	Growth Coefficient

# **PENGARUH HALANGAN FIZIKAL TERHADAP PERKUMPULAN IKAN SEPANJANG SUNGAI ENAM DI TAKUNGAN TEMENGOR, PERAK**

## **ABSTRAK**

Satu kajian mengenai pengaruh halangan fizikal terhadap perkumpulan ikan (bilangan spesis ikan, relatif faktor keadaan ikan dan pergerakan ikan) telah dijalankan dari Oktober 2014 hingga Jun 2015 di sembilan ruas sepanjang Sungai Enam di Takungan Temengor, Perak. Sembilan ruas tersebut termasuk dua halangan semula jadi (air terjun) dan dua halangan buatan manusia (empangan kecil). Persampelan telah dijalankan sebulan sekali dengan memilih satu kolam secara rawak dari setiap ruas (kaedah persampelan rawak berstrata). Ikan ditangkap menggunakan beg gelas pengejut ikan (Smith-Root LR-24 Electrofisher) dengan jaring. Selepas pengecaman ikan dan pengukuran, sirip ikan yang tertentu dipotong berdasarkan ruas sebagai kaedah penandaan untuk menentukan pergerakan mereka di sepanjang sungai. Hubungan antara pembolehubah persekitaran dan ketiga-tiga parameter ikan tersebut juga telah dinilai. Hasil kajian menunjukkan bahawa 13 spesis ikan (3,643 spesimen ikan) yang terdiri daripada enam famili telah dikenal pasti di sepanjang sembilan ruas di sungai ini. Kajian ini juga merekodkan *P. smedleyi* sebagai spesis ikan yang paling kerap dan banyak ditemui di sembilan ruas. *Mastacembelus favus* telah direkodkan buat kali pertama di sungai ini. Bilangan spesis ikan meningkat dengan ketara dari hulu ke hilir sungai. Ikan telah ditemui di kawasan atas air terjun yang tinggi di sungai ini. Berdasarkan pada 515 spesimen ikan yang ditangkap semula selepas proses penandaan, pergerakan enam spesis (28 spesimen) ke hulu sungai dan pergerakan lima spesis (39 spesimen ikan) ke hilir sungai telah direkodkan dan *Poropuntius smedleyi* merupakan spesis ikan yang

paling banyak bergerak di sepanjang sungai ini. Relatif faktor keadaan ( $K_n$ ) *P. smedleyi* menunjukkan bahwa kehadiran halangan fizikal tidak mempengaruhi keadaan *P. smedleyi* di sepanjang sungai ini. Hubungan antara pembolehubah persekitaran dan parameter ikan adalah lemah. Kajian ini telah mengesan bahawa air terjun yang mempunyai tahap kecerunan yang tinggi lebih cenderung mempengaruhi bilangan spesis ikan berbanding dengan air terjun yang mempunyai tahap kecerunan yang rendah. Manakala, halangan buatan manusia yang terdapat di sungai ini tidak memberikan pengaruh yang bererti terhadap bilangan spesis ikan. Walau bagaimanapun, halangan fizikal di Sungai Enam tidak memberikan pengaruh terhadap relatif faktor keadaan *P. smedleyi* dan pergerakan ikan di sepanjang sungai ini.



**THE INFLUENCE OF PHYSICAL BARRIERS TO FISH ASSEMBLAGE  
ALONG SUNGAI ENAM IN TEMENGOR RESERVOIR, PERAK**

**ABSTRACT**

A study on the influence of physical barriers to fish assemblage (i.e., the number of fish species, relative condition factor of fish and fish movements) was carried out from October 2014 to June 2015 at nine reaches along Sungai Enam in Temengor Reservoir, Perak. The nine reaches include two natural barriers (i.e., waterfalls) and two man-made barriers (i.e., small dams). Sampling excursions were carried out once a month by selecting one pool randomly from each reach (stratified random sampling method). Fish were captured using a backpack electrofishing (Smith-Root LR-24 Electrofisher) with scoop nets. After fish identification and measurements, selected fish fin were clipped according to reach as a marking method to determine their movement along the stream. Relationships between environmental variables and the three fish parameters were also assessed. Results showed that 13 fish species (3,643 fish specimens) comprising of six families were identified along the nine reaches of this stream. This study also recorded *P. smedleyi* as frequently and greatest fish species found in nine reaches. *Mastacembelus favus* was recorded for the first time in this stream. The number of fish species significantly increased from upstream to downstream. Fish were present above high waterfalls in this stream. Based on 515 fish specimens that were recaptured after marking process, upstream movement by six species (28 specimens) and downstream movement by five species (39 specimens) were recorded and *Poropuntius smedleyi* was the most travelled fish species along this stream. Relative condition factor ( $K_n$ ) of *P. smedleyi* indicated that the presence of physical barriers has no influence towards the

condition of *P. smedleyi* along this stream. The relationships between environmental variables and fish parameters were weak. This study detected that a waterfall with greatest slope is more likely to influence the number of fish species compared to the waterfall with least slope. Meanwhile, man-made barriers that present in this stream did not have significant influence to the number of fish species. However, physical barriers in Sungai Enam have no influence on relative condition factor of *P. smedleyi* and fish movements along the stream.

# CHAPTER I

## INTRODUCTION

### 1.1 Background

Malaysia is enriched with natural beauty and resources which are to be proud of. One of the attractions that are well-known in Malaysia is its waterfalls. Waterfalls attract people by their beautiful landscapes and refreshing air, since these area possess greater concentration of negative ions which increase serotonin level (mood chemical) that helps to reduce depression, stress and improve daytime energy (Mann & Nazario, 2002). Besides attraction, waterfall is also one of the natural habitats for aquatic biodiversity. Many endangered and rare species could be found in such secluded habitat compared to other aquatic habitats located near human hotspots, because most of the waterfalls are located deeply in the forest which ensures and secures the presence of those species.

Waterfall is a watercourse that is created when the water falls from high elevation over a cliff or rock. This eventually erodes the river bed and forms a pool. Formation of waterfall often occurs at the upper reach of a river where it flows over different band of rocks (Corrigan, 2007). The presence of waterfall in the river network affects stream communities (Gilliam *et al.*, 1993; Mazzoni *et al.*, 2006), changes the prey-predator dynamics (Reznick & Ghalambor, 2005), and longitudinal abundance of species (Greathouse & Pringle, 2006).

Waterfalls are categorised as natural barriers in a river system. This natural barrier is a dominant factor in determining the composition of regional faunas and in promoting endemism (Ricklefs & Schluter, 1993; Cox & Moore, 2000; Rahel, 2007).

The dominance and abundance of species in a fish assemblage could change with the changes of river gradient. Waterfall which is called as a bio-geographic barrier is able to prevent species exchange among regions and disrupts the movement of an aquatic organism (Rahel, 2007). For example, waterfalls in the Snake River (Idaho, USA) are restricted to certain fish species in upstream colonisation (McPhail & Lindsey, 1986).

Sungai Enam is one of the prominent headwaters that flow into Temengor Reservoir. Several studies have been conducted by ecologists in Sungai Enam such as fish fauna by Zakaria-Ismail and Sabariah (1995), Mohd- Akhir (1999), Berryhill-Jr. and Ahyaudin (2002), Hashim *et al.* (2012), Ismail *et al.* (2013) and Mohd Shafiq *et al.* (2013), hydrology and water quality by Hashim *et al.* (2011) and Wan Ruslan *et al.* (2013), flowering plants by Siti Munirah *et al.* (2013), lianas by Zakaria *et al.* (2013), macrofungal by Kin (2013) and mammals by Shahfiz *et al.* (2013) which proved that Sungai Enam solely able to possess great number of biodiversity and is a crucial headwater for research in the reservoir.

Nevertheless, these studies were only carried out at certain elevation within a short time frame. Therefore, no studies have yet been conducted on the influence of physical barriers to fish assemblages especially in the reaches above and below the waterfalls along Sungai Enam. Since the understanding of the influence of waterfalls to freshwater fish assemblages, especially in the tropical regions are still incomplete, Sungai Enam was selected to determine the influence of physical barriers to fish assemblage in this stream. At the same time, an experiment was carried out to observe fin regeneration.

## 1.2 Problem statement

Some definitive evidence showed that diversity and distribution of freshwater fish in Malaysia has been disrupted rapidly due to the developmental pressure and modification of fish habitat (Zakaria-Ismail, 1991; Bowen *et al.*, 2003; Chong *et al.*, 2010). One of the reasons that jeopardize freshwater fish diversity and its distribution in Malaysia is through habitat modification of dam constructions or any barriers that possibly will limit fish movement from downstream to upstream reaches or vice versa. The presence of barriers, either man-made or natural, will affect the fish assemblage in the fluvial ecosystem.

Waterfalls may disrupt the stream connectivity and influence the spatial distribution of freshwater fish. Fast-flowing water washes away fish species from upstream to downstream and prevents them from moving upstream as they have no adaptations either in the aspect of morphology or behaviour. It is clearly observed that only a few fish species can survive above the waterfall and move across the barrier. Therefore, the species, which are robust and tolerant against such physical and geographical obstacles hardly face biological competition and predation risks.

It is evident that the fish movement varies between taxa and life-history stages (Baras & Lucas, 2001; Crook, 2004). But complications and difficulties are common phenomenon in monitoring fish movement due to picking out the unfavourable type of tagging/marketing method. Some tags or marks such as external tags (e.g., T-bar tag) (Collins *et al.*, 1994), external marks (e.g., dyes) (Dussault & Rodríguez, 1997), internal tags (e.g., PIT tags) (Zydlewski *et al.*, 2006), internal marks (e.g., alizarine) (Simon *et al.*, 2009) and visible implant elastomer (VIE) tags (Younk *et al.*, 2010; Kozłowski *et al.*, 2016), have certain limitations to certain number of species as well as individuals and are only applicable to large sizes of fish

species. To reduce bias, tagging/marking techniques should be applied to a wide range of size. Therefore, fin clipping was selected and applied in this study because of its easiness and simplicity. Moreover, this marking method can be used to different types of fin. However, this type of marking method is considered harmful by a few scientists due to the lack of knowledge and practical understanding about the fin regeneration process (Akimenko *et al.*, 2003).

### **1.3 Significance of study**

Fish require unimpeded access along waterways to survive and reproduce. Both types of fishes, either fresh or saltwater, move within the water at different times to access food and shelter, to avoid predators, and to seek mates. Some fish species have to migrate at least once as a part of their life cycle. However, the presence of physical barriers such as dams and waterfalls can disrupt the continuity of river or stream. At the same time, these barriers may limit fish movements, resulting in distinct assemblages on both side of the barriers (Barbosa *et al.*, 2015).

Studies of spatiotemporal patterns of fish diversity, distribution, and composition are beneficial to determine the factors that influence the structure of fish community (Belliard *et al.*, 1997; Galacatos *et al.*, 2004; Khairul Adha *et al.*, 2009). Fish distribution in the habitats is influenced by various factors such as the availability of food, breeding sites, water velocity, depth, topography and physico-chemical properties of water (Harris, 1995).

Disclosing fish-habitat interaction in a habitat is also crucial to understand fish adaptations. Besides, by studying the fish that live in altitudinal variation habitats could provide new insight into how fishes respond to environmental gradients and directly help on freshwater conservation efforts. However, high-

altitude streams, such as above the waterfall is one of the habitats that have depauperate fish faunas of less than 15 species (Samat, 1990; Martin-Smith & Laird, 1998). These depauperate areas are hardly investigated by ecologists due to their physical isolation (Matthews, 1998).

To identify the location of any impacts arising in the future, information on upstream progress rates, and longitudinal changes in species movements is important (Bishop *et al.*, 1995). Observation towards the formation of groups between the waterfalls is necessary to confirm that waterfalls are able to produce dissimilar fish community resulting from different capability of fish to move against it.

#### **1.4 Limitations of study**

In this study, there were some limitations that may have affected the outcomes of this study. The first limitation is the sampling gears. In this study, only one active gear, electrofishing, was used to capture the fish. The combination of sampling gears either in passive or active gears, are generally able to collect more fish species because of the gear selectivity. Some species like to swim and forage on the surface of water (e.g., danios and rasboras), whereas some species like to swim and forage at the river bed (e.g., catfishes and eels). However, in this study, backpack electrofisher was the only suitable gear, due to shallow and narrow stream reaches.

The second limitation was the lack of manpower for such hard sampling excursions. In the field, sampling activities such as fish measurements and fish markings are needed to be carried out as fast as possible to reduce fish mortality. Along with hiking and other related procedures, the captured fish could be released earlier if there were enough manpower. In such case, a lot of manpower are needed as it is very difficult to manage with only three or four people.

## 1.5 Objectives

The objectives of this study can be summarized as:

- a) To determine fish assemblages and their similarities among reaches in Sungai Enam, primarily between above and below the barriers.
- b) To determine and to compare the relative condition factor of selected fish species among reaches in Sungai Enam, primarily between above and below the barriers.
- c) To determine fish movement along reaches in Sungai Enam, primarily between above and below the barriers.
- d) To correlate environmental variables with fish parameters in Sungai Enam.



## CHAPTER II

### LITERATURE REVIEW

#### 2.1 Running water

Running waters range from a small stream to large rivers (Allan, 2012). Running waters harbour a diverse and unique collection of species, habitats, and ecosystems, including some of the most threatened species and ecosystems (Allan & Flecker, 1993). However, nowadays, running waters are perhaps the most impacted ecosystems on the planet (Malmqvist & Rundle, 2002; Aura, 2008) due to human settlements and heavy exploitation for water supplies, irrigation, electricity, and waste disposal.

##### 2.1.1 River and stream

Rivers are mostly understood as lotic habitats where the water continuously flow. A river begins at a source (or most often several sources) and ends at a river mouth. Rivers and streams are always being misidentified due to their similar characteristics. Even though both are fast moving water bodies, but in general, a river is larger, deeper, and longer than a stream and is also formed from a collection of many streams (Chiotti, 2004). On the other hand, a stream is much shallower than a river, which sometimes can even be walked across.

Studies on rivers and streams started in Europe (Cushing *et al.*, 2006). Early studies were concerned with the distribution, abundance, and taxonomic composition of lotic organisms. Then, these studies advanced with the development of technology and expertise, as summarized in Figure 2.1.




Figure 2.1: Chronology on the studies of river and stream ecosystem (adapted from Cushing *et al.*, 2006)

Generally, rivers and streams are classified by their size according to the Strahler Stream Order (Strahler, 1957). A first-order stream is usually the smallest stream with round flow and no tributaries. Second-order streams are formed when two first-order streams flow into each other. Third order-streams are formed when two second-order streams flow into each other. A river starts to form when all streams are interconnected; this is also known as a fourth-order stream (Chiotti, 2004).

### **2.1.2 River Continuum Concept**

River Continuum Concept (RCC), is a framework that has been proposed by Vannote *et al.* (1980), that is important for the study of running waters. RCC, which has been tested in many countries, is able to describe the function of lotic ecosystems from the source to the river mouth (Allan, 2012). In this concept, geomorphological and hydrological performance changes as a stream order changed (Allan & Castillo, 2007). Allochthonous material was mentioned to be more abundant in the headwaters, whereas autochthonous input plays a greater role in the mid reaches.

According to Johnson *et al.* (1995), River Continuum Concept (RCC) was debated with different views regarding to several issues such as border issue, stochastic issue, human disturbance issue and spatio-temporal variation issue. To make it more realistic, several other concepts have been integrated into the RCC, including on the influence of riparian zone (Cummins *et al.*, 1989) and the Flood Pulse Concept (Junk *et al.*, 1989).

Flood Pulse Concept highlighted the importance of flood pulse, which is responsible for the existence, productivity, and interactions between biota in river and the floodplain systems. This concept was assumed as one of the most innovative concepts in order to adjust some weaknesses in the RCC. However, Ta Fang (2010)

concluded that the application of RCC is still valid in order to provide an important model for proper management in riverine functional enhancement.

### **2.1.3 Malaysian river system with special emphasis on Sungai Enam**

Malaysia, a tropical country, has a diverse collection of rivers and tributaries. Out of 150 major rivers, 100 are located in Peninsular Malaysia and another 50 are located in Sabah and Sarawak (Hashim *et al.*, 2011). Sungai Rajang with 51,000 km<sup>2</sup> of catchment area is the largest river in Malaysia, whereas in Peninsular Malaysia, it is Sungai Pahang with 29,000 km<sup>2</sup> of the catchment areas (Chan *et al.*, 2003). Most Malaysian rivers are short, steep (Abdullah, 2002) and located in remote areas (Berryhill-Jr. & Ahyaudin, 2002).

In Malaysia, low order streams are tropical mountain streams which are located deep within the forest and are difficult to be accessed. Basically, this low order stream is classified as oligotrophic with low fish diversity. Sungai Enam is one of the low order and high gradient streams in Malaysia. This stream is located within the Temengor Reservoir watershed, which is enriched with waterfalls at different heights and two small man-made dams. Water quality in Sungai Enam has recovered and was classified as Class I (Hashim *et al.*, 2011).

## **2.2 Assessment of freshwater fishes in streams**

Fish is a crucial inhabitant of an aquatic ecosystem. Fish constitute almost half from the total number of vertebrates in the world due to their ability to live virtually in every conceivable aquatic habitat (Nelson, 2006). Out of 39,900 species of vertebrates, there are 21,723 living fish species recorded (Jayaram, 1999), which is comprised of 8,411 freshwater species and 11,650 marine species (Kar *et al.*, 2006).

### **2.2.1 Fish assemblages in streams**

Fish assemblages play an important role in aquatic ecosystems. Fish assemblage is the term used to describe the collection of fish species in a habitat or fishing ground. Research on freshwater fish in Malaysia had begun in the mid-19<sup>th</sup> century (Zakaria-Ismail, 1991). However, studies on entire river stretches are still limited and need to be further discovered, especially in the upper reaches of rivers because this river section is the least studied, among other fluvial sections (Kottelat & Whitten, 1996), and possibly has many undiscovered endemic species (Shah *et al.*, 2009).

### **2.2.2 Previous studies of fish assemblages in the Temengor Reservoir**

Temengor Reservoir is one of the four consecutive hydroelectric reservoirs located along Sungai Perak after Chenderoh, Bersia, and Kenering Reservoir. Temengor Reservoir, which has approximately 30 m of mean depth and 152 km<sup>2</sup> of mean surface area (Khairul-Rizal, 2004) receives water from its surrounding water catchment areas that have countless headwaters.

Previous studies on fish diversity of Temengor Reservoir (lotic and lentic) include Zakaria-Ismail and Lim (1995), Zakaria-Ismail and Sabariah (1995), Mohd-Akhir (1999), Berryhill-Jr. (2003), Hashim *et al.* (2012), Hamid *et al.* (2012), Kaviarasu *et al.* (2013), Ismail *et al.* (2013), Mohd-Shafiq *et al.* (2013) and Amiruddin and Zakaria-Ismail (2015). To date, Amiruddin and Zakaria-Ismail (2015) reported that the total number of fish species found in the reservoir and streams that flow into Temengor Reservoir is 42 species compared to the first study on fish fauna in Temengor Reservoir carried out by Zakaria-Ismail and Lim (1995) with only 23 species. However, Hashim *et al.* (2012) recorded only 21 species in Temengor

Reservoir. This finding showed that there were some missing species in comparison with Mohd-Akhir (1999) who found 37 species in the same reservoir.

Meanwhile, in Sungai Enam, the number of fish species recorded for the first study in 1994 was only nine species (Zakaria-Ismail & Sabariah, 1995) followed by 21 species in 1998 (Mohd-Akhir, 1999). Fish species recorded in Sungai Enam declined afterwards to 14 species in 2002 (Berryhill-Jr., 2003), 11 species in 2004 (Hashim, 2006) and ten species in 2012 (Ismail *et al.*, 2013).

### **2.2.3 Electrofishing**

Electrofishing is a common technique used by fish biologists to catch fish in freshwater bodies. It is an active fishing method that establishes an electric field in the water to capture fish. Generally, there are three types of electrofishing unit used: (1) backpack, (2) stream-side or shore, and (3) electrofishing boats. An electrofishing operator typically moves through habitats accompanied by assistants who collect shocked fish with scoop nets (Portt *et al.*, 2006). A backpack unit is carried by an operator and may be powered by either a battery or a small gasoline powered generator. The efficiency of electrofishing in attracting and immobilizing fish is influenced by the characteristics of the electric field that is established. The effectiveness of electrofishing is lower in larger streams than in small streams (Paller, 1995).

Electrofishing is easy to be used in the field compared to other sampling gears that depend on the user's skills which results in different amounts of captured fish for different operators. However, this gear is limited to a certain depth and is only efficient in a specific water condition (Table 2.1).

Table 2.1: Advantages and disadvantages of electrofishing (adapted from Mohd-Akhir, 1999).

Advantages	Disadvantages
<ul style="list-style-type: none"> <li>• It is selective, whereby unwanted fish can be released back to the habitat alive without any physical injuries.</li> <li>• It is more efficient to be used compared to other sampling gears.</li> <li>• It is safe to be used if operators wear the wader and is also easy to use.</li> </ul>	<ul style="list-style-type: none"> <li>• Most fish will sink, and it is possible that the fish will not leave their habitats when shocked.</li> <li>• It is quite difficult to detect fish that were shocked in cloudy water.</li> <li>• The effectiveness of this gear is limited to a certain depth and certain distance.</li> </ul>

#### **2.2.4 Altitude as a key factor in limiting fish distribution in streams**

In fish study, most of the changes in diversity are related to altitude (Lomolino, 2001). In Malaysia, studies that show the trend of fish diversity and distribution in high altitude zones include studies conducted at Taman Kinabalu Sabah (Samat, 1990), Danum Valley Field Centre in Sabah (Martin-Smith & Laird, 1998), Taman Negara Pahang (Yahya & Singh, 2012), and Sungai Enam in Perak (Berryhill-Jr. & Ahyaudin, 2002; Hashim *et al.*, 2012; Ismail *et al.*, 2013). Other studies include Madeira River of Amazon basin (Torrente-Vilara *et al.*, 2011), and Agbokum waterfalls in South Eastern Nigeria (Offem & Ikpi, 2012).

The richness of freshwater fishes declines with the increase in altitude (Wootton, 1992; Jaramillo-Villa *et al.*, 2010). This statement is in concordance with a study by Agarwal *et al.* (2011) who found a low number of fish species at the upper stream stretches of Bhilagna river, Uttarakhand, India. Similar results of fish richness and diversity indices that were shown to be greater in downstream than upstream regions were obtained by several researchers such as Chang *et al.* (1999), Offem and Ikpi (2011), and Singh and Agarwal (2013).

Spatial change in fish community resulted from the shifts in the physical gradient of the stream from a shallow and temporally variable area (e.g., upstream and riffles) to a deeper and more stable area (e.g., downstream and pools) (Isaac, 1982). From upstream to downstream, and from riffle to pool, habitat diversity and water volume increased. Fish community also changes from headwaters to downstream, which due to the progressing addition of new species (Rahel & Hubert, 1991).



## **2.3 Assessment of the impacts of barriers to fish species**

In ecology, a barrier is a physical or biological factor that tends to limit the migration, interbreeding or free movement of individuals or populations. Migratory fish require different environments to complete their life cycle phases such as reproduction, juvenile production, growth and sexual maturation (Larinier, 2001). For instance, the life cycle of diadromous fish species partly takes place in freshwater, and partly in seawater, whereas the reproduction of anadromous species, takes place in freshwater. The presence of structures such as dams, weirs, culverts, waterfalls and cascades, fish movement could be restricted and at worst, can disrupt their sustaining habitats (Rahel, 2007).

### **2.3.1 The influence of man-made barriers to fish species**

Most rivers throughout the world have been dammed. Dam construction has its own pros and cons, but usually provides negative effects to fish distribution (Table 2.2). Dam construction affects nutrient dynamics along the system and fisheries in downstream reservoirs (Welcomme, 1985) and river channels (Hess *et al.*, 1982). However, a dam also enhances some riverine fisheries, particularly with respect to tailwater fisheries which are located below the dam. For example, tailwater fisheries below Table Rock and Taneycomo dams on the White River (Missouri, USA) received seven to ten times greater value of catch per unit effort (CPUE) than upstream reservoir (Fry, 1965).

Table 2.2: Physical and biological impacts of dams

<b>Impacts</b>	
Physical	<ul style="list-style-type: none"> <li>• Modifies the flow, temperature regimes, and water turbidity (Chang <i>et al.</i>, 1999).</li> <li>• Changes water quality (Chang <i>et al.</i>, 1999).</li> <li>• Blocks the passage of young migratory fishes in downstream and dams act as a barrier to upstream migrations (Larinier, 2001).</li> <li>• Breaks the upstream-downstream connectivity for a series of dams (Hashim, 2013).</li> <li>• Changes water discharge and current in some distance off downstream with the presence of impoundment and is able to alter a river's natural periodicity of flow for a series of large dams (Choi <i>et al.</i>, 2005).</li> </ul>
Biological	<ul style="list-style-type: none"> <li>• Causes permanent biological change when migratory fish species are eliminated (Kanehl <i>et al.</i>, 1997).</li> <li>• Reduces the productivity in the rivers with its floodplain in response to reduced and altered river flow (Chen, 2000).</li> <li>• Much more likely to eliminate flora and fauna (Pereira <i>et al.</i>, 2010) due to high flows that are being released from dams, especially for the hydroelectric dam.</li> <li>• Blocks the upstream passage of anadromous fish such as salmon, catadromous fish such as eels, and some stream resident fishes, but likely induces migration to spawn, such as the case for catfish (Godinho &amp; Kynard, 2009).</li> <li>• In some cases, water pressures encountered in turbines and spillways damage fishes migrating downstream (Cada, 1990).</li> </ul>

### 2.3.2 The influence of natural barriers to fish species

The blockage of fish passage, resulting from barriers, either man-made or natural, can have negative implications to fish population. The deleterious effects of dams on fish assemblage are commonly noticed because most fish are unable to jump or move against the dams unless fish ladder is provided (Noonan *et al.*, 2012; McKay *et al.*, 2013; Ottburg & Blank, 2015). Surprisingly, a natural barrier such as a waterfall is also able to interrupt fish movements along the stream.

A waterfall is one of the obvious natural barriers present in river ecosystem. Waterfalls are generally assumed as a vertical or very steep descent of a watercourse. Waterfalls can be found in the upper course of a river and are created by erosion process (Lamb *et al.*, 2007; Fuller, 2012). The steepness and height of waterfalls are varied between places (Kano *et al.*, 2012).

Waterfall increases speciation rates (Dias *et al.*, 2013) through geographical isolation created by this barrier (Rahel, 2007; Cote *et al.*, 2009). High waterfall can be a barrier to the upstream fish movement (Rahel, 2007; Torrente-Vilara *et al.*, 2011; Kano *et al.*, 2012) and prevent species exchange among regions (Rahel, 2007).

Fish diversity decreases above waterfalls (Martin-Smith & Laird, 1998; Carlsson & Nilsson, 2001; Costello *et al.*, 2003) as waterfalls block the upstream movement of some fish species. Above waterfall, the populations will only persist if the fish have special behavioural or morphological adaptations. Adaptability of swimming pattern in order to move against water current may reduce downstream losses (Northcote, 1978; Wootton, 1992). Resident species above waterfalls generally have dispersal mechanisms such as united pelvic fins in some species under genus *Gastromyzon*, which allows them to ascend these upstream barriers (Martin-Smith & Laird, 1998). Thus, a waterfall, which acts as a bio-geographic

barrier, is expected to produce different assemblage structures in riverine ecosystems (Torrente-Villara *et al.*, 2011) and generate freshwater biodiversity in river basins (Dias *et al.*, 2013).

## **2.4 Assessment of fish condition on the basis of length-weight relationships and relative condition factor**

Biometric relationships are often required in research field to transform collected data into appropriate indices (Ecoutin & Albaret, 2003). Both analyses (*i.e.*, length-weight relationships and relative condition factor) are important in fisheries and fish biology to determine the condition or the well-being of fish in their habitat. Generally, the fish well-being indicates their health and suitability with the environment. This is because fish is easily affected by any changes that occur in the environment. Heavier fishes at a specific length are said to be in a better condition (Jones *et al.*, 1999).

Most fish species tend to grow in length and weight after they have reached sexual maturity, even though the growth rate declines as the fish get bigger (Wootton, 1992). Mainly, food availability and temperature always play an important role in affecting their growth. Fish growth depends on the efficiency of fish in obtaining sufficient food and the ranges of prey they have as the fish grows.

### **2.4.1 Length-weight relationships (LWR)**

Many researchers use length-weight relationship (LWR) in their studies to observe the growth pattern for fish species. LWR provides the beneficial information of the growth patterns of fish (Bagenal & Tesch, 1978). LWR of fishes is necessary to evaluate the general well-being of fish species in terms of fatness, breeding, and feeding states and how the environment is suitable enough for them (Schneider *et al.*,

2000; Farzana & Saira, 2008; Mohd Shafiq *et al.*, 2012). LWR can also be an indicator of environmental changes and sustainable stock management (Samat *et al.*, 2008; Mohd-Shafiq *et al.*, 2012). The  $b$  value in the length-weight relationship of fish depends on biotic and abiotic factors such as water temperature, food availability, and habitat type (Wootton, 1992; Chu *et al.*, 2012).

The  $b$  value of isometric growth is 3, and values greater or lower than 3 are considered as allometric growth. The  $b$  value originated from the slope of regression from the length-weight equation. The formula of length-weight relationships of most fish can be described as:

$$W = aL^b$$

Where  $W$  = weight,  $L$  = length,  $a$  is a constant and  $b$  is an exponent that usually ranges between 2.5 to 4.0 (Le Cren, 1951).

#### **2.4.2 Relative condition factor ( $Kn$ )**

Suitability of habitat for fish growth could be determined by the value of relative condition factor ( $Kn$ ) (Samat *et al.*, 2008; Abowei, 2009; Mohd-Shafiq *et al.*, 2012). Expected weight of fish in the formula of relative condition factor ( $Kn$ ) could be identified from the equation of LWR (Isa *et al.*, 2010). Based on Pope and Kruse (2007), formula for the relative condition factor that is widely used such as by Le Cren (1951), Blackwell *et al.* (2000), Mohd-Shafiq *et al.* (2012) and Amani *et al.* (2015) is:

$$Kn = (W/W') \cdot 100$$

where  $W$ =observed weight of fish individual (g) and  $W'$ =expected weight of fish individual (g) from LWR.

Fish are assumed to be in a good condition when their relative condition factor ( $Kn$ ) is greater than 100, as 100 is used as a benchmark value in this parameter

(Pope & Kruse, 2007). The greater the value of relative condition factor, the better the habitat suitability in terms of food availability for optimum fish growth (Samat *et al.*, 2008; Abowei, 2009; Isa *et al.*, 2010; Mohd-Shafiq *et al.*, 2012). This *Kn* value also can be used to compare spatial and temporal condition between or among fish species (Samat *et al.*, 2008). The state of their sexual maturity and appetite condition also contributes to different values of the relative condition factor (Williams, 2000; Pervin & Mortuza, 2008; Isa *et al.*, 2010).

## **2.5 Fish movement over barriers**

Fish come and go within habitats (Roff, 1988; Wootton, 1992), and the spatial position of sites within stream provides a chance for fish to move freely to access resources or colonized new habitats (Hitt, 2007). Knowledge of fish movement is particularly important to managers as it provides information on fish's spatial and temporal distribution (Lucas & Baras, 2000). Fish movement studies would also provide a resource to determine specific spawning area and nursery habitat used by offspring and are appropriate for future fisheries management activities to be planned (Landsman *et al.*, 2011).

### **2.5.1 Importance of fish movement**

Movement is a necessary organisms' behaviour, which not only brings beneficial encounters, but also exposes the organism to dangerous encounters (Bianco *et al.*, 2014). Movements to a new habitat may increase individual fitness in promoting their reproductive success, which sometimes cannot be obtained from their previous habitat (Hitt, 2007). This movement could also reduce the rate of local extinction (Jackson *et al.*, 2001; Offem & Ikpi, 2011). Fish movement brings the fish into more suitable habitats than the habitats they left (Wootton, 1992). Commonly,