

# Technische Universität Chemnitz-Zwickau

DFG-Forschergruppe "SPC" · Fakultät für Mathematik

Sergej V. Nepomnyaschikh

## **Domain Decomposition and Multilevel Techniques for Preconditioning Operators**

Computing Center  
Siberian Branch of Russian Academy of Sciences  
Novosibirsk, 63090  
Russia  
e-mail: [svnep@comcen.nsk.su](mailto:svnep@comcen.nsk.su)

**Preprint-Reihe der Chemnitzer DFG-Forschergruppe  
"Scientific Parallel Computing"**

**SPC 95\_30**

**November 1995**

# 1 Introduction

In recent years, domain decomposition methods have been used extensively to efficiently solve boundary value problems for partial differential equations in complex-form domains [4, 13, 16]. On the other hand, multilevel techniques on hierarchical data structures also have developed into an effective tool for the construction and analysis of fast solvers [2, 5, 15, 17]. But direct realization of multilevel techniques on a parallel computer system for the global problem in the original domain involves difficult communication problems. In this paper, we present and analyze a combination of these two approaches: domain decomposition and multilevel decomposition on hierarchical structures to design optimal preconditioning operators.

Let  $\Omega \subset R^2$  be a polygon. In the domain  $\Omega$  we consider the boundary value problem

$$\left\{ \begin{array}{ll} - \sum_{i,j=1}^2 \frac{\partial}{\partial x_i} \left( a_{ij}(x) \frac{\partial u}{\partial x_j} \right) + a_0(x)u = f(x), & x \in \Omega \\ u(x) = 0, & x \in \Gamma_0 \\ \frac{\partial u}{\partial N} + \sigma(x)u = 0, & x \in \Gamma_1. \end{array} \right. \quad (1.1)$$

where

$$\frac{\partial u}{\partial N} = \sum_{i,j=1}^2 a_{ij}(x) \frac{\partial u}{\partial x_j} \cos(n, x_i)$$

is the conormal derivative,  $n$  denotes the outward normal to  $\Gamma$ , and  $\Gamma_0$  is a union of a finite number of curvilinear segments,  $\Gamma = \Gamma_0 \cup \Gamma_1$ ,  $\Gamma_0 = \bar{\Gamma}_0$ . Here  $\bar{\Gamma}_0$  denotes the closure of  $\Gamma_0$ .

By  $H^1(\Omega, \Gamma_0)$  we denote the subspace of the Sobolev space  $H^1(\Omega)$

$$H^1(\Omega, \Gamma_0) = \{v \in H^1(\Omega) \mid v(x) = 0, \quad x \in \Gamma_0\}.$$

We introduce the bilinear form  $a(u, v)$  and the linear functional  $l(v)$ :

$$a(u, v) = \int_{\Omega} \left( \sum_{i,j=1}^2 a_{ij}(x) \frac{\partial u}{\partial x_j} \frac{\partial v}{\partial x_i} + a_0(x)uv \right) dx + \int_{\Gamma_1} \sigma(x)uv dx$$

$$l(v) = \int_{\Omega} f(x)v dx.$$

Let us suppose that the operator coefficients and the right-hand side of the problem (1.1) are such that the bilinear form  $a(u, v)$  is symmetric, elliptic, and continuous on  $H^1(\Omega, \Gamma_0) \times H^1(\Omega, \Gamma_0)$ , i.e.

$$a(u, v) = a(v, u) \quad \forall u, v \in H^1(\Omega, \Gamma_0)$$

$$\alpha_0 \|u\|_{H^1(\Omega)}^2 \leq a(u, u) \leq \alpha_1 \|u\|_{H^1(\Omega)}^2 \quad \forall u \in H^1(\Omega, \Gamma_0)$$

and the linear functional  $l(v)$  is continuous on  $H^1(\Omega, \Gamma_0)$ :

$$|l(u)| \leq \alpha \|u\|_{H^1(\Omega)} \quad \forall u \in H^1(\Omega, \Gamma_0).$$

The generalized solution  $u \in H^1(\Omega, \Gamma_0)$  of (1.1) is, by definition, a solution to the projection problem [1]

$$u \in H^1(\Omega, \Gamma_0) : a(u, v) = l(v) \quad \forall v \in H^1(\Omega, \Gamma_0). \quad (1.2)$$

We know that under these assumptions for  $a(u, v)$  and  $l(v)$  there exists a unique solution of (1.2).

Let  $\Omega$  be a union of  $n$  nonoverlapping subdomains  $\Omega_i$ ,

$$\bar{\Omega} = \bigcup_{i=1}^n \bar{\Omega}_i, \quad \Omega_i \cap \Omega_j = \emptyset, \quad i \neq j,$$

where  $\Omega_i$  are polygons with diameters on the order of  $H$ . Let us consider a coarse grid triangulation of  $\Omega$

$$\Omega_0^h = \bigcup_{i=1}^n \Omega_{i,0}^h, \quad \Omega_{i,0}^h = \bigcup_{l=1}^{M_i^{(0)}} \bar{\tau}_{i,l}^{(0)},$$

$$\text{diam}(\tau_{i,l}^{(0)}) = O(H)$$

and we refine  $\Omega_{i,0}^h$  several times. This results in a sequence of nested triangulations

$$\Omega_{i,0}^h, \Omega_{i,1}^h, \dots, \Omega_{i,J}^h$$

such that

$$\bar{\Omega}_{i,k}^h = \bigcup_{l=1}^{M_i^{(k)}} \bar{\tau}_{i,l}^{(k)}, \quad k = 0, 1, \dots, J,$$

where the triangles  $\tau_{i,l}^{(k+1)}$  are generated by subdividing triangles  $\tau_{i,l}^{(k)}$  into four congruent subtriangles by connecting the midpoints of the edges.

Introduce the spaces

$$\begin{aligned} W_{i,0} \subset W_{i,1} \subset \dots \subset W_{i,J} &= H_h(\Omega_i), \\ V_{i,0} \subset V_{i,1} \subset \dots \subset V_{i,J} &= H_h(\Gamma_i), \end{aligned} \quad (1.3)$$

$$\Gamma_i = \partial\Omega_i, \quad i = 1, 2, \dots, n.$$

Here the space  $W_{i,k}$  consists of real-valued functions which are continuous on  $\Omega$  and linear on the triangles in  $\Omega_{i,k}^h$ . The space  $V_{i,k}$  is the space of traces on  $\Gamma_i$  of functions from  $W_{i,k}$ :

$$V_{i,k} = \left\{ \varphi^h \mid \varphi^h = u^h|_{\Gamma_i}, \text{ with } u^h \in W_{i,k} \right\}.$$

We define the space  $H_h(\Omega)$  of real continuous functions which are linear on each triangle of  $\Omega^h$  and vanish at  $\Gamma_0$ .

Let us consider the projection problem

$$u^h \in H_h(\Omega) : a(u^h, v^h) = l(v^h) \quad \forall v^h \in H_h(\Omega) \quad (1.4)$$

which is an approximation of the problem (1.2).

Each function  $u^h \in H_h(\Omega)$  is put in correspondence with a real column vector  $u \in R^N$  whose components are values of the function  $u^h$  at the corresponding nodes of the triangulation  $\Omega^h$ . Then (1.4) is equivalent to the system of mesh equations

$$\begin{aligned} Au &= f, \\ (Au, v) &= a(u^h, v^h) \quad \forall u^h, v^h \in H_h(\Omega), \\ (f, v) &= l(v^h) \quad \forall v^h \in H_h(\Omega), \end{aligned} \quad (1.5)$$

where  $u^h$  and  $v^h$  are the respective interpolations of vectors  $u$  and  $v$ ;  $(f, v)$  is the Euclidean scalar product in  $R^N$ .

The goal of this work is to construct a symmetric positive definite preconditioning operator  $B$  for (1.5) so as to satisfy the inequalities

$$c_1(Bu, u) \leq (Au, u) \leq c_2(Bu, u) \quad (1.6)$$

where the positive constants  $c_1$  and  $c_2$  are independent of  $h$  and  $H$ , the multiplication of a vector by  $B^{-1}$  should be easy to implement.

Using a combination of Additive Schwarz and Fictitious Space Methods, optimal preconditioning operators have been constructed in [11, 12, 13] for the case of arbitrary (unstructured) grids. However, that construction involves explicit extension operators whose implementation for three dimensional problems is optimal from the arithmetic cost and the condition number points of view but difficult for practice realization. The main goal of this work is to construct, using the hierarchical structure (1.3), a robust optimal preconditioning operator. One of the crucial points in [11, 12, 13] and this paper is using of non-exact solvers in subdomains and explicit extension operators. It means, to construct optimal preconditioning operators, we can design norm preserving operators of functions given at  $\Gamma_i$  into  $\Omega_i$  with the optimal arithmetic cost (a number of arithmetic operations should be proportional to a number degrees of freedom) and then, instead of exact solvers in subdomains, we can use any spectrally equivalent preconditioning operators. Optimal extension operators have been presented in [8, 9, 11] for unstructured grids and robust explicit extension operators on hierarchical data structures in [5, 14].

The paper is organized as follows. In the Section 2, using Additive Schwarz Method, we describe general construction of a preconditioning operator with local multilevel preconditioning operators. In the Section 3, we present an optimal multilevel extension of grid functions from boundaries subdomains into inside subdomains. In the Section 4, we propose an optimal interface preconditioning operator at the boundaries of the subdomains which involves a multilevel decomposition and corresponding explicit extension operators at interfaces.

## 2 Domain decomposition – additive Schwarz-Method

To design the preconditioning operator for the system (1.5), we use the additive Schwarz-Method [7] and realize the main idea of the construction of preconditioners from [13] for the hierarchical grids. Denote by  $\mathring{H}_h(\Omega_i)$  the subspace of  $H_h(\Omega_i)$

$$\mathring{H}_h(\Omega_i) = \{u^h \in H_h(\Omega_i) \mid u^h(x) = 0, \quad x \in \Gamma_i\}$$

and define the local preconditioning operators  $B_i$  such that

$$B_i : \mathring{H}_h(\Omega_i) \rightarrow \mathring{H}_h(\Omega_i),$$

$$c_3 \|u^h\|_{H^1(\Omega_i)}^2 \leq (B_i u, u) \leq c_4 \|u^h\|_{H^1(\Omega_i)}^2 \quad \forall u^h \in \mathring{H}_h(\Omega_i),$$

where  $c_3, c_4$  are independent of  $h$  and  $H$ . We hereafter use the same designation for an operator and its matrix representation. For instance, to define  $B_i$ , we can use the so-called BPX-preconditioners [3]. To do it, denote by  $\{f_l^{(k)}\}$  nodal basis functions from the  $k$ -th level and define

$$B_i^{-1} u^h = \sum_{k=0}^J \sum_{f_l^{(k)} \in \mathring{H}_h(\Omega_i)} (u^h, f_l^{(k)})_{L_2(\Omega_i)} f_l^{(k)}. \quad (2.1)$$

Let us assume that we can define the extension operators  $t_i$

$$t_i : V_{i,J} \longrightarrow W_{i,J}$$

such that

$$\begin{aligned} t_i \varphi^h &= u^h, \\ u^h(x) &= \varphi^h(x), \quad x \in \Gamma_i, \end{aligned} \tag{2.2}$$

$$\|t_i \varphi^h\|_{H^1(\Omega_i)} \leq c_5 \|\varphi^h\|_{H^{1/2}(\Gamma_i)} \quad \forall \varphi^h \in V_{i,J},$$

with  $c_5$  independent of  $h$  and  $H$ . Here  $\|\varphi^h\|_{H^{1/2}(\Gamma_i)}$  is the norm [10] in the Sobolev space  $H^{1/2}(\Gamma_i)$

$$\|\varphi^h\|_{H^{1/2}(\Gamma_i)}^2 = H \int_{\Gamma_i} (\varphi^h(x))^2 dx + \int_{\Gamma_i} \int_{\Gamma_i} \frac{(\varphi^h(x) - \varphi^h(y))^2}{|x - y|^2} dx dy.$$

Then, we can define the extension operator  $t$

$$t : H_h(S) \rightarrow H_h(\Omega),$$

where  $H_h(S)$  is the space of traces of functions from  $H_h(\Omega)$  at  $S$

$$S = \bigcup_{i=1}^n \Gamma_i$$

and for any  $\varphi^h \in H_h(S)$

$$\begin{aligned} t \varphi^h &= u^h, \\ u^h(x) &= \varphi^h(x), \quad x \in S, \\ \|t \varphi^h\|_{H^1(\Omega)} &\leq c_5 \|\varphi^h\|_{H^{1/2}(S)}. \end{aligned}$$

Here

$$\|\varphi^h\|_{H^{1/2}(S)}^2 = \sum_{i=1}^n \|\varphi^h\|_{H^{1/2}(\Gamma_i)}^2.$$

The operator  $t_i$  from (2.2) is constructed in the Section 3.

Let  $\Sigma$  satisfies to the following inequalities

$$c_6 \|\varphi^h\|_{H^{1/2}(S)}^2 \leq (\Sigma \varphi, \varphi) \leq c_7 \|\varphi^h\|_{H^{1/2}(S)}^2 \quad \forall \varphi^h \in H_h(S), \tag{2.3}$$

where  $c_6, c_7$  independent of  $h$  and  $H$ . Then, according to [11], we can define the preconditioning operator  $B$  as follows

$$B^{-1} = \begin{bmatrix} 0 & & & \\ & B_1^{-1} & & \\ & & \ddots & \\ & & & B_n^{-1} \end{bmatrix} + t \Sigma^{-1} t^*. \tag{2.4}$$

Here  $0$  is the null-matrix which corresponds to nodes of the triangulation  $\Omega^h$  at  $S$  and  $B_i$  is from (2.1).

The following theorem is valid

**Theorem 2.1** *If the operator  $B$  is from (2.4), then the constants  $c_1, c_2$  from (1.6) are independent of  $h$  and  $H$ .*

### 3 Multilevel explicit extension operators

The main goal of this section is to construct the robust operator  $t_i$  from (2.2). During this section, we omit the subscript  $i$ .

To design the extension operator

$$t : V_J \rightarrow W_J,$$

we follow to [5, 14]. Denote by  $\varphi_i^{(k)}, i = 1, 2, \dots, N_k$ , the nodal basis of  $V_k$  and denote by  $\Phi_i^{(k)}$  the one-dimensional subspace spanned by this function  $\varphi_i^{(k)}$ . Define

$$Q_i^{(k)} : L_2(\Gamma) \rightarrow \Phi_i^{(k)}$$

the  $L_2$ -orthoprojection from  $L_2(\Gamma)$  on to  $\Phi_i^{(k)}$  and denote

$$\tilde{Q}_k = \sum_{i=1}^{N_k} Q_i^{(k)}, \quad k = 0, 1, \dots, J-1.$$

For  $k = J$  we define  $\tilde{Q}_J$  as the  $L_2$ -orthoprojection from  $L_2(\Gamma)$  on to  $V_J$ .

The following lemmas are valid [14].

**Lemma 3.1** *There exists a positive constant  $c_8$ , independent of  $h$  and  $H$ , such that for any  $\varphi^h \in V_J$*

$$\|\varphi_0^h\|_{H^{1/2}(\Gamma)}^2 + \frac{1}{H} \|\varphi_1^h\|_{L_2(\Gamma)}^2 + |\varphi_1^h|_{H^{1/2}(\Gamma)}^2 \leq c_8 \|\varphi^h\|_{H^{1/2}(\Gamma)}^2,$$

where

$$\varphi_0^h = \tilde{Q}_0 \varphi^h, \quad \varphi_1^h = \varphi^h - \varphi_0^h. \quad (3.1)$$

Here

$$|\varphi^h|_{H^{1/2}(\Gamma)}^2 = \int_{\Gamma} \int_{\Gamma} \frac{(\varphi^h(x) - \varphi^h(y))^2}{|x - y|^2} dx dy.$$

**Lemma 3.2** *There exists a positive constant  $c_9$ , independent of  $h$  and  $H$ , such that*

$$\|\varphi_0^h\|^2 + \frac{1}{H} \left( \|\tilde{Q}_0 \varphi_1^h\|_{L_2(\Gamma)}^2 + \sum_{k=1}^J 2^k \|(\tilde{Q}_k - \tilde{Q}_{k-1}) \varphi_1^h\|_{L_2(\Gamma)}^2 \right) \leq c_9 \|\varphi^h\|_{H^{1/2}(\Gamma)}^2,$$

where  $\varphi_0^h, \varphi_1^h$  from (3.1).

The construction of the operator  $t$  is based on the decomposition from the Lemma 3.2. Denote by  $x_i^{(k)}, i = 1, 2, \dots, L_k$ , the nodes of the triangulation  $\Omega_k^h$  (we assume that nodes  $x_i^{(k)}$  are enumerated first on  $\Gamma$  and then inside  $\Omega$ ) and define the extension operator  $t$  in the following way. For any  $\varphi^h \in V_J$  set

$$\begin{aligned} \psi_0^h &= \tilde{Q}_0 \varphi^h, \\ \psi_k^h &= (\tilde{Q}_k - \tilde{Q}_{k-1}) \varphi^h, \quad k = 1, 2, \dots, J. \end{aligned} \quad (3.2)$$

Then

$$\varphi^h = \psi_0^h + \psi_1^h + \dots + \psi_J^h.$$

Define the extension  $u_k^h \in W_k$  as follows

$$\begin{aligned} u_0^h(x_i^{(0)}) &= \begin{cases} \psi_0^h(x_i^{(0)}), & x_i^{(0)} \in \Gamma, \\ \bar{\psi}, & x_i^{(0)} \notin \Gamma, \end{cases} \\ u_k^h(x_i^{(k)}) &= \begin{cases} \psi_k^h(x_i^{(k)}), & x_i^{(k)} \in \Gamma, \\ 0, & x_i^{(k)} \notin \Gamma, \end{cases} \\ &k = 1, 2, \dots, J. \end{aligned} \tag{3.3}$$

Here  $\bar{\psi}$  is, for instance, the meanvalue of the function  $\psi_0^h$  on  $\Gamma$

$$\bar{\psi} = \frac{1}{N_0} \sum_{i=1}^{N_0} \psi_0^h(x_i^{(0)}).$$

Define

$$t\varphi^h = u^h \equiv u_0^h + u_1^h + \dots + u_J^h \tag{3.4}$$

**Remark 3.1** *We can use the  $L_2$ -orthoprojections from  $L_2(\Gamma)$  on to  $V_k$  instead of  $\tilde{Q}_k$ ,  $k = 0, 1, \dots, J-1$ . But in this case the cost of the decomposition (3.2) is expensive (especially for three dimensional problems).*

**Theorem 3.1** *There exists a positive constant  $c_{10}$ , independent of  $h$  and  $H$ , such that*

$$\|t\varphi^h\|_{H^1(\Omega)} \leq c_{10} \|\varphi^h\|_{H^{1/2}(\Gamma)} \quad \forall \varphi^h \in V_J.$$

Here the operator  $t$  is from (3.2)–(3.4).

**Remark 3.2** *It is obvious that*

$$Q_i^{(k)} \varphi^h = \frac{(\varphi^h, \varphi_i^{(k)})_{L_2(\Gamma)}}{(\varphi_i^{(k)}, \varphi_i^{(k)})_{L_2(\Gamma)}} \varphi_i^{(k)}$$

and the cost of the action of  $t$  and  $t^*$  is proportional to the number of nodes of the grid domain.

## 4 Interface preconditioning operators

In this section, we construct an optimal interface preconditioner in the space  $H_h(S)$  which satisfies (2.3). To do it, we use the idea of Additive Schwarz Method at interface  $S$  from [13]. Let  $S$  be a union of  $K$  nonoverlapping edges  $E_i$  of the triangulation  $\Omega_0^h$

$$S = \bigcup_{j=1}^K \bar{E}_j, \quad E_j \cap E_i = \emptyset, \quad i \neq j.$$

Split  $H_h(S)$  into a vector sum of subspaces

$$H_h(S) = U_0 + U_1 + \dots + U_K, \tag{4.1}$$

where  $U_0$  is the coarse space which consists of continuous functions linear on the edges  $E_j$ ,  $j = 1, 2, \dots, K$ , and  $U_j$ ,  $j = 1, 2, \dots, K$ , correspond to  $E_j$  and are defined below.

Denote by

$$\begin{aligned}\mathring{U}_j &= \{\varphi^h \in H_h(S) \mid \varphi^h(x) = 0, x \notin E_j\}, \\ \tilde{U}_j^{(k)} &= V_k|_{E_j}, \quad k = 0, 1, \dots, J.\end{aligned}$$

For any edge  $E_j$  we define the explicit extension operator  $\tau_j$

$$\tau_j : \tilde{U}_j^{(J)} \rightarrow H_h(S)$$

as follows. Denote by  $\varphi_{j,i}^{(k)}$ ,  $i = 1, 2, \dots, I_j^{(k)}$ , the nodal basis of  $\tilde{U}_j^{(k)}$  (the functions  $\varphi_{j,i}^{(k)}$  are differ from the functions  $\varphi_i^{(k)}$  from the Section 3 only at the end points of  $E_j$ ) and denote by  $\Phi_{j,i}^{(k)}$  the one-dimensional subspace spanned by this function  $\varphi_{j,i}^{(k)}$ . Define

$$Q_{j,i}^{(k)} : L_2(E_j) \rightarrow \Phi_{j,i}^{(k)}$$

corresponding  $L_2$ -orthoprojection. Set

$$\tilde{Q}_j^{(k)} = \sum_{i=1}^{I_j^{(k)}} Q_{j,i}^{(k)}, \quad k = 0, 1, \dots, J-1,$$

and define  $\tilde{Q}_J^{(k)}$  as the  $L_2$ -orthoprojection from  $L_2(E_j)$  onto  $\tilde{U}_j^{(J)}$ . Now we can define the extension operator  $\tau_j$  according to (3.2)–(3.4). For any  $\varphi^h \in \tilde{U}_j^{(J)}$  set

$$\begin{aligned}\psi_0^h &= \tilde{Q}_j^{(0)}\varphi^h, \\ \psi_k^h &= (\tilde{Q}_j^{(k)} - \tilde{Q}_j^{(k-1)})\varphi^h, \quad k = 1, 2, \dots, J\end{aligned}\tag{4.2}$$

and

$$u_k^h = \begin{cases} \psi_k^h(x_i^{(k)}), & x_i^{(k)} \in E_j \\ 0, & x_i^{(k)} \notin E_j, \end{cases} \quad k = 0, 1, \dots, J\tag{4.3}$$

$$\tau_j\varphi^h = u_0^h + u_1^h + \dots + u_J^h.$$

Define

$$U_j = \mathring{U}_j + \tau_j\tilde{U}_j.$$

Then from the Theorem 3.1 and [13] for the decomposition (4.1) we have the following theorem.

**Theorem 4.1** *There exists a positive constant  $c_{11}$ , dependent of  $h$  and  $H$ , such that for any function  $\varphi^h \in H_h(S)$  there exist  $\varphi_j^h \in U_j$ ,  $j = 0, 1, \dots, K$ , such that*

$$\varphi_0^h + \varphi_1^h + \dots + \varphi_k^h = \varphi^h,$$

$$\|\varphi_0^h\|_{H^{1/2}(S)}^2 + \|\varphi_1^h\|_{H^{1/2}(S)}^2 + \dots + \|\varphi_k^h\|_{H^{1/2}(S)}^2 \leq c_{11}\|\varphi^h\|_{H^{1/2}(S)}^2$$

Let the operator  $\Sigma_0$  generates an equivalent norm in  $U_0$ .

$$c_{12}\|\varphi^h\|_{H^{1/2}(S)}^2 \leq (\Sigma_0\varphi, \varphi) \leq c_{13}\|\varphi^h\|_{H^{1/2}(S)}^2 \quad \forall \varphi^h \in U_0,\tag{4.4}$$



where  $c_{12}$ ,  $c_{13}$  independent of  $h$  and  $H$ . Define local preconditioners for  $U_j$ ,  $j = 1, 2, \dots, K$ . Denote by  $\overset{\circ}{\Sigma}_j$  and  $\tilde{\Sigma}_j$  the BPX-like preconditioners in the spaces  $\overset{\circ}{U}_j$  and  $\tilde{U}_j$ , respectively

$$\begin{aligned}\overset{\circ}{\Sigma}_i^{-1}\varphi^h &= \sum_{k=0}^J \sum_{\text{supp } \varphi_{j,i}^{(k)} \subset E_j} (\varphi^h, \varphi_{j,i}^{(k)})_{L_2(E_j)} \varphi_{j,i}^{(k)} \quad \forall \varphi^h \in \overset{\circ}{U}_j, \\ \Sigma_i^{-1}\varphi^h &= \sum_{k=0}^J \sum_{\text{supp } \varphi_{j,i}^{(k)} \cap E_j \neq \emptyset} (\varphi^h, \varphi_{j,i}^{(k)})_{L_2(E_j)} \varphi_{j,i}^{(k)} \quad \forall \varphi^h \in \tilde{U}_j.\end{aligned}$$

Then, define the interface preconditioning operator  $\Sigma$  in the following way

$$\Sigma^{-1} = \Sigma_0^+ + \sum_{j=1}^K (\overset{\circ}{\Sigma}_j^{-1} + \tau_j \tilde{\Sigma}_j^{-1} \tau_j^*). \quad (4.5)$$

Here  $\Sigma_0^+$  is a pseudo-inverse of  $\Sigma_0$  from (4.4),  $\tau_j$  is from (4.2), (4.3), and we extend the operator  $\overset{\circ}{\Sigma}_j^{-1}$  by zero outside  $E_j$ . The following theorem is valid.

**Theorem 4.2** *If the operator  $\Sigma$  is from (4.5) then the constants  $c_6, c_7$  from (2.3) are independent of  $h$  and  $H$ .*

**Remark 4.1** *The method suggested in this paper can be generalized evidently for three dimensional problems.*

**Remark 4.2** *Using combination of presented technique and technique from [10], effective preconditioning operators for elliptic problems with jump coefficients can be constructed.*

### Acknowledgment

The author wish to acknowledge the DFG–Forschergruppe “SPC” Chemnitz and the ISF for supporting this research under grants La 767/3 and NPB 000.

### References

- [1] J.-P. Aubin. *Approximation of elliptic boundary–value problems*. Wiley–Interscience, New York, London, Sydney, Toronto, 1972.
- [2] J. H. Bramble. *Multigrid Methods*. Research Notes in Mathematics Series. Pitman, Boston-London-Melbourne, 1993.
- [3] J. H. Bramble, J. E. Pasciak, and J. Xu. Parallel multilevel preconditioners. *Math. Comput.*, 55(191):1–22, 1990.
- [4] M. Dryja and O. B. Widlund. Multilevel additive methods for elliptic finite element problems. In W. Hackbusch, editor, *Parallel Algorithms for Partial Differential Equations*, pages 58–69, Braunschweig, 1991. Vieweg–Verlag. Proc. of the Sixth GAMM–Seminar, Kiel, January 19–21, 1990.
- [5] G. Haase, U. Langer, A. Meyer, and S. V. Nepomnyaschikh. Hierarchical extension and local multigrid methods in domain decomposition preconditioners. *East–West J. Numer. Math.*, 2(3):173–193, 1994.

- [6] W. Hackbusch. *Multi-Grid Methods and Applications*, volume 4 of *Springer Series in Computational Mathematics*. Springer-Verlag, Berlin, 1985.
- [7] A. M. Matsokin and S. V. Nepomnyaschikh. A Schwarz alternating method in a subspace. *Soviet Mathematics*, 29(10):78–84, 1985.
- [8] A. M. Matsokin and S. V. Nepomnyaschikh. Norms in the space of traces of mesh functions. *Sov. J. Numer. Anal. Math. Modelling*, 3:199–216, 1988.
- [9] S. V. Nepomnyaschikh. *Domain decomposition and Schwarz methods in a subspace for the approximate solution of elliptic boundary value problems*. PhD thesis, Computing Center of the Siberian Branch of the USSR Academy of Sciences, Novosibirsk, 1986.
- [10] S. V. Nepomnyaschikh. Domain decomposition methods for elliptic problems with discontinuous coefficients. In R. Glowinski, Y. A. Kuznetsov, G. A. Meurant, and J. Periaux, editors, *Domain decomposition methods for partial differential equations*, pages 242–251, Philadelphia, 1991. SIAM. Proceedings of the 4th International Symposium, Moscow, 1990.
- [11] S. V. Nepomnyaschikh. Method of splitting into subspaces for solving elliptic boundary value problems in complex-form domains. *Sov. J. Numer. Anal. Math. Modelling*, 6:151–168, 1991.
- [12] S. V. Nepomnyaschikh. Mesh theorems on traces, normalization of function traces and their inversion. *Sov. J. Numer. Anal. Math. Modelling*, 6:1–25, 1991.
- [13] S. V. Nepomnyaschikh. Decomposition and Fictitious Domains Methods for Elliptic Boundary Value Problems. In T. F. Chan, D. E. Keyes, G. A. Meurant, T. S. Scroggs, and R. G. Voigt, editors, *5th Conference on Domain Decomposition Methods for PDE*, pages 62–72, Philadelphia, 1992. SIAM.
- [14] S. V. Nepomnyaschikh. Optimal multilevel extension operators. Preprint SPC 95\_3, Technische Universität Chemnitz-Zwickau, Fakultät für Mathematik, 1995.
- [15] P. Oswald. *Multilevel Finite Element Approximation: Theory and Applications*. Teubner Skripten zur Numerik. B. G. Teubner Stuttgart, 1994.
- [16] P. Le Tallec. Domain Decomposition Methods in Computational Mechanics. *Computational Mechanics Advances (North Holland)*, 1(2):121–220, Feb. 1994.
- [17] J. Xu. Iterative methods by space decomposition and subspace correction. *SIAM Review*, 34:581–613, 1992.

Other titles in the SPC series:

- 95\_1 T. Apel, G. Lube. Anisotropic mesh refinement in stabilized Galerkin methods Januar 1995.
- 95\_2 M. Meisel, A. Meyer. Implementierung eines parallelen vorkonditionierten Schur-Komplement CG-Verfahrens in das Programmpaket FEAP. Januar 1995.
- 95\_3 S. V. Nepomnyaschikh. Optimal multilevel extension operators. January 1995
- 95\_4 M. Meyer. Grafik-Ausgabe vom Parallelrechner für 3D-Gebiete. Januar 1995
- 95\_5 T. Apel, G. Haase, A. Meyer, M. Pester. Parallel solution of finite element equation systems: efficient inter-processor communication. Februar 1995
- 95\_6 U. Groh. Ein technologisches Konzept zur Erzeugung adaptiver hierarchischer Netze für FEM-Schemata. Mai 1995
- 95\_7 M. Bollhöfer, C. He, V. Mehrmann. Modified block Jacobi preconditioners for the conjugate gradient method. Part I: The positive definit case. January 1995
- 95\_8 P. Kunkel, V. Mehrmann, W. Rath, J. Weickert. GELDA: A Software Package for the Solution of General Linear Differential Algebraic Equation. February 1995
- 95\_9 H. Matthes. A DD preconditioner for the clamped plate problem. February 1995
- 95\_10 G. Kunert. Ein Residuenfehlerschätzer für anisotrope Tetraedernetze und Dreiecksnetze in der Finite-Elemente-Methode. März 1995
- 95\_11 M. Bollhöfer. Algebraic Domain Decomposition. March 1995
- 95\_12 B. Nkemzi. Partielle Fourierdekomposition für das lineare Elastizitätsproblem in rotationssymmetrischen Gebieten. März 1995
- 95\_13 A. Meyer, D. Michael. Some remarks on the simulation of elasto-plastic problems on parallel computers. March 1995
- 95\_14 B. Heinrich, S. Nicaise, B. Weber. Elliptic interface problems in axisymmetric domains. Part I: Singular functions of non-tensorial type. April 1995
- 95\_15 B. Heinrich, B. Lang, B. Weber. Parallel computation of Fourier-finite-element approximations and some experiments. May 1995
- 95\_16 W. Rath. Canonical forms for linear descriptor systems with variable coefficients. May 1995
- 95\_17 C. He, A. J. Laub, V. Mehrmann. Placing plenty of poles is pretty preposterous. May 1995
- 95\_18 J. J. Hench, C. He, V. Kučera, V. Mehrmann. Dampening controllers via a Riccati equation approach. May 1995
- 95\_19 M. Meisel, A. Meyer. Kommunikationstechnologien beim parallelen vorkonditionierten Schur-Komplement CG-Verfahren. Juni 1995
- 95\_20 G. Haase, T. Hommel, A. Meyer and M. Pester. Bibliotheken zur Entwicklung paralleler Algorithmen. Juni 1995.
- 95\_21 A. Vogel. Solvers for Lamé equations with Poisson ratio near 0.5. June 1995.

- 95\_22 P. Benner, A. J. Laub, V. Mehrmann. A collection of benchmark examples for the numerical solution of algebraic Riccati equations I: Continuous-time case. October 1995.
- 95\_23 P. Benner, A. J. Laub, V. Mehrmann. A collection of benchmark examples for the numerical solution of algebraic Riccati equations II: Discrete-time case. to appear: December 1995.
- 95\_24 P. Benner, R. Byers. Newton's method with exact line search for solving the algebraic Riccati equation. October 1995.
- 95\_25 P. Kunkel, V. Mehrmann. Local and Global Invariants of Linear Differential-Algebraic Equations and their Relation. July 1995.
- 95\_26 C. Israel. NETGEN69 - Ein hierarchischer paralleler Netzgenerator. August 1995.
- 95\_27 M. Jung. Parallelization of multi-grid methods based on domain decomposition ideas. November 1995.
- 95\_28 P. Benner, H. Faßbender. A restarted symplectic Lanczos method for the Hamiltonian eigenvalue problem. October 1995.
- 95\_29 G. Windisch. Exact discretizations of two-point boundary value problems. October 1995.
- 95\_30 S. V. Nepomnyaschikh. Domain decomposition and multilevel techniques for preconditioning operators. November 1995.
- 95\_31 H. Matthes. Parallel preconditioners for plate problems. November 1995.
- 95\_32 V. Mehrmann, H. Xu. An analysis of the pole placement problem. I. The single input case. November 1995.
- 95\_33 Th. Apel. SPC-PM Po3D — User's manual. December 1995.
- 95\_34 Th. Apel, F. Milde, M. Theß. SPC-PM Po3D — Programmer's manual. December 1995.
- 95\_35 S. A. Ivanov, V. G. Korneev. On the preconditioning in the domain decomposition technique for the p-version finite element method. Part I. December 1995.
- 95\_36 S. A. Ivanov, V. G. Korneev. On the preconditioning in the domain decomposition technique for the p-version finite element method. Part II. December 1995.

Some papers can be accessed via anonymous ftp from server `ftp.tu-chemnitz.de`, directory `pub/Local/mathematik/SPC`. (Note the capital L in Local!)  
The complete list of current and former preprints is available via  
`http://www.tu-chemnitz.de/~pester/sfb/spc95pr.html`.