



A guide to studying human dermal adipocytes in situ

DOI:

[10.1111/exd.13549](https://doi.org/10.1111/exd.13549)

Document Version

Accepted author manuscript

[Link to publication record in Manchester Research Explorer](#)

Citation for published version (APA):

Nicu, C., Pople, J., Bonsell, L., Bhogal, R., Ansell, D., & Paus, R. (2018). A guide to studying human dermal adipocytes in situ. *Experimental Dermatology*. <https://doi.org/10.1111/exd.13549>

Published in:

Experimental Dermatology

Citing this paper

Please note that where the full-text provided on Manchester Research Explorer is the Author Accepted Manuscript or Proof version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version.

General rights

Copyright and moral rights for the publications made accessible in the Research Explorer are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Takedown policy

If you believe that this document breaches copyright please refer to the University of Manchester's Takedown Procedures [<http://man.ac.uk/04Y6Bo>] or contact uml.scholarlycommunications@manchester.ac.uk providing relevant details, so we can investigate your claim.



A guide to studying human dermal adipocytes *in situ*

Journal:	<i>Experimental Dermatology</i>
Manuscript ID	EXD-17-0432.R1
Manuscript Type:	Methods Review (invited)
Date Submitted by the Author:	12-Mar-2018
Complete List of Authors:	Nicu, Carina; University of Manchester, Centre for Dermatology Research Pople, Jennifer; Unilever R&D Colworth, Bonsell, Laura; University of Manchester, Centre for Dermatology Research Ansell, David; University of Manchester, Centre for Dermatology Research Bhogal, Ranjit; Unilever R&D, Colworth Discover, Tony James Building Paus, Ralf; University of Manchester, Centre for Dermatology Research
Keywords:	white adipose tissue, immunohistochemistry, immunofluorescence

1
2
3 **Prof. Thomas Luger**
4
5 **Editorial Board Member**
6
7 **Experimental Dermatology**
8
9

10
11 Manchester, March 12, 2018
12

13 **RE: EXD-17-0432 2**
14
15

16
17
18
19 Dear Prof. Luger, lieber Thomas,
20

21 We thank you and your referees for the time and effort spent reviewing our methods review
22 by Nicu et al. entitled 'A guide to studying human dermal adipocyte *in situ*'.
23
24

25 We are pleased to submit a revised manuscript that fully implements the valuable suggestions
26 made by your expert reviewers. All the changes have been acknowledged and detailed in our
27 point-by-point reply.
28
29

30
31 As explained below, we are in full agreement with both reviewers and their helpful feedback.
32 We hope that you shall consider our revised manuscript acceptable for publication, and we
33 look forward to hearing back from you soon.
34
35
36

37 Kind regards,
38

39
40 Ralf
41
42
43

44 ***Ralf Paus, MD, FRSB***

45 *Professor of Cutaneous Medicine*

46 *The Centre for Dermatology Research,*

47 *University of Manchester,*

48 *Manchester, United Kingdom*

49 *ralf.paus@manchester.ac.uk*
50
51
52
53
54
55
56
57
58
59
60

Nicu et al – Point-by-point Reply

Reviewer: 1

Suggestions for Authors

In this method review, Nicu et al well summarized the available techniques for studying human dermal fat, and also provide detailed guidelines for dermal fat study. This review would be of great help to the investigators of dermal fat study.

We are pleased that the reviewer finds this review to be of “great help to investigators” in the field.

Minor suggestions are listed below:

Human dermal fat is different from murine one, no panniculus carnosus layer exist in human skin to separate the dermal fat layer and subcutaneous fat layer. The primary adipocytes that mentioned in the "Dermal Adipocyte Culture" Section (Page 11), the Lonza derived adipocytes, was isolated from the whole subcutaneous fat of human skin. As a result, Lonza preadipocytes are the mixture of preadipocytes from both dermal fat and subcutaneous fat. The author should make it clear in this review.

Thanks a lot – we absolutely agreed that this needs to be made clearer. We already discuss the distinction between human and mouse skin (page 2, section ‘Human DWAT anatomy and biology in a nutshell’, and mention that, perhaps the platysma colli of human neck skin may be somewhat comparable to the *panniculus carnosus* that divides the subcutis of murine skin) (Larson et al, 2014). As requested, we now further clarify that the Lonza pre-adipocytes used in the current study are derived from whole subcutaneous human fat.

As Magnetic resonance imaging (MRI) is a good way for fat imaging. MRI method could be included in this review.

We have incorporated the reviewer’s helpful suggestion into our manuscript by adding a section on imaging adipocytes using MRI after the ‘Transmission Electron Microscopy’ paragraph, as well as adding MRI as an imaging technique in Table 1, along with appropriate references.

Reviewer: 2

Suggestions for Authors

Over the last years, several research groups have focus their activities on the study of subcutaneous and visceral fat, trying to understand the role of these adipose tissues in the development of metabolic diseases (insulin resistance, type 2 diabetes, cardiovascular diseases, liver diseases). Interestingly, adipose cells are found in several other compartments of the body and are now known to participate in the development of other diseases. This is the case for dermal adipocytes, that are now known to play key roles in regulating several biological processes in the skin. In this manuscript, Carina et al. review the common techniques used to study dermal adipocytes. The timing for this review is good considering the growing interest in dermal fat cells. Overall, this review is interesting and well written. I liked the enthusiasm of the authors toward the study of this fat depot. I only have minor comments that should be addressed before the publication of the manuscript.

We thank the reviewer for this encouraging feedback.

1
2
3 1) P5, line 38. There is an emerging literature on bright fat cells. The authors should at least mention that these
4 cells exist.

5
6 The reviewer makes an excellent point. As requested, we have now discussed bright fat cells in our 'Human
7 DWAT anatomy and biology in a nutshell' section, along with appropriate references, in which details can be
8 further explored.
9

10
11
12 2) P5. Please define what HF refers to. I presume that this means hair follicle but I could not find the
13 information.

14 We apologize for this omission, which has now been corrected. Yes, indeed: HF is an abbreviation of hair
15 follicle.
16
17

18
19 3) Table 1. This table is interesting but several changes should be made. First, the section 'adipocyte size and
20 number' could be fused with the 'Mature adipocyte' section. The information presented in these section is the
21 same.
22

23 We thank the reviewer for their valuable comments on Table 1. Based on this feedback we have made extensive
24 changes to Table 1. Among these, we have amalgamated "adipocyte size and number" with "mature adipocyte"
25 as suggested.
26
27

28
29 4) Table 1 (and reference in the text, p.9-10). I was surprised to see that CD68, PLIN1, and caveolin are used as
30 marker of apoptosis. I agree that adipose cell death often produces necrotic sections and crown like structures.
31 However, dying adipocytes will not necessarily be associated with CD68 staining. Moreover, the presence of
32 macrophages (CD68) in fat does not mean that adipose cells are dying. This section should be corrected as it can
33 be misleading.
34

35 Thank you for encouraging us to clarify this. We meant that adipocyte death could be measured using a double
36 stain of CD68 coupled with either PLIN1 or Caveolin1 to mark both macrophages and dermal adipocytes –
37 namely those that are completely surrounded, and thereby supposedly phagocytosed, by CD68+ macrophages.
38 We have now added this information since it features prominently in several mouse adipocyte research papers
39 (Berry et al, 2014; Martinez-Santibanez et al, 2014; Cinti et al, 2005; Giordano et al, 2013), which are now cited
40 in the revised review. However, we understand that dying adipocytes can express CD68+ crown-like structures
41 only during necrosis and pyroptosis (and have phrased this accordingly in the core text under 'Apoptosis
42 markers'. We have made the appropriate changes in both Table 1 and the core text under 'Apoptosis markers' to
43 avoid any confusion with the readers.
44
45
46
47
48
49

50 5) Table 1. It would be important to add some staining in the table. In the text, the authors refer to Masson
51 Trichrome and Picrosirius Red to stain fibrosis. Unfortunately, we do not find any reference to these staining in
52 Table 1.
53

54 6) Table 1. The same applies for Methylene blue.

55 Thank you for helping us to improve the table regarding Points 5) and 6). These markers have now been added
56 to Table 1.
57

58 7) Table 1. The importance of including the sections 'lipid synthesis' and 'lipolysis' in Table 1 is debatable. It is
59 not accurate to measure lipolysis by looking at ATGL, MGL and HSL protein levels by IF. Lipolysis is a
60

1
2
3 complex biological process that involves rapid phosphorylation /dephosphorylation steps and looking at the total
4 levels of enzyme does not mean anything in terms of enzymatic activity. The same comment applies for lipid
5 synthesis. The levels of FAS, DGAT and LPL can be a marker of differentiation (mature vs immature fat cells)
6 but these proteins cannot be used by IF to measure lipogenesis.
7

8
9 The reviewer raises an excellent point. ATGL, MGL and HSL are regulated at the post-transcriptional level,
10 meaning that there can be a lack of correlation between their levels and the process of lipolysis. Thus, we have
11 eliminated ATGL and MGL from Table 1 as markers of lipolysis, and instead have included phospho-Perilipin 1
12 and phospho-HSL which, in their phosphorylated form, constitute a direct readout of PKA activity (Schweiger
13 et al, 2014) and could be used to semi-quantitatively study levels of lipolysis by IF. Furthermore, we have
14 transferred FAS, DGAT and LPL markers to the mature adipocyte section in Table 1 and, instead, have added *ex*
15 *vivo* BODIPY C1/C12 uptake as an efficient measure of DWAT lipogenesis to Table 1; we have also added
16 BODIPY C1/C12 uptake assay in human abdominal organ culture as an additional supplementary figure (Suppl.
17 Fig. 6a). We also include free glycerol released by adipocytes into media as a readout parameter for lipolysis *ex*
18 *vivo* alongside an example of quantification for this assay (Suppl. Fig 6b). Moreover, in the
19 lipogenesis/lipolysis section of Table 1 we highlight a simple method to measure these parameters *in situ* via
20 quantifying changes in adipocyte size/volume. In addition, we have referred the readers to reviews which
21 explain how to efficiently measure lipogenesis/lipolysis in activity-based assays (Schweiger et al, 2014)
22 (Section ‘Studying lipogenesis/lipolysis in human DWAT’) and now explicitly acknowledge that lipolysis is a
23 complex biological process that involves rapid phosphorylation /dephosphorylation steps so that looking at the
24 total protein levels of enzymes involved in this complex process is uninformative in terms of assessing enzymatic
25 activity (Lass et al, 2011; Nielsen et al, 2014) (section ‘Examining lipogenesis/lipolysis *in situ* and using
26 activity-based techniques’).
27
28
29
30
31
32
33
34
35

36 8) P.7 It would be great to say few words about the timing for the fixation steps. Many people over or under fix
37 their samples, which greatly impairs the staining process.

38 Yes, that makes perfect sense, as fixation time can be critical. We now recommend fixation times in the core
39 text in the ‘Fixation’ section for both paraffin-embedded and cryopreserved tissue. Thank you!

40 9) Generally, it would be important to carefully review the abbreviations used throughout the review. I have
41 seen many abbreviations that are never fully described in the text. FABP4, DAG, MGL, LPL, PPARG, CEBP,
42 CD68, HF, DGAT, FASN, LD, NEFA, PKA...

43 Thank you for pointing out this error. We have now surveyed the entire text and made the appropriate changes.
44 Where dispensable, abbreviations were deleted and words spelled-out in full.
45

46 10) Table 2. The line that describes pCEBPB. Please mention that this protein promotes terminal adipocyte
47 differentiation together with CEBPA (not CEBPD).
48

49 Thank you for drawing attention to this mistake in Table 2 – this has now been amended.
50

51 11) The lipidomics, proteomics and metabolomics sections should be either i) removed or ii) improved. These
52 sections are extremely short and underdeveloped. These sections also appear out of place. Personally, I would
53 take them out.
54
55
56

57
58 The aim of these sections was to present the readers with alternative approaches to study dermal fat in addition
59 to immunohistochemistry, ISH and TEM. We agree that these sections were underdeveloped and we have now
60

1
2
3 removed them from the manuscript, as recommended. In addition, we clarify in the introduction that the review
4 will not discuss techniques associated with lipidomics and metabolomics – instead, we point the readers to a few
5 useful reviews in case they wish to delve into these complex topics after reading our review.
6
7

8 We are genuinely grateful to this expert reviewer for the excellent advice we have received on making our
9 review maximally useful – thank you very much!
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Methods Review

A guide to studying human dermal adipocytes *in situ*

Carina Nicu^{1,2}, Jennifer Pople³, Laura Bonsell^{1,2}, Ranjit Bhogal³ David M Ansell^{1,2} and Ralf Paus^{1,2}

¹ Centre for Dermatology Research, The University of Manchester, Manchester, UK

² NIHR Manchester Biomedical Research Centre and Manchester Academic Health Science Centre, UK

³ Unilever R&D Colworth, Colworth Science Park, Bedford, U.K

Corresponding author: ralf.paus@manchester.ac.uk

Abstract

Dermal white adipose tissue (DWAT) is a main component of human skin, composed of individual lipid-laden mesenchymal cells known as dermal adipocytes (DAs). Besides their well-known role in lipid storage and release, DAs also promote skin immunity, wound healing and hair follicle cycling and are important players in cutaneous neuro-endocrinology. The ever-growing insights into DWAT functions, albeit mostly in mice, have invited speculation that it may be involved in multiple skin diseases ranging from fibrosis to alopecia and psoriasis, thus designating human DWAT a clinically relevant, but as yet insufficiently investigated skin compartment. Therefore, this practical, user-friendly guide aims to introduce the techniques available to study human DWAT *in situ* and *ex vivo*, including immunohistochemistry, immunofluorescence microscopy and analysis via quantitative immunohistomorphometry. Here we provide information on a collection of stains comprising pre-adipocyte (Pref1) and mature adipocyte markers (Perilipin1, Caveolin1), as well as various lipid (OilRedO, BODIPY) and histochemical stains (H&E, Trichrome) available for use on human DWAT. We offer the reader guidelines on fixation, tissue processing and staining human DAs and highlight caveats and solutions to common problems that one may encounter when studying this fascinating skin compartment. We also suggest standard methods for conducting quantitative immunohistomorphometry on human DWAT and its individual adipocytes in order to quantify cell size, number, lipid content and fluorescence intensity of adipose-specific markers. Finally, we briefly introduce *in situ* hybridisation, transmission electron microscopy, and essentials of MRI imaging as additional tools for instructively interrogating this largest, but still least-known compartment of human skin.

Key Words

white adipose tissue; immunohistochemistry; immunofluorescence

Introduction

Historically often viewed as just an inert, energy-storing and thermo-insulating spacer material [1], dermal white adipose tissue (DWAT) is making a resounding scientific comeback as increasingly more studies are emerging which highlight its important and diverse functions in cutaneous biology and beyond [2–9]. It is now recognized that the dermal adipose depot is involved in multiple additional activities, including protection against bacterial infections [4], controlling hair cycle wave patterns [6, 7, 10], thermogenesis [1], promoting wound healing [1, 11–13] and hypothesized neuroendocrine activities [14, 15]. Human DWAT as well as deeper subcutaneous fat layers also harbour heterogeneous, but multi-potent adipose-derived stem cell populations that have been used in the context of cell-based autologous cell therapy, thus finding widespread entry into clinical practice [3, 16–21].

Multiple reviews have appeared on the functions and biology of DWAT in mice [2, 8, 9, 22–25], including reviews on how to image murine adipose tissue [26, 27]. However, there is no comprehensive guide available on staining and imaging techniques of human DAs *in situ*. This constitutes a major open question since the anatomy of human and murine skin adipose layers (and quite likely their physiology) are distinct, calling for independent characterisation.

The present Methods Review attempts to close the literature gap by firstly providing a brief introduction into DA biology in humans. We then present a compendium of histological stains and immunohistochemistry markers available to study human DWAT functions, ranging from lipid droplet biology to distinguishing between pre- and mature adipocytes. We also present data from a primary pre-adipocyte cell line that has potential to differentiate and act as a positive control for DWAT markers. Although we do not discuss key adipocyte techniques associated with lipidomics, metabolomics or proteomics, we direct the readers towards specialized reviews on the subject [28–37]. We aim to provide advice to both enthusiasts and newcomers in the skin adipocyte field.

Human DWAT anatomy and biology in a nutshell

Adipose tissue is found within multiple body sites and can be divided into brown and white fat, the first thought to be involved predominantly in thermogenesis whereas the latter is recognized as a key player in whole-body metabolism [1, 8, 9, 38–40]. An additional type of fat cell exists i.e. beige/bright adipocytes that undergo transdifferentiation from white fat [41–43]. Recently, there has been an important paradigm shift in how “fat” and what used to be called the “subcutis” or “hypodermis” [44] is viewed, since it has turned out that functionally and embryologically distinct layers of adipose tissue are distinguishable in mammalian skin (Fig. 1a) [45].

Whilst murine DWAT is clearly divided from the underlying subcutaneous white adipose tissue by a distinctive muscle layer, the *panniculus carnosus* [46–48], it remains unclear how to morphologically distinguish between human DWAT and the subcutaneous fat layer (Fig. 1a,b) [49]. However, one notable exception described on human cadavers is that of the platysma colli muscle that separates the subcutis into two fat compartments in the neck, the distal (upper) which may be related to human DWAT [50]. Interestingly, human DWAT exhibits a unique cone geometry, clustering around the pilosebaceous units and forming so-called ‘dermal cones’ (Fig. 1c).

1
2
3 DAs typically envelope the connective tissue sheath of hair follicles (HFs) in scalp and abdominal skin, and can
4 stretch from underneath the HF matrix up to eccrine sweat glands and the bulge area of HFs [51, 52] (Fig. 1c).

7 ***Dermal Adipocytes***

8
9 DWAT contains a variety of components including fibroblast-like progenitor adipocytes, mature lipid-laden
10 adipocytes and other cell types such as fibroblasts, macrophages, pericytes, mast cells, endothelial cells,
11 adipose-derived stem cells, as well as an extensive vascular network (Fig 1f) [27]. The main building blocks of
12 DWAT are dermal adipocytes (DAs) – unilocular spherical cells which function to store and release energy via
13 lipogenesis and lipolysis [53]. Their typical shape is the result of the resident lipid droplet, the main site of
14 triglyceride and cholesterol storage which occupies >90% of the cell volume [24]. With their characteristic
15 honeycomb-shape, adipocytes can oscillate in size from 50µm up to 150µm, thus representing the largest cells
16 in the skin. Their cytoplasmic structures and nucleus are compactly located in a thin layer on the inner rim of the
17 cell (Fig 1.d,e).

23 ***Clinical relevance of DWAT***

24
25 At a whole-body level, adipose tissue is a complex endocrine organ system involved in regulating immunity,
26 inflammation, metabolism, insulin sensitivity and overall tissue homeostasis via the secretion of hormones,
27 metabolites and signalling molecules referred to as adipokines [38, 39, 54, 55].

28
29 DAs stimulate wound repair by their own regeneration from myofibroblasts and by promoting fibroblast
30 migration and vessel regeneration [3, 11, 13, 56]. DWAT also defends against *S. aureus* skin infections by
31 producing the antimicrobial peptide Cathelicidin [4], and DAs are thought to play a key role in skin fibrosis via
32 adipocyte-myofibroblast transition [5, 57]. This mechanism is speculated to promote development of fibrotic
33 structures associated with hair follicles in androgenetic alopecia [40], supported by higher plasma leptin levels
34 being associated with an increased incidence of androgenetic alopecia [58].

35
36 Apart from classical dermatoses that affect the adipocyte layers of human skin (e.g., panniculitis,
37 lipogranulomatoses, lipomata, lipoatrophy, lipodystrophy) [59, 60] DAs are likely involved in other cutaneous
38 diseases. Studies have shown fluctuating levels of adipokines in psoriasis patients [61, 62], and the use of
39 therapeutics in this condition has been associated with fluctuations in DWAT thickness [63–65]. Moreover, a
40 recent study supports the role of DAs in promoting melanoma aggressiveness [19].

48 **Overview of visualisation methods to consider**

49
50 When studying DWAT, there are a number of methods that one can choose from. Standard stains such as
51 Haematoxylin and Eosin (H&E) and Masson's Trichrome allow the visualization of adipocyte morphology and
52 any changes in tissue or individual cell expansion or contraction. Lipid stains such as Oil Red O (ORO) and
53 BODIPY can be used to visualize and analyse lipid droplet content, size and accumulation within DAs. Lastly,
54 immunohistochemical/immunofluorescence techniques are widely employed to localize and detect changes in
55 proteins of interest within DWAT by immunohistomorphometry.

Key considerations prior to human DA visualization in situ

The predominant approach to assessing DWAT is via histology/immunohistochemistry, however, prior to staining there are some key considerations to be made (Table 1).

Paraffin-embedded versus cryopreserved tissue

Although paraffin embedding offers excellent morphological detail of DWAT, lipids are displaced during paraffin processing rendering lipid staining protocols impossible. In contrast, cryosectioning offers many advantages including time-efficient protocols and the ability to conduct lipid stains. However, if tissue is scarce and multiple sections are required, sectioning via the microtome is preferable to obtain sufficient thin sections.

Section thickness

When the primary aim is to study lipid content, it is advisable to cut cryopreserved samples at a thickness of 15 – 30 μ m to maintain the cellular lipid content (Suppl Fig. 1). Sections may be quickly visualized using a microscope, though must not be allowed to thaw before storage at -80°C in order to preserve the lipids. For immunohistochemical-based approaches thinner sections are preferred, however, when collecting sections from cryopreserved DWAT, maintaining a temperature equal to or lower than -30°C within the cryostat is crucial to obtain thin (7 μ m) sections of high quality.

Fixation

Choice of fixative and its duration can greatly affect the quality of tissue sections and subsequent staining. Though neutral-buffered formalin (NBF) is the universal fixative used for tissues aimed to undergo paraffin processing, our observations suggest that it is a harsh fixative for DWAT. One alternative fixative is Zinc Formalin which better preserves DA morphology as recommended by [26]. Alternative fixation methods include 4% paraformaldehyde (PFA), 20% normal-buffered formalin (NBF) and 99% ethanol (EtOH), however these have very different effects upon RNA quality, morphology and staining intensities [66]. For fatty samples that are to be embedded in paraffin we recommend 24hr fixation prior to processing.

For cryopreserved tissue, multiple fixatives can be employed including 4% PFA, ethanol, acetone and methanol; however, alcohol-based fixatives should be avoided when the aim is to examine lipid droplet content as these solvents displace lipids [67]. Instead, 4%PFA or ethanol:glacial acetic acid can be successfully used to obtain high-quality lipid stains (Suppl Fig. 2). We recommend performing the fixation step in 4%PFA immediately after thawing the tissue. For cryosections containing fat we have found 20-30min to be an optimal fixation time. Following fixation and sectioning, DWAT can be stained using a range of techniques including immunohistochemistry and immunofluorescence stains. To help our readers, we have compiled a list of suggestions to quickly and easily combat common problems when working with human DWAT (Box 1).

Histochemistry

Standard *haematoxylin and eosin* (H&E) staining may be employed to visualize DWAT structure within skin, and around pilosebaceous units (Fig 1b,c; Fig 2a). H&E is useful when assessing human biopsies because it

1
2
3 shows gross changes in morphology and may point towards further areas for closer examination [68]. For
4 example, one can infer whether human DWAT thickness or the amount of fibroblast networks surrounding DAs
5 changes in samples from healthy, alopecia areata and sebaceous nevus patients [69]. Moreover, H&E can be
6 used to pinpoint whether expansion or reduction occurs within DWAT as highlighted in *S. aureus* infection in
7 mice [4]. Using H&E, the exact distribution and changes in individual cell size and number can be assessed
8 using Adiposoft and Cell Counter plugins in ImageJ (Suppl. Fig 3 a,b).
9
10

11
12 Another histological stain that proves useful when studying adipocytes is *Masson's Trichrome* (Fig 2b) which
13 stains fibrillary collagens I and III surrounding DWAT. Masson's Trichrome can highlight ectopic collagen
14 deposition between individual DAs, a marker of DWAT fibrosis [70], and also infers collagen deposition
15 following organ culture of skin biopsies (Fig. 2b), a simple method to study fibrosis in DWAT [57, 71–73]. An
16 alternative stain for collagen fibrils is *Picrosirius Red* (Fig. 2c).
17
18
19

20
21 *Methylene Blue* (MB) is a vital dye that can be used to stain the skin as a whole [74]. MB stains human skin
22 compartments such as the epidermis, follicular units and the dermis but is not absorbed within DAs apart from
23 in the nucleus [75, 76]. An advantage of MB compared to H&E is that it can be performed as a wholemount
24 technique, allowing the dissection and isolation of DWAT by providing contrast to the surrounding dermis and
25 follicular unit (Fig. 2d).
26
27
28

29 *Lipid stains*

30
31 *Oil red O* (ORO) is a hydrophobic dye used to visualize lipid droplet localization, morphology and
32 accumulation within adipose tissue [67, 77]. ORO stains neutral lipids (triacylglycerols, diacylglycerols) and
33 cholesteryl esters (Fig 2e), but not polar lipids on the lipid droplet membrane such as ceramides, sphingolipids
34 and phospholipids [78]. ORO content in DAs can be quantified using ImageJ as detailed in Suppl. Fig 3c. ORO
35 staining can prove difficult to quantify if sections are too thin (Suppl. Fig 2) or not stored and fixed
36 appropriately following cryosectioning. Precautions should be taken when choosing the reagent in which to
37 dissolve ORO as certain chemicals may result in staining of non-adipogenic cells [79].
38
39
40

41
42 Other chemicals that stain lipid within DAs include Nile Red [80], Sudan Red [81], and Osmium Tetroxide
43 which is routinely used in electron microscopy, but may result in adipocyte swelling [82].
44
45

46
47 In order to quantify changes in lipid accumulation in DAs, fluorescent lipid stains on cryosections such as
48 *BODIPY 493/503* (Fig. 2f) or *LipidTOX* (Suppl. Fig. 1) can be used as an alternative to ORO. BODIPY
49 493/503 and LipidTOX are cell-permeable lipophilic neutral dyes that can be employed for both microscopy and
50 flow cytometry [2, 83–90]. In our studies BODIPY (Fig 2f) appears superior in terms of fluorescence intensity
51 given that LipidTOX rapidly bleaches and does not display the lipid droplet morphology described in other
52 studies (Suppl. Fig. 1) [2, 7].
53
54

55
56 *Cell Mask Plasma Membrane* (CPM) stain (Suppl. Fig. 1) is an amphipathic molecule that anchors into, and
57 histochemically defines, the plasma membrane of human DAs and can aid in visualizing DAs in conjunction
58 with lipid stains. This double stain also allows the correct identification of adipocyte nuclei, located between the
59 lipid droplet and the adipocyte plasma membrane [26].
60

Immunohistochemistry and immunofluorescence microscopy

When staining adipocytes using IHC or immunofluorescence (IF) techniques, various protein markers such as pC/EBP β , PPAR γ 2, Caveolin1, Perilipin1 and Pref1 can be employed to stain progenitor and mature adipocytes (Table 2). Overall, markers including Perilipin1, Pref1 and PPAR γ 2 exclusively stain human DAs. Although sebocytes are cells of the epithelial lineage derived from outer root sheath keratinocytes [91], and embryologically distinct from DAs, they share common antigens. In our stains human sebocytes provide an internal positive control within the skin for adipocyte markers such as phospho-C/EBP β (pC/EBP β), Caveolin1, DGAT1 and FASN.

Pre-adipocyte and mature adipocyte markers

Perilipin1, the ubiquitous marker of mature DAs in mice, has been used in multiple studies to demarcate the lipid droplet and cytoplasm of DWAT [4, 7, 92]. Perilipin1 also marks human DAs and can be assessed in both cryosections and paraffin sections (Fig 2g). Caveolin1, an intracellular scaffolding protein, is abundant in murine adipocytes where it occupies 30% of their plasma membrane, and is involved in lipid uptake [93, 94]. In human skin, Caveolin1 shows the expected pattern in IF and IHC, staining the cell membrane of DAs (Fig. 2h).

Two markers that stain both differentiating pre-adipocytes and mature DAs are Peroxisome proliferator-activated receptor gamma 2 (PPAR γ 2) and CCAAT-enhancer-binding protein beta (C/EBP β) [95]. PPAR γ 2 is the master regulator of adipogenesis [96, 97], and exhibits a nuclear staining pattern (Fig. 2i). C/EBP β is an early marker of pre-adipocytes, and its phosphorylation is crucial for the expression of downstream adipogenic markers such as adiponectin and CCAAT-enhancer-binding protein alpha (C/EBP α) [98]. Phosphorylated C/EBP β (pC/EBP β) shows nuclear staining in human adipocytes (Fig 2j). In contrast, Preadipocyte Factor 1 (Pref1) (Fig. 2k) is a well-known inhibitor of adipocyte differentiation, thus acting as a *bone fide* human pre-adipocyte marker [99–102].

Overall, it is crucial to utilize appropriate positive and negative controls for each marker in order to ensure the staining is reliable (see Table 1 for details on internal and tissue positive controls). For detailed reviews on imaging adipose tissue using fluorescence and confocal microscopy, the reader is directed towards specialized reviews [26, 27].

Analysis by quantitative immunohistomorphometry

Depending on the scientific question asked, various staining techniques (Table 1) as well as adipocyte-specific functional markers (Table 2) can be utilized as described above.

For example, changes in adipocyte size/volume as well as cell number can be quantified via H&E or by using mature adipocyte markers such as Perilipin1 and Caveolin1 using ImageJ (Fig. 2g-h; Suppl Fig. 3a). Alternatively, lipid droplet stains can be employed to quantify adipocyte size, although they require tissue processing via cryopreservation and sectioning at approx. 30 μ m (Table 1). The reader is also directed towards automated techniques of quantifying size and number of individual cells in adipose tissue using software such as MetaMorph, Adiposoft and ImageJ [103, 104].

1
2
3 Changes in lipid content can be inferred from ORO/NileRed/Sudan Black stains or fluorescent
4 BODIPY/LipidTOX stains, quantified also using ImageJ (Fig. 2, Suppl Fig 1,2,3b). An alternative option for
5 measuring adipocyte differentiation in cell lines is by using AdipoRed – a Nile Red kit (Lonza) that is quantified
6 using a microplate reader (Suppl. Fig. 4) [105].
7
8

9 ***Apoptosis markers***

10
11 Cell death in skin adipocytes has historically proven difficult to capture since classical apoptosis markers such
12 as TUNEL and Cleaved Caspase-3 routinely escape detection in murine studies [106, 107]. Given their
13 considerable size and relatively small cytoplasmic area, most DAs in one section will not have a visible nucleus,
14 making reliable quantification difficult (Suppl Fig. 5 a-d). Moreover, distinguishing between nuclei pertaining to
15 mature adipocytes or surrounding cells can also complicate TUNEL staining and its subsequent quantification.
16 We also confirm the lack of TUNEL+ or Cleaved Caspase3+ adipocytes in control human scalp skin (Suppl.
17 Fig. 4 a-d).
18
19

20
21 In obesity adipocyte cell death is driven by necrosis [108] and pyroptosis [109] with macrophages forming
22 crown-like structures around Perilipin1 –ve adipocytes [26, 110]. Thus, loss of Perilipin1 or Caveolin1 (Suppl.
23 Fig. 4e,f) can be used as a marker of adipocyte necrosis, since dying adipocytes lose Perilipin1 and Caveolin1
24 and are surrounded by macrophages [108, 111–113].
25
26

27 ***Proliferation markers***

28
29 Mature adipocytes are post-mitotic under physiological conditions [114]. Any newly-produced adipocytes
30 normally undergo differentiation from resident progenitor cells [115]. Therefore, proliferation markers such as
31 Ki-67 (Fig. 2l) or Proliferating cell nuclear antigen (PCNA) can be employed to quantify proliferation of all
32 resident cells within DWAT, and only estimate the amount of pre-adipocyte proliferation within DWAT (Fig.
33 2l). Rather, a more specific approach is measuring co-localization of pre-adipocyte markers such as Pref1 and
34 Zinc finger protein 521 (Zfp521) (the human homologue of murine Zfp423) with a proliferative marker as
35 described in murine DWAT [4].
36
37

38
39 Other techniques such as EdU and BrdU incorporation can also be utilized to study proliferation within DWAT
40 [7]. To do so in preclinical studies of human skin *ex vivo*, one can resort to full-thickness human skin organ
41 culture [116]. The same techniques and markers for cell cycle analysis *ex vivo* can be employed for human
42 DWAT that we previously described for human hair follicle organ culture [117, 118].
43
44

45 ***Examining lipogenesis/lipolysis in situ and using activity-based techniques***

46
47 Metabolic processes are central to white adipocytes which function to store and release energy [24]. Various key
48 enzymes involved in lipid synthesis and breakdown such as Diacylglycerol O-acyltransferase 1/2 (DGAT1/2)
49 (Fig. 2m), Fatty acid Synthase (FASN) (Fig 2n), lipoprotein lipase (LPL) [119, 120], adipose triglyceride lipase
50 (ATGL), hormone-sensitive lipase (HSL) and monoglyceride lipase (MGL) [121–123] can be used to stain
51 human DAs *in situ* (Fig. 2b). However, lipolysis is a complex biological process that involves rapid
52 phosphorylation/dephosphorylation steps [124–126] during which ATGL, MGL and HSL are regulated at the
53
54
55
56
57
58
59
60

1
2
3 post-transcriptional level [127]. Therefore, quantifying total protein levels of enzymes involved in this complex
4 process is uninformative in terms of assessing enzymatic activity. Furthermore, when studying adipocyte
5 differentiation, the simple presence of lipid cannot attest to the full metabolic capacity of the cell, which can be
6 tested *in situ* using the activity of other enzymes including NADH, NADP, esterase, succinate dehydrogenase
7 and α -glycerol phosphate dehydrogenase [128–130].
8
9

10
11 A number of activity-based assays can be used to study lipogenesis/lipolysis in a more quantitative manner. For
12 instance, lipogenesis can be measured in human DWAT *ex vivo* by using BODIPY C1/C12, a fluorescent fatty
13 acid analogue that is taken up by cells undergoing lipogenesis (Suppl. Fig. 6a) [131–134]. Lipolysis can also be
14 assessed *ex vivo* by measuring levels of glycerol and free fatty acids released into the culture media (Suppl. Fig
15 6b). For more detailed measurements of lipogenesis and lipolysis the reader is directed towards specialised
16 reviews on these topics [127, 135–138].
17
18

19 20 ***In situ* hybridisation**

21
22 When antibody stains for secreted factors prove difficult to optimize, *in situ* hybridization (ISH) can be used to
23 measure messenger RNA concentration within adipose tissue e.g. [6] (Fig. 2o). In addition, when one is not
24 certain whether an immunostain is specific, if ISH detects no corresponding mRNA, then immunoreactivity is
25 likely non-specific. The reader is directed towards the methods review by [139] for detailed protocols on
26 conducting ISH on adipose tissue. A commercially available kit is the RNAscope assay (ACD CellBio) which
27 can be used to conduct ISH on human DAs (Fig. 2o).
28
29
30
31

32 33 **Confocal microscopy**

34
35 In addition to fluorescence and brightfield microscopy, confocal microscopy is a widespread tool used to image
36 adipose tissue either in wholemount, thick sections or adipose cell lines [26, 27, 140]. Confocal microscopy is
37 an advantageous tool because of its enhanced optical resolution and capacity to create serial images throughout
38 5-50 μ m-thick sections, allowing visual 3D tissue reconstruction [141]. Confocal imaging is particularly
39 advantageous to visualize features of interest within human DAs that may otherwise escape detection due to the
40 thin cytoplasm in relation to the lipid droplet (Fig. 2p) [142]. Furthermore, confocal imaging can be used in
41 conjunction with other techniques for a more in-depth study of human DWAT in large skin samples [143].
42
43
44

45 46 **Transmission electron microscopy**

47
48 Using transmission electron microscopy (TEM), one can observe unparalleled detail of intracellular structures
49 (Fig. 2q-s) and quantify changes in adipocytes such as general cellular degeneration, glycogen deposits,
50 cytoplasmic thickness, mitochondrial area and density, extracellular matrix, as well as the presence of crown-
51 like structures reminiscent of adipose pyroptosis [109]. Various transmission and scanning electron microscopy
52 studies have focused on human white adipocytes [144] following liposuction [145], facial autologous graft [146]
53 as well as functional studies of secreted factors [147], white adipocyte de-differentiation into a stem-cell-like
54 state [148] and cell behaviour following adrenergic stimulation [149].
55
56
57
58
59
60

Magnetic Resonance Imaging

Used widely both in clinic and lab-based settings, magnetic resonance imaging (MRI) is a versatile, non-invasive tool for 3D imaging extensively used for studying various fat depots including DWAT [150–155]. MRI could be employed to measure parameters such as thermogenic potential, thickness and true volume of DWAT or individual adipocytes [156, 157], since measuring adipocyte area using 2D methods is prone to error [158, 159]. MRI could also be used to determine the exact division between dermal and subcutaneous adipose compartments in human skin [160], which currently remains uncharacterized. An alternative to MRI would be computed tomography which has also been used for 3D imaging of adipose depots [151, 152, 161–164].

Studying DWAT in human skin organ culture

Although human DWAT tends to degenerate rapidly in standard skin organ culture, it can be kept vital for several days in serum-free William's E medium supplemented with insulin, hydrocortisone and L-glutamine [116]. DWAT viability depends on how freshly the tissue is obtained, how gently it is handled, how fast it is placed into culture and how much trauma is applied during tissue trimming and organ culture set-up. We recommend using 4-6 mm punch biopsies to facilitate tissue perfusion/oxygenation and standardization of the organ-cultured tissue volume (Suppl Fig. 7 a,b). Alternatively, human DWAT can be dissected from scalp/abdominal skin and cultured in the same conditions described above as a mini-organ on its own (Suppl. Fig. 7 c,d). Placing the biopsies onto a pre-hydrated extracellular matrix support structure, such as porcine skin-derived gelatine sponges [165] may prolong DWAT survival *ex vivo*. Tissue viability can be monitored by measuring LDH release into the medium [116].

Dermal adipocyte culture

An alternative to studying human DAs *in situ* is the *in vitro* culture of human white adipocytes [166–169]. However, this removes DAs from the physiological concert of diverse stimuli (systemic, cell-cell, cell-matrix, vascular and neural) that they are exposed to within their normal tissue habitat and subjects them to unphysiological stressors, including the typical hypoxic, reactive oxygen species-generating conditions of cell culture. Therefore, one is well-advised to assume that some results obtained with isolated DAs *in vitro* might represent cell culture artefacts under conditions of oxidative stress. The unphysiological nature of isolated DA studies can be partly overcome by adopting suitable 3D designs [170–175].

One advantage of adipose cell lines over *in situ* analysis is that they provide a model for studying adipogenesis, trans-differentiation [148], formation of lipid droplets [176] or their interaction with other cells in co-culture experiments [10, 177]. In addition, adipocyte cell lines facilitate manipulation of the tissue *ex vivo* that would not otherwise be possible *in situ* such as adding inhibitors/stimulators, performing siRNA knockdowns or measuring secretion of various factors into the culture media.

The most widely-used murine cell line for adipogenesis studies is 3T3-L1 [178], with human pre-adipocytes cell lines developed from various depots across the body including abdominal and gluteal tissue [179], peripheral blood mononuclear cells [180], stromal cell fraction of subcutaneous adipocytes [181, 182] and even liposarcomas (SW872 cells) [183, 184]. A widely used model for studying human fat biology are SGBS pre-

1
2
3 adipocytes [185–187]. Human DAs can also be studied by differentiating white adipocytes from human
4 pluripotent stem cells [188] or by reprogramming human dermal fibroblasts [189].
5

6
7 A useful tool to study human adipocytes *in vitro* are primary subcutaneous pre-adipocytes (Lonza) that are
8 isolated from whole human fat i.e. containing both dermal and subcutaneous pre-adipocytes. These cells have
9 the capacity to differentiate into mature lipid-laden adipocytes (Lonza). Seven days following stimulation,
10 differentiated adipocytes express the same biomarkers seen in mature adipocyte cells such as Perilipin1,
11 Caveolin1 and lipids shown using ORO and BODIPY (Fig. 3). In primary cells, Perilipin1 is expressed seven
12 days following maturation, with no expression observed in pre-adipocytes (Fig. 3). Primary pre-adipocytes
13 commence expressing Caveolin1 after three days in culture, while phospho-CEBP β stains nuclei of both pre-
14 and differentiated adipocytes [95] (Fig. 3). The *bone fide* pre-adipocyte marker Pref1, appears to stain both pre-
15 and mature adipocytes (Fig. 3), highlighting the artificial conditions of adipocyte culture *in vitro*.
16
17
18
19

20
21 Overall, we confirm that these commercially available primary subcutaneous pre-adipocytes are an instructive
22 tool for studying human DAs given that they express the expected markers and can be easily differentiated as an
23 alternative to the other human adipose cell lines available.
24
25

26 **Summary and perspective**

27
28 In this practical guide, we have reviewed the routine methods available to study the human DWAT *in situ* and
29 *ex vivo* including immunohistochemistry, immunofluorescence microscopy and analysis via quantitative
30 immunohistomorphometry. We have discussed some particularly useful stains/markers that cover wide-ranging
31 aspects of human adipocyte biology (e.g. OilRedO, Trichrome, Perilipin1, Caveolin1, BODIPY) and
32 histochemical stains (H&E, Trichrome) available for use on human DWAT, along with guidelines on fixation,
33 tissue processing and quantitative immuno-histomorphometry of human DAs *in situ*. We have also highlighted
34 caveats and solutions to common problems frequently encountered when studying DWAT, including the
35 challenges one faces when assessing adipocyte proliferation and apoptosis. For more advanced analyses, we
36 have briefly covered ISH, TEM, DWAT *ex vivo* and *in vitro* culture, and essentials of MRI imaging of
37 adipocytes”.
38
39
40
41
42

43
44 It is our intention that this guide greatly facilitates future DWAT analysis in human skin and encourages
45 newcomers to this field to join the growing number of investigators that have begun to (re)-appreciate DWAT
46 as one of the last great frontiers in human skin research. Thus, the current guide is meant to complement the rich
47 clinical dermatology and pathology literature on diseases that affect the human subcutis [59, 60].
48
49

50
51 While the existing textbook literature documents that clinicians have long appreciated the clinical importance of
52 subcutaneous disease, skin research into the physiology of human DWAT has yet to be fully dissected.
53 Currently, medical therapy only deals with the clinical consequences of DWAT pathology, without attempting
54 to target DWAT functions and thus impact on the function of skin and its appendages. One notable exception
55 remains the use of adipose-derived autologous stem cells [18, 177, 190]. Hopefully, stimulated by the recent
56 progress that several landmark papers have swept into both murine [3, 191, 192] and human DWAT research
57 [19, 20, 193], this will change in future, with the current guide serving as encouragement and a laboratory
58 companion for furthering these almost entirely mouse-based concepts in the human system.
59
60

1
2
3 **Acknowledgements**
4

5 Writing of the review was supported by a Biotechnology and Biological Science Research Council (BBSRC)
6 Case Studentship (awardee: RP, recipient: C.N) and the NIHR Manchester Biomedical Research Centre,
7 Dermatology Theme.
8
9

10 **Author contributions**
11

12 Photographs and images were provided by C.N. and L.B. The review was written by C.N. with full scientific
13 and writing contributions from all co-authors. The review was designed and edited by RP.
14
15

16 **Conflict of interests**
17

18 The authors declare no conflict of interest. However, for the record, JP and RB are employees of Unilever Plc.,
19 and RP is founder and owner of Monasterium Laboratory, Münster/Germany.
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only

Methods Review

A guide to studying human dermal adipocytes *in situ*

Carina Nicu^{1,2}, Jennifer Pople³, Laura Bonsell^{1,2}, Ranjit Bhogal³ David M Ansell^{1,2} and Ralf Paus^{1,2}

¹ Centre for Dermatology Research, The University of Manchester, Manchester, UK

² NIHR Manchester Biomedical Research Centre and Manchester Academic Health Science Centre, UK

³ Unilever R&D Colworth, Colworth Science Park, Bedford, U.K

Corresponding author: ralf.paus@manchester.ac.uk

Abstract

Dermal white adipose tissue (DWAT) is a main component of human skin, composed of individual lipid-laden mesenchymal cells known as dermal adipocytes (DAs). Besides their well-known role in lipid storage and release, DAs also promote skin immunity, wound healing and hair follicle cycling and are important players in cutaneous neuro-endocrinology. The ever-growing insights into DWAT functions, albeit mostly in mice, have invited speculation that it may be involved in multiple skin diseases ranging from fibrosis to alopecia and psoriasis, thus designating the human DWAT a clinically relevant, but as yet insufficiently investigated skin compartment. Therefore, this practical, user-friendly guide aims to introduce the techniques available to study ~~the~~ human DWAT *in situ* and *ex vivo*, including immunohistochemistry, immunofluorescence microscopy and analysis via quantitative immunohistomorphometry. Here we provide information on a collection of stains comprising pre-adipocyte (Pref1) and mature adipocyte markers (Perilipin1, Caveolin1), as well as various lipid (OilRedO, BODIPY) and histochemical stains (H&E, Trichrome) available for use on human DWAT. We offer the reader guidelines on fixation, tissue processing and staining human DAs and highlight caveats and solutions to common problems that one may encounter when studying this fascinating skin compartment. We also suggest standard methods for conducting quantitative immunohistomorphometry on ~~the~~ human DWAT and its individual adipocytes in order to quantify cell size, number, lipid content and fluorescence intensity of adipose-specific markers. Finally, we also briefly introduce *in situ* hybridisation, transmission electron microscopy, and essentials of DWAT-related “-omics”, as additional tools for instructively interrogating this largest, but still least-known compartment of human skin.

Key Words

white adipose tissue; immunohistochemistry; immunofluorescence

Introduction

Historically often viewed as just an inert, energy-storing and thermo-insulating spacer material [1], dermal white adipose tissue (DWAT) is making a resounding scientific comeback as increasingly more studies are emerging which highlight its important and diverse functions in cutaneous biology and beyond [2–9]. It is now recognized that the dermal adipose depot is involved in multiple additional activities, including protection against bacterial infections [4], controlling hair cycle wave patterns [6, 7, 10], thermogenesis [1], promoting wound healing [1, 11–13] and hypothesized neuroendocrine activities [14, 15]. Human DWAT as well as deeper subcutaneous fat layers also harbour heterogeneous, but multi-potent adipose-derived stem cell populations that have been used in the context of cell-based autologous cell therapy, thus finding widespread entry into clinical practice [3, 16–21].

Multiple reviews have appeared on the functions and biology of DWAT in mice [2, 8, 9, 22–25], including reviews on how to image murine adipose tissue [26, 27]. However, there is no comprehensive guide available on staining and imaging techniques of human DAs *in situ*. This constitutes a major open question since the anatomy of human and murine skin adipose layers (and quite likely their physiology) are distinct, calling for independent characterisation.

The present Methods Review attempts to close the literature gap by firstly providing a brief introduction into DA biology in humans. We then present a compendium of histological stains and immunohistochemistry markers available to study human DWAT functions, ranging from lipid droplet biology to distinguishing between pre- and mature adipocytes. We also present data from a primary pre-adipocyte cell line that has potential to differentiate and act as a positive control for DWAT markers. [Although we do not discuss key adipocyte techniques associated with lipidomics, metabolomics or proteomics, we direct the readers towards specialized reviews on the subject](#) [28–37]. We aim to provide advice to both enthusiasts and newcomers in the skin adipocyte field.

Human DWAT anatomy and biology in a nutshell

Adipose tissue is found within multiple body sites and can be divided into brown and white fat, the first thought to be involved predominantly in thermogenesis whereas the latter is recognized as a key player in whole-body metabolism [1, 8, 9, 38–40]. [An additional type of fat cell exists i.e. beige/bright adipocytes that undergo transdifferentiation from white fat](#) [41–43]. Recently, there has been an important paradigm shift in how “fat” and what used to be called the “subcutis” or “hypodermis” [44] is viewed, since it has turned out that functionally and embryologically distinct layers of adipose tissue are distinguishable in mammalian skin (Fig. 1a) [45].

Whilst murine DWAT is clearly divided from the underlying subcutaneous white adipose tissue by a distinctive muscle layer, the *panniculus carnosus* [46–48], it remains unclear how to morphologically distinguish between human DWAT and the subcutaneous fat layer (Fig. 1a,b) [49]. [However, one notable exception described on human cadavers is that of the platysma colli muscle that separates the subcutis into two fat compartments in the neck, the distal \(upper\) which may be related to human DWAT](#) [50]. Interestingly, human DWAT exhibits a unique cone geometry, clustering around the pilosebaceous units and forming so-called ‘dermal cones’ (Fig. 1c).

1
2
3
4
5
6
7
8 | DAs typically envelope the connective tissue sheath of [hair follicles \(HFs\)](#) -in scalp and abdominal skin, and can
9 stretch from underneath the HF matrix up to eccrine sweat glands and the bulge area of HFs [51, 52] (Fig. 1c).

11 | ***Dermal Adipocytes***

Formatted: Indent: First line: 0 cm

12
13 | DWAT contains a variety of components including fibroblast-like progenitor adipocytes, mature lipid-laden
14 | adipocytes and other cell types such as fibroblasts, macrophages, pericytes, mast cells, endothelial cells,
15 | adipose-derived stem cells, as well as an extensive vascular network (Fig 1f) [27]. The main building blocks of
16 | DWAT are dermal adipocytes (DAs) – unilocular spherical cells which function to store and release energy via
17 | lipogenesis and lipolysis [53]. Their typical shape is the result of the resident lipid droplet, the main site of
18 | triglyceride and cholesterol storage which occupies >90% of the cell volume [24]. With their characteristic
19 | honeycomb-shape, adipocytes can oscillate in size from 50µm up to 150µm, thus representing the largest cells
20 | in the skin. Their cytoplasmic structures and nucleus are compactly located in a thin layer on the inner rim of the
21 | cell (Fig 1.d,e).

24 | ***Clinical relevance of DWAT***

Formatted: Indent: First line: 0 cm

26 | At a whole-body level, adipose tissue is a complex endocrine organ system involved in regulating immunity,
27 | inflammation, metabolism, insulin sensitivity and overall tissue homeostasis via the secretion of hormones,
28 | metabolites and signalling molecules referred to as adipokines [38, 39, 54, 55].

30 | DAs stimulate- wound repair- by their own regeneration from myofibroblasts and by promoting fibroblast
31 | migration and vessel regeneration [3, 11, 13, 56]. DWAT also defends against *S. aureus* skin infections by
32 | producing the antimicrobial peptide Cathelicidin [4], and DAs are thought to play a key role in skin fibrosis via
33 | adipocyte-myofibroblast transition [5, 57]. This mechanism is speculated to promote development of fibrotic
34 | structures associated with hair follicles in androgenetic alopecia [40], supported by higher plasma leptin levels
35 | being associated with an increased incidence of androgenetic alopecia [58].

38 | Apart from classical dermatoses that affect the adipocyte layers of human skin (e.g., panniculitis,
39 | lipogranulomatoses, lipomata, lipoatrophy, lipodystrophy) [59, 60] DAs are likely involved in other cutaneous
40 | diseases. Studies have shown fluctuating levels of adipokines in psoriasis patients [61, 62], and the use of
41 | therapeutics in this condition has been associated with fluctuations in DWAT thickness [63–65]. Moreover, a
42 | recent study supports the role of DAs in promoting melanoma aggressiveness [19].

45 | **Overview of visualisation methods to consider**

47 | When studying DWAT, there are a number of methods that one can choose from. Standard stains such as
48 | Haematoxylin and Eosin (H&E) and Masson's Trichrome allow the visualization of adipocyte morphology and
49 | any changes in tissue or individual cell expansion or contraction. Lipid stains such as Oil Red O (ORO) and
50 | BODIPY can be used to visualize and analyse lipid droplet content, size and accumulation within DAs. Lastly,
51 | immunohistochemical/immunofluorescence techniques are widely employed to localize and detect changes in
52 | proteins of interest within DWAT by immunohistomorphometry.
53 |
54 |
55 |
56 |
57 |
58 |
59 |
60 |

Key considerations prior to human DA visualization in situ

Formatted: Indent: First line: 0 cm

The predominant approach to assessing DWAT is via histology/immunohistochemistry, however, prior to staining there are some key considerations to be made (Table 1).

Paraffin-embedded versus cryopreserved tissue

Although paraffin embedding offers excellent morphological detail of DWAT, lipids are displaced during paraffin processing rendering lipid staining protocols impossible. In contrast, cryosectioning offers many advantages including time-efficient protocols and the ability to conduct lipid stains. However, if tissue is scarce and multiple sections are required, sectioning via the microtome is preferable to obtain sufficient thin sections.

Section thickness

Formatted: Indent: First line: 0 cm

When the primary aim is to study lipid content, it is advisable to cut cryopreserved samples at a thickness of 15 – 30µm to maintain the cellular lipid content (Suppl Fig. 1). Sections may be quickly visualized using a microscope, though must not be allowed to thaw before storage at -80°C in order to preserve the lipids. For immunohistochemical-based approaches thinner sections are preferred, ~~though~~ however, when collecting sections from cryopreserved DWAT, maintaining a temperature equal to or lower than -30°C within the cryostat is crucial to obtain thin (7µm) sections of high quality.

Fixation

Formatted: Indent: First line: 0 cm

Choice of fixative and its duration can greatly affect the quality of tissue sections and subsequent staining. Though neutral-buffered formalin (NBF) is the universal fixative used for tissues aimed to undergo paraffin processing, our observations suggest that it is a harsh fixative for DWAT. One alternative fixative is Zinc Formalin which better preserves DA morphology as recommended by [26]. Alternative fixation methods include 4% paraformaldehyde (PFA), 20% normal-buffered formalin (NBF) and 99% ethanol (EtOH), however these have very different effects upon RNA quality, morphology and staining intensities [66]. For fatty samples that are to be embedded in paraffin we recommend 24hr fixation prior to processing.

For cryopreserved tissue, multiple fixatives can be employed including 4% PFA, ethanol, acetone and methanol; however, alcohol-based fixatives should be avoided when the aim is to examine lipid droplet content as these solvents displace lipids [67]. Instead, 4%PFA or ethanol:glacial acetic acid can be successfully used to obtain high-quality lipid stains (Suppl Fig. 2). We recommend performing the fixation step in 4%PFA immediately after thawing the tissue. For cryosections containing fat we have found 20-30min to be an optimal fixation time.

Following fixation and sectioning, DWAT can be stained using a range of techniques including immunohistochemistry and immunofluorescence stains. To help our readers, we have compiled a list of suggestions to quickly and easily combat common problems when working with human DWAT (Box 1).

Histochemistry

1
2
3
4
5
6
7
8 Standard *haematoxylin and eosin* (H&E) staining may be employed to visualize DWAT structure within skin,
9 and around pilosebaceous units (Fig 1b,c; Fig 2a). H&E is useful when assessing human biopsies because it
10 shows gross changes in morphology and may point towards further areas for closer examination [68]. For
11 example, one can infer whether human DWAT thickness or the amount of fibroblast networks surrounding DAs
12 changes in samples from healthy, alopecia areata and sebaceous nevus patients [69]. Moreover, H&E can be
13 used to pinpoint whether expansion or reduction occurs within DWAT as highlighted in *S. aureus* infection in
14 mice [4]. Using H&E, the exact distribution and changes in individual cell size and number can be assessed
15 using Adiposoft and Cell Counter plugins in ImageJ (Suppl. Fig 3 a,b).
16
17

18 Another histological stain that proves useful when studying adipocytes is *Masson's Trichrome* (Fig 2b) which
19 stains fibrillary collagens I and III surrounding DWAT. Masson's Trichrome can highlight ectopic collagen
20 deposition between individual DAs, a marker of DWAT fibrosis [70], and also infers collagen deposition
21 following organ culture of skin biopsies (Fig. 2b), a simple method to study fibrosis in DWAT [57, 71–73]. An
22 alternative stain for collagen fibrils is *Picrosirius Red* (Fig. 2c).
23
24

25 *Methylene Blue* (MB) is a vital dye that can be used to stain the skin as a whole [74]. MB stains human skin
26 compartments such as the epidermis, follicular units and the dermis but is not absorbed within DAs apart from
27 in the nucleus [75, 76]. An advantage of MB compared to H&E is that it can be performed as a wholemount
28 technique, allowing the dissection and isolation of DWAT by providing contrast to the surrounding dermis and
29 follicular unit (Fig. 2d).
30
31

32 *Lipid stains*

33 *Oil red O* (ORO) is a hydrophobic dye used to visualize lipid droplet localization, morphology and
34 accumulation within adipose tissue [67, 77]. ORO stains neutral lipids (triacylglycerols, diacylglycerols) and
35 cholesteryl esters (Fig 2e), but not polar lipids on the lipid droplet membrane such as ceramides, sphingolipids
36 and phospholipids [78]. ORO content in DAs can be quantified using ImageJ as detailed in Suppl. Fig 3c. ORO
37 staining can prove difficult to quantify if sections are too thin (Suppl. Fig 2) or not stored and fixed
38 appropriately following cryosectioning. Precautions should be taken when choosing the reagent in which to
39 dissolve ORO as certain chemicals may result in staining of non-adipogenic cells [79].
40
41

42 Other chemicals that stain lipid within DAs include Nile Red [80], Sudan Red [81], and Osmium Tetroxide
43 which is routinely used in electron microscopy, but may result in adipocyte swelling [82].
44
45

46 In order to quantify changes in lipid accumulation in DAs, fluorescent lipid stains on cryosections such as
47 *BODIPY 493/503* (Fig. 2f) or *LipidTOX* (Suppl. Fig. 1) can be used as an alternative to ORO. BODIPY
48 493/503 and LipidTOX are cell-permeable lipophilic neutral dyes that can be employed for both microscopy and
49 flow cytometry [2, 83–90]. In our studies BODIPY (Fig 2f) appears superior in terms of fluorescence intensity
50 given that LipidTOX rapidly bleaches and does not display the lipid droplet morphology described in other
51 studies (Suppl. Fig. 1) [2, 7].
52
53

54 *Cell Mask Plasma Membrane* (CPM) stain (Suppl. Fig. 1) is an amphipathic molecule that anchors into, and
55 histochemically defines, the plasma membrane of human DAs and can aid in visualizing DAs in conjunction
56
57
58
59
60

with lipid stains. This double stain also allows the correct identification of adipocyte nuclei, located between the lipid droplet and the adipocyte plasma membrane [26].

Immunohistochemistry and immunofluorescence microscopy

When staining adipocytes using IHC or immunofluorescence (IF) techniques, various protein markers such as pC/EBP β , PPAR γ 2, Caveolin1, Perilipin1 and Pref1 can be employed to stain progenitor and mature adipocytes (Table 2). Overall, markers including Perilipin1, Pref1 and PPAR γ 2 exclusively stain human DAs. Although sebocytes are cells of the epithelial lineage derived from outer root sheath keratinocytes [91], and embryologically distinct from DAs, they share common antigens. In our stains human sebocytes provide an internal positive control within the skin for adipocyte markers such as phospho-C/EBP β (pC/EBP β), Caveolin1, DGAT1 and FASN.

Pre-adipocyte and mature adipocyte markers

Perilipin1, the ubiquitous marker of mature DAs in mice, has been used in multiple studies to demarcate the lipid droplet and cytoplasm of DWAT [4, 7, 92]. Perilipin1 also marks human DAs and can be assessed in both cryosections and paraffin sections (Fig 2g). Caveolin1, an intracellular scaffolding protein, is abundant in murine adipocytes where it occupies 30% of their plasma membrane, and is involved in lipid uptake [93, 94]. In human skin, Caveolin1 shows the expected pattern in IF and IHC, staining the cell membrane of DAs (Fig. 2h).

Two markers that stain both differentiating pre-adipocytes and mature DAs are [Peroxisome proliferator-activated receptor gamma 2 \(PPAR \$\gamma\$ 2\)](#) and [CCAAT-enhancer-binding protein beta \(C/EBP \$\beta\$ \)](#) [95]. PPAR γ 2 is the master regulator of adipogenesis [96, 97], and exhibits a nuclear staining pattern (Fig. 2i). C/EBP β is an early marker of pre-adipocytes, and its phosphorylation is crucial for the expression of downstream adipogenic markers such as adiponectin and [CCAAT-enhancer-binding protein alpha \(C/EBP \$\alpha\$ \)](#) [98]. [Phosphorylated C/EBP \$\beta\$ \(pC/EBP \$\beta\$ \)](#) shows nuclear staining in human adipocytes (Fig 2j). In contrast, [Preadipocyte Factor 1 \(Pref1\)](#) (Fig. 2k) is a well-known inhibitor of adipocyte differentiation, thus acting as a *bone fide* human pre-adipocyte marker [99–102].

Overall, it is crucial to utilize appropriate positive and negative controls for each marker in order to ensure the staining is reliable (see Table 1 for details on internal and tissue positive controls). For detailed reviews on imaging adipose tissue using fluorescence and confocal microscopy, the reader is directed towards [specialized the reviews of](#) [26, 27].

Analysis by quantitative immunohistomorphometry

Depending on the scientific question asked, various staining techniques (Table 1) as well as adipocyte-specific functional markers (Table 2) can be utilized as described above.

For example, changes in adipocyte size/volume as well as cell number can be quantified via H&E or by using mature adipocyte markers such as Perilipin1 and Caveolin1 using ImageJ (Fig. 2g-h; Suppl Fig. 3a). Alternatively, lipid droplet stains can be employed to quantify adipocyte size, although they require tissue processing via cryopreservation and sectioning at approx. 30 μ m (Table 1). The reader is also directed towards

Formatted: Indent: First line: 0 cm

automated techniques of quantifying size and number of individual cells in adipose tissue using software such as MetaMorph, Adiposoft and ImageJ [103, 104].

Changes in lipid content can be inferred from ORO/NileRed/Sudan Black stains or fluorescent BODIPY/LipidTOX stains, quantified also using ImageJ (Fig. 2, Suppl Fig 1,2,3b). An alternative option for measuring adipocyte differentiation in cell lines is by using AdipoRed – a Nile Red kit (Lonza) that is quantified using a microplate reader (Suppl. Fig. 4) [105].

Apoptosis markers

Cell death in skin adipocytes has historically proven difficult to capture since classical apoptosis markers such as TUNEL and Cleaved Caspase-3 routinely escape detection in murine studies [106, 107]. Given their considerable size and relatively small cytoplasmic area, most DAs in one section will not have a visible nucleus, making reliable quantification difficult (Suppl Fig. 5 a-d). Moreover, distinguishing between nuclei pertaining to mature adipocytes or surrounding cells can also complicate TUNEL staining and its subsequent quantification. We also confirm the lack of TUNEL+ or Cleaved Caspase3+ adipocytes in control human scalp skin (Suppl. Fig. 4 a-d).

In obesity adipocyte cell death in obesity is driven by necrosis [108] and pyroptosis [109] with macrophages forming crown-like structures around Perilipin1 –ve adipocytes [26, 110]. Therefore, the best method of visualizing DA cell death is by using macrophage markers such as CD68 or Mac 2 [27]. Alternatively, Thus, loss of Perilipin1 or Caveolin1 (Suppl. Fig. 4e,f) can be used as a marker of adipocyte necrosis, since dying adipocytes lose Perilipin1 and Caveolin1 and are surrounded by macrophages [108, 111–113].

Proliferation markers

Mature adipocytes are post-mitotic under physiological conditions [114]. Any newly-produced adipocytes normally undergo differentiation from resident progenitor cells [115]. Therefore, proliferation markers such as Ki-67 (Fig. 2l) or Proliferating cell nuclear antigen (PCNA) can be employed to quantify proliferation of all resident cells within DWAT, and only estimate the amount of pre-adipocyte proliferation within DWAT (Fig. 2l). Rather, a more specific approach is measuring co-localization of pre-adipocyte markers such as Pref1 and Zinc finger protein 521 (Zfp521) (the human homologue of murine Zfp423) with a proliferative marker as described in murine DWAT [4].

Other techniques such as EdU and BrdU incorporation can also be utilized to study proliferation within DWAT [7]. To do so in preclinical studies of human skin *ex vivo*, one can resort to full-thickness human skin organ culture [116]. The same techniques and markers for cell cycle analysis *ex vivo* can be employed for human DWAT that we previously described for human hair follicle organ culture [117, 118].

Enzyme histochemistry for important lipid metabolism/synthesis enzymes

Examining lipogenesis/lipolysis *in situ* and using activity-based techniques

Formatted: Indent: First line: 0 cm

Formatted: Indent: First line: 0 cm

Formatted: Font: Italic

Metabolic processes are central to white adipocytes which function to store and release energy [24]. Various key enzymes involved in lipid synthesis and breakdown such as [Diacylglycerol O-acyltransferase 1/2 \(DGAT1/2\)](#) (Fig. 2m), [Fatty acid Synthase \(FASN\)](#) (Fig 2n), [lipoprotein lipase \(LPL\)](#) [119, 120], ~~and~~ [adipose triglyceride lipase \(ATGL\)](#), [hormone-sensitive lipase \(HSL\)](#) and [monoglyceride lipase \(MGL\)](#) [121–123] can be used to stain human DAs *in situ* (Fig. 2b). ~~These stains are incredibly useful because they can semi-quantitatively determine enzyme activity *in situ*.~~ However, lipolysis is a complex biological process that involves rapid phosphorylation/dephosphorylation steps [124–126] during which ATGL, MGL and HSL are regulated at the post-transcriptional level [127]. ~~Therefore, quantifying total protein levels of enzymes involved in this complex process is uninformative in terms of assessing enzymatic activity.~~ Furthermore, ~~W~~hen studying adipocyte differentiation, the simple presence of lipid cannot attest to the full metabolic capacity of the cell, which can be tested *in situ* using the activity of other enzymes including NADH, NADP, esterase, succinate dehydrogenase and α -glycerol phosphate dehydrogenase [128–130].

Comment [CN1]: Re-write this section!

~~A number of activity-based assays can be used to study lipogenesis/lipolysis in a more quantitative manner. For instance, lipogenesis can be measured in human DWAT *ex vivo* by using BODIPY C1/C12, a fluorescent fatty acid analogue that is taken up by cells undergoing lipogenesis (Suppl. Fig. 6a) [131–134]. Lipolysis can also be assessed *ex vivo* by measuring levels of glycerol and free fatty acids released into the culture media (Suppl. Fig 6b). For more detailed measurements of lipogenesis and lipolysis the reader is directed towards specialised reviews on these topics [127, 135–138].~~

Formatted: Font: Italic

Formatted: Font: Italic

***In situ* hybridisation**

When antibody stains for secreted factors prove difficult to optimize, *in situ* hybridization (ISH) can be used to measure messenger RNA concentration within adipose tissue e.g. [6] (Fig. 2o). In addition, when one is not certain whether an immunostain is specific, if ISH detects no corresponding mRNA, then immunoreactivity is likely non-specific. The reader is directed towards the methods review by [139] for detailed protocols on conducting ISH on adipose tissue. A commercially available kit is the RNAscope assay (ACD CellBio) which can be used ~~to conduct for~~ ISH on human DAs (Fig. 2o).

Confocal microscopy

In addition to fluorescence and brightfield microscopy, confocal microscopy is a widespread tool used to image adipose tissue either in wholemount, thick sections or adipose cell lines [26, 27, 140]. Confocal microscopy is an advantageous tool because of its enhanced optical resolution and capacity to create serial images throughout 5-50 μ m-thick sections, allowing visual 3D tissue reconstruction [141]. Confocal imaging is particularly advantageous to visualize features of interest within human DAs that may otherwise escape detection due to the thin cytoplasm in relation to the lipid droplet (Fig. 2p) [142]. Furthermore, confocal imaging can be used in conjunction with other techniques for a more in-depth study of human DWAT in large skin samples [143].

Transmission electron microscopy

Using transmission electron microscopy (TEM), one can observe unparalleled detail of intracellular structures (Fig. 2q-s) and quantify changes in adipocytes such as general cellular degeneration, glycogen deposits, cytoplasmic thickness, mitochondrial area and density, extracellular matrix, as well as the presence of crown-like structures reminiscent of adipose pyroptosis [109]. Various transmission and scanning electron microscopy studies have focused on human white adipocytes [144] following liposuction [145], facial autologous graft [146] as well as functional studies of secreted factors [147], white adipocyte de-differentiation into a stem-cell-like state [148] and cell behaviour following adrenergic stimulation [149].

Magnetic Resonance Imaging

Used widely both in clinic and lab-based settings, magnetic resonance imaging (MRI) is a versatile, non-invasive tool for 3D imaging extensively used for studying various fat depots including DWAT [150–155]. MRI could be employed to measure parameters such as thermogenic potential, thickness and true volume of DWAT or individual adipocytes [156, 157], since measuring adipocyte area using 2D methods is prone to error [158, 159]. MRI could also be used to determine the exact division between dermal and subcutaneous adipose compartments in human skin [160], which currently remains uncharacterized. An alternative to MRI would be computed tomography which has also been used for 3D imaging of adipose depots [151, 152, 161–164].

Studying DWAT in human skin organ culture

Although human DWAT tends to degenerate rapidly in standard skin organ culture, it can be kept vital for several days in serum-free William's E medium supplemented with insulin, hydrocortisone and L-glutamine [116]. DWAT viability depends on how freshly the tissue is obtained, how gently it is handled, how fast it is placed into culture and how much trauma is applied during tissue trimming and organ culture set-up. We recommend using 4-6 mm punch biopsies to facilitate tissue perfusion/oxygenation and standardization of the organ-cultured tissue volume (Suppl Fig. 76 a,b). Alternatively, human DWAT can be dissected from scalp/abdominal skin and cultured in the same conditions described above as a mini-organ on its own (Suppl. Fig. 76 c,d). Placing the biopsies onto a pre-hydrated extracellular matrix support structure, such as porcine skin-derived gelatine sponges [165] may prolong DWAT survival *ex vivo*. Tissue viability can be monitored by measuring LDH release into the medium [116].

Dermal adipocyte culture

An alternative to studying human DAs *in situ* is the *in vitro* culture of human white adipocytes [166–169]. However, this removes DAs from the physiological concert of diverse stimuli (systemic, cell-cell, cell-matrix, vascular and neural) that they are exposed to within their normal tissue habitat and subjects them to unphysiological stressors, including the typical hypoxic, reactive oxygen species-generating conditions of cell culture. Therefore, one is well-advised to assume that some results obtained with isolated DAs *in vitro* might represent cell culture artefacts under conditions of oxidative stress. The unphysiological nature of isolated DA studies can be partly overcome by adopting suitable 3D designs [170–175].

One advantage of adipose cell lines over *in situ* analysis is that they provide a model for studying adipogenesis, trans-differentiation [148], formation of lipid droplets [176] or their interaction with other cells in co-culture

experiments [10, 177]. In addition, adipocyte cell lines facilitate manipulation of the tissue *ex vivo* that would not otherwise be possible *in situ* such as adding inhibitors/stimulators, performing siRNA knockdowns or measuring secretion of various factors into the culture media.

The most widely-used murine cell line for adipogenesis studies is 3T3-L1 [178], with human pre-adipocytes cell lines developed from various depots across the body including abdominal and gluteal tissue [179], peripheral blood mononuclear cells [180], stromal cell fraction of subcutaneous adipocytes [181, 182] and even liposarcomas (SW872 cells) [183, 184]. A widely used model for studying human fat biology are SGBS pre-adipocytes [185–187]. Human DAs can also be studied by differentiating white adipocytes from human pluripotent stem cells [188] or by reprogramming human dermal fibroblasts [189].

A useful tool to study human adipocytes *in vitro* are primary subcutaneous pre-adipocytes (Lonza) that are isolated from whole human fat i.e. containing both dermal and subcutaneous pre-adipocytes. These cells which have the capacity to differentiate into mature lipid-laden adipocytes (Lonza). Seven days following stimulation, differentiated adipocytes express the same biomarkers seen in mature adipocyte cells such as Perilipin1, Caveolin1 and lipids shown using ORO and BODIPY (Fig. 3). In primary cells, Perilipin1 is expressed seven days following maturation, with no expression observed in pre-adipocytes (Fig. 3). Primary pre-adipocytes commence expressing Caveolin1 after three days in culture, while phospho-CEBP β stains nuclei of both pre- and differentiated adipocytes [95] (Fig. 3). The *bone fide* pre-adipocyte marker Pref1, appears to stain both pre- and mature adipocytes (Fig. 3), highlighting the artificial conditions of adipocyte culture *in vitro*.

Overall, we confirm that these commercially available primary subcutaneous pre-adipocytes are an instructive tool for studying human DAs given that they express the expected markers and can be easily differentiated as an alternative to the other human adipose cell lines available.

Summary and perspective

In this practical guide, we have reviewed the routine methods available to study the human DWAT *in situ* and *ex vivo* including immunohistochemistry, immunofluorescence microscopy and analysis via quantitative immunohistomorphometry. We have discussed some particularly useful stains/markers that cover wide-ranging aspects of human adipocyte biology (e.g. OilRedO, Trichrome, Perilipin1, Caveolin1, BODIPY) and histochemical stains (H&E, Trichrome) available for use on human DWAT, along with guidelines on fixation, tissue processing and quantitative immuno-histomorphometry of human DAs *in situ*. We have also highlighted caveats and solutions to common problems frequently encountered when studying DWAT, including the challenges one faces when assessing adipocyte proliferation and apoptosis. For more advanced analyses, we have briefly covered I¹H, TEM, DWAT *ex vivo* and *in vitro* culture, and essentials of MRI imaging of adipocytes of DWAT-related “-omics”.

It is our intention that this guide greatly facilitates future DWAT analysis in human skin and encourages newcomers to this field to join the growing number of investigators that have begun to (re-)appreciate the DWAT as one of the last great frontiers in human skin research. Thus, the current guide is meant to complement the rich clinical dermatology and pathology literature on diseases that affect the human subcutis [59, 60].

Formatted: Font: Not Italic

1
2
3
4
5
6
7
8 While the existing textbook literature documents that clinicians have long appreciated the clinical importance of
9 subcutaneous disease, skin research into the physiology of ~~the~~ human DWAT has yet to be fully dissected.
10 Currently, medical therapy only deals with the clinical consequences of DWAT pathology, without attempting
11 to target DWAT functions and thus impact on the function of skin and its appendages. One notable exception
12 remains the use of adipose-derived autologous stem cells [18, 177, 190]. Hopefully, stimulated by the recent
13 progress that several landmark papers have swept into both murine [3, 191, 192] and human DWAT research
14 [19, 20, 193], this will change in future, with the current guide serving as encouragement and a laboratory
15 companion for furthering these almost entirely mouse-based concepts in the human system.
16
17
18
19

20 **Acknowledgements**

21
22 Writing of the review was supported by a Biotechnology and Biological Science Research Council (BBSRC)
23 Case Studentship (awardee: RP, recipient: C.N) and the NIHR Manchester Biomedical Research Centre,
24 Dermatology Theme.
25

26 **Author contributions**

27
28 Photographs and images were provided by C.N. and L.B. The review was written by C.N. with full scientific
29 and writing contributions from all co-authors. The review was designed and edited by RP.
30

31 **Conflict of interests**

32
33 The authors declare no conflict of interest. However, for the record, JP and RB are employees of Unilever Plc.,
34 and RP is founder and owner of Monasterium Laboratory, Münster/Germany.
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1. Structure of dermal white adipose tissue and its components. Whilst murine DWAT is divided from subcutaneous adipocytes by the *panniculus carnosus*, human DWAT is found as a continuous layer in the human dermis (a) with no equivalent muscle structure separating it from subcutaneous white adipose tissue (SWAT) (a). DWAT surrounds the bottom half of hair follicles in the form of so-called 'dermal cones' (b-c). At the ultrastructural level (d-e), the cellular components of dermal adipocytes become clearer – most of the cell volume is occupied by the centrally-located lipid droplet (LD) which is surrounded by a thin sheath of cytoplasm (represented between the dotted lines); the cytoplasm is enveloped by the plasma membrane on the outside. DWAT is composed of various cell types including mature adipocytes, fibroblast-like pre-adipocytes, macrophages, pericytes, endothelial cells and a multitude of blood vessels (f). Scale bar is 2μm in (d) and 500nm in (e).




Figure 2. Stains for observing and quantifying human dermal white adipose tissue *in situ*. Adipocytes can be examined using histological stains such as (a) Haematoxylin and Eosin (H&E), (b) Masson's Trichrome, (b) Picrosirius Red and (d) Methylene blue (MB). The lipid within DAs can be measured using (e) Oil Red O (ORO), or fluorescently via (f) BODIPY493/503. Furthermore, various markers are available to study pre- and mature adipocytes including (g) Perilipin1 (PLIN1), (h) Caveolin1 (CAV1), (i) PPAR γ 2, (j) p-C/EBP β and (k) PREF-1. In addition, proliferation (l) via Ki67 as well as lipid synthesis enzymes such as (m) diacylglycerol acetyl transferase (DGAT1) and (n) fatty acid synthase (FASN) can be used to stain DAs *in situ*. Alternative methods of studying adipocyte *in situ* include (o) *in situ* hybridization (ISH) to study synthesis of individual strands of mRNA, and various cellular processes at the ultrastructural level (q-s) using transmission electron microscopy (TEM). Scale bar is 50 μ m across all images except the last two TEM images.




Figure 3. Human primary subcutaneous adipocytes as a model for studying human DAs *in vitro*. Primary pre-adipocytes can be differentiated over 7 days into differentiated adipocytes that express PLIN1, p-CEBP β , CAV1 and PREF1. Lipid stains such as Oil Red O and BODIPY also stain differentiated adipocytes. Thus, primary cells can also serve as positive and negative controls for antibody stains conducted *in situ*. Scale bar is 50 μ m.

Table 1. A guide to choosing the right processing route and stain for analysing dermal adipocytes.

LD – lipid droplet; BC – breast carcinoma; BF – brightfield; CFM –confocal microscopy; DA – dermal adipocyte; FB – fibroblasts; FFPE – formalin fixed paraffin embedded; H&E – haematoxylin and eosin; IFM – immunofluorescence microscopy; KC - keratinocyte; LD – lipid droplet; MP – macrophage; PC- prostate carcinoma; SC – stratum corneum; SG – sebaceous gland; TEM – transmission electron microscopy

<u>What should I choose when studying...?</u>	<u>How to process the tissue</u>	<u>Stains/Markers</u>	<u>Target / Antigen Localization</u>	<u>Positive Control Tissue</u>	<u>Imaging and Analysis</u>	<u>References</u>
<u>DWAT Morphology</u>	<u>FFPE Cryo</u>	<u>H&E</u>	<u>n/a</u>	<u>n/a</u>	<u>BF</u>	[68, 69, 194]
<u>Lipid droplet content and Morphology</u>	<u>Cryo</u>	<u>Oil Red O, Nile Red</u>	<u>LD</u>	<u>SG, SC, Breast</u>	<u>BF</u>	[3, 7, 105, 115, 140, 148, 183, 194–196]
		<u>BODIPY, LipidTOX</u>	<u>LD</u>	<u>SG, SC, Breast</u>	<u>IFM CFM</u>	
<u>Mature Adipocyte Adipocyte size and number</u>	<u>FFPE Cryo</u>	<u>PLIN1</u>	<u>LD Membrane</u>	<u>SC, BC</u>		[7, 101, 108, 163-166, 169-173]
		<u>CAV1</u>	<u>Cell membrane</u>	<u>KCs, FBs, HeLa Cells</u>	<u>BF/IFM</u>	
		<u>FABP4</u>	<u>Cytoplasm</u>	<u>KCs, Lymph node</u>		
		<u>CPM</u>	<u>Cell membrane</u>	<u>All cells, Any tissue</u>	<u>IFM</u>	
		<u>H&E</u>	<u>n/a</u>	<u>n/a</u>	<u>BF</u>	
		<u>DGAT1/2</u>	<u>Cytoplasm - ER</u>	<u>KCs, Colon</u>		
		<u>FASN</u>	<u>Cytoplasm</u>	<u>KCs, PC, BC</u>	<u>BF/IFM</u>	
		<u>LPL</u>	<u>Membrane</u>	<u>Intestine</u>		
<u>Adipocyte differentiation</u>	<u>FFPE Cryo</u>	<u>pC/EBPβ</u>	<u>Nucleus</u>	<u>SG, BC</u>	<u>BF/IFM</u>	[7, 195]
		<u>PPARγ</u>	<u>Nucleus</u>	<u>SG, BC</u>		
<u>Pre-adipocytes</u>	<u>FFPE Cryo</u>	<u>Pref1</u>	<u>Membrane / Nucleus</u>	<u>Placenta</u>	<u>BF/IFM</u>	[195, 197]
		<u>Zfp521</u>	<u>Nucleus</u>	<u>Bone</u>		
<u>Adipocyte apoptosis</u>	<u>FFPE Cryo</u>	<u>PLIN1</u>	<u>LD Membrane</u>	<u>SC, BC</u>	<u>Loss of Perilipin1</u>	[27, 194, 198]
		<u>CAV1</u>	<u>Cell membrane</u>	<u>KCs, FBs, HeLa Cells</u>	<u>Loss of Caveolin1</u>	
<u>Pre-adipocyte proliferation</u>	<u>FFPE Cryo</u>	<u>Pref1 + Ki67/ BrdU/ EdU</u>	<u>Nucleus</u>	<u>Placenta</u>	<u>BF/IFM</u>	[7, 115, 195]
<u>Lipid synthesis (lipogenesis)</u>	<u>Unfixed tissue</u>	<u>BODIPY C1/C12</u>	<u>Cytoplasm - ER</u>	<u>Adipocytes</u>	<u>IFM</u>	[132, 134, 199, 200]
	<u>FFPE Cryo</u>	<u>Mature DA stain to measure change in size (see above)</u>	<u>Various</u>	<u>Dependent on stain chosen</u>	<u>BF/IFM</u>	<u>See section above</u>

<u>Lipid metabolism (lipolysis)</u>	<u>FFPE</u> <u>Cryo</u>	<u>phospho-Perilipin1</u>	<u>Cytoplasm</u>	<u>Adipocyte cell line stimulated to undergo lipolysis</u>	<u>BF/IFM</u>	[127, 201, 202]
		<u>phospho-HSL</u>	<u>Cytoplasm</u>			
<u>Adipocyte Ultrastructure</u>	<u>EM-specific</u>	<u>OsO4 Uranyl Acetate</u>	<u>Only for immuno-EM</u>	<u>n/a</u>	<u>TEM</u>	[203],[109, 144, 148, 193, 204],[176]
<u>3D Volume and Thickness</u>	<u>As per MRI/μCT protocol</u>	<u>Various</u>	<u>n/a</u>	<u>n/a</u>	<u>MRI</u> <u>μCT</u>	[150–157, 160–164]
<u>mRNA</u>	<u>FFPE</u> <u>Cryo</u>	<u>Fluorescent or chromogenic ISH</u>	<u>Cytoplasm</u>	<u>Dependent upon gene of interest</u> <u>PPIB; HeLa cells</u>	<u>ISH</u>	[139] [205]
<u>Fibrosis</u>	<u>FFPE</u>	<u>Masson's Trichrome/ Picrosirius Red</u>	<u>Fibrillary Collagens I and III</u>	<u>n/a</u>	<u>BF</u>	[70, 206, 207]
		<u>Methylene Blue</u>	<u>Dermal components other than DAs</u>	<u>n/a</u>		

Review Only

Table 2. Markers of human dermal adipocytes and their functional importance.

<u>Marker</u>	<u>Target Localization</u>	<u>Function</u>	<u>Murine References</u>	<u>Human References</u>
<u>Perilipin1</u>	<u>LD membrane ER</u>	<u>Master regulator of lipolysis</u> <u>Upon phosphorylation by PKA, Perilipin1 allows HSL to translocate to the adipocyte lipid droplet and commence lipolysis</u>	[7, 69, 140, 195, 196, 207, 208] [209]	[102, 170, 176, 210, 211]
<u>Caveolin1</u>	<u>Plasma membrane</u>	<u>Endo- and exocytosis</u> <u>Component of caveolae which are sites of triacylglycerol synthesis</u>	[212, 213]	[214]
<u>PPARγ2</u>	<u>Nucleus</u>	<u>Master regulator of adipogenesis</u>	[95]	[102, 215]
<u>pC/EBPβ</u>	<u>Nucleus</u>	<u>Promotes terminal adipocyte differentiation together with C/EBPα</u>	[216]	[217, 218]
<u>Pref1/Dik1</u>	<u>Plasma Membrane</u>	<u>Inhibits pre-adipocyte proliferation in its membrane-bound form</u> <u>Prevents adipocyte differentiation (adipogenesis)</u>	[99, 219–223]	[100–102]
<u>Zfp423/ Znf521</u>	<u>Nucleus</u>	<u>Induces PPARγ expression thus initiating preadipocyte commitment</u>	[224–226]	[226]
<u>DGAT 1/2</u>	<u>Cytoplasm - ER</u>	<u>Catalyses addition of third fatty acid to diacylglycerol</u>	[119, 227, 228]	[120]
<u>FASN</u>	<u>Cytoplasm</u>	<u>Uses acetylCoA, malonyl-CoA, and NADPH for <i>de novo</i> synthesis of long chain saturated fatty acids.</u>	[229, 230]	[120, 231, 232]
<u>LPL</u>	<u>Adipocyte plasma membrane</u>	<u>Hydrolyses lipoproteins at capillary endothelium that are triglyceride-rich to generate NEFA</u>	[233, 234]	[120, 234]
<u>ATGL</u>	<u>LD membrane</u>	<u>Hydrolyses TAG to DAG</u>	[235–237]	[236, 237]
<u>HSL</u>	<u>Cytosol, LD membrane</u>	<u>Hydrolyses TAG to DAG, and DAG to MAG</u>	[237–240]	[237, 241]
<u>MGL</u>	<u>Cytosol</u>	<u>Hydrolyses MAG to glycerol and and a fatty acid.</u>	[236, 237]	[237]

Box 1. Pitfalls and tricks when working with dermal adipose tissue.

<u>Issue</u>	<u>Solution(s)</u>
<u>Poor quality DWAT in cryosections</u>	<ul style="list-style-type: none"> • <u>Cut thicker sections</u> • <u>Lower the cryostat temperature to -35 degrees</u> • <u>Gently apply coverslips upon tissue sections</u>
<u>Lipid is displaced from adipocytes in cryosections</u>	<ul style="list-style-type: none"> • <u>Lower the cryostat temperature to -35 degrees</u> • <u>Store slides in cryostat without allowing them to thaw</u> • <u>Cut sections of up to 30µm thickness</u> • <u>Change fixative to 4%PFA or Ethanol:Glacial Acetic Acid</u>
<u>Lipid is absent in paraffin sections</u>	<ul style="list-style-type: none"> • <u>Switch to cryosections – lipids are damaged during processing</u>
<u>Poor quality of adipocytes in paraffin sections</u>	<ul style="list-style-type: none"> • <u>Set waterbath temperature to 35 degrees (adipocytes tend to burst at higher temperatures)</u> • <u>Switch to a more gentle fixative (Zinc Formalin instead of 10% NBF)</u> • <u>Apply coverslips gently upon tissue sections</u> • <u>Ensure the tissue is surrounded by enough fixative</u>
<u>Uncertainty as to whether adipocyte stain worked</u>	<ul style="list-style-type: none"> • <u>Check staining within positive control (human tissue distinct to skin)</u> • <u>Check whether there is staining in sebaceous gland (there are many shared markers/lipid stains between sebocytes and DAs)</u>
<u>Antibody optimization of secreted factors repeatedly fails</u>	<ul style="list-style-type: none"> • <u>Try Tyramide Signal Amplification method</u> • <u>Resort to detecting mRNA of adipose-derived factors using ISH</u>
<u>Low resolution of target protein in immunofluorescence</u>	<ul style="list-style-type: none"> • <u>Try confocal microscopy instead (also available for tissue that is 20µm or thicker)</u> • <u>Attempt immuno-EM (as a last resort as the technique is very difficult)</u>
<u>Identifying adipocyte nuclei within sections</u>	<ul style="list-style-type: none"> • <u>Cut thicker sections (the volume of adipocyte renders few nuclei pertaining to mature adipocytes)</u> • <u>Double stain with a lipid stain (e.g. BODIPY) and plasma membrane stain (e.g. CPM) to identify adipocyte-specific nuclei</u>
<u>Difficulty in co-localising target antigen with actual adipose cells</u>	<ul style="list-style-type: none"> • <u>Conduct double stain with an adipocyte marker e.g. Perilipin1/BODIPY</u> • <u>Use IHC instead of IF stain</u>
<u>Difficulty in detecting dying adipocytes <i>in situ</i></u>	<ul style="list-style-type: none"> • <u>Stain for a macrophage marker (CD68/Mac2) to detect crown-like structures around adipocytes</u> • <u>Quantify loss of Perilipin1/CAVI</u> • <u>Do an ultrastructure test for apoptosis using TEM</u>

References

1. Alexander CM, Kasza I, Yen C-LE, et al (2015) Dermal white adipose tissue: a new component of the thermogenic response. *J Lipid Res* 56:2061–9 . doi: 10.1194/jlr.R062893
2. Shook B, Rivera Gonzalez G, Ebmeier S, et al (2016) The Role of Adipocytes in Tissue Regeneration and Stem Cell Niches. *Annu Rev Cell Dev Biol*. doi: 10.1146/annurev-cellbio-111315-125426
3. Plikus M V., Guerrero-Juarez CF, Ito M, et al (2017) Regeneration of fat cells from myofibroblasts during wound healing. *Science* (80-) 355:748–752 . doi: 10.1126/science.aai8792
4. Zhang L -j., Guerrero-Juarez CF, Hata T, et al (2015) Dermal adipocytes protect against invasive *Staphylococcus aureus* skin infection. *Science* (80-) 347:67–71 . doi: 10.1126/science.1260972
5. Marangoni RG, Lu TT (2017) The roles of dermal white adipose tissue loss in scleroderma skin fibrosis. *Curr Opin Rheumatol* 1 . doi: 10.1097/BOR.0000000000000437
6. Plikus M V., Mayer JA, de la Cruz D, et al (2008) Cyclic dermal BMP signalling regulates stem cell activation during hair regeneration. *Nature* 451:340–344 . doi: 10.1038/nature06457
7. Festa E, Fretz J, Berry R, et al (2011) Adipocyte lineage cells contribute to the skin stem cell niche to drive hair cycling. *Cell* 146:761–71 . doi: 10.1016/j.cell.2011.07.019
8. Guerrero-Juarez CF, Plikus M V. (2018) Emerging nonmetabolic functions of skin fat. *Nat Rev Endocrinol* 14:163–173 . doi: 10.1038/nrendo.2017.162
9. Zwick RK, Guerrero-Juarez CF, Horsley V, Plikus M V (2018) Anatomical, Physiological, and Functional Diversity of Adipose Tissue. *Cell Metab* 27:68–83 . doi: 10.1016/j.cmet.2017.12.002
10. Yang C-C, Sheu H-M, Chung P-L, et al (2015) Leptin of dermal adipose tissue is differentially expressed during the hair cycle and contributes to adipocyte-mediated growth inhibition of anagen-phase vibrissa hair. *Exp Dermatol* 24:57–60 . doi: 10.1111/exd.12566
11. Fu X, Fang L, Li H, et al (2007) Adipose tissue extract enhances skin wound healing. *Wound Repair Regen* 15:540–8 . doi: 10.1111/j.1524-475X.2007.00262.x
12. Kim W-S, Park B-S, Sung J-H, et al (2007) Wound healing effect of adipose-derived stem cells:

- 1
2
3 a critical role of secretory factors on human dermal fibroblasts. *J Dermatol Sci* 48:15–24 . doi:
4 10.1016/j.jdermsci.2007.05.018
5
6
7
8 13. Schmidt BA, Horsley V (2013) Intradermal adipocytes mediate fibroblast recruitment during
9 skin wound healing. *Development* 140:1517–27 . doi: 10.1242/dev.087593
10
11
12 14. Procaccini C, La Rocca C, Carbone F, et al (2017) Leptin as immune mediator: Interaction
13 between neuroendocrine and immune system. *Dev Comp Immunol* 66:120–129 . doi:
14 10.1016/j.dci.2016.06.006
15
16
17 15. Greenhill C (2017) Adipose tissue: Crosstalk between adipocytes and neurons. *Nat Rev*
18 *Endocrinol* 13:438–438 . doi: 10.1038/nrendo.2017.79
19
20
21
22 16. Klar AS, Zimoch J, Biedermann T (2017) Skin Tissue Engineering: Application of Adipose-
23 Derived Stem Cells. *Biomed Res Int* 2017:9747010
24
25
26 17. Owczarczyk-Saczonek A, Wociór A, Placek W, et al (2017) The Use of Adipose-Derived Stem
27 Cells in Selected Skin Diseases (Vitiligo, Alopecia, and Nonhealing Wounds). *Stem Cells Int.*
28 2017
29
30
31
32 18. Pak J, Lee JH, Park KS, et al (2017) Current use of autologous adipose tissue-derived stromal
33 vascular fraction cells for orthopedic applications. *J Biomed Sci* 24:9 . doi: 10.1186/s12929-
34 017-0318-z
35
36
37
38 19. Lazar I, Clement E, Dauvillier S, et al (2016) Adipocyte Exosomes Promote Melanoma
39 Aggressiveness through Fatty Acid Oxidation: A Novel Mechanism Linking Obesity and Cancer.
40 *Cancer Res* 76:4051–4057 . doi: 10.1158/0008-5472.CAN-16-0651
41
42
43
44 20. Shih BB, Nirmal AJ, Headon DJ, et al (2017) Derivation of marker gene signatures from human
45 skin and their use in the interpretation of the transcriptional changes associated with
46 dermatological disorders. *J Pathol* 241:600–613 . doi: 10.1002/path.4864
47
48
49
50 21. Wang Y, Zhang L, Yu J, et al (2017) A Co-Drug of Butyric Acid Derived from Fermentation
51 Metabolites of the Human Skin Microbiome Stimulates Adipogenic Differentiation of
52 Adipose-Derived Stem Cells: Implications in Tissue Augmentation. *J Invest Dermatol* 137:46–
53 56 . doi: 10.1016/j.jid.2016.07.030
54
55
56
57 22. Rivera-Gonzalez G, Shook B, Horsley V (2014) Adipocytes in skin health and disease. *Cold*
58 *Spring Harb Perspect Med* 4: . doi: 10.1101/cshperspect.a015271
59
60

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
23. Driskell RR, Watt FM (2015) Understanding fibroblast heterogeneity in the skin. *Trends Cell Biol* 25:92–9 . doi: 10.1016/j.tcb.2014.10.001
24. Rutkowski JM, Stern JH, Scherer PE (2015) The cell biology of fat expansion. *J Cell Biol* 208:501–12 . doi: 10.1083/jcb.201409063
25. Kruglikov IL, Scherer PE, Wollina U (2016) Are dermal adipocytes involved in psoriasis? *Exp Dermatol* n/a-n/a . doi: 10.1111/exd.12996
26. Berry R, Church CD, Gericke MT, et al (2014) Imaging of adipose tissue. *Methods Enzymol* 537:47–73 . doi: 10.1016/B978-0-12-411619-1.00004-5
27. Martínez-Santibañez G, Cho KW, Lumeng CN (2014) Imaging white adipose tissue with confocal microscopy. *Methods Enzymol* 537:17–30 . doi: 10.1016/B978-0-12-411619-1.00002-1
28. Stanley EG, Jenkins BJ, Walker CG, et al (2017) Lipidomic profiling of human adipose tissue identifies a pattern of lipids associated with fish oil supplementation. *J Proteome Res* acs.jproteome.7b00161 . doi: 10.1021/acs.jproteome.7b00161
29. Pietiläinen KH, Røgg T, Seppänen-Laakso T, et al (2011) Association of Lipidome remodeling in the Adipocyte membrane with acquired obesity in humans. *PLoS Biol* 9: . doi: 10.1371/journal.pbio.1000623
30. Baker RC, Nikitina Y, Subauste AR (2014) Analysis of adipose tissue lipid using mass spectrometry. *Methods Enzymol* 538:89–105 . doi: 10.1016/B978-0-12-800280-3.00006-2
31. López-Bascón MA, Calderón-Santiago M, Sánchez-Ceinos J, et al (2017) Influence of sample preparation on lipidomics analysis of polar lipids in adipose tissue. *Talanta*. doi: 10.1016/j.talanta.2017.09.017
32. Camera E, Ludovici M, Tortorella S, et al (2016) Use of lipidomics to investigate sebum dysfunction in juvenile acne. *J Lipid Res* 57:1051–1058 . doi: 10.1194/jlr.M067942
33. Shields KJ, Wu C (2017) Differential Adipose Tissue Proteomics. In: *Methods in molecular biology* (Clifton, N.J.)
34. Kirkwood JS, Miranda CL, Bobe G, et al (2016) 18O-Tracer metabolomics reveals protein turnover and CDP-choline cycle activity in differentiating 3T3-L1 pre-adipocytes. *PLoS One* 11: . doi: 10.1371/journal.pone.0157118

- 1
2
3 35. Suárez-Zamorano N, Fabbiano S, Chevalier C, et al (2015) Microbiota depletion promotes
4 browning of white adipose tissue and reduces obesity. *Nat Med* 21:1497–1501 . doi:
5
6 10.1038/nm.3994
7
8
- 9 36. Sanmiguel C, Gupta A, Mayer EA (2015) Gut Microbiome and Obesity: A Plausible Explanation
10 for Obesity. *Curr. Obes. Rep.* 4:250–261
11
12
- 13 37. Schugar RC, Shih DM, Warriar M, et al (2017) Erratum: The TMAO-Producing Enzyme Flavin-
14 Containing Monooxygenase 3 Regulates Obesity and the Beiging of White Adipose Tissue
15 (Cell Reports (2017) 19(12) (2451–2461) (S2211124717307519)
16 (10.1016/j.celrep.2017.05.077)). *Cell Rep.* 20:279
17
18
19
- 20 38. Carobbio S, Pellegrinelli V, Vidal-Puig A (2017) Adipose tissue function and expandability as
21 determinants of lipotoxicity and the metabolic syndrome. In: *Advances in Experimental*
22 *Medicine and Biology.* pp 161–196
23
24
25
- 26 39. Reilly SM, Saltiel AR (2017) Adapting to obesity with adipose tissue inflammation. *Nat Rev*
27 *Endocrinol.* doi: 10.1038/nrendo.2017.90
28
29
- 30 40. Kruglikov IL, Scherer PE (2017) Adipocyte-myofibroblast transition as a possible
31 pathophysiological step in androgenetic alopecia. *Exp. Dermatol.* 26:522–523
32
33
34
- 35 41. Wu J, Boström P, Sparks LM, et al (2012) Beige adipocytes are a distinct type of thermogenic
36 fat cell in mouse and human. *Cell* 150:366–76 . doi: 10.1016/j.cell.2012.05.016
37
38
- 39 42. Giralt M, Villarroya F (2013) White, Brown, Beige/Brite: Different Adipose Cells for Different
40 Functions? *Endocrinology* 154:2992–3000 . doi: 10.1210/en.2013-1403
41
42
- 43 43. Wang QA, Scherer PE (2016) Human Beige Adipocytes: Epiphenomenon or Drivers of
44 Metabolic Improvements? *Trends Endocrinol Metab* 27:244–246 . doi:
45 10.1016/j.tem.2016.03.008
46
47
48
- 49 44. Driskell RR, Jahoda CAB, Chuong CM, et al (2014) Defining dermal adipose tissue. *Exp.*
50 *Dermatol.* 23:629–631
51
52
- 53 45. Driskell RR, Lichtenberger BM, Hoste E, et al (2013) Distinct fibroblast lineages determine
54 dermal architecture in skin development and repair. *Nature* 504:277–81 . doi:
55 10.1038/nature12783
56
57
58
- 59 46. DURWARD A, RUDALL KM (1949) Studies on hair growth in the rat. *J Anat* 83:325–35, 4 pl
60

- 1
- 2
- 3
- 4 47. Hausman GJ, Champion DR, Richardson RL, Martin RJ (1981) Adipocyte development in the rat
- 5 hypodermis. *Am J Anat* 161:85–100 . doi: 10.1002/aja.1001610107
- 6
- 7
- 8 48. Driskell RR, Jahoda CAB, Chuong C-M, et al (2014) Defining dermal adipose tissue. *Exp*
- 9 *Dermatol* 23:629–631 . doi: 10.1111/exd.12450
- 10
- 11
- 12 49. Schneider MR (2014) Coming home at last: Dermal white adipose tissue. *Exp. Dermatol.*
- 13 23:634–635
- 14
- 15
- 16 50. Larson JD, Tierney WS, Ozturk CN, Zins JE (2014) Defining the Fat Compartments in the Neck:
- 17 A Cadaver Study. *Aesthetic Surg J* 34:499–506 . doi: 10.1177/1090820X14526406
- 18
- 19
- 20 51. Poblet E, Jiménez-Acosta F, Hardman JA, et al (2016) Is the eccrine gland an integral,
- 21 functionally important component of the human scalp pilosebaceous unit? *Exp Dermatol*
- 22 25:149–150 . doi: 10.1111/exd.12889
- 23
- 24
- 25
- 26 52. Poblet E, Jimenez-Acosta F, Escario-Travesedo E, et al (2018) Eccrine sweat glands associate
- 27 with the human hair follicle within a defined compartment of dermal white adipose tissue. *Br*
- 28 *J Dermatol.* doi: 10.1111/bjd.16436
- 29
- 30
- 31
- 32 53. Saely CH, Geiger K, Drexel H (2012) Brown versus White Adipose Tissue: A Mini-Review.
- 33 *Gerontology* 58:15–23 . doi: 10.1159/000321319
- 34
- 35
- 36 54. Kwok KHM, Lam KSL, Xu A (2016) Heterogeneity of white adipose tissue: molecular basis and
- 37 clinical implications. *Exp Mol Med* 48:e215 . doi: 10.1038/emm.2016.5
- 38
- 39
- 40 55. Sun K, Kusminski CCM, Scherer PEP (2011) Adipose tissue remodeling and obesity. *J Clin ...*
- 41 121:2094–2101 . doi: 10.1172/JCI45887.2094
- 42
- 43
- 44 56. Horsley V, Watt F (2017) Repeal and Replace: Adipocyte Regeneration in Wound Repair. *Cell*
- 45 *Stem Cell* 20:424–426 . doi: 10.1016/j.stem.2017.03.015
- 46
- 47
- 48 57. Divoux A, Tordjman J, Lacasa D, et al (2010) Fibrosis in Human Adipose Tissue : Composition ,
- 49 Distribution , and Link With Lipid Metabolism and Fat. *Diabetes* 59:2817–2825 . doi:
- 50 10.2337/db10-0585.A.D.
- 51
- 52
- 53
- 54 58. Yang C-C, Chung P-L, Lin L-Y, et al (2017) Higher plasma leptin is associated with higher risk of
- 55 androgenetic alopecia in men. *Exp Dermatol* 26:524–526 . doi: 10.1111/exd.13369
- 56
- 57
- 58 59. Chen G (2016) Chapter 100: “Other acquired disorders of subcutaneous fat.” *Rook’s Textb*
- 59 *dermatology* Wiley-Blackwell 9:67–82
- 60

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
60. Fung (2017) Chapter 2 "Panniculitis." *Dermatopathology* 2nd editio:Elsevier/Saunders, Philadelphia
61. Toussirost É, Aubin F, Dumoulin G (2014) Relationships between adipose tissue and psoriasis, with or without arthritis. *Front. Immunol.* 5
62. Kruglikov I, Wollina U (2017) Local effects of adipose tissue in psoriasis and psoriatic arthritis. *Psoriasis Targets Ther* Volume 7:17–25 . doi: 10.2147/PTT.S122959
63. Kruglikov IL (2017) Interfacial Adipose Tissue in Systemic Sclerosis. *Curr Rheumatol Rep* 19:1–8 . doi: 10.1007/s11926-017-0627-y
64. Lynch M, Ahern T, Sweeney CM, et al (2017) Adipokines, psoriasis, systemic inflammation, and endothelial dysfunction. *Int. J. Dermatol.* 56:1103–1118
65. Cheung L, Fisher RM, Kuzmina N, et al (2016) Psoriasis Skin Inflammation-Induced microRNA-26b Targets NCEH1 in Underlying Subcutaneous Adipose Tissue. *J Invest Dermatol* 136:640–648 . doi: 10.1016/j.jid.2015.12.008
66. Matsuda Y, Fujii T, Suzuki T, et al (2011) Comparison of Fixation Methods for Preservation of Morphology, RNAs, and Proteins From Paraffin-Embedded Human Cancer Cell-Implanted Mouse Models. *J Histochem Cytochem* 59:68–75 . doi: 10.1369/jhc.2010.957217
67. Mehlem A, Hagberg CE, Muhl L, et al (2013) Imaging of neutral lipids by oil red O for analyzing the metabolic status in health and disease. *Nat Protoc* 8:1149–1154 . doi: 10.1038/nprot.2013.055
68. Ezure T, Amano S (2015) Increment of subcutaneous adipose tissue is associated with decrease of elastic fibres in the dermal layer. *Exp Dermatol* 24:924–929 . doi: 10.1111/exd.12816
69. Donati G, Proserpio V, Lichtenberger BM, et al (2014) Epidermal Wnt/ -catenin signaling regulates adipocyte differentiation via secretion of adipogenic factors. *Proc Natl Acad Sci* 111:E1501–E1509 . doi: 10.1073/pnas.1312880111
70. Halberg N, Khan T, Trujillo ME, et al (2009) Hypoxia-inducible factor 1alpha induces fibrosis and insulin resistance in white adipose tissue. *Mol Cell Biol* 29:4467–4483 . doi: 10.1128/MCB.00192-09
71. Fleischmajer R, Damiano V, Nedwich A (1971) Scleroderma and the subcutaneous tissue.

- 1
2
3 Science 171:1019–21
4
5
6 72. Huang C, Murphy GF, Akaishi S, Ogawa R (2013) Keloids and hypertrophic scars: update and
7 future directions. *Plast Reconstr surgery Glob open* 1:e25 . doi:
8 10.1097/GOX.0b013e31829c4597
9
10
11 73. Marangoni RG, Korman BD, Wei J, et al (2015) Myofibroblasts in Murine Cutaneous Fibrosis
12 Originate From Adiponectin-Positive Intradermal Progenitors. *Arthritis Rheumatol* 67:1062–
13 1073 . doi: 10.1002/art.38990
14
15
16
17 74. HARRIS JE, Peters A (1953) Experiments on vital staining with methylene blue. *Q J Microsc Sci*
18 3:113–124
19
20
21 75. Genina EA, Bashkatov AN, Tuchin V V. (2004) Methylene blue diffusion in skin tissue. In:
22 ALT'03 International Conference on Advanced Laser Technologies: Biomedical Optics. pp
23 315–323
24
25
26
27 76. Tchoukalova YD, Koutsari C, Karpyak M V, et al (2008) Subcutaneous adipocyte size and body
28 fat distribution. *Am J Clin Nutr* 87:56–63
29
30
31 77. Ramírez-Zacarías JL, Castro-Muñozledo F, Kuri-Harcuch W (1992) Quantitation of adipose
32 conversion and triglycerides by staining intracytoplasmic lipids with oil red O. *Histochemistry*
33 97:493–497 . doi: 10.1007/BF00316069
34
35
36
37 78. Fowler SD, Greenspan P (1985) Application of Nile red, a fluorescent hydrophobic probe, for
38 the detection of neutral lipid deposits in tissue sections: comparison with oil red O. *J*
39 *Histochem Cytochem* 33:833–836 . doi: 10.1177/33.8.4020099
40
41
42
43 79. Kinkel AD, Fernyhough ME, Helterline DL, et al (2004) Oil red-O stains non-adipogenic cells: A
44 precautionary note. *Cytotechnology* 46:49–56 . doi: 10.1007/s10616-004-3903-4
45
46
47
48 80. Greenspan P, Mayer EP, Fowler SD (1985) Nile red: a selective fluorescent stain for
49 intracellular lipid droplets. *J Cell Biol* 100:965–973 . doi: 10.1083/jcb.100.3.965
50
51
52 81. Mendel D (1912) Behavior of Fat-Soluble Dyes
53
54
55 82. Hirsch J, Gallian E (1968) Methods for the determination of adipose cell size in man and
56 animals. *J Lipid Res* 9:110–119
57
58
59 83. Spangenburg EE, Pratt SJP, Wohlers LM, Lovering RM (2011) Use of BODIPY (493/503) to
60 visualize intramuscular lipid droplets in skeletal muscle. *J Biomed Biotechnol* 2011: . doi:

- 1
2
3 10.1155/2011/598358
4
5
6 84. Qiu B, Simon M (2016) BODIPY 493/503 Staining of Neutral Lipid Droplets for Microscopy and
7 Quantification by Flow Cytometry. *BIO-PROTOCOL* 6: . doi: 10.21769/BioProtoc.1912
8
9
10 85. Jiang M, Gu X, Lam JWY, et al (2017) Two-photon AIE bio-probe with large Stokes shift for
11 specific imaging of lipid droplets. *Chem Sci* 8:5440–5446 . doi: 10.1039/C7SC01400G
12
13
14 86. Tang J, Zhang Y, Yin H-Y, et al (2017) Precise Labeling and Tracking of Lipid Droplets in
15 Adipocytes Using a Luminescent ZnSalen Complex. *Chem - An Asian J* 12:2533–2538 . doi:
16 10.1002/asia.201701010
17
18
19
20 87. Jester J V., Potma E, Brown DJ (2016) PPAR γ Regulates Mouse Meibocyte Differentiation and
21 Lipid Synthesis. *Ocul Surf* 14:484–494 . doi: 10.1016/j.jtos.2016.08.001
22
23
24 88. Dutta P, Haller E, Sharp A, Nanjundan M (2016) MIR494 reduces renal cancer cell survival
25 coinciding with increased lipid droplets and mitochondrial changes. *BMC Cancer* 16:33 . doi:
26 10.1186/s12885-016-2053-3
27
28
29
30 89. Ding J, Liu Y, Sullivan DA (2015) Effects of insulin and high glucose on human meibomian
31 gland epithelial cells. *Investig Ophthalmol Vis Sci* 56:7814–7820 . doi: 10.1167/iovs.15-18049
32
33
34 90. Wang H, Sreenivasan U, Sreenevasan U, et al (2011) Perilipin 5, a lipid droplet-associated
35 protein, provides physical and metabolic linkage to mitochondria. *J Lipid Res* 52:2159–68 .
36 doi: 10.1194/jlr.M017939
37
38
39
40 91. Hinde E, Haslam IS, Schneider MR, et al (2013) A practical guide for the study of human and
41 murine sebaceous glands in situ. *Exp. Dermatol.* 22:631–637
42
43
44 92. Bayati V, Abbaspour MR, Dehbashi FN, et al (2016) A dermal equivalent developed from
45 adipose-derived stem cells and electrospun polycaprolactone matrix: an in vitro and in vivo
46 study. *Anat Sci Int* 1–12 . doi: 10.1007/s12565-016-0352-z
47
48
49
50 93. Fan JY, Carpentier JL, van Obberghen E, et al (1983) Morphological changes of the 3T3-L1
51 fibroblast plasma membrane upon differentiation to the adipocyte form. *J Cell Sci* 61:219–30
52
53
54 94. Kwok KHM, Lam KSL, Xu A (2016) Heterogeneity of white adipose tissue: molecular basis and
55 clinical implications. *Exp Mol Med* 48:e215 . doi: 10.1038/emm.2016.5
56
57
58 95. Lefterova MI, Zhang Y, Steger DJ, et al (2008) PPAR and C/EBP factors orchestrate adipocyte
59 biology via adjacent binding on a genome-wide scale. *Genes Dev* 22:2941–2952 . doi:
60

- 1
2
3 10.1101/gad.1709008
4
5
6 96. Shao X, Wang M, Wei X, et al (2016) Peroxisome Proliferator-Activated Receptor- γ : Master
7 Regulator of Adipogenesis and Obesity. *Curr Stem Cell Res Ther* 11:282–9 . doi:
8 10.2174/1574888X10666150528144905
9
10
11 97. Tontonoz P, Hu E, Devine J, et al (1995) PPAR gamma 2 regulates adipose expression of the
12 phosphoenolpyruvate carboxykinase gene. *Mol Cell Biol* 15:351–357 . doi:
13 10.1128/MCB.15.1.351
14
15
16
17 98. Park B-H, Qiang L, Farmer SR (2004) Phosphorylation of C/EBPbeta at a consensus
18 extracellular signal-regulated kinase/glycogen synthase kinase 3 site is required for the
19 induction of adiponectin gene expression during the differentiation of mouse fibroblasts into
20 adipocytes. *Mol Cell Biol* 24:8671–80 . doi: 10.1128/MCB.24.19.8671-8680.2004
21
22
23
24
25 99. Hudak CS, Sul HS (2013) Pref-1, a gatekeeper of adipogenesis. *Front Endocrinol (Lausanne)*
26 4:79 . doi: 10.3389/fendo.2013.00079
27
28
29 100. Mitterberger MC, Lechner S, Mattesich M, et al (2012) DLK1(PREF1) is a negative regulator of
30 adipogenesis in CD105+/CD90+/CD34+/CD31–/FABP4– adipose-derived stromal cells from
31 subcutaneous abdominal fat pats of adult women. *Stem Cell Res* 9:35–48 . doi:
32 10.1016/j.scr.2012.04.001
33
34
35
36 101. Zwierzina ME, Ejaz A, Bitsche M, et al (2015) Characterization of DLK1(PREF1)+/CD34+ cells in
37 vascular stroma of human white adipose tissue. *Stem Cell Res* 15:403–418 . doi:
38 10.1016/j.scr.2015.08.004
39
40
41
42 102. Marques AP, Rosmaninho-Salgado J, Estrada M, et al (2017) Hypoxia mimetic induces lipid
43 accumulation through mitochondrial dysfunction and stimulates autophagy in murine
44 preadipocyte cell line. *Biochim Biophys Acta - Gen Subj* 1861:673–682 . doi:
45 10.1016/j.bbagen.2016.12.005
46
47
48
49 103. Galarraga M, Campi3n J, Mu3noz-Barrutia A, et al (2012) Adiposoft: automated software for
50 the analysis of white adipose tissue cellularity in histological sections. *J Lipid Res* 53:2791–
51 2796 . doi: 10.1194/jlr.D023788
52
53
54
55 104. Parlee SD, Lentz SI, Mori H, MacDougald OA (2014) Quantifying size and number of
56 adipocytes in adipose tissue. *Methods Enzymol* 537:93–122 . doi: 10.1016/B978-0-12-
57 411619-1.00006-9
58
59
60

- 1
2
3 105. Gadupudi G, Gourronc FA, Ludewig G, et al (2015) PCB126 inhibits adipogenesis of human
4 preadipocytes. *Toxicol Vitro* 29:132–141 . doi: 10.1016/j.tiv.2014.09.015
5
6
- 7 106. Wojciechowicz K, Gledhill K, Ambler CA, et al (2013) Development of the mouse dermal
8 adipose layer occurs independently of subcutaneous adipose tissue and is marked by
9 restricted early expression of FABP4. *PLoS One* 8:e59811 . doi:
10 10.1371/journal.pone.0059811
11
12
- 13 107. Prins JB, Walker NI, Winterford CM, Cameron DP (1994) Human adipocyte apoptosis occurs in
14 malignancy. *Biochem Biophys Res Commun* 205:625–630
15
16
- 17 108. Cinti S (2005) Adipocyte death defines macrophage localization and function in adipose tissue
18 of obese mice and humans. *J Lipid Res* 46:2347–2355 . doi: 10.1194/jlr.M500294-JLR200
19
20
- 21 109. Giordano A, Murano I, Mondini E, et al (2013) Obese adipocytes show ultrastructural features
22 of stressed cells and die of pyroptosis. *J Lipid Res* 54:2423–2436 . doi: 10.1194/jlr.M038638
23
24
- 25 110. Cinti S, Mitchell G, Barbatelli G, et al (2005) Adipocyte death defines macrophage localization
26 and function in adipose tissue of obese mice and humans. *J Lipid Res* 46:2347–2355 . doi:
27 10.1194/jlr.M500294-JLR200
28
29
- 30 111. Feng D, Tang Y, Kwon H, et al (2011) High-fat diet-induced adipocyte cell death occurs
31 through a cyclophilin D intrinsic signaling pathway independent of adipose tissue
32 inflammation. *Diabetes* 60:2134–2143 . doi: 10.2337/db10-1411
33
34
- 35 112. Lumeng CN, Delproposto JB, Westcott DJ, Saltiel AR (2008) Phenotypic switching of adipose
36 tissue macrophages with obesity is generated by spatiotemporal differences in macrophage
37 subtypes. *Diabetes* 57:3239–3246 . doi: 10.2337/db08-0872
38
39
- 40 113. Lumeng CN, Bodzin JL, Saltiel AR (2007) Obesity induces a phenotypic switch in adipose tissue
41 macrophage polarization. *J Clin Invest* 117:175–184 . doi: 10.1172/JCI29881
42
43
- 44 114. Simon G (1965) Adipose Tissue. *Handb Physiol* 5:101–107
45
46
- 47 115. Rodeheffer MS, Birsoy K, Friedman JM (2008) Identification of white adipocyte progenitor
48 cells in vivo. *Cell* 135:240–9 . doi: 10.1016/j.cell.2008.09.036
49
50
- 51 116. Lu Z, Hasse S, Bodo E, et al (2006) Towards the development of a simplified long-term organ
52 culture method for human scalp skin and its appendages under serum-free conditions. doi:
53 10.1111/j.1600-0625.2006.00510.x
54
55
56
57
58
59
60

- 1
2
3 117. Langan EA, Philpott MP, Kloepper JE, Paus R (2015) Human hair follicle organ culture: theory,
4 application and perspectives. *Exp. Dermatol.* 24:903–911
5
6
7 118. Purba TS, Haslam IS, Shahmalak A, et al (2015) Mapping the expression of epithelial hair
8 follicle stem cell-related transcription factors LHX2 and SOX9 in the human hair follicle. *Exp*
9 *Dermatol* 24:462–7 . doi: 10.1111/exd.12700
10
11
12
13 119. Yen CL, Stone SJ, Koliwad S, et al (2000) DGAT enzymes and triacylglycerol biosynthesis. *J*
14 *Lipid Res* . doi: 10.1194/jlr.R800018-JLR200
15
16
17 120. Ranganathan G, Unal R, Pokrovskaya I, et al (2006) The lipogenic enzymes DGAT1, FAS, and
18 LPL in adipose tissue: effects of obesity, insulin resistance, and TZD treatment. *J Lipid Res*
19 47:2444–2450 . doi: 10.1194/jlr.M600248-JLR200
20
21
22
23 121. Girusse A, Langin D (2012) Adipocyte lipases and lipid droplet-associated proteins: insight
24 from transgenic mouse models. *Int J Obes* 36:581–594 . doi: 10.1038/ijo.2011.113
25
26
27 122. Yang X, Lu X, Lombès M, et al (2010) The G(0)/G(1) switch gene 2 regulates adipose lipolysis
28 through association with adipose triglyceride lipase. *Cell Metab* 11:194–205 . doi:
29 10.1016/j.cmet.2010.02.003
30
31
32
33 123. Morimoto C, Kameda K, Tsujita T, Okuda H (2001) Relationships between lipolysis induced by
34 various lipolytic agents and hormone-sensitive lipase in rat fat cells. *J Lipid Res* 42:120–7
35
36
37 124. Lass A, Zimmermann R, Oberer M, Zechner R (2011) Lipolysis - a highly regulated multi-
38 enzyme complex mediates the catabolism of cellular fat stores. *Prog Lipid Res* 50:14–27 . doi:
39 10.1016/j.plipres.2010.10.004
40
41
42
43 125. Nielsen TS, Jessen N, Jørgensen JOL, et al (2014) Dissecting adipose tissue lipolysis: molecular
44 regulation and implications for metabolic disease. *J Mol Endocrinol* 52:R199-222 . doi:
45 10.1530/JME-13-0277
46
47
48
49 126. Martin S, Okano S, Kistler C, et al (2009) Spatiotemporal regulation of early lipolytic signaling
50 in adipocytes. *J Biol Chem* 284:32097–107 . doi: 10.1074/jbc.M109.002675
51
52
53 127. Schweiger M, Eichmann TO, Taschler U, et al (2014) Measurement of lipolysis. *Methods*
54 *Enzymol* 538:171–93 . doi: 10.1016/B978-0-12-800280-3.00010-4
55
56
57 128. Society RM (1965) Enzyme histochemistry of developing human skin in tissue culture.
58 84:317–322
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
129. Danse HJC (1974) 脂褐素是氧化终产物 *Enzyme Histochemical Studies of Adipose Tissue in Porcine Yellow Fat Disease*. 476:465–476
 130. Hausman GJ (1984) Histochemistry of connective tissue cells in subcutaneous adipose tissue of normal and decapitated pig fetuses. *Acta Anat (Basel)* 118:147–52
 131. Li H, Black PN, DiRusso CC (2005) A live-cell high-throughput screening assay for identification of fatty acid uptake inhibitors. *Anal Biochem* 336:11–19 . doi: 10.1016/J.AB.2004.09.025
 132. Li H, Black PN, Chokshi A, et al High-throughput screening for fatty acid uptake inhibitors in humanized yeast identifies atypical antipsychotic drugs that cause dyslipidemias. doi: 10.1194/jlr.D700015-JLR200
 133. Liao J, Sportsman R, Harris J, Stahl A (2005) Real-time quantification of fatty acid uptake using a novel fluorescence assay. *J Lipid Res* 46:597–602 . doi: 10.1194/jlr.D400023-JLR200
 134. Kolahi K, Louey S, Varlamov O, Thornburg K (2016) Real-Time Tracking of BODIPY-C12 Long-Chain Fatty Acid in Human Term Placenta Reveals Unique Lipid Dynamics in Cytotrophoblast Cells. *PLoS One* 11:e0153522 . doi: 10.1371/journal.pone.0153522
 135. Iglesias J, Lamontagne J, Erb H, et al (2015) Simplified assays of lipolysis enzymes for drug discovery and specificity assessment of known inhibitors. *J Lipid Res* 57:131–141 . doi: 10.1194/jlr.D058438
 136. Previs SF, McLaren DG, Wang SP, et al (2014) New methodologies for studying lipid synthesis and turnover: Looking backwards to enable moving forwards. *Biochim. Biophys. Acta - Mol. Basis Dis.* 1842:402–413
 137. Murphy EJ (2006) Stable isotope methods for the in vivo measurement of lipogenesis and triglyceride metabolism. *J. Anim. Sci.* 84 Suppl
 138. Akie TE, Cooper MP (2015) Determination of Fatty Acid Oxidation and Lipogenesis in Mouse Primary Hepatocytes. *J Vis Exp* 2–7 . doi: 10.3791/52982
 139. Grino M (2008) Study of adipose tissue gene expression by in situ hybridization. *Methods Mol Biol* 456:271–283 . doi: 10.1007/978-1-59745-245-8_20
 140. Moore HPH, Silver RB, Mottillo EP, et al (2005) Perilipin targets a novel pool of lipid droplets for lipolytic attack by hormone-sensitive lipase. *J Biol Chem* 280:43109–43120 . doi: 10.1074/jbc.M506336200

- 1
2
3 141. Jonkman J, Brown CM (2015) Any Way You Slice It - A Comparison of Confocal Microscopy
4 Techniques. *J Biomol Tech* 26:54–65 . doi: 10.7171/jbt.15-2602-003
5
6
- 7 142. Malide (2001) Confocal Microscopy of Adipocytes. *Adipose Tissue Protoc Methods Mol Biol*
8 155:
9
10
- 11 143. Wilke K, Wick K, Keil FJ, et al (2008) A strategy for correlative microscopy of large skin
12 samples: Towards a holistic view of axillary skin complexity. *Exp. Dermatol.* 17:73–80
13
14
- 15 144. Sbarbati A, Accorsi D, Benati D, et al (2010) Subcutaneous adipose tissue classification. *Eur J*
16 *Histochem* 54:48 . doi: 10.4081/ejh.2010.e48
17
18
- 19 145. Conti G, Benati D, Bernardi P, et al (2014) The post-adipocytic phase of the adipose cell cycle.
20 *Tissue Cell* 46:520–526 . doi: 10.1016/j.tice.2014.09.005
21
22
- 23 146. Bertossi D (2003) Lipofilling of the lips: ultrastructural evaluation by transmission electron
24 microscopy of injected adipose tissue. *Arch facial ...* 5:392–398 . doi: 10.1001/archfaci.5.5.392
25
26
- 27 147. Kranendonk MEG, Visseren FLJ, Van Balkom BWM, et al (2014) Human adipocyte extracellular
28 vesicles in reciprocal signaling between adipocytes and macrophages. *Obesity* 22:1296–1308 .
29 doi: 10.1002/oby.20679
30
31
- 32 148. Maurizi G, Poloni A, Mattiucci D, et al (2017) Human White Adipocytes Convert Into
33 “Rainbow” Adipocytes In Vitro. *J Cell Physiol* 232:2887–2899 . doi: 10.1002/jcp.25743
34
35
- 36 149. Frontini A, Cinti S (2010) Distribution and Development of Brown Adipocytes in the Murine
37 and Human Adipose Organ. *Cell Metab.* 11:253–256
38
39
- 40 150. Mirrashed F, Sharp JC, Krause V, et al (2004) Pilot study of dermal and subcutaneous fat
41 structures by MRI in individuals who differ in gender, BMI, and cellulite grading. *Ski Res*
42 *Technol* 10:161–168 . doi: 10.1111/j.1600-0846.2004.00072.x
43
44
- 45 151. Juan Y-H, Saboo SS, Tirumani SH, et al (2014) Malignant Skin and Subcutaneous Neoplasms in
46 Adults: Multimodality Imaging With CT, MRI, and ¹⁸F-FDG PET/CT. *Am J Roentgenol*
47 202:W422–W438 . doi: 10.2214/AJR.13.11424
48
49
- 50 152. Kudawara I, Ueda T, Yoshikawa H, et al (1999) Granular cell tumor of the subcutis: CT and MRI
51 findings. *Skelet Radiol* 28:
52
53
- 54 153. Deng J, Schoeneman SE, Zhang H, et al (2015) MRI characterization of brown adipose tissue in
55 obese and normal-weight children. *Pediatr Radiol* 45:1682–1689 . doi: 10.1007/s00247-015-
56
57
58
59
60

- 1
2
3 3391-z
4
5
6 154. Poonawalla AH, Sjoberg BP, Rehm JL, et al (2013) Adipose tissue MRI for quantitative
7 measurement of central obesity. *J Magn Reson Imaging* 37:707–716 . doi:
8 10.1002/jmri.23846
9
10
11 155. Lindenberg KS, Weydt P, Müller H-P, et al (2014) Two-Point Magnitude MRI for Rapid
12 Mapping of Brown Adipose Tissue and Its Application to the R6/2 Mouse Model of
13 Huntington Disease. *PLoS One* 9:e105556 . doi: 10.1371/journal.pone.0105556
14
15
16
17 156. Kasza I, Hernando D, Roldán-Alzate A, et al (2016) Thermogenic profiling using magnetic
18 resonance imaging of dermal and other adipose tissues. *JCI insight* 1:e87146 . doi:
19 10.1172/jci.insight.87146
20
21
22
23 157. Gronemeyer SA, Steen RG, Kauffman WM, et al (2000) Fast adipose tissue (FAT) assessment
24 by MRI. *Magn Reson Imaging* 18:815–818 . doi: 10.1016/S0730-725X(00)00168-5
25
26
27 158. Nerlekar N, Baey Y-W, Brown AJ, et al (2018) Poor Correlation, Reproducibility, and
28 Agreement Between Volumetric vs. Linear Epicardial Adipose Tissue Measurement: A 3D
29 Computed Tomography vs. 2D Echocardiography Comparison. *JACC Cardiovasc Imaging*. doi:
30 10.1016/J.JCMG.2017.10.019
31
32
33
34 159. Hu HH, Nayak KS, Goran MI (2011) Assessment of abdominal adipose tissue and organ fat
35 content by magnetic resonance imaging. *Obes Rev* 12:e504-15 . doi: 10.1111/j.1467-
36 789X.2010.00824.x
37
38
39
40 160. Hu HH, Chen J, Shen W (2016) Segmentation and quantification of adipose tissue by magnetic
41 resonance imaging. *MAGMA* 29:259–76 . doi: 10.1007/s10334-015-0498-z
42
43
44
45 161. Kim WH, Kim CG, Kim D-W (2012) Optimal CT Number Range for Adipose Tissue When
46 Determining Lean Body Mass in Whole-Body F-18 FDG PET/CT Studies. *Nucl Med Mol Imaging*
47 (2010) 46:294–9 . doi: 10.1007/s13139-012-0175-3
48
49
50
51 162. Judex S, Luu YK, Ozcivici E, et al (2010) Quantification of adiposity in small rodents using
52 micro-CT. *Methods* 50:14–9 . doi: 10.1016/j.ymeth.2009.05.017
53
54
55 163. Marchadier A, Vidal C, Tafani J-P, et al (2011) Quantitative CT imaging for adipose tissue
56 analysis in mouse model of obesity Quantitative CT imaging for adipose tissue analysis in
57 mouse model of obesity Quantitative CT imaging for adipose tissue analysis in mouse model
58 of obesity. *SPIE Med Imaging*
59
60

- 1
2
3 164. Kullberg J, Hedström A, Brandberg J, et al (2017) Automated analysis of liver fat, muscle and
4 adipose tissue distribution from CT suitable for large-scale studies. *Sci Rep* 7:10425 . doi:
5 10.1038/s41598-017-08925-8
6
7
8
9 165. Li L, Margolis LB, Paus R, Hoffman RM (1992) Hair shaft elongation, follicle growth, and
10 spontaneous regression in long-term, gelatin sponge-supported histoculture of human scalp
11 skin. *Proc Natl Acad Sci U S A* 89:8764–8 . doi: 10.1073/pnas.89.18.8764
12
13
14
15 166. Gesta S, Lolmède K, Daviaud D, et al (2003) Culture of human adipose tissue explants leads to
16 profound alteration of adipocyte gene expression. *Horm Metab Res = Horm und*
17 *Stoffwechselforsch = Horm métabolisme* 35:158–63 . doi: 10.1055/s-2003-39070
18
19
20
21 167. Carswell KA, Lee M-J, Fried SK (2012) Culture of isolated human adipocytes and isolated
22 adipose tissue. *Methods Mol Biol* 806:203–14 . doi: 10.1007/978-1-61779-367-7_14
23
24
25 168. Skurk T, Hauner H (2012) Primary culture of human adipocyte precursor cells: expansion and
26 differentiation. *Methods Mol Biol* 806:215–26 . doi: 10.1007/978-1-61779-367-7_15
27
28
29 169. Ruiz-Ojeda F, Rupérez A, Gomez-Llorente C, et al (2016) Cell Models and Their Application for
30 Studying Adipogenic Differentiation in Relation to Obesity: A Review. *Int J Mol Sci* 2016, Vol
31 17, Page 1040 17:1040 . doi: 10.3390/IJMS17071040
32
33
34
35 170. Huber B, Link A, Linke K, et al (2016) Integration of Mature Adipocytes to Build-Up a
36 Functional Three-Layered Full-Skin Equivalent. *Tissue Eng Part C Methods* 22:756–764 . doi:
37 10.1089/ten.tec.2016.0141
38
39
40
41 171. Lequeux C, Auxenfans C, Thépot A, et al (2011) A simple way to reconstruct a human 3-D
42 hypodermis: A useful tool for pharmacological functionality. *Skin Pharmacol Physiol* 25:47–55
43 . doi: 10.1159/000330904
44
45
46
47 172. Miyamoto Y, Ikeuchi M, Noguchi H, et al (2017) Enhanced Adipogenic Differentiation of
48 Human Adipose-Derived Stem Cells in an in vitro Microenvironment: The Preparation of
49 Adipose-Like Microtissues Using a Three-Dimensional Culture. *Cell Med* 9:35–44 . doi:
50 10.3727/215517916X693096
51
52
53
54 173. Turner PA, Gurumurthy B, Bailey JL, et al (2017) Adipogenic differentiation of human adipose-
55 derived stem cells grown as spheroids. *Process Biochem* 59:312–320 . doi:
56 10.1016/j.procbio.2017.02.003
57
58
59
60 174. Furuhata Y, Kikuchi Y, Tomita S, Yoshimoto K (2016) Small spheroids of adipose-derived stem

- 1
2
3 cells with time-dependent enhancement of IL-8 and VEGF-A secretion. *Genes to Cells*
4 21:1380–1386 . doi: 10.1111/gtc.12448
5
6
7
8 175. Feng J, Mineda K, Wu S-H, et al (2017) An injectable non-cross-linked hyaluronic-acid gel
9 containing therapeutic spheroids of human adipose-derived stem cells. *Sci Rep* 7:1548 . doi:
10 10.1038/s41598-017-01528-3
11
12
13 176. Heid H, Rickelt S, Zimbelmann R, et al (2014) On the formation of lipid droplets in human
14 adipocytes: The organization of the perilipin-vimentin cortex. *PLoS One* 9: . doi:
15 10.1371/journal.pone.0090386
16
17
18 177. Paino F, La Noce M, Di Nucci D, et al (2017) Human adipose stem cell differentiation is highly
19 affected by cancer cells both in vitro and in vivo: implication for autologous fat grafting. *Cell*
20 *Death Dis* 8:e2568 . doi: 10.1038/cddis.2016.308
21
22
23
24 178. Poulos SP, Dodson M V, Hausman GJ (2010) Cell line models for differentiation: preadipocytes
25 and adipocytes. *Exp Biol Med* 235:1185–1193 . doi: 10.1258/ebm.2010.010063
26
27
28
29 179. Todorčević M, Hilton C, McNeil C, et al (2017) A cellular model for the investigation of depot
30 specific human adipocyte biology. *Adipocyte* 6:40–55 . doi: 10.1080/21623945.2016.1277052
31
32
33
34 180. Torii I, Morikawa S, Nakano A, Morikawa K (2003) Establishment of a Human Preadipose Cell
35 Line, HPB-AML-I: Refractory to PPAR γ -Mediated Adipogenic Stimulation. *J Cell Physiol*
36 197:42–52 . doi: 10.1002/jcp.10326
37
38
39 181. Wabitsch M, Brenner RE, Melzner I, et al (2001) Characterization of a human preadipocyte
40 cell strain with high capacity for adipose differentiation. *Int J Obes* 25:8–15 . doi:
41 10.1038/sj.ijo.0801520
42
43
44 182. Iwata T, Kuribayashi K, Nakasono M, et al (2017) The AMPK/mTOR pathway is involved in D-
45 dopachrome tautomerase gene transcription in adipocytes differentiated from SGBS cells, a
46 human preadipocyte cell line. *Cytokine* 96:195–202 . doi: 10.1016/j.cyto.2017.04.017
47
48
49
50 183. Guennoun A, Kazantzis M, Thomas R, et al (2015) Comprehensive molecular characterization
51 of human adipocytes reveals a transient brown phenotype. *J Transl Med* 13:135 . doi:
52 10.1186/s12967-015-0480-6
53
54
55 184. Gibellini L, De Biasi S, Pinti M, et al (2017) The protease inhibitor atazanavir triggers
56 autophagy and mitophagy in human preadipocytes. *Aids* 26:2017–2026 . doi:
57 10.1097/QAD.0b013e328359b8be
58
59
60

- 1
2
3 185. Fischer-Posovszky P, Newell FS, Wabitsch M, Tornqvist HE (2008) Human SGBS cells - A
4 unique tool for studies of human fat cell biology. *Obes. Facts* 1:184–189
5
6
7 186. Zhou Y, Robciuc MR, Wabitsch M, et al (2012) OSBP-Related Proteins (ORPs) in Human
8 Adipose Depots and Cultured Adipocytes: Evidence for Impacts on the Adipocyte Phenotype.
9 *PLoS One* 7: . doi: 10.1371/journal.pone.0045352
10
11
12
13 187. Do M-S, Jeong H-S, Choi B-H, et al (2006) Inflammatory gene expression patterns revealed by
14 DNA microarray analysis in TNF-alpha-treated SGBS human adipocytes. *Yonsei Med J* 47:729–
15 736 . doi: 10.3349/ymj.2006.47.5.729
16
17
18
19 188. Lee Y-K, Cowan C a (2014) Differentiation of white and brown adipocytes from human
20 pluripotent stem cells.
21
22
23 189. Chen J-H, Goh KJ, Rocha N, et al (2017) Evaluation of human dermal fibroblasts directly
24 reprogrammed to adipocyte-like cells as a metabolic disease model. *Dis Model Mech*
25 *dmm.030981* . doi: 10.1242/dmm.030981
26
27
28
29 190. Choi S, Ryoo S-B, Park KJ, et al (2017) Autologous adipose tissue-derived stem cells for the
30 treatment of complex perianal fistulas not associated with Crohn’s disease: a phase II clinical
31 trial for safety and efficacy. *Tech Coloproctol* 21:345–353 . doi: 10.1007/s10151-017-1630-z
32
33
34
35 191. Dranse HJ, Muruganandan S, Fawcett JP, Sinal CJ (2016) Adipocyte-secreted chemerin is
36 processed to a variety of isoforms and influences MMP3 and chemokine secretion through an
37 NFkB-dependent mechanism. *Mol Cell Endocrinol* 436:114–129 . doi:
38 10.1016/j.mce.2016.07.017
39
40
41
42 192. Forni MF, Pelligia J, Braga TT, et al (2017) Caloric Restriction Promotes Structural and
43 Metabolic Changes in the Skin. *Cell Rep* 20:2678–2692 . doi: 10.1016/j.celrep.2017.08.052
44
45
46
47 193. Wang YY, Attané C, Milhas D, et al (2017) Mammary adipocytes stimulate breast cancer
48 invasion through metabolic remodeling of tumor cells. *JCI Insight* 2: . doi:
49 10.1172/jci.insight.87489
50
51
52
53 194. Berry R, Church CD, Gericke MT, et al (2014) Chapter Four – Imaging of Adipose Tissue. In:
54 *Methods in Enzymology*. pp 47–73
55
56
57 195. Zhang L -j., Guerrero-Juarez CF, Hata T, et al (2015) Dermal adipocytes protect against
58 invasive *Staphylococcus aureus* skin infection. *Science* (80-) 347:67–71 . doi:
59 10.1126/science.1260972
60

- 1
2
3 196. Shook B, Rivera Gonzalez G, Ebmeier S, et al (2016) The Role of Adipocytes in Tissue
4 Regeneration and Stem Cell Niches. *Annu Rev Cell Dev Biol*. doi: 10.1146/annurev-cellbio-
5 111315-125426
6
7
- 8
9 197. Kang S, Akerblad P, Kiviranta R, et al (2012) Regulation of Early Adipose Commitment by
10 Zfp521. *PLoS Biol* 10: . doi: 10.1371/journal.pbio.1001433
11
12
- 13 198. Murano I, Barbatelli G, Parisani V, et al (2008) Dead adipocytes, detected as crown-like
14 structures, are prevalent in visceral fat depots of genetically obese mice. *J Lipid Res* 49:1562–
15 8 . doi: 10.1194/jlr.M800019-JLR200
16
17
- 18 199. Klapper M, Ehmke M, Palgunow D, et al (2011) Fluorescence-based fixative and vital staining
19 of lipid droplets in *Caenorhabditis elegans* reveal fat stores using microscopy and flow
20 cytometry approaches. *J Lipid Res* 52:1281–93 . doi: 10.1194/jlr.D011940
21
22
- 23 200. Jia Z, Moulson CL, Pei Z, et al (2007) Fatty Acid Transport Protein 4 Is the Principal Very Long
24 Chain Fatty Acyl-CoA Synthetase in Skin Fibroblasts *. doi: 10.1074/jbc.M700568200
25
26
- 27 201. McDonough PM, Maciejewski-Lenoir D, Hartig SM, et al (2013) Differential Phosphorylation
28 of Perilipin 1A at the Initiation of Lipolysis Revealed by Novel Monoclonal Antibodies and
29 High Content Analysis. *PLoS One* 8:e55511 . doi: 10.1371/journal.pone.0055511
30
31
- 32 202. Wang H, Hu L, Dalen K, et al (2009) Activation of hormone-sensitive lipase requires two steps,
33 protein phosphorylation and binding to the PAT-1 domain of lipid droplet coat proteins. *J Biol*
34 *Chem* 284:32116–25 . doi: 10.1074/jbc.M109.006726
35
36
- 37 203. Mirancea N, Mirancea D (2011) THE ULTRASTRUCTURE OF THE WHITE SUBCUTANEOUS
38 ADIPOSE TISSUE. *Proc Rom Acad, Ser B* 3:206–211
39
40
- 41 204. Kolokol'chikova EG, Pal'tsyn AA, Shchegolev AI, et al (2005) Proliferative activity of adipocytes
42 in adipose tissue tumors. *Bull Exp Biol Med* 140:122–126 . doi: doi [pii]
43
44
- 45 205. Plikus M V, Mayer JA, de la Cruz D, et al (2008) Cyclic dermal BMP signalling regulates stem
46 cell activation during hair regeneration. *Nature* 451:340–4 . doi: 10.1038/nature06457
47
48
- 49 206. Khan T, Muise ES, Iyengar P, et al (2009) Metabolic dysregulation and adipose tissue fibrosis:
50 role of collagen VI. *Mol Cell Biol* 29:1575–91 . doi: 10.1128/MCB.01300-08
51
52
- 53 207. Sun Z, Gong J, Wu H, et al (2013) Perilipin1 promotes unilocular lipid droplet formation
54 through the activation of Fsp27 in adipocytes. *Nat Commun* 4:1594 . doi:
55
56
57
58
59
60

- 1
2
3 10.1038/ncomms2581
4
5
6 208. Tansey JT, Sztalryd C, Hlavin EM, et al (2004) The central role of perilipin A in lipid metabolism
7 and adipocyte lipolysis. *IUBMB Life* 56:379–385
8
9
10 209. Sun Z, Gong J, Wu H, et al (2013) Perilipin1 promotes unilocular lipid droplet formation
11 through the activation of Fsp27 in adipocytes. *Nat Commun.* doi: 10.1038/ncomms2581
12
13
14 210. Stenson BM, Rydén M, Venteclef N, et al (2011) Liver X Receptor (LXR) regulates human
15 adipocyte lipolysis. *J Biol Chem* 286:370–379 . doi: 10.1074/jbc.M110.179499
16
17
18 211. Mastrogiannaki M, Lichtenberger BM, Reimer A, et al (2016) β -Catenin Stabilization in Skin
19 Fibroblasts Causes Fibrotic Lesions by Preventing Adipocyte Differentiation of the
20 Reticular Dermis. *J Invest Dermatol* 136:1130–1142 . doi: 10.1016/j.jid.2016.01.036
21
22
23
24 212. Öst A, Örtegren U, Gustavsson J, et al (2005) Triacylglycerol is synthesized in a specific
25 subclass of caveolae in primary adipocytes. *J Biol Chem* 280:5–8 . doi:
26 10.1074/jbc.C400429200
27
28
29
30 213. Briand N, Prado C, Mabileau G, et al (2014) Caveolin-1 expression and cavin stability regulate
31 caveolae dynamics in adipocyte lipid store fluctuation. *Diabetes* 63:4032–4044 . doi:
32 10.2337/db13-1961
33
34
35
36 214. Catalán V, Gómez-Ambrosi J, Rodríguez A, et al (2008) Expression of caveolin-1 in human
37 adipose tissue is upregulated in obesity and obesity-associated type 2 diabetes mellitus and
38 related to inflammation. *Clin Endocrinol (Oxf)* 68:213–219 . doi: 10.1111/j.1365-
39 2265.2007.03021.x
40
41
42
43 215. Giusti V, Verdumo C, Suter M, et al (2003) Expression of peroxisome proliferator-activated
44 receptor-gamma1 and peroxisome proliferator-activated receptor-gamma2 in visceral and
45 subcutaneous adipose tissue of obese women. *Diabetes* 52:1673–1676
46
47
48
49 216. Tanaka T, Yoshida N, Kishimoto T, Akira S (1997) Defective adipocyte differentiation in mice
50 lacking the C/EBPbeta and/or C/EBPdelta gene. *EMBO J* 16:7432–43 . doi:
51 10.1093/emboj/16.24.7432
52
53
54
55 217. Muruganandan S, Roman AA, Sinal CJ (2009) Adipocyte differentiation of bone marrow-
56 derived mesenchymal stem cells: Cross talk with the osteoblastogenic program. *Cell. Mol. Life*
57 *Sci.* 66:236–253
58
59
60

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
218. Yang HN, Park JS, Woo DG, et al (2011) C/EBP- β and C/EBP- δ -mediated adipogenesis of human mesenchymal stem cells (hMSCs) using PLGA nanoparticles complexed with poly(ethyleneimine). *Biomaterials* 32:5924–5933 . doi: 10.1016/j.biomaterials.2011.04.072
219. Garcés C, Ruiz-Hidalgo MJ, Bonvini E, et al (1999) Adipocyte differentiation is modulated by secreted delta-like (dlk) variants and requires the expression of membrane-associated dlk. *Differentiation* 64:103–114 . doi: 10.1007/s002580050265
220. Mei B, Zhao L, Chen L, Sul HS (2002) Only the large soluble form of preadipocyte factor-1 (Pref-1), but not the small soluble and membrane forms, inhibits adipocyte differentiation: role of alternative splicing. *Biochem J* 364:137–44
221. Mortensen SB, Jensen CH, Schneider M, et al (2012) Membrane-Tethered Delta-Like 1 Homolog (DLK1) Restricts Adipose Tissue Size by Inhibiting Preadipocyte Proliferation. *Diabetes* 61:
222. Sul HS (2009) Minireview: Pref-1: Role in Adipogenesis and Mesenchymal Cell Fate. *Mol Endocrinol* 23:1717–1725 . doi: 10.1210/me.2009-0160
223. Wang Y, Kim K-A, Kim J-H, Sul HS (2006) Pref-1, a Preadipocyte Secreted Factor That Inhibits Adipogenesis. *Journal Nutr* 136:2953–2956 . doi: 136/12/2953 [pii]
224. Gupta RK, Arany Z, Seale P, et al (2010) Transcriptional control of preadipocyte determination by Zfp423. *Nature* 464:619–623 . doi: 10.1038/nature08816
225. Gupta RK, Mepani RJ, Kleiner S, et al (2012) Zfp423 expression identifies committed preadipocytes and localizes to adipose endothelial and perivascular cells. *Cell Metab* 15:230–239 . doi: 10.1016/j.cmet.2012.01.010
226. Longo M, Raciti GA, Zatterale F, et al (2017) Epigenetic modifications of the Zfp / ZNF423 gene control murine adipogenic commitment and are dysregulated in human hypertrophic obesity. 4:
227. Meegalla RL, Billheimer JT, Cheng D (2002) Concerted elevation of acyl-coenzyme A:diacylglycerol acyltransferase (DGAT) activity through independent stimulation of mRNA expression of DGAT1 and DGAT2 by carbohydrate and insulin. *Biochem Biophys Res Commun* 298:317–323 . doi: 10.1016/S0006-291X(02)02466-X
228. Lopes JLS, Nobre TM, Cilli EM, et al (2014) Deconstructing the DGAT1 enzyme: Binding sites and substrate interactions. *Biochim Biophys Acta - Biomembr* 1838:3145–3152 . doi:

- 1
2
3 10.1016/j.bbamem.2014.08.017
4
5
6 229. Loftus TM (2000) Reduced Food Intake and Body Weight in Mice Treated with Fatty Acid
7 Synthase Inhibitors. *Science* (80-) 288:2379–2381 . doi: 10.1126/science.288.5475.2379
8
9
10 230. Padilla-Benavides T, Velez-delValle C, Marsch-Moreno M, et al (2016) Lipogenic Enzymes
11 Complexes and Cytoplasmic Lipid Droplet Formation During Adipogenesis. *J Cell Biochem*
12 2315–2326 . doi: 10.1002/jcb.25529
13
14
15 231. Mayas MD, Ortega FJ, Macías-González M, et al (2010) Inverse relation between FASN
16 expression in human adipose tissue and the insulin resistance level. *Nutr Metab (Lond)* 7:3 .
17 doi: 10.1186/1743-7075-7-3
18
19
20 232. Kusakabe T, Maeda M, Hoshi N, et al (2000) Fatty acid synthase is expressed mainly in adult
21 hormone-sensitive cells or cells with high lipid metabolism and in proliferating fetal cells. *J*
22 *Histochem Cytochem* 48:613–622 . doi: 10.1177/002215540004800505
23
24
25 233. Merkel M, Eckel RH, Goldberg IJ (2002) Lipoprotein lipase: genetics, lipid uptake, and
26 regulation. *J Lipid Res* 43:1997–2006 . doi: 10.1194/jlr.R200015-JLR200
27
28
29 234. Mead JR, Irvine SA, Ramji DP (2002) Lipoprotein lipase: structure, function, regulation, and
30 role in disease. *J Mol Med (Berl)* 80:753–69 . doi: 10.1007/s00109-002-0384-9
31
32
33 235. Zimmermann R (2004) Fat Mobilization in Adipose Tissue Is Promoted by Adipose Triglyceride
34 Lipase. *Science* (80-) 306:1383–1386 . doi: 10.1126/science.1100747
35
36
37 236. Bézaire V, Langin D (2009) Regulation of adipose tissue lipolysis revisited. *Proc Nutr Soc*
38 68:350–60 . doi: 10.1017/S0029665109990279
39
40
41 237. Brasaemle DL (2007) *Thematic review series: Adipocyte Biology* . The perilipin family of
42 structural lipid droplet proteins: stabilization of lipid droplets and control of lipolysis. *J Lipid*
43 *Res* 48:2547–2559 . doi: 10.1194/jlr.R700014-JLR200
44
45
46 238. Haemmerle G, Zimmermann R, Hayn M, et al (2002) Hormone-sensitive lipase deficiency in
47 mice causes diglyceride accumulation in adipose tissue, muscle, and testis. *J Biol Chem*
48 277:4806–4815 . doi: 10.1074/jbc.M110355200
49
50
51 239. Haemmerle G, Lass A, Zimmermann R, et al (2006) Defective lipolysis and altered energy
52 metabolism in mice lacking adipose triglyceride lipase. *Science* 312:734–7 . doi:
53 10.1126/science.1123965
54
55
56
57
58
59
60

- 1
2
3 240. Bezaire V, Mairal A, Ribet C, et al (2009) Contribution of adipose triglyceride lipase and
4 hormone-sensitive lipase to lipolysis in hMADS adipocytes. J Biol Chem 284:18282–91 . doi:
5 10.1074/jbc.M109.008631
6
7
8
9 241. Bezaire V, Langin D (2009) Regulation of adipose tissue lipolysis revisited. Proc Nutr Soc
10 68:350–360 . doi: S0029665109990279 [pii]\r10.1017/S0029665109990279
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only




Figure 1. Structure of dermal white adipose tissue and its components. Whilst murine DWAT is divided from subcutaneous adipocytes by the *panniculus carnosus*, human DWAT is found as a continuous layer in the human dermis (a) with no equivalent muscle structure separating it from subcutaneous white adipose tissue (SWAT) (a). DWAT surrounds the bottom half of hair follicles in the form of so-called ‘dermal cones’ (b-c). At the ultrastructural level (d-e), the cellular components of dermal adipocytes become clearer – most of the cell volume is occupied by the centrally-located lipid droplet (LD) which is surrounded by a thin sheath of cytoplasm (represented between the dotted lines); the cytoplasm is enveloped by the plasma membrane on the outside. DWAT is composed of various cell types including mature adipocytes, fibroblast-like pre-adipocytes, macrophages, pericytes, endothelial cells and a multitude of blood vessels (f). Scale bar is 2 μm in (d) and 500 nm in (e).





Figure 2. Stains for observing and quantifying human dermal white adipose tissue *in situ*. Adipocytes can be examined using histological stains such as (a) Haematoxylin and Eosin (H&E), (b) Masson's Trichrome, (b) Picrosirius Red and (d) Methylene blue (MB). The lipid within DAs can be measured using (e) Oil Red O (ORO), or fluorescently via (f) BODIPY493/503. Furthermore, various markers are available to study pre- and mature adipocytes including (g) Perilipin1 (PLIN1), (h) Caveolin1 (CAV1), (i) PPAR γ 2, (j) p-C/EBP β and (k) PREF-1. In addition, proliferation (l) via Ki67 as well as lipid synthesis enzymes such as (m) diacylglycerol acetyl transferase (DGAT1) and (n) fatty acid synthase (FASN) can be used to stain DAs *in situ*. Alternative methods of studying adipocyte *in situ* include (o) *in situ* hybridization (ISH) to study synthesis of individual strands of mRNA, and various cellular processes at the ultrastructural level (q-s) using transmission electron microscopy (TEM). Scale bar is 50 μ m across all images except the last two TEM images.



18 **Figure 3. Human primary subcutaneous adipocytes as a model for studying human DAs *in vitro*.** Primary
19 pre-adipocytes can be differentiated over 7 days into differentiated adipocytes that express PLIN1, p-CEBP β ,
20 CAV1 and PREF1. Lipid stains such as Oil Red O and BODIPY also stain differentiated adipocytes. Thus,
21 primary cells can also serve as positive and negative controls for antibody stains conducted *in situ*. Scale bar is
22 50 μ m.
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1. A guide to choosing the right processing route and stain for analysing dermal adipocytes.

LD – lipid droplet; BC – breast carcinoma; BF – brightfield; CFM – confocal microscopy; DA – dermal adipocyte; FB – fibroblasts; FFPE – formalin fixed paraffin embedded; H&E – haematoxylin and eosin; IFM – immunofluorescence microscopy; KC - keratinocyte; LD – lipid droplet; MP – macrophage; PC- prostate carcinoma; SC – stratum corneum; SG – sebaceous gland; TEM – transmission electron microscopy

What should I choose when studying...?	How to process the tissue	Stains/ Markers	Target / Antigen Localization	Positive Control Tissue	Imaging and Analysis	References
DWAT Morphology	FFPE Cryo	H&E	n/a	n/a	BF	[68, 69, 194]
Lipid droplet content and Morphology	Cryo	Oil Red O, Nile Red	LD	SG, SC, Breast	BF	[3, 7, 105, 115, 140, 148, 183, 194-196]
		BODIPY, LipidTOX	LD	SG, SC, Breast	IFM CFM	
Mature Adipocyte Adipocyte size and number	FFPE Cryo	PLIN1	LD Membrane	SC, BC	BF/IFM	[7, 101, 108, 163-166, 169-173]
		CAV1	Cell membrane	KCs, FBs, HeLa Cells		
		FABP4	Cytoplasm	KCs, Lymph node		
		CPM	Cell membrane	All cells, Any tissue	IFM	
		H&E	n/a	n/a	BF	
		DGAT1/2	Cytoplasm - ER	KCs, Colon	BF/IFM	
		FASN	Cytoplasm	KCs, PC, BC		
Adipocyte differentiation	FFPE Cryo	pC/EBP β	Nucleus	SG, BC	BF/IFM	[7, 195]
		PPAR γ	Nucleus	SG, BC		
Pre-adipocytes	FFPE Cryo	Pref1	Membrane / Nucleus	Placenta	BF/IFM	[195, 197]
		Zfp521	Nucleus	Bone		
Adipocyte apoptosis	FFPE Cryo	PLIN1	LD Membrane	SC, BC	Loss of Perilipin1	[27, 194, 198]
		CAV1	Cell membrane	KCs, FBs, HeLa Cells	Loss of Caveolin1	
Pre-adipocyte proliferation	FFPE Cryo	Pref1 + Ki67/ BrdU/ EdU	Nucleus	Placenta	BF/IFM	[7, 115, 195]
Lipid synthesis (lipogenesis)	Unfixed tissue	BODIPY C1/C12	Cytoplasm - ER	Adipocytes	IFM	[132, 134, 199, 200]
	FFPE Cryo	Mature DA stain to measure change in size (see above)	Various	Dependent on stain chosen	BF/IFM	See section above

Lipid metabolism (lipolysis)	FFPE Cryo	phospho-Perilipin1	Cytoplasm	Adipocyte cell line stimulated to undergo lipolysis	BF/IFM	[127, 201, 202]
		phospho-HSL	Cytoplasm			
Adipocyte Ultrastructure	EM-specific	OsO4 Uranyl Acetate	Only for immuno-EM	n/a	TEM	[203] [109, 144, 148, 193, 204] [176]
3D Volume and Thickness	As per MRI/ μ CT protocol	Various	n/a	n/a	MRI μ CT	[150–157, 160–164]
mRNA	FFPE Cryo	Fluorescent or chromogenic ISH	Cytoplasm	Dependent upon gene of interest PPIB; HeLa cells	ISH	[139] [205]
Fibrosis	FFPE	Masson's Trichrome/ Picrosirius Red	Fibrillary Collagens I and III	n/a	BF	[70, 206, 207]
		Methylene Blue	Dermal components other than DAs	n/a		

Review Only


Table 2. Markers of human dermal adipocytes and their functional importance.

Marker	Target Localization	Function	Murine References	Human References
Perilipin1	LD membrane ER	Master regulator of lipolysis Upon phosphorylation by PKA, Perilipin1 allows HSL to translocate to the adipocyte lipid droplet and commence lipolysis	[7, 69, 140, 195, 196, 207, 208] [209]	[102, 170, 176, 210, 211]
Caveolin1	Plasma membrane	Endo- and exocytosis Component of caveolae which are sites of triacylglycerol synthesis	[212, 213]	[214]
PPARγ2	Nucleus	Master regulator of adipogenesis	[95]	[102, 215]
pC/EBPβ	Nucleus	Promotes terminal adipocyte differentiation together with C/EBP α	[216]	[217, 218]
Pref1/Dlk1	Plasma Membrane	Inhibits pre-adipocyte proliferation in its membrane-bound form Prevents adipocyte differentiation (adipogenesis)	[99, 219–223]	[100–102]
Zfp423/ Znf521	Nucleus	Induces PPAR γ expression thus initiating preadipocyte commitment	[224–226]	[226]
DGAT 1/2	Cytoplasm - ER	Catalyses addition of third fatty acid to diacylglycerol	[119, 227, 228]	[120]
FASN	Cytoplasm	Uses acetylCoA, malonyl-CoA, and NADPH for <i>de novo</i> synthesis of long chain saturated fatty acids.	[229, 230]	[120, 231, 232]
LPL	Adipocyte plasma membrane	Hydrolyses lipoproteins at capillary endothelium that are triglyceride-rich to generate NEFA	[233, 234]	[120, 234]
ATGL	LD membrane	Hydrolyses TAG to DAG	[235–237]	[236, 237]
HSL	Cytosol, LD membrane	Hydrolyses TAG to DAG, and DAG to MAG	[237–240]	[237, 241]
MGL	Cytosol	Hydrolyses MAG to glycerol and a fatty acid.	[236, 237]	[237]


Box 1. Pitfalls and tricks when working with dermal adipose tissue.

Issue	Solution(s)
Poor quality DWAT in cryosections	<ul style="list-style-type: none"> • Cut thicker sections • Lower the cryostat temperature to -35 degrees • Gently apply coverslips upon tissue sections
Lipid is displaced from adipocytes in cryosections	<ul style="list-style-type: none"> • Lower the cryostat temperature to -35 degrees • Store slides in cryostat without allowing them to thaw • Cut sections of up to 30µm thickness • Change fixative to 4%PFA or Ethanol:Glacial Acetic Acid
Lipid is absent in paraffin sections	<ul style="list-style-type: none"> • Switch to cryosections – lipids are damaged during processing
Poor quality of adipocytes in paraffin sections	<ul style="list-style-type: none"> • Set waterbath temperature to 35 degrees (adipocytes tend to burst at higher temperatures) • Switch to a more gentle fixative (Zinc Formalin instead of 10% NBF) • Apply coverslips gently upon tissue sections • Ensure the tissue is surrounded by enough fixative
Uncertainty as to whether adipocyte stain worked	<ul style="list-style-type: none"> • Check staining within positive control (human tissue distinct to skin) • Check whether there is staining in sebaceous gland (there are many shared markers/lipid stains between sebocytes and DAs)
Antibody optimization of secreted factors repeatedly fails	<ul style="list-style-type: none"> • Try Tyramide Signal Amplification method • Resort to detecting mRNA of adipose-derived factors using ISH
Low resolution of target protein in immunofluorescence	<ul style="list-style-type: none"> • Try confocal microscopy instead (also available for tissue that is 20µm or thicker) • Attempt immuno-EM (as a last resort as the technique is very difficult)
Identifying adipocyte nuclei within sections	<ul style="list-style-type: none"> • Cut thicker sections (the volume of adipocyte renders few nuclei pertaining to mature adipocytes) • Double stain with a lipid stain (e.g. BODIPY) and plasma membrane stain (e.g. CPM) to identify adipocyte-specific nuclei
Difficulty in co-localising target antigen with actual adipose cells	<ul style="list-style-type: none"> • Conduct double stain with an adipocyte marker e.g. Perilipin1/BODIPY • Use IHC instead of IF stain
Difficulty in detecting dying adipocytes <i>in situ</i>	<ul style="list-style-type: none"> • Stain for a macrophage marker (CD68/Mac2) to detect crown-like structures around adipocytes • Quantify loss of Perilipin1/CAV1 • Do an ultrastructure test for apoptosis using TEM


Supplementary Figures




Suppl. Fig. 1. Section thickness determines lipid droplet content and morphology in dermal adipocytes. Although integrity of adipocyte cytoplasm and plasma membrane is maintained in samples sectioned at 7µm, most of the lipid droplet remains intact only when cutting the tissue at a thickness of 15-30µm as revealed by various lipid stains such as Lipid TOX (a-c), BODIPY (d-f) and OilRedO (g-i). BODIPY (d-f) appears a more robust fluorescent lipid stain in comparison with LipidTOX that fades fairly quickly (a-c). The advantage of fluorescent lipid stains (a-f) over chromogenic ones such as OilRedO (g-i) is that they can be paired with a fluorescent plasma membrane marker such as Cell Plasma Membrane stain (CPM). Scale bar = 50µm.




Suppl. Fig. 2. Effect of fixation upon lipid content of human dermal adipocytes. The choice of fixative has a great impact upon the lipid content of human dermal adipocytes. Whilst acetone displaces the lipid in the sebaceous gland (SG) and DWAT (a, b) thus making it an unsuitable choice for fixation prior to BODIPY or ORO staining, Ethanol:Glacial Acetic Acid (Eth:GlacAc) (c,d) and 4%PFA (e,f) are both suitable fixatives for subsequent lipid stains. Note how BODIPY also stains the lipid-filled sebocytes within the sebaceous gland (SG), rendering the SG as a positive control. Scale bar = 50 μ m.





Suppl. Fig. 3. Quantifying adipocyte size/volume and lipid content in ImageJ. Different tools can be employed to measure adipocyte size/volume in ImageJ (a) such as freehand selection, elliptical tool and polygon selection. We recommend utilizing the polygon selection tool (a) to measure adipocyte size/volume, which allows a more efficient tracing of the adipocyte plasma membrane, generating more consistent measurements of individual surface area/volume. Alternatively, adipocyte size can be measured automatically/manually using an ImageJ plugin called 'Adiposoft'. Quantifying adipocyte number (b) can be done using the Cell Counter Tool in ImageJ, whilst lipid stains such as OilRedO (c) can also be quantified using ImageJ. However, before quantifying staining intensity, the ORO stain needs to be inverted. Scale bar = 50 μ m.




Suppl. Fig. 4. Quantifying adipocyte differentiation in culture using the AdipoRed assay. Human primary subcutaneous adipocytes (Lonza) in culture can be differentiated in 7-10 days following addition of differentiation medium to display multiple bead-like lipid droplets within their cytoplasm (a). The level of differentiation can be measured using the AdipoRed kit (Lonza) which acts like a lipophilic dye similar to Nile Red that stains the lipid droplets of adipocytes. Maturation can be quantified using a plate reader. Displayed are the levels of adipocyte maturation at distinct passages (b,c) and also when cells are thawed and cultured following cryopreservation (d).



Suppl. Fig. 5. Cell death of skin adipocytes. Conventional methods for detecting apoptosis such as TUNEL (a,b) and Cleaved Caspase 3 (CASP3) (c,d) show no cell death within human DAs. Instead, apoptosis can be measured via different approaches, for example by measuring loss of Perilipin1 (c) or Caveolin1 (d) in dermal adipocytes within abdominal skin following 10 days in culture. Asterisks in the images denote the adipocytes that have lost Perilipin1 (c) and Caveolin1 (d) expression. The percentage of skin adipocytes that are negative for Perilipin1 (c) or Caveolin1 (d) out of total adipocytes per field of view (FOV) can be calculated in ImageJ using the Cell Counter. Results can then be represented in a graph style when comparing different methods of treatment. Scale bar = 50µm.



Suppl. Fig. 6. Measuring lipogenesis and lipolysis in human DWAT *ex vivo*. One of the methods that can be used to assess fatty acid uptake *ex vivo* is by adding BODIPY C1/C12 (green or red) to the media of abdominal skin organ culture (a). For instance, BODIPY C1/C12 can be added to adipocyte culture either as a single or double labelling with green or red BODIPY to observe the cells that are undergoing continuous or dynamic changes in lipid uptake over a certain period of time (a). Studying lipolysis in abdominal skin organ culture can be done by a colorimetric kit that detects free glycerol released into the media over a certain time period; such a test can be used to determine the effect of different medias or treatments upon adipocyte lipolysis (b). Scale bar in (a) is 50 μm .



Suppl. Fig. 7 Culturing human DWAT *ex vivo*. Several methods can be used to culture human DWAT *ex vivo* such as 4mm punch biopsies collected from human abdominal skin (a,b) or human DWAT carefully dissected from scalp skin. Depending on the depth of the punch biopsy, significant amounts of subcutaneous adipose tissue may be present, at the bottom of the punch, whereas DWAT is present mostly around follicular units in abdominal skin. Scale bar = 1mm