



Postoperative daily living activities of geriatric patients administered general or spinal anesthesia for hip fracture surgery: A retrospective cohort study

著者	Fukuda Taeko, Imai Shinobu, Nakadera Masaya, Wagatsuma Yukiko, Horiguchi Hiromasa
journal or publication title	Journal of orthopaedic surgery
volume	26
number	1
page range	230949901775410
year	2018-01
権利	(C) The Author(s) 2018 Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (http://www.creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (https://us.sagepub.com/en-us/nam/open-access-at-sage).
URL	http://hdl.handle.net/2241/00150878

doi: 10.1177/2309499017754106



Postoperative daily living activities of geriatric patients administered general or spinal anesthesia for hip fracture surgery: A retrospective cohort study

Taeko Fukuda^{1,2}, Shinobu Imai³, Masaya Nakadera³,
Yukiko Wagatsuma^{1,4} and Hiromasa Horiguchi³

Abstract

Purpose: Maintaining independence after hip fracture repair is important for geriatric patients and general welfare. We investigated the effects of anesthetic methods on postoperative activities of daily living (ADLs) following hip fracture surgery in elderly patients. **Methods:** The medical records of 12,342 patients aged ≥ 65 years who underwent typical surgeries for hip fracture using either general anesthesia or spinal anesthesia were reviewed. To adjust for baseline differences and minimize selection bias for the chosen method of anesthesia, patients were matched by propensity scores. Factors affecting the deterioration in ADLs during hospital stay were also investigated in all subjects using a multivariate logistic regression analysis. Eating, grooming, toileting, bathing, and walking were selected as the ADL parameters, as they are considered important for an independent life. **Results:** Of the 12,342 patients, 6918 (56.1%) received general anesthesia and 5424 (43.9%) received spinal anesthesia. After the propensity score matching, the anesthesia types were not associated with ADL scores except toileting at discharge. Results from the multivariate logistic regression analysis showed that the types of anesthesia were not associated with deterioration in ADL scores. Advanced age, male sex, high Charlson Comorbidity Index scores, psychiatric disease, no administration of nonsteroidal anti-inflammatory drugs, and short length of hospital stay were associated with deterioration in ADL scores. **Conclusion:** The anesthesia types were not associated with ADL dependency except toileting at discharge. Spinal anesthesia adversely affected toilet use at hospital discharge. However, anesthesia types were not factors that affected deterioration in ADL during hospital stay in elderly patients who underwent hip fracture surgery.

Keywords

activities of daily living, general anesthesia, geriatrics, hip fracture, spinal anesthesia

Date received: 11 August 2017; Received revised 22 November 2017; accepted: 16 December 2017

Introduction

Hip fractures are a global public health concern. Approximately 1.6 million hip fractures are reported to occur annually worldwide, and the incidence is expected to drastically increase several times over the next three decades due to the aging population.^{1,2} Most hip fractures are treated with surgical repair, which requires general or regional anesthesia.³ Several investigations have attempted to determine the type of anesthesia that is optimal for hip fracture, but the evidence remains conflicting.^{3–9} General anesthesia enables patients to be completely free from mental stress

¹ University of Tsukuba, Tsukuba, Ibaraki, Japan

² Faculty of Medicine, Department of Anesthesiology, Tsuchiura Center for Medical Education and Training (National Hospital Organization, Kasumigaura Medical Center), Tsuchiura, Ibaraki Prefecture, Japan

³ Department of Clinical Data Management and Research, Clinical Research Center, National Hospital Organization, Meguro-ku, Tokyo, Japan

⁴ Faculty of Medicine, Department of Clinical Trial and Clinical Epidemiology, University of Tsukuba, Tsukuba, Ibaraki, Japan

Corresponding author:

Taeko Fukuda, Faculty of Medicine, Department of Anesthesiology, University of Tsukuba, Tsukuba, Ibaraki 305-8575, Japan.
Email: taekof@md.tsukuba.ac.jp



during the surgeries. Covert and Fox reported that general anesthesia significantly reduced hypotensive episodes and blood loss when compared to regional anesthesia.¹⁰ Hypotension and/or blood loss may preclude adequate oxygen delivery to the central nervous system during surgery. Anemia is reported to be a factor related to a poor outcome after hip fracture surgery.^{11,12} On the contrary, spinal anesthesia can keep the patient at a consistent level of consciousness through the perioperative and operative periods. A systematic review reported that general anesthesia, compared to regional anesthesia, might increase the risk of developing postoperative cognitive dysfunction.¹³ It is widely known that cognitive impairment is a risk factor for poor functional outcomes in geriatric trauma patients.^{14–16}

Thus far, the main comparisons between general anesthesia and regional anesthesia methods have been mortality and morbidity, as it is difficult to investigate the changes in intellectual faculties, or in the ability to live independently, using large-scale retrospective studies. However, these factors are speculated to be associated with recovery, and poor outcomes require increased utilization of social financial assistance.¹⁷ Since 2002, data have been accumulated for 10 parameters of activities of daily living (ADLs) in many Japanese hospitals using the diagnosis procedure combination (DPC) system. We speculated that the parameters for ADLs within the nationwide administrative claims database are indicators of postoperative physical and mental status, both of which have serious effects on both quality of life and the social welfare budget.

The purpose of this study was to investigate the effects of anesthetic technique (general or spinal) on five ADL parameters (independent eating, grooming, toileting, bathing, and walking), at discharge, and on the changes in the ADL parameters during the hospital stay in elderly patients who had undergone surgery for hip fracture.

Materials and methods

This retrospective cohort study was approved by the ethics review board of the Kasumigaura Medical Center and National Hospital Organization (NHO).

Following the Japanese ethics guidelines for human medical research, this study protocol was made available to the general public via the NHO website to obtain patient objections. To protect patient privacy, all types of personal identification were encrypted in a security room of the NHO data bank. The requirement for individual informed consent was waived due to the anonymous nature of the data.

Data sources

The study period was from April 1, 2012, to March 31, 2016. We used data from the Japanese DPC administrative claims database that was obtained from 69 hospitals of the NHO group. The DPC database is a diagnosis-dominant, case-mix system administered by the Ministry of Health, Labor, and

Welfare of Japan and linked with a lump-sum payment system. We obtained hospital characteristics data from the 2016 resident recruitment guidebook of the NHO and the website of the Japanese Society of Anesthesiologists.

Selection of patients and variables

The inclusion criteria were that all patients had to be aged 65 years or older and should have undergone surgery for hip fracture (transcervical, intertrochanteric, and subtrochanteric femoral fracture) during the study period. The surgery type included total or hemiarthroplasty and other typical hip fracture procedures (plate/screw or intramedullary implant). Only patients undergoing either general anesthesia or spinal anesthesia were selected. Patients receiving unknown, mixed spinal and general, or other anesthesia types were excluded. Furthermore, we excluded 90 cases of atypical surgeries (repair of previous implanted plate, etc.) and 2347 cases for missing baseline characteristics data. This left 12,342 patients of an initial 14,779 for the analysis.

Data from the DPC database included age, sex, height, weight, fracture type, surgical procedure, comorbidities at admission, complications after admission, medications during admission, and functional status, which was based on the patient's ability to perform tasks related to ADL both at admission and at discharge. Body mass index (BMI) was calculated from height and weight. Functional status included the following 10 parameters: self-feeding, transfer, grooming, toilet use, bathing, mobility, use of stairs, dressing, defecation, and micturition.¹⁸ We chose 5 parameters (eating, grooming, toileting, bathing, and walking) from the 10 ADL abilities as indicators of independence. ADL scores for eating and toileting were defined as totally dependent (score 0), partially dependent (score 1), and independent (score 2). ADL scores for grooming and bathing were defined as dependent (score 0) and independent (score 1). ADL scores for walking were defined as totally dependent (score 0), partially dependent (wheelchair using; score 1), partially dependent (walking; score 2), and independent (score 3). The change in ADL was determined by deducting the baseline ADL score (at admission) from the ADL score at hospital discharge. A negative number (–3, –2, or –1) indicated deterioration, zero indicated no change, and a positive number (3, 2, or 1) was indicative of improvement.

A modified Charlson Comorbidity Index (CCI) was calculated for each patient based on the Quan coding algorithms.¹⁹ Hospitals authorized by the Japanese Society of Anesthesiologists have full-time, certified staff as well as residents in training. Data regarding the use of anesthetics and postoperative medication for analgesics and sedation were evaluated. Of the analgesics, 20 kinds of nonsteroidal anti-inflammatory drugs (NSAIDs) were included: aspirin, celecoxib, diclofenac, dimetotiazine mesilate, emorfazone, etodolac, flurbiprofen, ibuprofen,

indomethacin, isopropylantipyrene, ketoprofen, lornoxicam, loxoprofen, mefenamic acid, meloxicam, naproxen, salicylate, sulindac, tiaramide, and zaltoprofen. Haloperidol was used as an antipsychotic drug. The postoperative hospital length of stay (LOS) was defined as the number of calendar days from operation to discharge.

Propensity score matching

The propensity score-matched analysis was used for reducing the selection bias and potential baseline differences between the general and the spinal anesthesia groups. Scores were calculated using a binary logistic regression analysis with type of anesthesia as the dependent variable and 23 preoperative factors as predictor variables. The independent variables were age, sex, BMI, the modified CCI, length of hospital stay after surgery, fracture type, surgery type, hospital type (scale and level of anesthesiology), five ADL parameters (eating, grooming, toileting, bathing, and walking) at hospital admission, and baseline comorbidities (malignancy, anemia, endocrine/metabolism, mental, neurologic, circulatory, respiratory, digestive, and renal/urinary). The score was defined as probability of a patient receiving general anesthesia based on covariates. After adjustment with the scores, none of the preoperative variables remained statistically different between the two groups. The multivariate regression model of propensity for a patient given general anesthesia had a *C*-statistic of 0.67. (*C*-statistic is a standard measure of the predictive accuracy of a logistic regression model.)

Statistical analysis

Patient and operating characteristics were compared between the general and the spinal anesthesia groups using the *t*-test or the paired *t*-test for continuous variables and the χ^2 test for categorical variables. The ADL scores at hospital discharge were assessed using the Pearson χ^2 test after the propensity score matching. Multivariate logistic regression analysis was performed between the deterioration and the nondeterioration (no change and improvement) groups to analyze the effects of various factors. In addition to anesthesia methods, the following were selected for the model: age, sex, modified CCI, psychiatric disease (F00), NSAIDs, antipsychotic drugs, and LOS. These variables have been previously suggested to affect the prognosis.^{20–29} Odds ratios and 95% confidence intervals were determined. A *p* value of <0.05 was considered significant for all statistical tests. All statistical analyses were carried out using SPSS software version 24 (IBM, Armonk, New York).

Results

Of the 12,342 patients who underwent hip surgery during the 4-year observation period, 6918 (56.1%) received

general anesthesia and 5424 (43.9%) received spinal anesthesia. Table 1 compares the characteristics of the two groups. Before propensity score matching, 18 of the 23 baseline factors differed significantly between the two anesthesia groups. After 1:1 matching, there was a total of 3949 patients in each anesthetic group, with no significant baseline differences.

Table 2 summarizes the ADL score for independence at hospital discharge in the matched sample. Anesthesia types were not associated with ADL scores for eating, grooming, bathing, and walking. Patients in the spinal anesthesia group had a significantly higher percentage of total dependence for toileting than did patients in the general anesthesia group (spinal vs. general: 40.3% vs. 36.7%; *p* < 0.003). A total of 174 patients died and were excluded from the analysis: 109 (1.6%) in the general anesthesia group and 65 (1.2%) in the spinal anesthesia group. This difference in mortality rates was not significant (*p* = 0.078).

Results from the multivariate regression models showed that the type of anesthesia was not associated with deterioration in the ADL scores for eating, grooming, toileting, bathing, and walking after hip surgery. Nonadministration of NSAIDs was associated with deterioration in all ADL scores. In addition, advanced age, male sex, high CCI scores, psychiatric disease, and short length of hospital stay were associated with deterioration in four ADL (eating, grooming, toileting, and walking) scores (Table 3).

Discussion

Using the DPC database of the NHO group and a propensity-matched analysis, we studied the impact of anesthesia type on ADL (eating, grooming, toileting, bathing, and walking) scores for independence at hospital discharge and changes in the ADL scores during the hospital stay following surgery for hip fracture. After the propensity score matching, anesthesia types were not associated with ADL functions except toileting. Results from the multivariate logistic regression analysis showed that anesthesia types were not factors affecting ADL deterioration during hospital stay.

There are many studies evaluating levels of ADL after hip fracture.^{30–35} However, to our knowledge, only one group investigated the effect of anesthetic technique on ADL variables in their small-scale study.³⁶ They reported that no differences were observed in ADL (feeding, dressing, bathing, toileting, walking, etc.) scores between general and regional anesthesia groups at 3, 6, and 12 months after hip fracture. Although we used data for ADL scores at the time of hospital discharge, almost all of our results were in accordance with that of the long-term outcome study.

In the present study, patients in the spinal anesthesia group had a significantly higher percentage of total dependence for toileting than did patients in the general anesthesia group. Spinal anesthesia is reported to cause a clinically

Table 1. Demographic and clinical characteristics of patients who underwent hip fracture surgery.

	Before propensity score matching				After propensity score matching			
	General	Spinal	p Value	Standardized difference	General	Spinal	p Value	Standardized difference
<i>n</i>	6918	5424			3949	3949		
Age, mean (SD)	83.5 (7.5)	84.5 (7.5)	<0.001	0.03	83.8 (7.5)	84.0 (7.5)	0.242	<0.01
65–79	1940 (28.0%)	1285 (23.7%)	<0.001	0.10	1051 (26.6%)	1035 (26.2%)	0.718	<0.01
80–84	1667 (24.1%)	1231 (22.7%)		0.03	918 (23.2%)	959 (24.3%)		0.03
85–89	1794 (25.9%)	1480 (27.3%)		0.03	1045 (26.5%)	1019 (25.8%)		0.02
90+	1517 (21.9%)	1428 (26.3%)		0.10	935 (23.7%)	936 (23.7%)		0
Sex								
Male	1502 (21.7%)	1119 (20.6%)	0.145	0.03	822 (20.8%)	835 (21.1%)	0.719	<0.01
Female	5416 (78.3%)	4305 (79.4%)			3127 (79.2%)	3114 (78.9%)		
BMI, mean (SD)	20.59 (3.63)	20.25 (3.56)	<0.001	<0.01	20.49 (3.61)	20.35 (3.59)	0.082	<0.01
Charlson Comorbidity Index								
0	3087 (44.6%)	2516 (46.4%)	0.045	0.04	1799 (45.6%)	1787 (45.3%)	0.767	<0.01
1	2191 (31.7%)	1718 (31.7%)		0	1250 (31.7%)	1235 (31.3%)		<0.01
2+	1640 (23.7%)	1190 (21.9%)		0.04	900 (22.8%)	927 (23.5%)		0.02
Length of hospital stay (days)								
<21	2793 (40.4%)	3127 (57.7%)	<0.001	0.35	2017 (51.1%)	2058 (52.1%)	0.356	0.02
21+	4125 (59.6%)	2297 (42.3%)			1932 (48.9%)	1891 (47.9%)		
Fracture type								
Transcervical	3578 (51.7%)	2674 (49.3%)	0.001	0.05	2021 (51.2%)	2019 (51.1%)	0.922	<0.01
Intertrochanteric	3160 (45.7%)	2639 (48.7%)		0.06	1841 (46.6%)	1848 (46.8%)		<0.01
Subtrochanteric	180 (2.6%)	111 (2.0%)		0.04	87 (2.2%)	82 (2.1%)		<0.01
Surgery type								
Arthroplasty	2620 (37.9%)	1721 (31.7%)	<0.001	0.13	1369 (34.7%)	1351 (34.2%)	0.670	0.01
Other	4298 (62.1%)	3703 (68.3%)			2580 (65.3%)	2598 (65.8%)		
Hospital scale (bed#)								
400+	4974 (71.9%)	3899 (71.9%)	0.985	0	2864 (72.5%)	2883 (73.0%)	0.631	0.01
<400	1944 (28.1%)	1525 (28.1%)			1085 (27.5%)	1066 (27.0%)		
Anesthesiologist society authorized hospital								
Yes	6240 (90.2%)	4485 (82.7%)	<0.001	0.22	3422 (86.7%)	3426 (86.8%)	0.895	<0.01
No	678 (9.8%)	939 (17.3%)			527 (13.3%)	523 (13.2%)		
Activity of daily living (eating) at hospital admission								
Total dependent	2612 (37.8%)	1795 (33.1%)	<0.001	0.10	1407 (35.6%)	1373 (34.8%)	0.666	0.02
Partial dependent	2706 (39.1%)	2174 (40.1%)		0.02	1542 (39.0%)	1548 (39.2%)		<0.01
Independent	1600 (23.1%)	1455 (26.8%)		0.09	1000 (25.3%)	1028 (26.0%)		0.02
Activity of daily living (grooming) at hospital admission								
Dependent	5965 (86.2%)	4551 (83.9%)	<0.001	0.06	3375 (85.5%)	3351 (84.9%)	0.447	0.02
Independent	953 (13.8%)	873 (16.1%)			574 (14.5%)	598 (15.1%)		
Activity of daily living (toileting) at hospital administration								
Total dependent	5691 (82.3%)	4271 (78.7%)	<0.001	0.09	3184 (80.6%)	3172 (80.3%)	0.928	<0.01
Partial dependent	688 (9.9%)	623 (11.5%)		0.05	411 (10.4%)	421 (10.7%)		<0.01
Independent	539 (7.8%)	530 (9.8%)		0.07	354 (9.0%)	356 (9.0%)		0
Activity of daily living (bathing) at hospital administration								
Dependent	6432 (93.0%)	4931 (90.9%)	<0.001	0.08	3630 (91.9%)	3624 (91.8%)	0.805	<0.01
Independent	486 (7.0%)	493 (9.1%)			319 (8.1%)	325 (8.2%)		
Activity of daily living (walking) at hospital administration								
Total dependent	5940 (85.9%)	4488 (82.7%)	<0.001	0.09	3328 (84.3%)	3327 (84.2%)	0.961	<0.01
Partial dependent (wheelchair)	235 (3.4%)	267 (4.9%)		0.08	160 (4.1%)	162 (4.1%)		0
Partial dependent (walking)	260 (3.8%)	256 (4.7%)		0.04	163 (4.1%)	155 (3.9%)		0.01
Independent	483 (7.0%)	413 (7.6%)		0.02	298 (7.5%)	305 (7.7%)		<0.01
Malignancy								
–	6541 (94.6%)	5174 (95.4%)	0.035	0.04	3737 (94.6%)	3744 (94.8%)	0.725	<0.01
+	377 (5.4%)	250 (4.6%)			212 (5.4%)	205 (5.2%)		

(continued)

Table 1. (continued)

	Before propensity score matching				After propensity score matching			
	General	Spinal	p Value	Standardized difference	General	Spinal	p Value	Standardized difference
Anemia								
–	6430 (92.9%)	4754 (87.6%)	<0.001	0.18	3585 (90.8%)	3564 (90.3%)	0.420	0.02
+	488 (7.1%)	670 (12.4%)			364 (9.2%)	385 (9.7%)		
Endocrine/metabolism								
–	5075 (73.4%)	4116 (75.9%)	0.001	0.06	2975 (75.3%)	2938 (74.4%)	0.337	0.02
+	1843 (26.6%)	1308 (24.1%)			974 (24.7%)	1011 (25.6%)		
Mental								
–	5837 (84.4%)	4153 (76.6%)	<0.001	0.20	3175 (80.4%)	3166 (80.2%)	0.799	<0.01
+	1081 (15.6%)	1271 (23.4%)			774 (19.6%)	783 (19.8%)		
Neurologic								
–	5869 (84.8%)	4768 (87.9%)	<0.001	0.09	3437 (87.0%)	3422 (86.7%)	0.618	<0.01
+	1049 (15.2%)	656 (12.1%)			512 (13.0%)	527 (13.3%)		
Circulatory								
–	3202 (46.3%)	2906 (53.6%)	<0.001	0.15	2005 (50.8%)	1977 (50.1%)	0.529	0.01
+	3716 (53.7%)	2518 (46.4%)			1944 (49.2%)	1972 (49.9%)		
Respiratory								
–	6460 (93.4%)	4952 (91.3%)	<0.001	0.08	3621 (91.7%)	3632 (92.0%)	0.651	0.01
+	458 (6.6%)	472 (8.7%)			328 (8.3%)	317 (8.0%)		
Digestive								
–	5741 (83.0%)	4637 (85.5%)	<0.001	0.07	3333 (84.4%)	3311 (83.8%)	0.498	0.02
+	1177 (17.0%)	787 (14.5%)			616 (15.6%)	638 (16.2%)		
Renal/urinary								
–	6418 (92.8%)	5003 (92.2%)	0.262	0.02	3672 (93.0%)	3651 (92.5%)	0.363	0.02
+	500 (7.2%)	421 (7.8%)			277 (7.0%)	298 (7.5%)		

BMI: body mass index.

Table 2. ADL at hospital discharge.

Outcomes	No (%) of general anesthesia (N = 3877)	No (%) of spinal anesthesia (N = 3903)	p Value
Eating			
Total dependent	490 (12.6)	559 (14.3)	0.077
Partial dependent	1596 (41.2)	1550 (39.7)	
Independent	1791 (46.2)	1794 (46.0)	
Grooming			
Dependent	2633 (67.9)	2640 (67.6)	0.797
Independent	1244 (32.1)	1263 (32.4)	
Toileting			
Total dependent	1421 (36.7)	1574 (40.3)	0.003
Partial dependent	1525 (39.3)	1428 (36.6)	
Independent	931 (24.0)	901 (23.1)	
Bathing			
Dependent	3232 (83.4)	3278 (84.0)	0.457
Independent	645 (16.6)	625 (16.0)	
Walking			
Total dependent	1803 (46.5)	1790 (45.9)	0.124
Partial dependent (wheelchair)	565 (14.6)	524 (13.4)	
Partial dependent (walking)	830 (21.4)	916 (23.5)	
Independent	679 (17.5)	673 (17.2)	

ADL: activity of daily living.

Table 3. Multivariate analysis: Factors affecting ADL deterioration during hospital stay.

Variable	Eating				Grooming			
	β	OR	(95% CI)	p Value	β	OR	(95% CI)	p Value
Age								
(80–84/<79)	0.052	1.053	(0.850 to 1.305)	0.636	0.128	1.136	(0.862 to 1.498)	0.365
(85–90/<79)	0.322	1.380	(1.131 to 1.683)	0.001	0.294	1.341	(1.033 to 1.742)	0.028
(90+/<79)	0.502	1.652	(1.354 to 2.017)	<0.001	0.037	1.038	(0.780 to 1.380)	0.800
Sex (m/f)	0.299	1.349	(1.144 to 1.590)	<0.001	0.090	1.094	(0.869 to 1.376)	0.445
Charlson Comorbidity Index								
(1/0)	0.124	1.132	(0.958 to 1.339)	0.146	0.091	1.096	(0.868 to 1.383)	0.441
(2+/0)	0.249	1.283	(1.073 to 1.533)	0.006	0.388	1.475	(1.163 to 1.869)	0.001
Psychiatric disease (yes/no)	0.335	1.398	(1.181 to 1.655)	<0.001	0.108	1.114	(0.876 to 1.417)	0.379
Anesthesia (spinal/general)	–0.066	0.936	(0.809 to 1.083)	0.374	–0.035	0.965	(0.791 to 1.178)	0.728
NSAIDs (no/yes)	0.299	1.349	(1.169 to 1.557)	<0.001	0.291	1.338	(1.100 to 1.628)	0.004
Antipsychotic drugs (yes/no)	0.046	1.047	(0.893 to 1.227)	0.572	–0.032	0.969	(0.776 to 1.210)	0.781
Length of hospital stay (days) (<21/21+)	0.194	1.214	(1.054 to 1.398)	0.007	0.095	1.099	(0.906 to 1.333)	0.337
Variable	Toileting				Bathing			
	β	OR	(95% CI)	p Value	β	OR	(95% CI)	p Value
Age								
(80–84/<79)	0.289	1.336	(0.997 to 1.790)	0.053	–0.056	0.945	(0.697 to 1.281)	0.717
(85–90/<79)	0.584	1.794	(1.366 to 2.355)	<0.001	–0.014	0.986	(0.735 to 1.322)	0.926
(90+/<79)	0.540	1.716	(1.293 to 2.277)	<0.001	–0.218	0.804	(0.585 to 1.107)	0.181
Sex (m/f)	0.293	1.340	(1.077 to 1.668)	0.009	0.041	1.042	(0.800 to 1.358)	0.761
Charlson Comorbidity Index								
(1/0)	0.207	1.230	(0.982 to 1.541)	0.072	0.080	1.084	(0.835 to 1.407)	0.546
(2+/0)	0.363	1.438	(1.134 to 1.823)	0.003	0.240	1.271	(0.965 to 1.675)	0.088
Psychiatric disease (yes/no)	0.134	1.144	(0.908 to 1.440)	0.253	–0.032	0.968	(0.729 to 1.286)	0.824
Anesthesia (spinal/general)	0.064	1.066	(0.878 to 1.294)	0.519	0.181	1.199	(0.955 to 1.504)	0.117
NSAIDs (no/yes)	0.334	1.396	(1.153 to 1.691)	0.001	0.284	1.328	(1.061 to 1.663)	0.013
Antipsychotic drugs (yes/no)	0.106	1.112	(0.901 to 1.372)	0.323	0.055	1.057	(0.823 to 1.357)	0.664
Length of hospital stay (days) (<21/21+)	0.195	1.216	(1.006 to 1.470)	0.043	0.117	1.124	(0.901 to 1.403)	0.300
Variable	Walking							
	β	OR	(95% CI)	p Value				
Age								
(80–84/<79)	0.117	1.124	(0.865 to 1.461)	0.382				
(85–90/<79)	0.291	1.338	(1.045 to 1.714)	0.021				
(90+/<79)	0.269	1.308	(1.012 to 1.692)	0.040				
Sex (m/f)	0.176	1.192	(0.966 to 1.473)	0.102				
Charlson Comorbidity Index								
(1/0)	0.031	1.031	(0.834 to 1.276)	0.775				
(2+/0)	0.222	1.249	(0.999 to 1.561)	0.051				
Psychiatric disease (yes/no)	0.154	1.167	(0.935 to 1.456)	0.173				
Anesthesia (spinal/general)	–0.034	0.966	(0.804 to 1.162)	0.716				
NSAIDs (no/yes)	0.249	1.283	(1.070 to 1.538)	0.007				
Antipsychotic drugs (yes/no)	–0.014	0.986	(0.804 to 1.209)	0.892				
Length of hospital stay (days) (<21/21+)	0.238	1.269	(1.061 to 1.517)	0.009				

NSAIDs: nonsteroidal anti-inflammatory drugs; CI: confidence interval; OR: odds ratio; ADL: activity of daily living.

significant disturbance in bladder function.^{37,38} Intrathecal local anesthetics cause an interruption in the micturition reflex by blocking both afferent nerves and efferent fibers. Since the micturition reflex is a very delicate system, spinal anesthesia might confuse the reflex and slow the recovery of bladder function in this study.

Alarcon et al. reported that the probabilities of recovering the previous functional performance level in grooming and eating at the time of hospital discharge were 41.6% and 47.98%, respectively.³² Since they excluded nonambulatory patients with serious conditions from their study, the previous functional level is speculated to be independent.

In our study, the probabilities of independence in grooming and eating were about 32% and 46%, respectively. Although the probabilities determined from our data are slightly lower than those of Alarcon et al.'s study, we think the results are compatible with their data as we did not exclude patients with serious conditions.

Deterioration in a patient's ability to eat independently was not associated with the type of anesthesia in our multivariate logistic regression analysis. Sandman et al. stated that elderly people who were totally dependent on assisted feeding were more disabled with regard to other ADL functions.³⁹ Feeding problems might be a terminal phenomenon in the hierarchical loss of ADL functions. Furthermore, Siebens et al. reported that loss of independence in eating was associated with an increased risk of mortality within 6 months.⁴⁰ Thus, self-feeding may have to be considered as the most important item in the Barthel's ADL Index. We speculate that the type of anesthesia might have no impact on the overall deterioration in ADL, because it is not related to deterioration in eating independently.

A common factor for deterioration in independent eating, grooming, and toileting was a higher CCI score. An increase of more than two points in the CCI increases the rates of deterioration by 28% for eating, 48% for grooming, and 44% for toileting. This result is in accord with the finding of previous studies.^{31,35} Advanced age, male sex, and psychiatric disease were associated with deterioration in some ADL scores. Penrod et al. reported that advanced age and psychiatric disease (dementia and stroke) had a negative impact on ADL functions after hip fracture.³⁴ Advanced age is commonly recognized to be a determinant factor for loss of independence,^{30,31,35} whereas opinions are divided on the gender effect on independence as measured by ADL functions.^{20,41}

The probability of regaining prior levels of autonomy 6 months after surgery is reported to be only 33.6% even in previously independent elderly patients.⁴² Bertram et al. reported in their review that 29% of elderly patients did not reach their prefracture levels 1 year postfracture and went on to experience lifelong disability.⁴³ It is not easy for elderly patients to regain prefracture functional levels. Patients lose their confidence and independence due to long-term disability after a hip fracture. Thus, medical factors related to postoperative deterioration in ADL functions are important. In the present study, no administration of NSAIDs was associated with deterioration in all ADL scores. Hip pain was reported to be a risk factor related to functional dependence.^{28,35} The present study reconfirms that postoperative analgesia is important. Kawano et al. reported that ketoprofen prevented the development of surgery-associated memory deficits via its pain-relieving effects.⁴⁴ Delirium is also a risk factor for failure to regain prefracture levels of function.⁴⁵ In human and animal studies, postoperative cognitive dysfunction has been reported to be associated with a neuroinflammatory response triggered by surgery.⁴⁶⁻⁴⁸ NSAIDs might reduce the risk of

poor functional recovery via their anti-inflammatory mechanisms.

Our study had some limitations. Our data did not include several important data such as physical status as categorized by the American Society of Anesthesiologists, details of intraoperative hemodynamics, bleeding volume, and oxygen saturation. Our ADL data were collected only from hospitals. The ADL scores at hospital administration might be underestimated, because they are easily influenced by environment and/or social lifestyle. Additionally, not all our results may be generalizable to other races and countries because this study population was mainly Japanese.

Conclusion

In the propensity score matching analysis, anesthesia types were not associated with ADL (eating, grooming, bathing, and walking) scores at discharge. Results from the multivariate logistic regression analysis showed that the type of anesthesia was not associated with deterioration in the ADL scores. Spinal anesthesia adversely affected the toilet use at hospital discharge. However, it was not a deterioration factor for toileting. Advanced age, male sex, high CCI scores, psychiatric disease, no administration of NSAIDs, and short length of hospital stay were associated with deterioration in ADL scores. In addition to the patient's factors (CCI, age, sex, and coexisting disease), further investigation of additional medical factors should be considered in order for elderly patients to maintain independence after hip fracture surgery.

Acknowledgment

The authors thank Chiharu Okada, MD, PhD, MBA (National Hospital Organization Headquarters, Department of Hospital Management Assistance) for arranging this study.

Declaration of conflicting interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

1. Johnell O and Kanis JA. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporos Int* 2006; 17: 1726-1733.
2. Gullberg B, Johnell O and Kanis JA. World-wide projections for hip fracture. *Osteoporos Int* 1997; 7: 407-413.
3. Guay J, Parker MJ, Gajendragadkar PR, et al. Anaesthesia for hip fracture surgery in adults. *Cochrane Database Syst Rev* 2016; 2: CD000521.
4. Basques BA, Bohl DD, Golinvaux NS, et al. General versus spinal anaesthesia for patients aged 70 years and older

- with a fracture of the hip. *Bone Joint J* 2015; 97-B(5): 689–695.
5. Fields AC, Dieterich JD, Buterbaugh K, et al. Short-term complications in hip fracture surgery using spinal versus general anaesthesia. *Injury* 2015; 46: 719–723.
 6. Whiting PS, Molina CS, Greenberg SE, et al. Regional anaesthesia for hip fracture surgery is associated with significantly more peri-operative complications compared with general anaesthesia. *Int Orthop* 2015; 39: 1321–1327.
 7. Chu CC, Weng SF, Chen KT, et al. Propensity score-matched comparison of postoperative adverse outcomes between geriatric patients given a general or a neuraxial anesthetic for hip surgery. *Anesthesiology* 2015; 123: 136–147.
 8. Newman MD, Silber JH, Elkassabany NM, et al. Comparative effectiveness of regional versus general anesthesia for hip fracture surgery in adults. *Anesthesiology* 2012; 117: 72–92.
 9. Paterno E, Neuman MD, Schneeweiss S, et al. Comparative safety of anesthetic type for hip fracture surgery in adults: retrospective cohort study. *BMJ* 2014; 348: g4022.
 10. Covert CR and Fox GS. Anaesthesia for hip surgery in the elderly. *Can J Anaesth* 1989; 36: 311–319.
 11. Gregersen M. Postoperative red blood cell transfusion strategy in frail anemic elderly with hip fracture. A randomized controlled trial. *Dan Med J* 2016; 63(4): pii: B5221.
 12. Hagino T, Ochiai S, Sato E, et al. The relationship between anemia at admission and outcome in patients older than 60 years with hip fracture. *J Orthop Traumatol* 2009; 10: 119–122.
 13. Mason SE, Noel-Storr A and Ritchie CW. The impact of general and regional anesthesia on the incidence of postoperative cognitive dysfunction and post-operative delirium: a systematic review with meta-analysis. *J Alzheimers Dis* 2010; 22: 67–79.
 14. Maxwell CA, Mion LC, Mukherjee K, et al. Preinjury physical frailty and cognitive impairment among geriatric trauma patients determine postinjury functional recovery and survival. *J Trauma Acute Care Surg* 2016; 80: 195–203.
 15. Hakkinen A, Heinonen M, Kautiainen H, et al. Effect of cognitive impairment on basic activities of daily living in hip fracture patients: a 1-year follow-up. *Aging Clin Exp Res* 2007; 19: 139–144.
 16. Gruber-Baldini AL, Zimmerman S, Morrison RS, et al. Cognitive impairment in hip fracture patients: timing of detection and longitudinal follow-up. *J Am Geriatr Soc* 2003; 51: 1227–1236.
 17. Rundshagen I. Postoperative cognitive dysfunction. *Dtsch Arztebl Int* 2014; 111: 119–125.
 18. Collin C, Wade DT, Davies S, et al. The Barthel Index: a reliability study. *Int Disabil Stud* 1988; 10: 61–63.
 19. Quan H, Sundararajan V, Halfon P, et al. Coding algorithms for defining comorbidities in ICD-9-CM and ICD-10 administrative data. *Med Care* 2005; 43: 1130–1139.
 20. Arinzon Z, Shabat S, Peisakh A, et al. Gender differences influence the outcome of geriatric rehabilitation following hip fracture. *Arch Gerontol Geriatr* 2010; 50: 86–91.
 21. Ekstrom W, Samuelsson B, Ponzer S, et al. Sex effects on short-term complications after hip fracture: a prospective cohort study. *Clin Interv Aging* 2015; 10: 1259–1266.
 22. Smith T, Pelpola K, Ball M, et al. Pre-operative indicators for mortality following hip fracture surgery: a systematic review and meta-analysis. *Age Ageing* 2014; 43: 464–471.
 23. Shoda N, Yasunaga H, Horiguchi H, et al. Risk factors affecting in-hospital mortality after hip fracture: retrospective analysis using the Japanese diagnosis procedure combination database. *BMJ Open* 2012; 2(3): e000416.
 24. Radcliff TA, Henderson WG, Stoner TJ, et al. Patient risk factors, operative care, and outcomes among older community-dwelling male veterans with hip fracture. *J Bone Joint Surg Am* 2008; 90: 34–42.
 25. Holmes J and House A. Psychiatric illness predicts poor outcome after surgery for hip fracture: a prospective cohort study. *Psychol Med* 2000; 30: 921–929.
 26. Seitz DP, Gill SS, Gruneir A, et al. Effects of dementia on postoperative outcomes of older adults with hip fractures: a population-based study. *J Am Med Dir Assoc* 2014; 15: 334–341.
 27. Gulin T, Kruljac I, Kirigin L, et al. Advanced age, high β -CTX levels, and impaired renal function are independent risk factors for all-cause one-year mortality in hip fracture patients. *Calcif Tissue Int* 2016; 98: 67–75.
 28. Feldt KS and Oh HL. Pain and hip fracture outcomes for older adults. *Orthop Nurs* 2000; 19: 35–44.
 29. Brown CH, Azman AS, Gottschalk A, et al. Sedation depth during spinal anesthesia and survival in elderly patients undergoing hip fracture repair. *Anesth Analg* 2014; 118: 977–980.
 30. Gonzalez-Zabaleta J, Pita-Fernandez S, Seoane-Pillado T, et al. Dependence for basic and instrumental activities of daily living after hip fractures. *Arch Gerontol Geriatr* 2015; 60: 66–70.
 31. Vergara I, Vrotsou K, Orive M, et al. Factors related to functional prognosis in elderly patients after accidental hip fractures: a prospective cohort study. *BMC Geriatr* 2014; 14: 124.
 32. Alarcon T, Gonzalez-Montalvo JI, Gotor P, et al. Activities of daily living after hip fracture: profile and rate of recovery during 2 years of follow-up. *Osteoporos Int* 2011; 22: 1609–1613.
 33. Bentler SE, Liu L, Obrizan M, et al. The aftermath of hip fracture: discharge placement, functional status change, and mortality. *Am J Epidemiol* 2009; 170: 1290–1299.
 34. Penrod JD, Litke A, Hawkes WG, et al. The association of race, gender, and comorbidity with mortality and function after hip fracture. *J Gerontol A Biol Sci Med Sci* 2008; 63: 867–872.
 35. Cree M, Carriere KC, Soskolne CL, et al. Functional dependence after hip fracture. *Am J Phys Med Rehabil* 2001; 80: 736–743.
 36. Koval KJ, Aharonoff GB, Rosenberg AD, et al. Functional outcome after hip fracture. Effect of general versus regional anesthesia. *Clin Orthop Relat Res* 1998; 348: 37–41.
 37. Zanfini BA, Paradisi G, Savone R, et al. Bladder function after spinal anesthesia for cesarean section: an urodynamic

- evaluation. *Eur Rev Med Pharmacol Sci* 2012; 16: 1525–1529.
38. Lanz E and Grab BM. Micturition disorders following spinal anesthesia of different durations of action (lidocaine 2% versus bupivacaine 0.5%). *Anaesthetist* 1992; 41: 231–234.
 39. Sandman PO, Norberg A, Adolfsson R, et al. Prevalence and characteristics of persons with dependency on feeding at institutions for elderly. *Scand J Caring Sci* 1990; 4: 121–127.
 40. Siebens H, Trupe E, Siebens A, et al. Correlates and consequences of eating dependency in institutionalized elderly. *J Am Geriatr Soc* 1986; 34: 192–198.
 41. Radosavljevic N, Nikolic D, Lazovic M, et al. Estimation of functional recovery in patients after hip fracture by Berg balance scale regarding the sex, age and comorbidity of participants. *Geriatr Gerontol Int* 2013; 13: 365–371.
 42. Candel-Parra E, Corcoles-Jimenez MP, Del Egado-Fernandez MA, et al. Independence in activities of daily living 6 months after surgery in previously independent elderly patients with hip fracture caused by a fall. *Enferm Clin* 2008; 18: 309–316.
 43. Bertram M, Norman R, Kemp L, et al. Review of the long-term disability associated with hip fractures. *Inj Prev* 2011; 17: 365–370.
 44. Kawano T, Takahashi T, Iwata H, et al. Effects of ketoprofen for prevention of postoperative cognitive dysfunction in aged rats. *J Anesth* 2014; 28: 932–936.
 45. Vochteloo AJH, Moerman S, Tuinebreijer WE, et al. More than half of hip fracture patients do not regain mobility in the first postoperative year. *Geriatr Gerontol Int* 2013; 13: 334–341.
 46. Xu Z, Dong Y, Wang H, et al. Peripheral surgical wounding and age-dependent neuroinflammation in mice. *PLoS One* 2014; 9(5): e96752.
 47. Beloosesky Y, Hendel D, Weiss A, et al. Cytokines and C-reactive protein production in hip-fracture-operated elderly patients. *J Gerontol A Biol Sci Med Sci* 2007; 62: 420–426.
 48. Buvanendran A, Kroin JS, Berger RA, et al. Upregulation of prostaglandin E2 and interleukins in the central nervous system and peripheral tissue during and after surgery in humans. *Anesthesiology* 2006; 104: 403–410.