



Open Archive TOULOUSE Archive Ouverte (OATAO)


OATAO is an open access repository that collects the work of Toulouse researchers and makes it freely available over the web where possible.

This is an author-deposited version published in : <http://oatao.univ-toulouse.fr/>
Eprints ID : 16279

To link to this article :

URL : <https://prodinra.inra.fr/?locale=en#!ConsultNotice:348677>

To cite this version :

Kraus, Daniel and Büttler, Rita and Krumm, Franck and Lachat, Thibault and Larrieu, Laurent  and Mergner, Ulrich and Paillet, Yoan and Schuck, Andreas and Winter, Susanne *Catalogue of tree microhabitats : Reference field list*. (2016) [Report]

Any correspondence concerning this service should be sent to the repository administrator: staff-oatao@listes-diff.inp-toulouse.fr

Catalogue of tree microhabitats

Reference field list



This catalogue is available for download:
integrateplus.org

Recommended citation: Kraus, D., Bütler, R., Krumm, F., Lachat, T., Larrieu, L., Mergner, U., Paillet, Y., Rydkvist, T., Schuck, A., and Winter, S. 2016. Catalogue of tree microhabitats – Reference field list. Integrate+ Technical Paper. 16p.

Illustrations: Lisa Apfelbacher

Photos: Daniel Kraus

Disclaimer: The present catalogue is one of the outputs of the demonstration project '*Establishing a European network of demonstration sites for the integration of biodiversity conservation into forest management*' supported by the German Federal Ministry of Food and Agriculture (BMEL). The views expressed in this publication are those of the authors and do not necessarily represent those of the European Forest Institute.

European Forest Institute, 2016

Large quantities of deadwood and a high density of old microhabitat-bearing trees are characteristic elements of natural forests, especially of the old-growth phases. These are often absent or rare in managed forests, even in forests under close-to-nature management. Yet, an important share of forest biodiversity is strictly or primarily dependent on such elements for their survival, especially 'saproxylic' species, those are species depending on deadwood.

Tree related microhabitats are therefore recognised as important substrates and structures for biodiversity in forests. The retention of both existing and future tree microhabitats is thus one important aspect to take in to consideration in forest management. Giving tree microhabitats increased attention will help sustain and increase the habitat value for biodiversity also in managed forests .

This reference field list is developed to support training exercises conducted in Integrate+ Marteloscope sites. It aims at supporting forest managers, inventory personnel and other groups in identifying and describing tree microhabitats in the course of such exercises. It can also find use as illustrative material in forest education and as background documentation for other training events and field excursions.



Illustrations	Code	Type	Description	Saprophytic microhabitats
---------------	------	------	-------------	---------------------------

CV1

Woodpecker cavities



CV11 $\varnothing = 4 \text{ cm}$

Cavity entrance about $\varnothing = 4 \text{ cm}$ with a larger interior diameter of the cavity. The cavity of *Dendrocopos minor* is built in crown branches.



CV12 $\varnothing = 5 - 6 \text{ cm}$

Cavity entrance about $\varnothing = 5 - 6 \text{ cm}$ with a larger interior diameter of the cavity. *Picus viridis* builds its cavity into the trunk into the insertion of broken-off branches. The entrance is round in accordance with the round form of the branch insertion.



The cavities of the medium-sized woodpeckers such as *Dendrocopos major* are built into decaying branch insertions, dead branches or snags.



CV13 $\varnothing > 10 \text{ cm}$

Woodpecker hole in the trunk that indicates a cavity of *Dryocopus martius*. The cavity entrance width is about $\varnothing > 10 \text{ cm}$ with a larger interior diameter. Black woodpeckers build cavities with an oval entrance at trunk free of branches. The cavity trees have mostly a minimum average of 40 cm dbh thus it can remain for a longer decaying time (2 - 3 decades).



CV14 $\varnothing \geq 10 \text{ cm}$ feeding hole)

The excavation is conical: the entrance is larger than the interior.


Cavities

Saprophytic microhabitats	Description	Type	Code	Illustrations
---------------------------	-------------	------	------	---------------

At least three in the trunk connected woodpecker breeding cavities. If this cannot be checked: three cavity openings within two meters.

Woodpecker "flute"
/ cavity string

CV15



Trunk and mould cavities

Trunk cavity with mould, cavity bottom has ground contact thus soil humidity enters the cavity hole. Note that the cavity entrance can be higher at the trunk.

$\varnothing \geq 10$ cm (ground contact)

CV21




$\varnothing \geq 30$ cm (ground contact)

CV22

Mould containing trunk cavity without ground contact.

$\varnothing \geq 10$ cm

CV23



$\varnothing \geq 30$ cm

CV24



Semi-open trunk cavity with or without mould, cavity chamber is not completely protected from surrounding microclimate and precipitation may enter the interior. Note that the cavity entrance can be higher at the trunk.

$\varnothing \geq 30$ cm / semi-open

CV25



Large trunk cavity with open top and with or without ground contact.

$\varnothing \geq 30$ cm / open top

CV26



Cavities



Illustrations	Code	Type	Description	Saprophylic microhabitats
---------------	------	------	-------------	---------------------------

CV3

Branch holes



CV31

$\varnothing \geq 5 \text{ cm}$

Rot-holes originating from branch breakage at trunk when fungal decay of wood is progressing faster than occlusion of wound.



CV32

$\varnothing \geq 10 \text{ cm}$



CV33

Hollow branch,
 $\varnothing \geq 10 \text{ cm}$

Hollow more or less horizontal branch developing from breakage. Provides tubular shelter from surrounding microclimate.

Cavities

CV4

Dendrotelms and water-filled holes



CV41

$\varnothing \geq 3 \text{ cm}$ / trunk base

The entrance diameter is the same as the chamber's interior. Cup-shaped concavities that, due to their form, retain precipitation for a certain period of time (until it dries out).

CV42

$\varnothing \geq 15 \text{ cm}$ / trunk base



CV43

$\varnothing \geq 5 \text{ cm}$ / crown

The entrance diameter is the same as the chamber's interior. Cup-shaped concavities that, due to their form, retain precipitation for a certain period of time (until it dries out).

CV44

$\varnothing \geq 15 \text{ cm}$ / crown



Saproxyllic microhabitats	Description	Type	Code	Illustrations
---------------------------	-------------	------	------	---------------

Cavities

The entrance or exit diameter is the same as the interior's hole diameter. A bore hole net of xylophagous insects indicates a wood hole system. An insect gallery is a complex system of holes and chambers created by one or more insect species within the trunk.

Insect galleries and bore holes

Gallery with single small bore holes	CV51
Large bore hole $\varnothing \geq 2$ cm	CV52



CV5

Bark loss / exposed sapwood

Loss of trunk bark thus sapwood is exposed; caused e.g. by felling, natural falling of trees, rock fall. At the trunk base, bark loss may also be caused by skidding of logs, rodents, woodpecker sloughing.

Bark loss 25- 600 cm ² , decay stage < 3	IN 11
Bark loss > 600 cm ² , decay stage < 3	IN12
Bark loss 25- 600 cm ² , Decay stage = 3	IN13
Bark loss > 600 cm ² , decay stage = 3	IN14



IN1




Injuries and wounds

The tree has broken off at the trunk level, in a living tree. The tree is still alive and is developing a secondary crown with parts of the trunk decaying near the injury: the tree combines large decaying wood with xylem and phloem flux.

Exposed heartwood / trunk and crown breakage

Broken trunk, $\varnothing \geq 20$ cm at the broken end	IN21
--	------



IN2

Illustrations	Code	Type	Description	Saprophytic microhabitats
---------------	------	------	-------------	---------------------------



IN22

Broken tree crown / fork
Exposed wood ≥ 300 cm²

Exposed heartwood through the fork insertion breakage into the trunk - the rot initiates decaying substrate on the living tree.



IN23

Broken limb, $\varnothing \geq 20$ cm at the broken end


A 1st order branch has broken off. The tree is still alive. The injury provides a large entry gate for organisms and may develop into a cavity (rot hole) with xylem and phloem flux.



IN24

Splintered stem, $\varnothing \geq 20$ cm at the broken end

At wind breakage, trunk has splintered with several long splinters due to high force: splintered wound provides specific ecological conditions.



Cracks and scars



IN31

Length ≥ 30 cm;
width > 1 cm;
depth > 10 cm






Line-shaped injury (cleft) through the bark into the sapwood, exposing cambium and sapwood (not to be recorded if injury has occluded).

IN32

Length ≥ 100 cm;
width > 1 cm;
depth > 10 cm

Injuries and wounds

IN3

Saprophytic microhabitats	Description	Type	Code	Illustrations
Injuries and wounds	Bark loss and crack caused by lightning strike exposing the sapwood (not recorded when new bark has closed the scar).	Lightning scar	IN33	
	Fire scars at the lower trunk usually have a triangular shape and are located at the base of the tree on the leeward trunk side. Fire scars are associated with charred wood and eventually resin flow on exposed sapwood or bark.	Fire scar, $\geq 600 \text{ cm}^2$	IN34	
Bark	Space between bark and sapwood forming a shelter (open at the bottom).	Bark shelter, width > 1 cm; depth > 10 cm; height > 10 cm	Bark BA11	
	Space between bark and sapwood forming a pocket (open at the top), eventually containing mould.	Bark pocket, width > 1 cm; depth > 10 cm; height > 10 cm	BA12	
	Coarse and fissured bark, sometimes tree species specific.	Coarse bark	BA21	



Illustrations	Code	Type	Description	Saprophytic microhabitats
---------------	------	------	-------------	---------------------------

DE1



Dead branches and limbs / crown deadwood

DE11	\varnothing 10 - 20 cm, \geq 50 cm, sun exposed	Smaller sized (> 10 cm diameter) decaying wood, often horizontal or at a skewed angle, often in the shadow of the remaining canopy; in contact with living wood (xylem and phloem flow).
DE12	\varnothing > 20 cm, \geq 50 cm, sun exposed	
DE13	\varnothing 10 - 20 cm, \geq 50 cm, not sun exposed	
DE14	\varnothing > 20 cm, \geq 50 cm, not sun exposed	
DE15	Dead top $\varnothing \geq$ 10 cm	

Deadwood

Illustrations	Code	Type	Description	Epixylic microhabitats
---------------	------	------	-------------	------------------------

GR1



Root buttress cavities

GR11	$\varnothing \geq$ 5 cm	Natural cavity at the base of the tree trunk formed by the tree roots. May be densely covered with bryophytes. No wound or rot-hole.
GR12	$\varnothing \geq$ 10 cm	
GR13	Trunk cleavage length \geq 30 cm	Cleft formed by tree growth, no wound or open crack. Enclosure located higher at the trunk and therefore not part of the root buttress.

Deformation / growth form

Epixylic microhabitats	Description	Type	Code	Illustrations
------------------------	-------------	------	------	---------------

Deformation / growth form

Witches broom

Dense agglomeration of twigs caused by a parasite (such the fungi *Melampsorella caryophylacerum* or *Taphrina betulina*) or hemiparasite (genus *Arceuthobium*, *Viscaceae*).

Witches broom,
ø > 50 cm

GR21




GR2

Dense agglomeration of shoots on the trunk or branches of a tree. They originate from latent buds visible on the tree or can be submerged under the bark as epicormic buds.

Water sprout

GR22



Cankers and burrs

Proliferation of cell growth with rough bark and bark damage at the canker surface.

Cancerous growth,
ø > 20 cm

GR31



GR3

Decayed canker exposing necrotic tissue, e.g. caused by *Nectria spec.* on beech.

Decayed canker,
ø > 20 cm


GR32



Illustrations	Code	Type	Description	Epixylic microhabitats
---------------	------	------	-------------	------------------------

EP1

Fruiting bodies fungi



EP11 Annual polypores, $\varnothing > 5\text{cm}$

Fruiting bodies of polypores at the tree trunk lasting some weeks. European polypores only have one layer of tubes and are mostly of tough elastic soft consistency (no woody parts). Many species do not develop fruiting bodies each year. Main annual genera are *Abortiporus*, *Amylocystis*, *Bjerkandera*, *Bondar-zewia*, *Cerrena*, *Climacocystis*, *Fistulina*, *Gloeophyllum*, *Grifola*, *Haploporus*, *Inonotus*, *Ischnoderma*, *Laetiporus*, *Leptoporus*, *Meripilus*, *Oligoporus*, *Oxyporus*, *Perenniporia* pp, *Phaeolus*, *Piptoporus*, *Podofomes*, *Polyporus*, *Pycnoporus*, *Spongipellis*, *Stereum*, *Trametes*, *Trichaptum*, *Tyromyces* (underlined genera known to host a wide diversity / rare invertebrates).



EP12 Perennial polypores, $\varnothing > 10\text{cm}$

Woody, or at least tough fruiting bodies, showing distinct annual layers in the tube layer. Perennial fruiting bodies of the fungi indicating trunk decay caused by white rot (e.g. *Fomes fomentarius* (L. ex Fr.) Fr.) and brown rot (e.g. *Fomitopsis pinicola* (Swartz ex Fr.) Karst.). Main perennial genera are *Fomitopsis* pp, *Fomes*, *Perreniporia* pp, *Oxyporus*, *Ganoderma* pp, *Phellinus*, *Daedalea*, *Haploporus*, *Heterobasidion*, *Hexagonia*, *Laricifomes*, *Daedaleopsis* (underlined genera known to host a wide diversity / rare invertebrates).



EP13 Pulpy agaric, $\varnothing > 5\text{cm}$

Large, thick and pulpy or rather fleshy fruiting body of gill-bearing fungus (order Agaricales) - an agaric is a type of fungal fruiting body characterized by the presence of a pileus (cap) that is clearly differentiated from the stipe (stalk), with lamellae (gills) on the under-side of the pileus. "Agaric" can also refer to a basidiomycete species characterized by an agaric-type fruiting body. Examples: *Armillaria*, *Pleurotus*, *Megacollybia*, *large Pluteus* bear many arthropods and also parasitic fungi. The fruiting body remains generally several weeks.

Epiphytes

Epixylic microhabitats	Description	Type	Code	Illustrations
------------------------	-------------	------	------	---------------

Fungi cover of large tough hemispheric dark fungus looking like a lump of coal. Genus examples are *Daldinia* and *Hypoxylon*.

Large ascomycetes,
 $\varnothing > 5 \text{ cm}$

EP14




Myxomycetes

Amoeboid slime mould which forms moving plasmodium looking like gelatinous mass when fresh.

Myxomycetes,
 $\varnothing > 5 \text{ cm}$

EP21



Epiphytic crypto- and phanerogams

Epiphytes

Tree trunk covered by mosses and liverworts.

Epiphytic bryophytes
 coverage > 25 %

EP31



Tree trunk covered by foliose and fruticose lichens (lichens in combination with bryophytes).

Epiphytic foliose and fruticose lichens,
 coverage > 25 %

EP32



Lianas and other climbing plants cover the trunk surface (e.g. *Hedera helix*, *Clematis vitalba*).

Lianas,
 coverage > 25 %

EP33



Illustrations	Code	Type	Description	Epixylic microhabitats
---------------	------	------	-------------	------------------------



EP34

Epiphytic ferns,
> 5 fronds

Epiphytic ferns on trunk and large branches, often associated with bryophytes.



EP35

Mistletoe

Occurrence of these epiphytic and hemiparasitic plant species in the tree crown (*Viscum spp.*, *Arceuthobium spp.*, *Amyena spp.*, *Loranthus spp.*).

Epiphytes

Nests



NE11

Large vertebrate nest,
ø > 80 cm

Structures built by big raptors (eagles, black or white stork, grey heron) to hold eggs, offspring, or occasionally the animal itself. They may be composed of organic material such as twigs, grass, and leaves, and are located on branches, forks or witch brooms.



NE12

Small vertebrate nest,
ø > 10 cm

Nests built by small bird species, dormouse, mouse or squirrel.

Nests



NE21

Invertebrate nest

Larval nest of the pine processionary moth *Thaumetopoea pityocampa*, nest of the wood ant *Lasius fuliginosus* and of feral bees in tree trunk.

Sap and resin run



OT11



Sap flow,
> 50 cm

Fresh significant flow of sap, mainly at deciduous tree species.

Other

NE1

OT1


Epixylic microhabitats	Description	Type	Code	Illustrations
Other	Fresh significant flow of resin, at coniferous tree species.	Resin flow and pockets, > 50 cm	OT12	
	Result of micro-pedogenesis from epiphytic mosses, lichens or algae and necrosed old bark, it also depends on debris and litter fall from the crowns and bark.	Crown microsoil	OT21	<p style="text-align: center;">Microsoil</p> 
		Bark microsoil	OT22	

OT2



Integrate+ is a demonstration project funded by the German BMEL to establish a European network of demonstration sites for the integration of biodiversity conservation into forest management.

The Integrate+ project runs from December 2013 to December 2016 and builds on a partner network from research and practice with a focus on implementation of integrative management and enhancing transnational exchange of experiences.



European Forest Institute
Regional Office EFICENT
Wonnhaldestr. 4
79100 Freiburg, Germany

www.integrateplus.org
info@integrateplus.org