

**Bibliografia**

- AGUIAR, C., 2000. *Flora e vegetação da Serra da da Nogueira e do Parque Natural de Montesinho*. Tese de Doutoramento. Instituto Superior de Agronomia. Lisboa.
- COSTA, J.C., AGUIAR, C., CAPELO, J., LOUSÃ, M., NETO, C., 1998. Biogeografia de Portugal continental. *Quercetea* **0** : 5-56.
- COSTA, J.C., CAPELO, J., LOUSÃ, M., AGUIAR, C., 1993. Communautés de *Juniperus* au Portugal. *Colloques Phytosociologiques* **22** : 499-526.
- COSTA, J.C., CAPELO, J., LOUSÃ, M., ESPÍRITO SANTO, M.D., 1998. Sintaxonomia da vegetação halocasmofítica das falésias marítimas portuguesas (*Crithmo-Staticetea* Br.-Bl. 1947). *Itinera Geobotanica* **11** : 227-247.
- COSTA, J.C., CAPELO, J., ESPÍRITO SANTO, M.D., LOUSÃ, M., 1998. Aditamentos à vegetação do Sector Divisório Português. *Silva Lusitana* **10**(1) : 119-128.
- RIVAS-MARTÍNEZ, S., DÍAZ, T.E., FERNÁNDEZ-GONZÁLEZ, F., IZCO, J., LOIDI, J., LOUSÃ, M., PENAS, A., 2002. Vascular Plant Communities of Spain and Portugal. Adenda to the syntaxonomical checklist of 2001. Part I. *Itinera Geobotanica* **15**(1) : 5-432.

Ao longo do texto e nas tabelas, os nomes das plantas vasculares estão maioritariamente de acordo com a *Flora Iberica* (CASTROVIEJO *et al.*, 1986-2001) já publicada; AMARAL FRANCO & ROCHA AFONSO (1998) para as gramíneas e *Flora Europaea* (TUTIN *et al.*, 1968-1980, 1993) para os restantes grupos. A checklist de RIVAS-MARTÍNEZ, DÍAZ, F. DEZ-GONZÁLEZ, IZCO, LOIDI, LOUSA & PENAS (2002) *Itinera Geobotanica* 15(2): 697-813 foi também considerada. Por vezes os nomes estão abreviados até ao último epíteto infra-específico.

**Carlos Aguiar**, Escola Superior Agrária de Bragança. Inst. Polit. Bragança. [cfaguiar@ipb.pt](mailto:cfaguiar@ipb.pt); **José Carlos Costa**; Depto. Protecção de Plantas e Fitoeco-

logia. Instituto Superior de Agronomia. [jccosta@isa.utl.pt](mailto:jccosta@isa.utl.pt); **Jorge Capelo**; Depto. de Ecologia, Recursos Naturais e Ambiente. Estação Florestal Nacional. [jorge.capelo@efn.com.pt](mailto:jorge.capelo@efn.com.pt); **Anabela Amado**; Parque Natural do Douro Internacional, [xoa99@yahoo.com](mailto:xoa99@yahoo.com); **J. Honrado**; UGEV-CBIO/ICETA & Depto. Botânica, Universidade do Porto. [jhonrado@fc.up.pt](mailto:jhonrado@fc.up.pt); **M. Dalila Espírito Santo**; Depto. Protecção de Plantas e Fitoecologia. Instituto Superior de Agronomia.; [dalilaesanto@isa.utl.pt](mailto:dalilaesanto@isa.utl.pt) & **Mário Lousã**; Depto. Protecção de Plantas e Fitoecologia. Instituto Superior de Agronomia. [mariolousa@isa.utl.pt](mailto:mariolousa@isa.utl.pt)

**XXXIV: The vegetation of Madeira: II - woody caulirosetted communities of evergreen forest clearings: *Euphorbia melliferae* all. nova.**

The endemic caulirosetted microphanerophytes [with a rosette of leaves on top of a long woody few-branched naked stem] – *e.g.* *Euphorbia mellifera*, *Isoplexis sceptrum*, *Melanoselinum decipens*, *Musschia wollastonii* and *Sonchus fruticosus* – are among the most striking plants of Madeira Island. They are often found in the "levadas" artificial system of channels that runs through the *Ocotea foetens* forest [*Clethro arborea*-*Ocoteetum foetentis*]. These plants organize themselves in a particular phytocoenosis – *Isoplexido sceptri-Euphorbietum melliferae* ass. nova. – that reflects a worldwide recurrent phenomenon in forest ecosystems: the presence of plants adapted to cuts in the continuous crown layer of dense forest, such as dry ravines, forest clearings produced by tempests, landslides and other natural

disturbances. "Levadas" are artificial simulations of these natural habitats.

Since irregular streams, with rocky ground, found amidst the *Pruno-Lauretalia* forest share some of the above described habitat features, some caulirosetted elements are also found there. Thus, the *Isoplexido-Euphorbietum melliferae* phytocoenosis is somewhat floristically and physiognomically similar to the *Rhamno glandulosi-Sambucetum lanceolati* [riparian higrophyllous leafy shrub community, see note XXXV], but can easily be differentiated from it by the absence of *Rhamnus glandulosa* and *Sambucus lanceolata*, the scarcity of escio-higrophyllous ferns and the abundance of *Trifolio-Geranietae* characteristics.

The caulirosetted microphanerophytes share a common physiognomy [caulirosetted] and ecology [heliophyllous], belonging to the same functional group [C-strategist in the primary strategies of GRIME, 2001, *Plant Strategies, Vegetation Processes and Ecosystem Properties*] and many of them evolved from herbaceous ancestors (e.g. *Sonchus*: KIM *et al.*, 1996, *Proc. Natl. Acad. Sci. USA* 93: 7743-7748) that managed to adapt to an island without indigenous mammal herbivores, once almost totally covered by dense forest vegetation. A coarse comparison can be made with the *Trifolio-Geranietae* herbaceous vegetation, of which the *caulirosetulati* would be a woody evolutionary equivalent.

Physiognomical, ecological and floristical independence of this vegetation, as a special type among forest groupings [*Pruno-Lauretalia*]

suggests that the rank of *alliance* should be used: *Euphorbion melliferae*. The biogeographical scope of the proposed alliance should include, most probably, the vicariant *Euphorbia mellifera* dominated communities also found in the Canary Islands [c.f. RIVAS-MARTÍNEZ *et al.* (1993), *Itin. Geobot.* 7: 169-364].

\* ***Euphorbion melliferae*** Capelo, J.C. Costa, Jardim, Sequeira, Aguiar & Lousã *alliancia nova hoc loco*

[Typus: *Isoplexido sceptri-Euphorbion melliferae* Capelo, J.C. Costa, Jardim, Sequeira, Aguiar & Lousã *ass. nova*; characteristic taxa: *Euphorbia mellifera*, *Isoplexis sceptrum*\*, *Melanoselinum decipens*\*, *Musschia wollastonii*\*, *Sonchus fruticosus*\*; \*territorial of Madeiran Province]. Affiliated to the *Pruno hixae-Lauretalia novocanariensis* Oberdorfer ex Rivas-Martínez *et al.* 1977 *corr.* Rivas-Martínez *et al.* 2002; *PRUNO - LAURETEA NOVOCANARIENSIS* Oberdorfer 1965 *corr.* Rivas-Martínez *et al.* 2002].

1 - ***Isoplexido sceptri-Euphorbietum melliferae*** Capelo, J.C. Costa, Jardim, Sequeira, Aguiar & Lousã *associatio nova hoc loco*

[typus: relevé #2 table 1]

[taxonomical nomenclature follows: PRESS & SHORT (1994) *Flora of Madeira*. BM. London; and also the checklist of taxa of RIVAS-MARTÍNEZ, DÍAZ, F.DEZ-GONZÁLEZ, IZCO, LOIDI, LOUSA & PENAS (2002) - *Itinera Geobotanica* 15(2): 697-813. Sometimes names are shortened to the last infra-specific rank].

**Acknowledgment:** the authors would like to thank the most valuable support of the *Direcção Regional de Florestas da Secretaria Regional do Ambiente e Recursos Naturais da Região Autónoma da Madeira*.

**Table 1** - *Isoplexido sceptri-Euphorbietum melliferae*

# of relevé	1	2	3
m.s.m. (1=10m)	102	102	.
Area (m <sup>2</sup> )	400	400	300
Aspect	NW	NW	NE
<b>Characteristic combination</b>			
<i>Euphorbia mellifera</i>	2	3	3
<i>Isoplexis sceptrum</i>	1	2	1
<i>Sonchus fruticosus</i>	3	2	1
<i>Erysimum bicolor</i>	1	1	1
<i>Melanoselinum decipiens</i>	2	1	.
<i>Musschia wollastonii</i>	1	2	.
<b>Characteristic of higher syntaxa</b>			
<i>Clethra arborea</i>	.	1	+
<i>Teline maderensis</i>	+	.	.
<i>Erica maderinicola</i>	+	+	.
<i>Rubus bollei</i>	+	.	.
<i>Rubia agostinhoi</i>	+	.	.
<i>Vaccinium padifolium</i>	.	+	+
<i>Phyllis nobla</i>	1	2	2
<i>Sibthorpia peregrina</i>	.	.	1
<i>Rosa mandonii</i>	.	+	.
<i>Ocotea foetens - pl.</i>	.	.	1
<i>Succisa pratensis</i>	+	1	.
<i>Origanum virens</i>	.	+	.
<i>Geranium palmatum</i>	1	2	1
<i>Pericallis aurita</i>	+	.	+
<i>Bystropogon punctatus</i>	+	1	+
<i>Dactylorhiza foliosa</i>	+	.	.
<i>Ageratina adenophora</i>	1	.	.
<i>Rubus bollei</i>	.	+	+
<i>Argyranthemum pinnatifidum</i>	+	+	.
<i>Erigeron karvinskianus</i>	+	1	+
<i>Hypochaeris radicata</i>	.	+	.
<i>Festuca donax</i>	+	1	.
<i>Pteris incompleta</i>	.	+	1
<i>Cirsium latifolium</i>	.	+	+
<i>Tolpis macrorhiza</i>	.	1	.
<i>Cedronella canariensis</i>	+	.	.
<i>Stegnogramma pozoi</i>	.	+	+
<i>Arachniodes webbiana</i>	.	+	.
<i>Aichryson divaricatum</i>	+	.	+
<i>Ranunculus cortusifolius</i>	.	+	+
<i>Carex peregrina</i>	.	+	.
<i>Blechnum spicant</i>	.	+	.
<i>Carex lowei</i>	.	+	+
<i>Dryopteris aitoniana</i>	.	+	.
<i>Davallia canariensis</i>	.	+	.
<i>Brachypodium sylvaticum</i>	.	1	+
<i>Doodia caudata</i>	.	+	.

**Sites:** 1,2 Folhadal; 3: Fajã da Nogueira.

**Jorge Capelo;** Depto. de Ecologia, Recursos Naturais e Ambiente. Estação Florestal Nacional, jorge.capelo@efn.com.pt. **José C. Costa;** Depto. Protecção de Plantas e Fitoecologia, Instituto Superior de Agronomia, Lisboa, jccosta@isa.utl.pt. **Roberto Jardim;** Jardim

Botânico da Madeira, Funchal, robertojardim.sra@gov-madeira.pt.

**Miguel Sequeira;** Depto. Biologia, Universidade da Madeira, Funchal, sequeira@uma.pt. **Carlos Aguiar;** Escola Superior Agrária de Bragança, cfaguiar@ipb.pt. **Mário Lousã;** Depto. Protecção de Plantas e Fitoecologia, Instituto Superior de Agronomia, Lisboa, mariolousa@isa.utl.pt.

**XXXV: The vegetation of Madeira: III - *Diplazio caudati-Perseetum indicum* ass. nova and *Rhamno glandulosi-Sambucetum lanceolati* ass. nova: two new hygrophillic forest associations from Madeira Island .**

Descriptions of forest vegetation of Madeira Island included, so far, two types of climatophylous broadleaf forest vegetation – *Clethro arboreae-Ocoteetum foetentis* and *Semele androgynae-Apollonietum barbujanae* – with an arboreal *stratum* respectively dominated by two *Lauraceae* trees: *Ocotea foetens* [til] and *Apollonia barbujana* [barbusano]. Recently, we discovered that the other *Lauraceae* Madeira's tree – *Persea indica* [vinhático] – is the dominant tree in a third type of broadleaf forest - *Diplazio caudati-Perseetum indicum* ass. nova. This is an edaphohygrophylous forest, proper of middle and lower-course permanent streams, over fairly thick colluvial (rarely alluvial) substrata. It ranges from the termomediterranean sub-humid stage to the mesomediterranean humid stage. Its characteristic combination involves: *Persea indica*, *Dryopteris aitoniana*, *Woodwardia radicans*, *Diplazium caudatum*, *Pteris incompleta*, etc. Nowadays, the *Diplazio-Perseetum indicum* is a rare phytocoenosis because *Persea indica* is a source of noble wood and many of its