

ANTIOXIDANT AND ANTI-INFLAMMATORY ACTIVITIES OF *CYTISUS MULTIFLORUS*



Susana C.S.^{ab}, Olívia R.P.^{ac}, Joana L.^{dg}, Maria T.B.^{bd}, Maria M.R.D.^e, Maria T.C.^{bg}, Susana M.C.^{af}

^aCERNAS, Escola Superior Agrária, Instituto Politécnico de Coimbra, Coimbra, Portugal

^bFaculdade de Farmácia, Universidade de Coimbra, Azinhaga de Santa Comba, 3000-548 Coimbra, Portugal

^cDepartamento de Tecnologias de Diagnóstico e Terapêutica, Escola Superior de Saúde, Instituto Politécnico de Bragança, Bragança, Portugal

^dCentro de Estudos Farmacêuticos, Azinhaga de Santa Comba, 3000-548 Coimbra, Portugal

^eCentro de espectrometria de massa, Departamento de Química, Universidade de Aveiro, 3810-193 Aveiro, Portugal

^fCIMO, Escola Superior Agrária, Instituto Politécnico de Bragança, Bragança, Portugal

^gCentro de Neurociências e Biologia Celular, Departamento de Zoologia da Universidade de Coimbra, 3004-517 Coimbra, Portugal



INTRODUCTION

Many traditional medicinal plants are potential candidates for finding new therapeutic and supplementary health products. *Cytisus multiflorus*, (White Spanish Broom), is used in folk medicine in the Iberian Peninsula, where it is claimed to have various health benefits, including diuretic, hypnotic, anxiolytic, antiparasitic, antidiabetic, antioxidant and anti-inflammatory properties¹. The usage of this plant is however, totally based on the available ethnopharmacological information, as no scientific data regarding its biological effects has been delivered. In this sense, is the aim of this work to contribute to the scientific knowledge of the antioxidant and anti-inflammatory properties of *C. multiflorus*.

METHODS

The ethanolic extract from flowers of *C. multiflorus* was prepared by extraction with an 80% ethanolic solution (v/v), as previously described¹. The total phenolic content of the extract was determined following the Folin-Ciocalteu procedure and the main phenolic constituents were identified and quantified by combined HPLC-DAD and ESI-MSⁿ analysis¹. The antioxidant abilities of the *C. multiflorus* extract were evaluated through the DPPH scavenging² and reducing power³ assays. The assessment of cell viability in the presence of distinct concentrations of the extract was performed by the MTT reduction colorimetric assay⁴ and the anti-inflammatory activity of a non-toxic extract concentration was assessed by its nitric oxide inhibition ability, as measured by the Griess assay, on lipopolysaccharide-stimulated Raw 264.7 macrophages.

Fig. 1- *Cytisus multiflorus*



RESULTS AND DISCUSSION

Tab.1- Phenolic content and antioxidant activity of *C. multiflorus* ethanolic extract

Mass (%)	^a Total phenolic (mg GAE/g)	^b DPPH (EC ₅₀) (µg/ml)	^c Reducing Power (EC ₅₀) (µg/ml)
22.96%	140.39 ± 11.67	13.4 ± 1.0	11.4 ± 2.1

Mean Values ± standard derivations of three replicate analyses

^a Data expressed as milligrams of gallic acid equivalents (GAE) per gram of extract; ^b EC₅₀ – concentration for a 50 % inhibition of the 60 µM radical 2,2-diphenyl-1-picrylhydrazyl (DPPH); ^c Amount of extract able to providing 0.5 of absorbance by reducing of 3.5 µM Fe³⁺ to Fe²⁺.

Tab.2- Effect of *C. multiflorus* ethanolic extracts in HepG2 viability

Condition	Cell Viability (%)
Control	100
LPS (1µg/mL)	82.25 ± 1.14
<i>C. multiflorus</i> (325 µg/mL)	90.67 ± 14
<i>C. multiflorus</i> (325 µg/mL) + LPS (1µg/mL)	86.93 ± 4.1
<i>C. multiflorus</i> (161 µg/mL)	104.78 ± 4.1
<i>C. multiflorus</i> (161 µg/mL) + LPS (1µg/mL)	94.01 ± 13

The main phenolic constituents of *C. multiflorus* were chrysin-7-*O*- glycopyranoside and a dihydroxyflavone isomer of chrysin, which accounted for 49.4±7.3 mg/g and 21.8±3.8 mg/g, respectively. As indicated by its low EC₅₀ values (13.4±1.0 and 11.4±2.1 µg/mL for DPPH scavenging potential and reducing power, respectively), the *C. multiflorus* ethanolic extract has a high antioxidant capacity. Moreover, the extract did not cause cytotoxicity against RAW 264.7 macrophages for concentrations up to 325 µg/mL and the treatment of this cell line with 160,8 µg/ml and 325 µg/ml of the extract induced a decrease in the levels of NO of 23.9 and 32.1%, respectively.

REFERENCES

- [1] Pereira O. R. *et al.* (2012). Food Chem 131, 652-659
- [2] Ferreira A. *et al.* (2006). J Ethnopharmacol, 108, 31-37
- [3] Oyaizu M. *et al.* (1986). Jpn J Nutr, 44, 307-15
- [4] Mosmann T (1983). Immunol Methods 65(1-2), 55-63




Fig.2- Chromatographic profile of the *C. multiflorus* ethanolic extract at 280 nm




Fig.3- Effect of *C. multiflorus* extract (CME) in the nitrite production of macrophages stimulated with LPS 1 µg/mL

CONCLUSION

The gathered data suggests that *C. multiflorus* is in fact a good antioxidant and anti-inflammatory plant, as believed by the folk knowledge.

ACKNOWLEDGEMENTS

The authors acknowledge the financial support provided by the FCT to CERNAS (project PEst-OE/AGR/UI0681/CNC/CEF/2011). Olívia R Pereira was supported by a PhD grant (SFRH/PROTEC/49600/2009).