

# Cloning, characterization and *in vitro* and *in planta* expression of a glucanase inhibitor protein (GIP) of *Phytophthora cinnamomi*

Ivone M. Martins · Fátima Martins ·  
Hélio Belo · Madalena Vaz · Marisa Carvalho ·  
Alfredo Cravador · Altino Choupina

Received: 20 March 2013 / Accepted: 4 January 2014  
© Springer Science+Business Media Dordrecht 2014

**Abstract** Oomycetes from the genus *Phytophthora* are fungus-like plant pathogens that are devastating for agriculture and natural ecosystems. They are able to secrete a glucanase inhibitor protein (GIP) that inhibits the activity of endoglucanases (EGases) involved in defense responses against infection. One of the most widely distributed and aggressive *Phytophthora* species, with more than 1,000 host plants is *P. cinnamomi*. In this work we report the sequencing and characterization of a class of GIPs secreted by *Phytophthora cinnamomi*. The *gip* gene from *P. cinnamomi* has a 937 bp ORF encoding a putative peptide of 312 deduced amino acids. The expression of this gene was studied during growth in different carbon sources (glucose, cellulose and sawdust), by RT-qPCR and its level of expression was evaluated at five time points. The highest expression of *gip* gene occurred in sawdust at 8 h of induction. *In vivo* infection of *C. sativa* revealed an increase in *gip* expression from 12 to 24 h. At 36 h its expression decreased suggesting that a compensatory mechanism must occur in plant.

**Keywords** *Castanea sativa* · GIP · HE-TAIL PCR · *Phytophthora cinnamomi* · RT-qPCR

## Introduction

The class oomycetes or “water moulds” form one of several lineages within the Kingdom Chromista that independently evolved to a parasitic lifestyle and consequently are thought to have developed alternative mechanisms of pathogenicity. Oomycetes share with many bacterial, fungal and nematode plant pathogens the requirement for living host tissue for at least part of the infection cycle. To establish infection, these pathogens must evade, suppress or manipulate host defenses. Due to their particular physiological characteristics, no efficient treatments against diseases caused by these microorganisms are presently available. To develop such treatments, namely against *Phytophthora*, it appears essential to dissect the molecular mechanisms that determine the interaction between *Phytophthora* species and host plants.

*Phytophthora cinnamomi* Rands is a destructive, widespread soil-borne pseudofungus belonging to the class oomycetes that infects many plant hosts including forest, ornamental and fruit species. It has been often associated with the decline of forest species, as well as other woody perennial plant species [1]. For instance, the ink disease of *Castanea sativa* Mill, one of the most destructive diseases in *C. sativa* is caused by this oomycete. The most common symptoms are root necrosis and reduction in root growth, which invariably lead to the trees death [2].

*Phytophthora cinnamomi*, whose life cycle develops integrally in the soil, requires moist soil conditions and warm temperatures to be active, but visible damage caused by the disease most often occurs in summer when plants are drought stressed. This pathogen can grow saprophytically in the soil and persists in soil or infected plant material as chlamydozoospores and, to a lesser extent, as oospores. It is known to survive for as long as 6 years in

I. M. Martins · F. Martins · H. Belo · M. Vaz · M. Carvalho ·  
A. Choupina (✉)  
CIMO, Escola Superior Agrária de Bragança, Instituto  
Politécnico de Bragança, Campus de Santa Apolónia,  
Apartado 1172, 5301-855 Bragança, Portugal  
e-mail: albracho@ipb.pt

A. Cravador  
IBB/CGB, Universidade do Algarve, Campus de Gambelas,  
8005-139 Faro, Portugal

moist soil, and it is clear that moisture is a key factor in the establishment, spread and longevity of this pathogen [3].

Identification and diagnostic techniques for *P. cinnamomi* require expert attention. Preventative measures and chemical application are the typical forms of control for this pathogen [4]. By the moment there are no eradication methods available.

Several key cellular events, such as adhesion to the host surface, penetration, and colonization of host tissue, take place during plant infection by oomycetes that can also manipulate biochemical and physiological processes in their host plants through a diverse array of virulence or avirulence molecules, known as effectors [5–8]. In susceptible plants, these effectors promote infection by suppressing defense responses, enhancing susceptibility, or inducing disease symptoms. Alternatively, in resistant plants, effectors are recognized by the products of plant resistance genes, resulting in host cell death and effective defense response known as the hypersensitive response (HR) restricting the pathogen to an area of scorched earth [7, 9]. *Phytophthora* effectors that suppress host defense responses have been described in several pathosystems [10–12]. Suppression of host defenses can occur through the production of inhibitor proteins that target host enzymes [13].

Early observations that plants secrete inhibitor proteins that bind and inactivate microbial hydrolases, specifically the binding of plant polygalacturonase inhibitor proteins (PGIPs) to fungal polygalacturonases, spurred the search for analogous inhibitor proteins from microbes that might inactivate the  $\beta(1,3)$ endoglucanases or chitinases of the plant host [14]. Glucanase inhibitor protein (GIP), is thought to function as counter defensive molecule that inhibits the degradation of  $\beta(1,3)$  and  $\beta(1,6)$ glucans in the pathogen cell wall and/or the release of defense-eliciting oligosaccharides by host  $\beta(1,3)$ endoglucanases [9].

Pathogens secrete endo $\beta(1,4)$ glucanases, which hydrolyze a range of plant cell wall glucans, while the host counters by synthesizing endo $\beta(1,3)$ glucanase that specifically degrade the hyphal wall  $\beta(1,3)$ ,  $\beta(1,6)$ glucans, rendering the pathogen susceptible to cell lysis. The  $\beta(1,3)$ ,  $\beta(1,6)$ glucan elicitors that are released during this process are perceived by the plant, by a presumed plasma membrane-bound receptor, and a signaling cascade is initiated, resulting in the up-regulation of a number of plant defenses.

Since the recent identification of GIPs, studies have focused on their biochemical characterization, addressing features such as binding properties and ligand specificity. The observation that these proteins bind strongly to enzymes that degrade cell wall polysaccharides, together with various sets of expression data, suggests that they are likely to influence plant–pathogen interactions. However,

their biological significance has not yet been established. Many questions also remain at the molecular level, such as the identity of the domains and key residues of the inhibitor proteins that contribute to the recognition specificity and high avidity binding.

The objective of the present work was to clone and characterize a gene from *P. cinnamomi* homologous to *gip* genes, and analyze their expression when this species grows in cultures with different carbon sources and during host plant infection.

## Materials and methods

### Biological material

Isolate Pr120 of *P. cinnamomi* was kindly provided by Eugénia Gouveia from Instituto Politécnico de Bragança. It was isolated from soil samples associated with a *C. sativa* tree affected by the ink disease in Trás-os-Montes region (northeast Portugal). The strain was grown in the dark for 4–6 days at 22–25 °C in PDA (Potato-Dextrose Agar) medium.

Total genomic DNA from *P. cinnamomi* mycelium was obtained as described in [15].

*C. sativa* chestnuts were surface sterilized, germinated in sterile vermiculite and grown in a greenhouse until their root length reached 5–6 cm.

### Plasmids

pET-28a(+) (Novagen), is a 5369 bp *Escherichia coli* expression vector containing the kanamycin resistance gene. It is regulated by the T7 promoter (present as a 16 bp fragment) and by the T7 terminator (present as a 46 bp).

pIM3.1, a 6.3 kb construct [based on pET28a(+)], contains an expression cassette consisting of the GIP gene (*gip*), present as a 937 bp *SacI-HindIII* fragment. This was obtained by polymerase chain reaction (PCR) amplification of a 937 bp ORF of *gip* from *P. cinnamomi* strain 120 DNA, using as primers SacGIP (5'-**GAG CTC GAC ATG TTT ACT TCC GG**, insertion of restriction site *SacI* in bold) and HindGIP (5'-**CCA AGC TTG TCC TTA TGC CTT GAT G**, insertion of restriction point *HindIII* in bold), digesting this fragment with *SacI-HindIII* and finally cloning it into *SacI-HindIII* digested pET-28a(+). The DNA digested vector band was purified in a low melting point (LMP) agarose gel and Wizard® SV Gel and PCR Clean-Up System (Promega), following the manufacturer's instructions. Correctly oriented clones were selected by restriction analysis with *XhoI*. The positive clone obtained was sequenced to confirm the correct integration of the insert. The sequencing of the fragments of the DNA

obtained was carried through in an automatic sequencer ABI Prism 377TM from Applied Biosciences (Foster City, CA, USA).

#### Amplification of a short fragment of the *gip* gene

A PCR was used to amplify a 308 bp fragment of the *gip* gene. Degenerate oligonucleotide primers M1 (5'-GAG AAG CCC AGC AAG TTC SCC) and M3 (5'-GCS GAC ACG CGM GMG TAC ACA) were designed based on *gip*, sequences alignment from *P. sojae* and *P. infestans* [14]. The PCR cycling conditions were 94 °C/5 min, followed by 36 cycles of 94 °C/1 min; 63 °C/1 min; 72 °C/30 s, and ending with 72 °C/5 min. Each 50 µl PCR contained 0.8 mM dNTP, 0.2 mM of each primer, 100 ng genomic DNA, and 1 U Taq DNA polymerase in the appropriate buffer. Aliquots of the PCR reactions were separated on 0.8 % w/v agarose gel electrophoresis and stained with ethidium bromide, to check for the presence of the expected amplicon.

#### Amplification of unknown genomic DNA sequence of *gip*

The high-efficiency thermal asymmetric interlaced-PCR (HE-TAIL PCR) is an efficient method to amplify unknown genomic DNA sequences adjacent to short known regions by flanking the known sequence with asymmetric PCR. In this procedure gene-specific primers M1, M3, M2 (5'-GCC GTT YTC CTT GAT CAG CGG), S1 (5'-AGG CGT TGT CGC CCC AGA CC), S2 (5'-CGG CCG CGG TGA CGC TGA CG) and S4 (5'-GGT CTG GGG CGA CAA CGC CT) were used. Degenerated primers R1 (5'-NGT CGA SWG ANA WGA A), R2 (5'-GTN CGA SWC ANA WGT T), R3 (5'-WGT GNA GWA NCA NAG A) and R4 (5'-NCA GCT WSC TNT SCT T) were applied [16]. Three rounds of PCR were performed using the product of each previous PCR as a template for the next. A detailed cyler program is given in Table 1.

The primary PCR was performed in a 50 µl volume containing 80 ng genomic DNA, 0.2 mM of primers S1 or S3, 2 mM of a random primer (R1, R2, R3, R4), 0.2 mM of each dNTP and 1U Taq DNA polymerase in the appropriate buffer. The secondary PCR was performed with primers M1 or M3 (0.2 mM) and the same random primer R (2 mM) as used in the primary reaction. 1 µl 1/50 dilution of the primary PCR was used as a template. Single-step annealing-extension PCR consisting of a combined annealing and extension step at 65 or 68 °C was used in primary and secondary PCR reactions. The tertiary reaction was carried out with 1 µl of 1/10 dilution of the secondary reaction, 0.2 mM primers S1 and S2, 0.2 mM random primer R (the same as used in the previous cycles), 0.2 mM

**Table 1** HE-TAIL PCR cycle settings

Reaction	Number of cycles	Thermal settings
Primary	1	93 °C (1 min); 95 °C (5 min)
	5	94 °C (30 s); 62 °C (1 min); 72 °C (2 min 30 s)
	1	94 °C (30 s); 25 °C ramping 72 °C (3 min); 72 °C (2 min 30 s)
	15	94 °C (20 s); 65 °C (3 min 30 s); 94 °C (20 s); 65 °C (3 min 30 s); 94 °C (30 s); 42 °C (1 min); 72 °C (2 min 30 s)
Secondary	1	72 °C (5 min); 4 °C Hold
	12	94 °C (20 s); 65 °C (3 min 30 s); 94 °C (20 s); 65 °C (3 min 30 s); 94 °C (30 s); 42 °C (1 min); 72 °C (2 min 30 s)
	1	72 °C (5 min); 4 °C Hold
Tertiary	1	72 °C (5 min); 4 °C Hold
	30	94 °C (30 s); 42 °C (1 min); 72 °C (2 min 30 s)
	1	72 °C (5 min); 4 °C Hold

each dNTP, 1U DNA Taq polymerase in the appropriate buffer. To exclude nonspecific amplification, a tertiary control reaction R–R was set up without adding gene-specific primers.

#### Amplification of *gip* ORF

A PCR was used to amplify the 937 bp ORF of the *gip* gene. The PCR cycling conditions were 95 °C/3 min, followed by 40 cycles of 94 °C/1 min, 61 °C/1 min, 72 °C/3 min, and ending with 72 °C/10 min. Each 25 µl PCR contained 1.6 mM dNTP, 0.2 mM of each primer, 100 ng genomic DNA, 1.5 mM MgCl<sub>2</sub> and 0.05 U Taq DNA polymerase in the appropriate buffer. Aliquots of the PCR reactions were separated on 0.8 % w/v agarose gel electrophoresis and stained with ethidium bromide, to check for the presence of the expected amplicon.

#### Bacteria transformation and DNA extraction

Plasmids were propagated in *E. coli* (DH5α cells), following the procedure described by [17], and plasmid DNA was extracted-purified with the Wizard<sup>®</sup> Plus SV Minipreps DNA Purification System (Promega), following the manufacturer's instructions.

#### Analysis of GIP production

pIM3.1 clone was harvested in LB with kanamycin and protein expression was induced by adding 1 mM IPTG at 0, 2, 4, 6 and 8 h of growth. Secreted proteins were

concentrated by precipitation with trichloroacetic acid (TCA 8 % v/v), from the supernatant and with a lysis solution (50 mM Tris–HCl pH 7.5, 50 mM EDTA e 1 mM PMSF) from *E. coli* cells.

Protein quantification was performed by using the Protein quantification kit-general use (Fluka), following the manufacturer's instructions. After quantification, secreted proteins at all induction time points (0, 2, 4, 6 and 8 h) were separated by SDS-PAGE (15 % w/v). The membrane was stained with Coomassie brilliant blue BG-250 0.25 % (BioRad), and then was washed in a solution of methanol 45 % (v/v) and acetic acid 10 % (v/v).

#### Expression of *gip* gene

A Real-Time Quantitative PCR (RT-qPCR) assay was performed in order to analyze gene expression *in vitro* and *in vivo*.

#### *In vitro*

*P. cinnamomi* was grown in different carbon sources (2 % (w/v) glucose, 0.2 % (w/v) cellulose and 0.2 % (w/v) sawdust) and the expression of *gip* gene was evaluated at 2, 4, 6 and 8 days of growth. Control cells were grown in dextrose-containing medium (PDA). The expression level of the *gip* gene was expressed relative to the expression level of the housekeeping gene *actin2*, measured at 4 different times of induction with glucose, cellulose and sawdust.

#### *In vivo*

*C. sativa* roots were covered with fully colonized PDA and incubated in the dark at 25 °C for 12, 24 and 36 h. Negative controls were provided by roots in contact with non-colonized agar. After the incubation period, the agar was removed, along with all external mycelia growth. The roots were examined for the presence and extent of necrosis and then frozen to –80 °C. The assays were repeated three times. The expression level of the *gip* gene was expressed relative to the level measured in the reference gene *actin2*. Results were normalized to *actin2* gene and calculated using the  $2^{-\Delta\Delta Ct}$  method, as fold change relative to control cells.

Total RNA was isolated from *P. cinnamomi* mycelia using the Rneasy Plant Mini Kit (Qiagen), following the manufacturer's instructions. Residual DNA was removed by DNase I (Qiagen) treatment, following the

manufacturer's instructions. The integrity of the RNA was assessed by formaldehyde agarose gel electrophoresis (1.5 % agarose). 1 µg of RNA was reverse transcribed the iScript™ cDNA Synthesis Kit (BioRad) primed with oligo (dT), following the manufacturer's instructions. The qPCR was performed with IQ™ SYBR® Green Supermix (Bio-rad) Real Time using a MiniOpticon™ Real-Time PCR Detection System (Biorad). Each 25 µl reaction contained 100 ng RNA, as well as 12.5 µl IQ SYBR Green Supermix (100 mM KCl, 40 mM Tris–HCl pH 8.4, 0.4 mM of each dNTP, 50 U/ml iTaq DNA polymerase, 6 mM MgCl<sub>2</sub>, 20 nM SYBR Green I fluorescein, and stabilizers) and 1.25 µM of each primer. The reactions were run in triplicate and incubated at 95 °C for 3 min, followed by 40 cycles of 95 °C/30 s, 61 °C/30 s and 72 °C/30 s. The endogenous control was normalized to the expression levels of *actin2* gene within the same sample. Amplification primers (Table 2) were targeted to the coding regions of *P. cinnamomi actin2* and *gip* genes.

#### Southern blot analysis

Genomic DNA of *P. cinnamomi* strain PR120 was isolated as described above. 12 µg of DNA was digested with *Apa*I, *Xho*I–*Cl*aI and *Eco*RI restriction enzymes and size-separated by electrophoresis on a 1 % agarose gel. DNA was blotted onto a Hybond-N<sup>+</sup> membrane (Amersham) by capillary transfer [18], and hybridized at high stringency at 65 °C, following the instructions for DIG Wash and Block Buffer Set (Roche Diagnostics, Basel, Switzerland).

DIG-labeled probes were generated by PCR using the primers SacGIP and HindGIP described above which amplify an ORF 937 bp fragment of the *gip* gene, following the protocol of the PCR DIG Probe Synthesis Kit (Roche Diagnostics, Basel, Switzerland). Blots were washed twice in 2× SSC and 0.5 % SDS at room temperature for 5 min, and subsequently washed twice in 0.5× SSC and 0.5 % SDS at 65 °C for 15 min. Detection was performed at room temperature following the instructions for DIG Wash and Block Buffer Set (Roche Diagnostics).

## Results and discussion

This work reports the sequencing, using the HE-TAIL PCR [16] of a class of GIPs, that are secreted by *P. cinnamomi* and that specifically inhibit the endoglucanase activity of

**Table 2** Sequences of primers used to amplify *actin2* and *gip* genes in the RT-qPCR experiments

Target gene	Forward primer sequence (5' → 3')	Reverse primer sequence (5' → 3')
<i>actin2</i>	GGCCTCGAGAAGAGCTACG	CTTCATGATGGTCTGGAACG
<i>gip</i>	TGGTGGTGCTCTCATCAGTC	CGGAAGCTTGCTGGATTCTC

their plant host, thereby suppressing the degradation of glucans in the oomycete cell wall and the release of oligoglucoside elicitors.

We obtained a small sequence of 308 bp by amplification using degenerated primers designed based on homology in the open reading frames of other GIPs (*P. sojae*), and the full length gene sequence (1301 bp) was obtained by flanking the known sequence with asymmetric PCR. *gip* Gene encodes a 312 aa protein that shares significant structural similarity with the chymotrypsin class of serine proteases (SP), but lack proteolytic activity due to the absence of an intact catalytic triad (His-Asp-Ser) and, thus, belong to a broader class of proteins called serine protease homologs (SPH).

GIPs show high homology with the S1A subfamily of SP; however questions remain about the expression patterns and potential roles of different GIPs during pathogenesis and their possible interaction with host EGases in the plant apoplast. In order to understand gene function, we analyzed gene expression during growth in different carbon sources and also performed a time course of *gip* expression by RT-qPCR procedure. The ORF (937 bp) of *gip* gene was cloned into a different expression plasmid in order to perform the analysis of heterologous expression, in *E. coli*.


FAST alignments revealed significant homologies (60–84 %) of the GIP sequence of *P. cinnamomi* with sequences of *P. sojae* and *P. infestans* including regulatory sequences as trypsin-like SP. Many of these are

synthesized by the precursor inactive zymogens and are cleaved during their limited proteolysis to generate active forms [14].

#### Amplification and characterization of the *gip* gene of *P. cinnamomi*

Starting with the amplification of a small 308 bp fragment of the *gip* gene obtained from *P. cinnamomi* strain 120 DNA, the application of the HE-TAIL PCR method led to the successful amplification of the entire sequence of the gene (GenBank: AM259384.1) with a total length of 1301 bp and a 937 bp ORF which encodes a 312 aa protein.

The genus *Phytophthora* is one of only a few eukaryotic genera for which at least three complete genome sequences are available, thereby allowing an unusually broad view of gene family evolution. The phylogeny provides eight examples of genes represented by at least one ortholog in each of the three species and two other cases in which an ortholog appears to have been duplicated, lost, or both in one of the three species. Examples include both GIPs (for example, GIP1 has a full set of orthologs and paralogs while GIP2 has orthologs but no paralogs) and apparently functional SP, such as PiSP1, which has a full set of orthologs. It is estimated that there are at least five GIP-related SP and at least five GIP-related SPH prior to the diversification of *Phytophthora* spp. The gene organization



**Fig. 1** Sequence alignment of GIP proteins and Ser Proteases. The N-terminal signal sequence is indicated by a horizontal bar. Conserved Cys residues involved in disulfide bond formation are indicated by asterisks. The position of the His57, Asp102, and Ser195

residues of the catalytic triad are indicated with arrows. Sites that are predicted to form SP surface loops 1 and 2 are boxed, and residues forming the walls of the substrate binding pocket are underlined boxes

of GIPs, SPH, and SP was assessed by searching their respective locations in each of the genome sequences. For all three species, the GIP genes and some of the GIP-like genes were clustered in a scaffold, or supercontig, while the remaining genes were distributed among other supercontigs [19].

A BLAST search of the databases revealed that the GIPs protein sequences are homologous with the chymotrypsin subclass of SP. The overall amino acid sequence homology of the GIPs with a number of chymotrypsins was similar to that typically observed among SP (~20–40%), and a number of characteristic features of SP were also present, including many sequence motifs and six cysteine residues (Fig. 1). A characteristic feature of SP is the catalytic triad charge relay system, referred to as His-57, Asp-102, and Ser-195 that are essential for the proteolytic function. However, interestingly none of the predicted GIPs possess this intact “catalytic triad” [14]. Therefore, while GIPs strongly resemble SP, they are not expected to function as proteases. The triad of *P. cinnamomi* GIP is Ser-57, Asn-102 and Thr-195.

Although GIPs are proteolytically inactive, several stretches of amino acids and motifs that are highly conserved among SP, are present in the GIPs. Firstly, an N-terminal signal sequence is present that targets the enzyme for secretion. Secondly, the GIPs possess three characteristically spaced pairs of Cys residues that are involved in the formation of disulfide bonds. Thirdly, key functional residues are present in the S1A specific substrate binding pocket and the two regions that flank the catalytic residues in the binding pocket form so-called variable surface loops 1 and 2. These surface loops contribute to the geometry of the SP specificity pocket and act synergistically to influence selective substrate binding and catalysis. A phylogenetic analysis of the GIP sequences aligned

with other S1A clan SP from a number of evolutionarily diverse organisms revealed that the GIPs form a distinct group.

The substantial amount of literature related to SP structure and function provides a valuable opportunity to make structural predictions about GIPs and to identify regions of the proteins that are involved in binding to the  $\alpha$ -endoglucanases ligands or that may dictate ligand specificity (Fig. 2).


The overall geometry of the two classes of proteins is clearly predicted to be very similar and the software analysis designated the structure as a high confidence prediction. The model in Fig. 5c assumes that the inhibitor uses a trypsin-like mechanism of recognition [20] whereby GIP identifies a Lys residue on the surface of the glucanase molecule.

GIPs represent the first example of proteolytically inactive SP homologs that function as enzyme inhibitors. It can be hypothesized that a major role for GIP is to suppress the release of glucan elicitors during *P. cinnamomi* infection, thereby reducing the effectiveness of the plant host’s surveillance system [19, 21].

#### Protein production assay


The expression of the *gip* gene of *P. cinnamomi* was induced in *E. coli* transformed with the plasmid pIM3.1 resulting from the insertion of the 937 bp *gip* ORF in the plasmid pET28a(+). The aim of this assay was to study the *gip* gene in a heterologous organism, *E. coli* in our case. For that have we used the pET-28a(+) expression system to clone the gene into the bacteria.

The band corresponding apparently to the 31 kDa GIP protein is shown with the arrow, 8 h after induction (Fig. 3).



**Fig. 2** **a** The crystal structure of R117 H mutant rat anionic trypsin complexed with bovine pancreatic trypsin inhibitor BPTI (PDB 1C07) was used as a template to predict the corresponding structure of *P. cinnamomi* GIP; in **b** using a computational approach (ESyPred3D). The catalytic triad of trypsin and the equivalent residues of GIP are

colored. *Red* catalytic triad, *blue* conserved Cys residues, *yellow* residues forming the walls of the substrate binding pocket, *orange* amino acids predicted to form surface loops1 and 2; **c** A model of an  $\beta$ -endoglucanase (PDB 1AQ0) docking with *P. cinnamomi* GIP



**Fig. 3** Separation of the expressed GIP protein in a polyacrylamide/SDS gel (15 %) at 8 h following induction. *M* precision plus protein marker; 1, 3 *E. coli* no induction; 2 GIP induction. (Color figure online)

#### Infection of *C. sativa* roots

Infection of the host roots was performed with mycelium of the *P. cinnamomi* isolate Pr120. The first necrotic lesions appeared after about 12 h, in those areas in direct contact with the inoculum. By 24 h, the original lesions had extended along the root. By 36 h post-inoculation, the root necrosis had spread, and was localized in the non-suberized region of the root (Fig. 4).

#### Quantification of *gip* transcripts

The expression of *gip* gene in response to glucose, cellulose and sawdust was studied in *P. cinnamomi* cells using primers specific to this gene. The expression levels of *actin2* gene were used as internal cDNA loading controls [22].

Mycelia of *P. cinnamomi* were incubated either with glucose 2 % (w/v), cellulose 0.2 % (w/v) or sawdust 0.2 % (w/v), for 2, 4, 6 and 8 days. Those three substrates were chosen considering the plant–pathogen system. The choice of actin mRNA as a stable endogenous control to normalize the amount of sample RNA was validated by evaluation of the oomycete actin mRNA levels in *in vitro* and *in planta* conditions. In RNA extracted from various *in vitro* cultures both bands (18S and 28S), had identical intensity, showing that an identical quantity of *Phytophthora* RNA was present, the intensity of actin mRNA bands was similar in all samples, showing that actin mRNA levels were also identical [23].

Total RNA was extracted from these cells, cDNA was synthesized and gene expression measured by qRT-PCR.




**Fig. 4** Necrotic effect of *P. cinnamomi* in *C. sativa* roots. *C. sativa* roots were covered by fully colonized V8 agar and incubated for 24 h in the dark at 25 °C. Necrotic tissue is indicated by arrows. **a** Control, non infected root; **b** Infected root

The results were normalized to *actin2* gene and calculated as fold change relative to control cells.

Analysis of the expression levels of the *gip* gene in mycelia of *P. cinnamomi* grown in glucose, cellulose and sawdust for 2, 4, 6 and 8 days shows that different carbon sources induce a small increase in the expression levels of *gip* gene from 2 to 4 days. A higher increase is observed after 6 and 8 days (Fig. 5). The gene expression level in sawdust tends to be higher than in glucose and in cellulose with increasing culture age.

The type of medium had a visible effect on the level of expression of the *gip* gene—it is significantly higher in sawdust than in glucose or cellulose and incremented in all media as the culture aged. This fact may be related with the adaptation of the pathogen at the culture media. Also, both glucose and cellulose are simple carbon sources compared with sawdust; thus, are easily degraded by the pathogen. In this work, sawdust is the substrate that present more similarity with the natural substrate used by the pathogen and protein expression is definitely related with this fact.

The expression levels of *gip* transcripts *in planta*, after infection of *C. sativa* roots, analyzed at different times, are represented in Fig. 6. Expression significantly increases from 12 to 24 h and then significantly decreases at 36 h.




**Fig. 5** Effect of glucose, cellulose and sawdust on the expression levels of the *gip* gene of *P. cinnamomi*. Mycelia of the wild type Pr120 isolate were exposed to glucose 2 % (w/v), cellulose 0,2 % (w/v) and sawdust 0,2 % (w/v) for 2, 4, 6 and 8 days. RNA was isolated from these cells, cDNA was synthesized and the expression levels of the *gip* gene were analyzed by qRT-PCR, using specific primers. Values were normalized to the expression levels of *actin2* gene determined in the same sample and are shown as relative mean values  $\pm$  standard deviation (SD) of 3 experiments done in triplicate

The expression values are generally lower than those found in growth media with different carbon sources (Fig. 5), which could be explained by an inhibitory factor secreted by the plant. Because GIPs are secreted by *P. cinnamomi* in the absence of the host plant, we do not observe as many variations of this protein expression as we observe in the presence of different carbon sources during infection (Fig. 5). This fact verifies the host-pathogens interactions *in planta*. This oscillation in values of expression during infection also suggests a complex mechanism of interaction involving the pathogen and a defensive response [14].

*In planta* results show that, for a pathogen to colonize a host successfully, it must develop mechanisms to evade detection or, failing that, to subvert defense responses. Models have been proposed in which pathogen-derived effector molecules, interfere with elicitor binding, signal transduction, gene activation, or the activities of the defense responses [2, 24].

The ability of plants to quickly and accurately perceive their biotic attackers is essential for mounting an effective defense. Plants must identify and respond to the chemical effectors and mechanical signals that accompany the attack. A number of structurally diverse “suppressors” have been identified that constrain active resistance in plants [25]. They are considered to be determinants of pathogenicity and are defined as factors that are produced at the infection site, participate in suppressing general



**Fig. 6** Expression levels of the *gip* gene in *P. cinnamomi* during infection of *C. sativa* roots. *C. sativa* roots were exposed to mycelia of the wild type isolate Pr120 for 12, 24 and 36 h. RNA was isolated from these cells, cDNA was synthesized and the expression levels of the *gip* gene were analyzed by qRT-PCR, using specific primers. Values were normalized to the expression levels of *actin2* gene determined in the same sample and are shown as relative mean values  $\pm$  standard deviation (SD) of 3 experiments done in triplicate


resistance, induce local susceptibility and that are not toxic to plants. Certain glycoproteins and glucans have been classified as suppressors (for a review, see [25]) and additional studies may reveal that GIPs also should be classified as suppressors. Unlike GIPs, none of the currently defined suppressor molecules has been shown to inhibit the generation of elicitors as well as the activity of a downstream defense response [14].

#### Southern blot analysis

Southern blot analysis of *P. cinnamomi* genomic DNA was performed to determine the existence of GIP homologous sequences. *ApaI* (which cuts at positions 26 and 1099), *XhoI-ClaI* (which cuts at positions 397 and 1019) and *EcoRI* (that does not cut in this sequence), restriction enzymes were used to digest the DNA.

The digestion with *ApaI* reveals a unique band corresponding to an expected 1073 bp fragment (Fig. 7, lane 1). However, the double digestion with the enzymes *XhoI-ClaI* (Fig. 7, lane 2) not only reveals the expected 622 bp band but also two more bands, longer than 1500 bp, suggesting





**Fig. 7** Southern blot hybridization using a DIG-labeled 937 bp long PCR-generated probe representing the ORF of *P. cinnamomi* *gip* gene. Genomic DNA was digested with *Apa*I (1), *Xho*I/*Cla*I (2) and *Eco*RI (3)

the possible existence of *gip* paralogs in *P. cinnamomi*. Despite the assessment of the double digestion efficiency, it is possible that the two bands longer than 1500 bp observed with the *Xho*I-*Cla*I enzymes could be partial digestions. The observation of a band larger than 1500 bp upon digestion with *Eco*RI shows that this enzyme cuts at restriction points flanking this gene (Fig. 7, lane 3).

This observation of GIP homologues in species of the *Phytophthora* genus has previously been documented. The phylogeny provides eight examples of genes represented by at least one ortholog in each of the three species and two other cases in which an ortholog appears to have been duplicated, lost, or both in one of the three species. Examples include both GIPs (*GIP1* has a full set of orthologs and paralogs while *GIP2* has orthologs but no paralogs) and apparently functional SP, such as *PiSP1*, which has a full set of orthologs [19].

In conclusion, HE-TAIL PCR is an efficient method to amplify unknown genomic DNA sequences adjacent to short known regions. This technique made possible to sequence *P. cinnamomi* *gip* gene that showed sequence homology with serine-proteases, and possessed a number of conserved sequence motifs, a similar geometric arrangement of the catalytic residues and a common reaction mechanism.

**Acknowledgments** This work was financially supported by the EC - III Framework Programme for Research and Technological Development, co-financed by the European Social Fund (ESF) and by national funding from the Portuguese Ministério da Ciência e do Ensino Superior (MCES) (PTDC/AGR-AAM/67628/2006).

## References

1. Erwin DC, Ribeiro OK (1996) *Phytophthora* diseases worldwide. American Phytopathological Society Press, St. Paul
2. Knogge W (1998) Fungal pathogenicity. *Curr Opin Plant Biol* 1(4):324–328
3. Hardham AR (2005) *Phytophthora cinnamomi*. *Mol Plant Pathol* 6(6):589–604
4. King M, Reeve W, Van der Hoek MB, Williams N, McComb J, O'Brien PA, Hardy GE (2010) Defining the phosphite-regulated transcriptome of the plant pathogen *Phytophthora cinnamomi*. *Mol Genet Genomics* 284(6):425–435
5. Birch PR, Rehmany AP, Pritchard L, Kamoun S, Beynon JL (2006) Trafficking arms: oomycete effectors enter host plant cells. *Trends Microbiol* 14(1):8–11
6. Ellis J, Catanzariti AM, Dodds P (2006) The problem of how fungal and oomycete avirulence proteins enter plant cells. *Trends Plant Sci* 11(2):61–63
7. Kamoun S (2007) Groovy times: filamentous pathogen effectors revealed. *Curr Opin Plant Biol* 10(4):358–365
8. Schornack S, Huitema E, Cano LM, Bozkurt TO, Oliva R, Van Damme M, Schwizer S, Raffaele S, Chaparro-Garcia A, Farrer R, Segretin ME, Bos J, Haas BJ, Zody MC, Nusbaum C, Win J, Thines M, Kamoun S (2009) Ten things to know about oomycete effectors. *Mol Plant Pathol* 10(6):795–803
9. Kamoun S (2003) Molecular genetics of pathogenic oomycetes. *Eukaryot Cell* 2(2):191–199
10. Attard A, Gourgues M, Galiana E, Panabieres F, Ponchet M, Keller H (2008) Strategies of attack and defense in plant-oomycete interactions, accentuated for *Phytophthora parasitica* Dastur (syn. *P. Nicotianae* Breda de Haan). *J Plant Physiol* 165(1): 83–94
11. Day B, Graham T (2007) The plant host pathogen interface: cell wall and membrane dynamics of pathogen-induced responses. *Ann N Y Acad Sci* 1113:123–134
12. Kamoun S, Huitema E, Vleeshouwers VG (1999) Resistance to oomycetes: a general role for the hypersensitive response? *Trends Plant Sci* 4(5):196–200
13. Valueva TA, Mosolov VV (2004) Role of inhibitors of proteolytic enzymes in plant defense against phytopathogenic microorganisms. *Biochemistry (Mosc)* 69(11):1305–1309
14. Rose JK, Ham KS, Darvill AG, Albersheim P (2002) Molecular cloning and characterization of glucanase inhibitor proteins: coevolution of a counterdefense mechanism by plant pathogens. *Plant Cell* 14(6):1329–1345
15. Raeder U, Broda P (1985) Rapid preparation of DNA from filamentous fungi. *Lett Appl Microbiol* 1(1):17–20
16. Michiels MT, Van den Ende W, Van Laere A (2003) Chromosomal walking of flanking regions from short known sequences in GC-rich plant genomic DNA. *Plant Mol Biol Rep* 21:295–302
17. Hanahan D (1983) Studies on transformation of *Escherichia coli* with plasmids. *J Mol Biol* 166(4):557–580
18. Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor
19. Damasceno CM, Bishop JG, Ripoll DR, Win J, Kamoun S, Rose JK (2008) Structure of the glucanase inhibitor protein (GIP) family from *Phytophthora* species suggests coevolution with

- plant endo-beta-1,3-glucanases. *Mol Plant Microbe Interact* 21(6):820–830
20. York WS, Qin Q, Rose JK (2004) Proteinaceous inhibitors of endo-beta-glucanases. *Biochim Biophys Acta* 1696(2):223–233
  21. Liu B, Xue X, Cui S, Zhang X, Han Q, Zhu L, Liang X, Wang X, Huang L, Chen X, Kang Z (2010) Cloning and characterization of a wheat beta-1,3-glucanase gene induced by the stripe rust pathogen *Puccinia striiformis* f. sp. tritici. *Mol Biol Rep* 37(2):1045–1052
  22. Murfett J, Wang XJ, Hagen G, Guilfoyle TJ (2001) Identification of *Arabidopsis histone* deacetylase HDA6 mutants that affect transgene expression. *Plant Cell* 13(5):1047–1061
  23. Horta M, Sousa N, Coelho AC, Neves D, Cravador A (2009) In vitro and in vivo quantification of elicitor expression in *Phytophthora cinnamomi*. *Physiol Mol Plant P* 73:48–57
  24. Staskawicz BJ (2001) Genetics of plant-pathogen interactions specifying plant disease resistance. *Plant Physiol* 125(1):73–76
  25. Shiraishi T, Yamada T, Ichinose Y, Kiba A, Toyoda K (1997) The role of suppressors in determining host-parasite specificities in plant cells. *Int Rev Cytol* 172:55–93