

Background

Diabetes mellitus type 2 is one of the non-communicable diseases with increasing prevalence, associated with strong physical, emotional and socio-economic repercussions. Therefore, it requires new integrated approaches to prevent and combat the disease, based on the evidence already being produced. Objective: This study aimed at assessing knowledge, empowerment and quality of life of people with type 2 diabetes, being followed in Alto Minho Region of Portugal.

Methods

This correlational descriptive study was conducted on a random sample of 537 individuals, 277 of them in primary care (clusters of health care centres, ACES). The questionnaire encompassed a sociodemographic and clinical characterization part, together with the DES-SF (empowerment), DKT (knowledge) and EQ-5D (quality of life) scales.

Results

Individuals' mean age was 64.7 ± 11.7 (ACES) and 62.3 ± 12.9 (hospital). Concerning empowerment and quality of life we found similar mean scores in both settings 3.7 ± 0.9 and 0.6 ± 0.3 in ACES, and 3.7 ± 0.7 and 0.6 ± 0.3 in hospital. On the other hand, knowledge was higher in the hospital (66.7 ± 12.4) vs. primary care (59.7 ± 16.3), especially in non-insulin treated individuals. Men followed in hospital also showed higher empowerment than women although having similar knowledge. The highest knowledge deficits were related to hypoglycaemia handling, meaning of haemoglobin HbA1C, ketoacidosis signs among the insulin treated, and the care of the feet. Younger and less educated individuals showed higher knowledge of quality of life and empowerment.

Conclusions

There is lack of knowledge in fundamental aspects of the disease's management, followed by an unrealistic feeling of empowerment, taking into account the evidence produced. Local intervention strategies should consider these findings.

Keywords

Type 2 diabetes, empowerment, knowledge, quality of life

O125**Antihypertensive therapy adherence among hypertensive patients from Bragança county, Portugal**

Isabel C. Pinto¹, Tânia Pires^{1,2}, Catarina Silva¹

¹Instituto Politécnico de Bragança, Bragança, 5300-253, Portugal; ²Centro de Investigação de Montanha, Escola Superior Agrária, 5300-253 Bragança, Portugal

Correspondence: Isabel C. Pinto (isabel.pinto@ipb.pt) – Instituto Politécnico de Bragança, Bragança, 5300-253, Portugal

BMC Health Services Research 2016, **16(Suppl 3)**:O125

Background

Hypertension increases the risk of cardiovascular diseases and is highly prevalent worldwide, reaching more than a quarter of the Portuguese population. Poor antihypertensive therapy adherence has been identified as the main cause of failure to control hypertension. Objective: To estimate the prevalence of antihypertensive therapy adherence and related factors.

Methods

This cross-sectional study was based on a questionnaire, with MAT scale (measure of adherence to therapy) validated for the Portuguese population [1], applied to 122 hypertensive patients from Bragança county, in northern Portugal. To assess therapy adherence, those whose average adherence levels were ≥ 5 were called adherent. Descriptive statistics were used, correlations were accessed using chi-square test, with a significance level of 5 %.

Results

The sample consisted mainly of females (59.1 % vs. 40.9 %), aged between 31 and 92 years old (mean 69.8). The participants show high antihypertensive therapy adherence level (82.8 %). Only marital status is related to therapy adherence, with married or widowed people being those who least adhered to antihypertensive treatment ($p = 0.04$).

Conclusions

This study shows that a high prevalence of hypertensive patients adhered to the antihypertensive therapy prescribed, the married or widowed being those who least adhered to treatment.

References

- Delgado AB, Lima ML. Contributo para validação concorrente de uma medida de adesão aos tratamentos. *Psicologia, Saúde & Doenças*. 2001; 2(2): 81-100.

Keywords

Antihypertensive therapy adherence, hypertensive patients, therapy adherence, therapy non-adherence

O126**Subjective perception of sexual achievement - An exploratory study on people with overweight**

Maria Ribeiro^{1,2}, Maria Viegas-Branco³, Filomena Pereira⁴, Ana Mª Pereira³

¹Escola Superior Agrária, Instituto Politécnico de Bragança, 5300-253 Bragança, Portugal; ²Centro de Estudos Transdisciplinares para o Desenvolvimento, Universidade de Trás-os-Montes e Alto Douro, 5000-801 Vila Real, Portugal; ³Escola Superior de Saúde, Instituto Politécnico de Bragança, 5300-121 Bragança, Portugal; ⁴Faculdade de Medicina, Universidade de Lisboa, 1649-028 Lisboa, Portugal

Correspondence: Maria Ribeiro (xilote@ipb.pt) – Escola Superior Agrária, Instituto Politécnico de Bragança, 5300-253 Bragança, Portugal

BMC Health Services Research 2016, **16(Suppl 3)**:O126

The literature reports that Body Mass Index (BMI) changes are directly related to sexual interest. Individuals with overweight and obesity self-report fewer sexual partners and are less likely to have a romantic relationship compared to their non-obese counterparts.

The objectives of this work are to know the level of perceived sexual achievement, in overweight people with the Binge Eating Scale (BES) and to study the association between age and the perception of quality of sex in people with overweight are the objectives of this work.

An exploratory and quantitative study was developed, based on a sample of 218 patients of both genders, aged between 18 and 65 years. Data was collected in various hospitals of northern and central Portugal with the Sexual Satisfaction Index (SSI) being used for this purpose.

Of all participants, 64.5 % expressed overweight. On a scale of 0 to 100, the SSI was, on average, 26.7 (SD = 14.6). On average, Individuals with severe poor scores on the BES registered the highest level of sexual dissatisfaction. There were, statistically, significant differences in the SSI, among individuals with and without overweight. Dissatisfaction is greater the more advanced the age of participants. Contrary to what is reported in the literature, there was no difference between individuals with and without overweight according to the number of sexual partners.

The results obtained with this research are consistent with the literature, except for those concerning the number of sexual partners.

Keywords

Perception, sexual achievement, overweight, Portugal

O127**Physical activity level and associated factors in hypertensive individuals registered in the family health strategy of a basic health unit from the city of Palhoça, Santa Catarina, Brazil**

Fabírcia M. Almeida, Gustavo L. Estevez, Sandra Ribeiro, Marcia R. Kretzer
Universidade do Sul de Santa Catarina, Palhoça, Santa Catarina, 88137-270 Brasil

Correspondence: Marcia R. Kretzer (marcia.kretzer1@gmail.com) – Universidade do Sul de Santa Catarina, Palhoça, Santa Catarina, 88137-270 Brasil

BMC Health Services Research 2016, **16(Suppl 3)**:O127