



Atención Primaria

www.elsevier.es/ap



PARALLEL SESSIONS: POSTERS

2nd World Congress of Health Research

Viseu – Portugal, 7-8 October 2014

1. CHILD AND ADOLESCENT HEALTH

MANDATORY REPORTING OF CHILD AND ADOLESCENT ABUSE BY GENERAL PRACTITIONERS IN A STATE OF BRAZILIAN NORTHEASTERN

Ana Carine Rolim^a, Gracyelle Moreira^b, Luiza Jane Vieira^b, Agediva Pordeus^b, Carlos Corrêa^a

^aDepartment of Public Health, Faculty of Medical Sciences, University of Campinas; São Paulo, Brazil; ^bUniversity of Fortaleza, Ceará, Brazil.

Contact details: anacarine.rolim@hotmail.com

Introduction: Violence against children and adolescents is an unequal power relationship. The Brazilian Child and Adolescent Act adopt a strategy of safety and protection of all suspected cases of abuse.

Objectives: The aim of this paper is to analyze the system of general practitioners (GPs) notification for child and adolescent mistreatment throughout Ceará State, Brazil.

Methods: A structured questionnaire was given to 227 general practitioners in 85 cities across the state. A quantitative analyze was done. The research was conducted in accordance with recognized ethical standards and national/international laws.

Results: The professional average age is 35.7 years (Standard Deviation = 12.2). The GPs have been working on primary healthcare less than 05 years (51.5%). The results illustrate that the majority of GP had no training on child/adolescent mistreatment notification (73.6%), but knew the Child and Adolescent Act (72.7%). GPs that trust in protection agencies of child and adolescent are the majority (70.8%) and do not feel under threat by the State (75.8%), for this reason they know how to forward the child/adolescent mistreatment cases (64.2%). Among 227 GPs participants, 53,1% affirmed that they had identified cases of child abuse in their careers (53.1%), nevertheless, just 52.6% of these occurrences of abuse had notified.

Conclusions: These findings show that as less training GPs are, less notification of abuse problem they do. This is under the Brazilian National Health System (SUS) expectation and an abuse notification system on the SUS must be done, looking for changes on the GPs notification.

Keywords: Child abuse. Notification. Primary healthcare.

INVESTIGATION AND ANALYSES OF THE CAUSES OF INFANTILE DEATHS IN THE CITY OF MANAUS, AMAZONAS-BRAZIL, IN 2012

Giane Santos-Melo^a, Adriana Taveira^a, Waldeyde Magalhães^a, João Sidney Azevedo^b, Ivone Candido^b, Rosane Melo^b, Lucinalva Souza^b, Ana Lucia Serrão^b, Aydelene Santos^b, Rosiandra Silva^c, Thiago Souza^c, Danielle Pinto^c, Dayan Hosokawa^c, Érica Borges^c, Hanna Santos^c, Júlio Nogueira^c, Lívia Perreira^c, Mayara Costa^c, Mariah Pimenta^c, Marlon Silva^c, Rafaela Oliveira^c, Sara Silva^c, Sonaya Andrade^c

^aTeacher of University of the State of Amazonas, Brazil; ^bPreceptors of Municipal General Office of Health of Manaus/ Amazonas, Brazil (SEMSA-Manaus); ^cAcademic of University of the State of Amazonas, Brazil.

Contact details: gzsantos3@hotmail.com

Introduction: Data of the United Nations for the Childhood Fund (UNICEF) demonstrate a reduction in the infant mortality taxes in Brazil in the last two decades. However the monitoring of this indicator is necessary, because this translates the situation of health of a população.

Objectives: In this work, academics and tutors of the nursing and medicine courses of the University of the State of Amazonas-Brazil (UEA) accompanied the professional work of the Municipal General office of Health of Manaus/Amazonas-Brazil (SEMSA-Manaus), with the objective of living actions related to the surveillance in health in the scope of the infant mortality.

define a typology of attitudes towards symptoms of disease as well as different practices in the consumption of drugs in each of the profiles.

Methods: This is a quantitative descriptive and explanatory study with a probability sample of the Portuguese population (2050 cases) based on the results of the European Social Survey (round 2).

Results: We have identified four groups of attitudes towards disease status. The comparison between the four groups showed the existence of statistically significant differences respecting the agreement with the use of drugs by healthy people (χ^2 KW(3) = 66.217; $p < 0.001$). We have also found the existence of statistically significant differences regarding the regular use of previously prescriptive drugs to herself/himself in each of the profiles of attitudes towards disease ($\chi^2(3) = 8.13$; $p = 0.043$). Respecting to self-medication practices (consumption of medication prescribed for others), we observed that it is most common among individuals who tend use informal support networks when facing disease symptoms (χ^2 KW(3) = 65.278; $p < 0.001$).

Conclusions: The results show that self-medication practices in Portugal are relatively common, although it may vary according to the profile of attitudes towards the disease symptoms. The results demonstrate the existence of an individual appropriation of expert logical therapeutic management. However, this trend assumes different proportions depending on how one interprets disease.

Keywords: Attitudes. Illness. Medication. Symptoms.

EVALUATION OF HEART FAILURE'S THERAPY IN THE PATIENTS OF OUR FAMILY HEALTH UNIT (FHU)

Carla Moreira, Martina Rocha, Jorge Santos, Rui Oliveira, Raquel Rodrigues, José Carvalho

USF Infante D. Henrique, Portugal.

Contact details: carlinhamoreira86@gmail.com

Background: The European Society of Cardiology created a guideline for the treatment of Heart Failure (HF). Due to this importance, was decided to study our patients' medication, based on this guideline.

Objectives: Patients with HF: 1-Under adequate therapy with ACEinhibitor/ARB+BB; 2-With ACEinhibitor+digoxin and AF; 3-With ACEinhibitors/ARB+BB+mineralocorticoid receptor antagonist (MRA); 4-With ACEinhibitor/ARB+BB+MRA+lvabradine.

Methods: Study observational, descriptive, cross-sectional in patients ≥ 18 years old from our FHU, with ≥ 1 consultation since 2012, HF in class II-IV and ejection fraction (EF) $\leq 40\%$. Exclusion criteria: patients without ECG, ionogram or renal function values.

Results: From 12224 adult patients, 1.12% have HF and we included 125 patients (64% ♀). Mean age of 77.59 years. 36% have Atrial Fibrillation (AF). 29.6% of the patients meet the 1st goal. The others (70.4%): 9.09% are without ACEinhibitor/ARB; 62.5% (n = 55) without BB (from these 36,36% have COPD); 20.45% are without ACEinhibitor/ARB and BB (from these 16.67% have COPD); 5.68% are medicated with ACEinhibitor, but have contraindication (CI) (20% have Creatinin > 2.5 , 80% have K^+ > 5); 2.27% are medicated with BB, despite CI (50% have asthma, 50% have FC < 60 bpm). 20% of the patients meet the 2nd goal. 5.6% meet the 3rd, but in 16% the MRA was introduced early. None of the patients meet the 4th objective, despite of 4.8% are medicated with lvabradine.

Conclusions: 29.6% have adequate therapy. In 20% of AF patients (7.2% of the sample), the therapy is also appropriate. From the patients without adequate therapy, 92.05% haven't started the

recommended therapy yet (ACEinhibitors/ARB+BB), the others started but with CI. In 20.8% of the sample the medication (MRA and lvabradine) was introduced early.

Keywords: Heart failure. Digoxin. lvabradine. Beta blocker.

INFORMAL CAREGIVING IN EXTREME OLD AGE: A UNIQUE EXPERIENCE?

Sara Alves^a, Daniela Brandão^a, Laetitia Teixeira^a, Maria João Azevedo^a, Constança Paúl^{a,b}, Oscar Ribeiro^{a,c}

^aResearch and Education Unit on Ageing (UNIFAI/ICBAS-UP);

^bInstitute of Biomedical Sciences Abel Salazar, University of Porto (ICBAS.UP), Portugal; ^cSuperior School of Health of the University of Aveiro (ESSUA.UA), Higher Institute of Social Service of Porto (ISSSP), Portugal.

Contact details: sara.alves@unifai.eu

Introduction: The increasing number of old people, particularly of centenarians, has enhanced the attention of health professionals to study their specific care needs. Caregiving can be a positive experience but also a straining one with negative impact on the caregivers' health.

Objectives: This study aims to explore differences between family carers of individuals aged 100+ yrs (group 1) and family carers of younger individuals aged < 90 yrs (group 2).

Methods: A sample of 106 informal caregivers was considered (48 caregivers of old people from the Caring at Home project and 58 caregivers of centenarians from the Oporto Centenarian Study). Information on sociodemographic aspects, on the caregiving experience (e.g. length of care, relationship with person), and the caregivers' mental health (selected items from the SF-12v2) were obtained.

Results: Informal caregivers of group 2 were younger (Mage 52,0 yrs; SD = 9.9 vs Mage 65.6 yrs; SD = 7.2), both groups were mostly constituted by women (93.1% in group 1, 91.7% in group 2), married (63.8% in group 1, 77.1% in group 2) and had the support of a secondary caregiver (91.4% in group 1, 70.2% in group 2). Significant differences between groups were only found for "vitality" ($\chi^2 = 13.267$; $p = 0.01$), with group 1 showing better results.

Conclusions: This is a preliminary comparison of informal caregivers of centenarians with informal caregivers of younger individuals. More research is needed to deepen these findings, namely the reasons that might be under the observed differences and its potential consequences for the care provided.

Keywords: Informal caregiving. Centenarians. Mental health.

SELF-MEDICATION IN A NORTHERN TOWN OF PORTUGAL

Isabel C. Pinto^a, Joana Coelho^b, Ana Teixeira^b, Carlos Bernardo^b, Cristina Vaz^b

^aNúcleo de Investigação e Intervenção no Idoso, Departamento de Tecnologias de Diagnóstico e Terapêutica, Escola Superior de Saúde do Instituto Politécnico de Bragança, Portugal;

^bDepartamento de Tecnologias de Diagnóstico e Terapêutica, Escola Superior de Saúde do Instituto Politécnico de Bragança, Portugal.

Contact details: isabel.pinto@ipb.pt

Introduction: Self-medication is defined as the consumption of a drug without guidance or prescription of competent professionals, in which the patient decides that the product will be used.

Objectives: To determine the prevalence and frequency of self-medication; characterize the therapy used, reasons, verify the occurrence of undesirable effects and knowledge about the risks; and to determine associated factors.

Methods: This cross-sectional and correlational study, had a sample of 330 individuals, 57.0% females and 43.0% males, aged between 18 and 88 years (mean 41.3). Data collection was carried out through a self-administered questionnaire. Statistical analyses were applied measures of central tendency and dispersion, and the chi-square considering a significance level of 5%.

Results: The prevalence of self-medication was 87.6%, of which 78.2% self-medicated in the last year, the majority only a few times per year (81.2%), with a minority that makes daily (0.9%). The drugs most frequently used are Paracetamol (75.8%) and Ibuprofen (61.2%), headaches (67.0%) and colds (65.2%) being the main reasons. Among individuals who self-medicate, 52.1% admit to inform physician of non-prescribed drugs used and 83.0% reported not having noticed the occurrence of undesirable effects. Of all individuals surveyed, 88.2% claim that self-medication is dangerous for health. Gender and educational level appear to be associated with self-medication ($p < 0.001$ and $p = 0.01$, respectively).

Conclusions: Self-medication is highly prevalent, but infrequent over the year. The analgesics and anti-inflammatories are the most used due to headaches and colds. Self-medication is considered risky to health and appears to be linked to gender and education level.

Keywords: Self-medication. Drugs not subject to prescription.

PERCEPTION OF USERS ABOUT THE PRESCRIBED DRUG THERAPY AFTER MEDICAL CONSULTATION AND AFTER PHARMACY DISPENSING

Isabel C. Pinto, Joana Coelho, Ana Braga, Paula Pereira, Tiffany Cardoso

Departamento de Tecnologias de Diagnóstico e Terapêutica, Escola Superior de Saúde do Instituto Politécnico de Bragança, Portugal.

Contact details: isabel.pinto@ipb.pt

Introduction: Promoting rational use of medicines is crucial to ensure therapeutic efficacy. Many users do not understand prescribed treatment, often for lack of information during the medical consultation and pharmacy dispensing, which results in difficulties for correct drug therapy.

Objectives: Determine the perception of knowledge about the drug therapy, after medical consultation and after Pharmacy dispensing, and identify related factors.

Methods: This cross-sectional and study, had a sample of 150 users of health centers and pharmacies in the north of Portugal, 64% females and 36% males, aged between 18 and 90 years (mean 57). A self-administered questionnaire was applied, including knowledge perception scale (Frohlich'10). In data analysis was used descriptive statistics and t-student test (significance level 5%).

Results: The perception of knowledge about drug therapy is insufficient either after medical consultation (70.7%) or after pharmacy dispensing (70.7%), only a minority of users had a good knowledge after medical consultation (5.3%) and after dispensing in community pharmacy (2.7%). The lowest knowledge was related with forgetting doses, drugs/food interactions and side effects. No differences were found between the perceptions of users of medical centers and pharmacy ($p = 0.191$), neither between the educational level (health center $p = 0.842$, $p = 0.307$ pharmacy).

Conclusions: The perception of knowledge about drug therapy is quite insufficient both after medical consultation and after pharmacy dispensing. There were no found differences between the perceptions of users of medical centers and pharmacy, probably due to the limited sample size. Contrary to expectation, the education level is not associated with the perception of knowledge about drugs therapy.

Keywords: Drugs therapy. Perception of knowledge. Pharmacy.

FLOWCHART FOR THE TREATMENT OF VENOUS LEG ULCERS – A BEST PRACTICE GUIDELINE

Eduardo Santos^a, Madalena Cunha^b, Fernando Pina^c

^aFundação Aurélio Amaro Diniz; ^bCenter of Studies in Education, Health and Technology, Research and Development Unit, Health School of the Polytechnic Institute of Viseu, Portugal; ^cGeneral Emergency Department, Hospital Center Tondela Viseu, Portugal.

Contact details: ejf.santos87@gmail.com

Context: Leg ulcers are a major health problem with a great impact on patients quality of life and the work of nurses, particularly at the level of primary health care and differentiated services. In the clinical practice health professionals should evaluate the etiology typology, aggravating risk factors and select the appropriate treatment that the evidence points as compression therapy. Thus, when combined with the appropriate dressings and therapeutic options, compression therapy improves the rate of healing. However, in the absence of best practice guidelines health professionals tends not to apply the best available evidence.

Objectives: To present the flowchart for the treatment of venous leg ulcers.

Methods: Based on a review of guidelines and systematic reviews performed on several scientific databases (CINAHL with Full Text, MedC Latina, MEDLINE with Full Text, Cochrane Database of Systematic Reviews, Cochrane Central Register of Controlled Trials and JBI Library), we created a best practice guideline for the Treatment of Venous Leg Ulcers. Prior to inclusion all studies were submitted to critical appraisal.

Results: Data collected from studies suggest that a correct evaluation of the patients with venous ulcers needs a focal physical exam, anamnesis and vascular assessment. Current knowledge shows that should be implemented the recommendations for treatment into clinical practice like the wound bed preparation, right dressings choose and correct compression therapy use.

Conclusions: The Flowchart for the Treatment of Venous Leg Ulcers allows a sustained practice on the best available evidence, ensuring the effectiveness of treatment, standardizing the practice of nursing teams and improving healing rates.

Keywords: Leg ulcer. Compression bandages. Wound healing.

INVESTIGATION AND ANALYSES OF THE CAUSES OF MATERNAL DEATHS IN MANAUS, AMAZONAS-BRAZIL, BETWEEN 2010 TO 2012

Giane Santos-Melo^a, Adriana Taveira^a, Waldeyde Magalhães^a, João Sidney Azevedo^b, Ivone Candido^b, Rosane Melo^b, Lucinalva Souza^b, Ana Lucia Serrão^b, Aydelene Santos^b, Rosiandra Silva^c, Thiago Souza^c, Danielle Pinto^c, Dayan Hosokawa^c, Érica Borges^c, Hanna Santos^c, Júlio Nogueira^c, Lívia Perreira^c, Mayara Costa^c, Mariah Pimenta^c, Marlon Silva^c, Rafaela Oliveira^c, Sara Silva^c, Sonaya Andrade^c

^aTeacher of University of the State of Amazonas, Brazil;

^bPreceptors of Municipal General Office of Health of Manaus/ Amazonas, Brazil, (SEMSA-Manaus); ^cAcademic of University of the State of Amazonas, Brazil.

Contact details: gzsantos3@hotmail.com

Introduction: The World Health Organization (WHO) estimates that in 2010 287.000 maternal deaths happened in the mundo¹. The maternal death is a concern all over the world for managers of health, because it translates the social inequalities of a country and the quality of services health offered in a place.