



Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

2-Bromo-4-(3,4-dimethyl-5-phenyl-1,3-oxazolidin-2-yl)-6-methoxyphenol

Maywan Hariono,^a Nurziana Ngah,^b Habibah A. Wahab^a
and Aisyah Saad Abdul Rahim^{a*}

^aSchool of Pharmaceutical Sciences, Universiti Sains Malaysia, 11800 USM Penang, Malaysia, and ^bKulliyah of Science, International Islamic University Malaysia, Bandar Indra Mahkota, 25200 Kuantan, Pahang, Malaysia

Correspondence e-mail: aisyah@usm.my


Received 24 November 2011; accepted 29 November 2011

Key indicators: single-crystal X-ray study; $T = 100\text{ K}$; mean $\sigma(\text{C}-\text{C}) = 0.003\text{ \AA}$; R factor = 0.023; wR factor = 0.056; data-to-parameter ratio = 14.3.

In the title compound, $\text{C}_{18}\text{H}_{20}\text{BrNO}_3$, the oxazolidine ring adopts an envelope conformation with the N atom at the flap position. The mean plane of oxazolidine ring makes dihedral angles of $82.96(13)$ and $70.97(12)^\circ$, respectively, with the phenyl and benzene rings. In the crystal, adjacent molecules are connected via $\text{O}-\text{H}\cdots\text{O}$ and $\text{C}-\text{H}\cdots\text{O}$ hydrogen bonds and $\text{C}-\text{H}\cdots\pi$ interactions into a zigzag chain along the b axis.

Related literature

For the synthesis and closely related structures, see: Asaruddin *et al.* (2010); Diwischek *et al.* (2003); Khruscheva *et al.* (1997); Duffy *et al.* (2004). For therapeutic properties of oxazolidine derivatives, see: Moloney *et al.* (1998); Wang *et al.* (2010); Nakano *et al.* (2010); Fülöp *et al.* (2004); Panneerselvam (2011). For standard bond lengths, see: Allen *et al.* (1987). For the low-temperature device used in the data collection, see: Cosier & Glazer (1986).



Experimental

Crystal data

$\text{C}_{18}\text{H}_{20}\text{BrNO}_3$

$M_r = 378.26$

Orthorhombic, $P2_12_12_1$

$a = 7.8056(4)\text{ \AA}$

$b = 11.9034(6)\text{ \AA}$

$c = 18.9109(9)\text{ \AA}$

$V = 1757.07(15)\text{ \AA}^3$

$Z = 4$

Mo $K\alpha$ radiation

$\mu = 2.35\text{ mm}^{-1}$

$T = 100\text{ K}$

$0.50 \times 0.36 \times 0.23\text{ mm}$

Data collection

Bruker SMART APEXII CCD

area-detector diffractometer

Absorption correction: multi-scan
(*SADABS*; Bruker, 2009)

$T_{\min} = 0.383$, $T_{\max} = 0.618$

10569 measured reflections

3074 independent reflections

2935 reflections with $I > 2\sigma(I)$

$R_{\text{int}} = 0.037$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.023$

$wR(F^2) = 0.056$

$S = 1.08$

3074 reflections

215 parameters

1 restraint

H atoms treated by a mixture of independent and constrained refinement

$\Delta\rho_{\max} = 0.32\text{ e \AA}^{-3}$

$\Delta\rho_{\min} = -0.26\text{ e \AA}^{-3}$

Absolute structure: Flack (1983),

1283 Friedel pairs

Flack parameter: 0.004 (7)

Table 1

Hydrogen-bond geometry (\AA , $^\circ$).

$Cg2$ is the centroid of the C1–C6 phenyl ring.

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O2—H2 \cdots O1 ⁱ	0.85 (1)	2.03 (1)	2.7853 (19)	148 (2)
C15—H15A \cdots O2 ⁱⁱ	0.95	2.46	3.232 (3)	138
C18—H18A \cdots Cg2 ⁱ	0.98	2.96	3.679 (3)	131

Symmetry codes: (i) $-x + 1, y + \frac{1}{2}, -z + \frac{3}{2}$; (ii) $-x + 1, y - \frac{1}{2}, -z + \frac{3}{2}$.

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXTL* (Sheldrick, 2008); program(s) used to refine structure: *SHELXTL*; molecular graphics: *SHELXTL*; software used to prepare material for publication: *SHELXTL* and *PLATON* (Spek, 2009).

MH, HAW and ASAR acknowledge the Malaysia Ministry of Science, Technology and Innovations (MOSTI) for funding the synthetic chemistry work under the R&D initiative grant Nos. 09-05-1fn-meb-004 and 304/PFARMASI/650544/I121. MH thanks Universiti Sains Malaysia for the award of a postgraduate fellowship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: IS5016).

References

- Allen, F. H., Kennard, O., Watson, D. G., Brammer, L., Orpen, A. G. & Taylor, R. (1987). *J. Chem. Soc. Perkin Trans. 2*, pp. S1–19.
- Asaruddin, M. R., Wahab, H. A., Mohamed, N., Goh, J. H. & Fun, H.-K. (2010). *Acta Cryst. E66*, o1452–o1453.
- Bruker (2009). *APEX2*, *SAINT* and *SADABS*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Cosier, J. & Glazer, A. M. (1986). *J. Appl. Cryst. 19*, 105–107.
- Diwischek, F., Heller, E. & Holzgrabe, U. (2003). *Monatsh. Chem. 134*, 1105–1111.
- Duffy, M., Gallagher, J. F. & Lough, A. J. (2004). *Acta Cryst. E60*, o234–o236.
- Flack, H. D. (1983). *Acta Cryst. A39*, 876–881.
- Fülöp, F., Lázár, L., Szakonyi, Z., Pihlavisto, M., Alaranta, S., Vainio, P. J., Juhakoski, A., Marjamäki, M. & Smith, D. J. (2004). *Pure Appl. Chem. 76*, 965–972.
- Khruscheva, N. S., Loim, N. M., Sokolov, V. I. & Makhaev, V. D. (1997). *J. Chem. Soc. Perkin Trans. 1*, pp. 2425–2427.
- Moloney, G. P., Craik, D. J., Iskander, M. N. & Nero, T. L. (1998). *J. Chem. Soc. Perkin Trans. 2*, pp. 199–206.

organic compounds

- Nakano, H., Osone, K., Takeshita, M., Kwon, E., Seki, C., Matsuyama, H., Takano, N. & Kohari, Y. (2010). *Chem. Commun.* **46**, 4827–4829.
Panneerelvam, T. (2011). *J. Adv. Pharm. Tech. Res.* **2**, 43–46.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
Spek, A. L. (2009). *Acta Cryst. D* **65**, 148–155.
Wang, Y., Chan, C., Su, Z. & Chen, C. (2010). *Biophys. Chem.* **147**, 74–80.

supporting information

Acta Cryst. (2012). E68, o35–o36 [doi:10.1107/S1600536811051269]

2-Bromo-4-(3,4-dimethyl-5-phenyl-1,3-oxazolidin-2-yl)-6-methoxyphenol

Maywan Hariono, Nurziana Ngah, Habibah A. Wahab and Aisyah Saad Abdul Rahim

S1. Comment

Oxazolidine compounds are important in understanding drug behaviour in medicinal chemistry (Duffy *et al.*, 2004). Derivatives of oxazolidine have shown inhibitory effects for several diseases or condition such as β -adrenoceptor antagonist (Moloney *et al.*, 1998), influenza antiviral (Wang *et al.*, 2010; Nakano *et al.*, 2010), antinflammatory agents (Fulop *et al.*, 2004) and antihyperglycemic (Panneerselvam, 2011). In this paper, we report the X-ray crystal structure of the title oxazolidine compound, (I).

The title compound, $C_{18}H_{20}BrNO_3$, consists of two aromatic rings which are connected through oxazolidine ring (Fig. 1). The molecule is similar with those reported by Asaruddin *et al.* (2010), in that only the present of Br atom at β position of 3-hydroxy-4-methoxyphenyl ring is different. The oxazolidine ring (O1/C7–C9/N1) adopts an envelope conformation with puckering parameters of $Q = 0.433$ (2) Å and $\varphi = 107.3$ (3) $^\circ$. The N1 atom is at the flap position and it deviates from the mean plane through the remaining four atoms by 0.651 (2) Å. The C1–C6 phenyl and C10–C15 benzene rings make dihedral angles of 82.96 (13) and 70.97 (12) $^\circ$, respectively, with the mean plane of oxazolidine ring. The bond lengths and angles are in normal ranges (Allen *et al.*, 1987) and in agreement with those reported by Asaruddin *et al.* (2010).


In the crystal structure, adjacent molecules are connected *via* intermolecular O2—H2 \cdots O1 and C15—H15A \cdots O2 hydrogen bonds and C18—H18A \cdots Cg2 interactions (Table 1) to form a chain along the [010] direction; Cg2 is the centroid of the C1–C6 ring.

S2. Experimental


Following a modified method (Asaruddin *et al.*, 2010; Diwischek *et al.*, 2003; Khruscheva *et al.*, 1997), (1*S*,2*S*)-2-methylamino-1-phenylpropan-1-ol (0.17 g, 1 mmol) was mixed with 3-bromo-4-hydroxy-5-methoxybenzaldehyde (0.23 g, 1 mmol) in a two-round neck bottom flask. The mixture was dissolved in methanol (4 ml) and molecular sieve 4 Å (0.1 g) was added to the reaction mixture then the solution was refluxed at 333 K for 6 h. The solution was filtered and the solvent was evaporated *in vacuo* to give a crude product which was then recrystallized three times from methanol to give colourless blocks with a yield 11%. These were washed with *n*-hexane and dried overnight to afford single crystals suitable for X-ray analysis.

S3. Refinement

X-ray data were collected at 100 K (Cosier & Glazer, 1986). The hydroxyl H atom was located in a difference map and refined freely [$O2—H2 = 0.8499$ (10) Å]. Other H atoms were positioned geometrically and refined using riding model with C—H = 0.95–1.00 Å and $U_{iso}(H)=1.2$ or $1.5U_{eq}(C)$. A rotating group model was applied for methyl group.

**Figure 1**

The molecular structure of the title compound, with displacement ellipsoids are drawn at the 50% probability level.

**Figure 2**

The molecular packing of the title compound viewed down the a axis.

2-Bromo-4-(3,4-dimethyl-5-phenyl-1,3-oxazolidin-2-yl)-6-methoxyphenol

Crystal data



$M_r = 378.26$

Orthorhombic, $P2_12_12_1$

Hall symbol: P 2ac 2ab

$a = 7.8056 (4)$ Å

$b = 11.9034 (6)$ Å

$c = 18.9109 (9)$ Å

$V = 1757.07 (15)$ Å 3

$Z = 4$

$F(000) = 776$

$D_x = 1.430 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 8965 reflections

$\theta = 2.0\text{--}24.9^\circ$

$\mu = 2.35 \text{ mm}^{-1}$

$T = 100$ K

Block, colourless

$0.50 \times 0.36 \times 0.23$ mm

Data collection

Bruker SMART APEXII CCD area-detector diffractometer
 Radiation source: fine-focus sealed tube
 Graphite monochromator
 Detector resolution: 83.66 pixels mm⁻¹
 φ and ω scan
 Absorption correction: multi-scan (*SADABS*; Bruker, 2009)
 $T_{\min} = 0.383$, $T_{\max} = 0.618$

10569 measured reflections
 3074 independent reflections
 2935 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.037$
 $\theta_{\max} = 24.9^\circ$, $\theta_{\min} = 2.0^\circ$
 $h = -9 \rightarrow 9$
 $k = -14 \rightarrow 14$
 $l = -22 \rightarrow 22$

Refinement

Refinement on F^2
 Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.023$
 $wR(F^2) = 0.056$
 $S = 1.08$
 3074 reflections
 215 parameters
 1 restraint
 Primary atom site location: structure-invariant direct methods
 Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites
 H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0197P)^2]$
 where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.32 \text{ e } \text{\AA}^{-3}$
 $\Delta\rho_{\min} = -0.26 \text{ e } \text{\AA}^{-3}$
 Absolute structure: Flack (1983), 1283 Friedel pairs
 Absolute structure parameter: 0.004 (7)

Special details

Experimental. The crystal was placed in the cold stream of an Oxford Cryosystems Cobra open-flow nitrogen cryostat (Cosier & Glazer, 1986) operating at 100.0 (1) K.

Geometry. All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

Refinement. Refinement of F^2 against ALL reflections. The weighted R -factor wR and goodness of fit S are based on F^2 , conventional R -factors R are based on F , with F set to zero for negative F^2 . The threshold expression of $F^2 > \sigma(F^2)$ is used only for calculating R -factors(gt) etc. and is not relevant to the choice of reflections for refinement. R -factors based on F^2 are statistically about twice as large as those based on F , and R -factors based on ALL data will be even larger.

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å²)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.58037 (3)	0.865565 (17)	0.942297 (10)	0.02402 (8)
O1	0.2217 (2)	0.40943 (12)	0.84900 (7)	0.0161 (3)
O2	0.6507 (2)	0.85865 (13)	0.78478 (7)	0.0187 (3)
O3	0.5513 (2)	0.71024 (11)	0.68917 (7)	0.0195 (3)
N1	0.0154 (3)	0.54199 (15)	0.83295 (9)	0.0179 (4)
C1	0.0051 (3)	0.1430 (2)	0.85221 (12)	0.0259 (5)
H1A	-0.0140	0.1381	0.8027	0.031*
C2	-0.0108 (4)	0.0473 (2)	0.89404 (16)	0.0362 (7)
H2A	-0.0416	-0.0224	0.8732	0.043*
C3	0.0182 (4)	0.0542 (2)	0.96587 (14)	0.0365 (7)
H3A	0.0072	-0.0110	0.9944	0.044*
C4	0.0633 (4)	0.1557 (2)	0.99672 (12)	0.0341 (6)

H4A	0.0831	0.1600	1.0462	0.041*
C5	0.0792 (4)	0.25110 (19)	0.95494 (11)	0.0257 (5)
H5A	0.1110	0.3205	0.9760	0.031*
C6	0.0490 (3)	0.24580 (18)	0.88251 (10)	0.0191 (5)
C7	0.0610 (3)	0.34994 (17)	0.83700 (10)	0.0170 (5)
H7A	0.0563	0.3265	0.7862	0.020*
C8	0.1793 (3)	0.52295 (17)	0.86771 (10)	0.0165 (5)
H8A	0.1630	0.5277	0.9201	0.020*
C9	-0.0781 (3)	0.43775 (18)	0.84969 (10)	0.0191 (5)
H9A	-0.1098	0.4379	0.9009	0.023*
C10	0.3140 (3)	0.60567 (17)	0.84574 (10)	0.0152 (5)
C11	0.3775 (3)	0.68157 (17)	0.89478 (11)	0.0167 (5)
H11A	0.3434	0.6764	0.9429	0.020*
C12	0.4899 (3)	0.76443 (18)	0.87389 (10)	0.0162 (5)
C13	0.5436 (3)	0.77558 (17)	0.80434 (10)	0.0153 (4)
C14	0.4837 (3)	0.69524 (17)	0.75520 (10)	0.0149 (4)
C15	0.3686 (3)	0.61278 (17)	0.77486 (10)	0.0153 (4)
H15A	0.3264	0.5611	0.7407	0.018*
C16	-0.2384 (3)	0.4204 (2)	0.80546 (13)	0.0302 (6)
H16A	-0.3266	0.4741	0.8202	0.045*
H16B	-0.2808	0.3437	0.8122	0.045*
H16C	-0.2112	0.4324	0.7554	0.045*
C17	-0.0712 (3)	0.64373 (18)	0.85614 (11)	0.0258 (5)
H17A	0.0013	0.7090	0.8461	0.039*
H17B	-0.0931	0.6394	0.9071	0.039*
H17C	-0.1802	0.6513	0.8309	0.039*
C18	0.5220 (3)	0.62188 (19)	0.63849 (10)	0.0224 (5)
H18A	0.5904	0.6361	0.5960	0.034*
H18B	0.4003	0.6199	0.6259	0.034*
H18C	0.5554	0.5496	0.6591	0.034*
H2	0.659 (3)	0.856 (2)	0.74000 (15)	0.022 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Br1	0.03139 (14)	0.02299 (12)	0.01767 (10)	-0.00772 (11)	-0.00524 (10)	-0.00103 (8)
O1	0.0127 (8)	0.0156 (7)	0.0200 (7)	-0.0001 (7)	0.0011 (6)	0.0004 (5)
O2	0.0203 (8)	0.0183 (8)	0.0176 (7)	-0.0032 (7)	0.0025 (6)	0.0005 (6)
O3	0.0228 (10)	0.0188 (8)	0.0171 (6)	-0.0038 (7)	0.0049 (7)	-0.0016 (5)
N1	0.0126 (10)	0.0177 (10)	0.0234 (8)	-0.0018 (8)	-0.0014 (8)	0.0019 (7)
C1	0.0216 (13)	0.0234 (12)	0.0328 (11)	-0.0021 (11)	-0.0001 (10)	-0.0009 (10)
C2	0.0318 (16)	0.0194 (13)	0.0574 (16)	-0.0070 (12)	0.0011 (14)	0.0022 (11)
C3	0.0310 (16)	0.0269 (14)	0.0517 (15)	-0.0027 (12)	0.0041 (13)	0.0184 (11)
C4	0.0358 (16)	0.0388 (15)	0.0278 (11)	0.0018 (14)	0.0042 (12)	0.0111 (10)
C5	0.0295 (14)	0.0222 (11)	0.0252 (10)	0.0001 (11)	0.0019 (12)	0.0020 (8)
C6	0.0132 (13)	0.0196 (11)	0.0245 (10)	0.0004 (10)	0.0034 (10)	0.0023 (8)
C7	0.0159 (12)	0.0190 (11)	0.0160 (8)	-0.0046 (10)	-0.0001 (9)	-0.0016 (8)
C8	0.0167 (13)	0.0180 (11)	0.0150 (9)	0.0035 (10)	-0.0007 (9)	-0.0005 (8)

C9	0.0143 (11)	0.0219 (11)	0.0211 (9)	-0.0030 (11)	0.0023 (10)	0.0026 (8)
C10	0.0110 (11)	0.0163 (11)	0.0184 (9)	0.0043 (9)	-0.0003 (8)	0.0024 (8)
C11	0.0165 (12)	0.0177 (10)	0.0159 (9)	0.0022 (9)	-0.0006 (9)	0.0031 (8)
C12	0.0146 (12)	0.0173 (11)	0.0167 (9)	0.0016 (9)	-0.0039 (9)	-0.0036 (8)
C13	0.0105 (12)	0.0149 (10)	0.0204 (9)	0.0012 (9)	-0.0012 (9)	0.0026 (7)
C14	0.0118 (11)	0.0167 (11)	0.0163 (9)	0.0040 (9)	0.0005 (9)	0.0015 (8)
C15	0.0149 (11)	0.0136 (11)	0.0174 (9)	0.0034 (9)	-0.0026 (9)	-0.0004 (7)
C16	0.0171 (13)	0.0353 (14)	0.0383 (12)	-0.0027 (12)	-0.0068 (12)	0.0025 (10)
C17	0.0178 (12)	0.0250 (12)	0.0345 (11)	0.0039 (14)	0.0010 (11)	0.0037 (9)
C18	0.0256 (12)	0.0229 (12)	0.0186 (9)	-0.0033 (11)	0.0040 (9)	-0.0058 (9)

Geometric parameters (\AA , $^\circ$)

Br1—C12	1.903 (2)	C7—H7A	1.0000
O1—C8	1.435 (3)	C8—C10	1.499 (3)
O1—C7	1.458 (3)	C8—H8A	1.0000
O2—C13	1.347 (3)	C9—C16	1.519 (4)
O2—H2	0.8499 (10)	C9—H9A	1.0000
O3—C14	1.367 (2)	C10—C11	1.386 (3)
O3—C18	1.441 (2)	C10—C15	1.409 (3)
N1—C17	1.455 (3)	C11—C12	1.378 (3)
N1—C8	1.456 (3)	C11—H11A	0.9500
N1—C9	1.474 (3)	C12—C13	1.387 (3)
C1—C2	1.392 (3)	C13—C14	1.413 (3)
C1—C6	1.394 (3)	C14—C15	1.382 (3)
C1—H1A	0.9500	C15—H15A	0.9500
C2—C3	1.379 (4)	C16—H16A	0.9800
C2—H2A	0.9500	C16—H16B	0.9800
C3—C4	1.387 (4)	C16—H16C	0.9800
C3—H3A	0.9500	C17—H17A	0.9800
C4—C5	1.389 (3)	C17—H17B	0.9800
C4—H4A	0.9500	C17—H17C	0.9800
C5—C6	1.391 (3)	C18—H18A	0.9800
C5—H5A	0.9500	C18—H18B	0.9800
C6—C7	1.512 (3)	C18—H18C	0.9800
C7—C9	1.526 (3)		
C8—O1—C7	107.31 (16)	N1—C9—H9A	109.2
C13—O2—H2	107.2 (18)	C16—C9—H9A	109.2
C14—O3—C18	116.80 (16)	C7—C9—H9A	109.2
C17—N1—C8	113.71 (18)	C11—C10—C15	119.30 (19)
C17—N1—C9	113.94 (18)	C11—C10—C8	119.56 (18)
C8—N1—C9	101.95 (17)	C15—C10—C8	120.99 (18)
C2—C1—C6	120.4 (2)	C12—C11—C10	120.15 (19)
C2—C1—H1A	119.8	C12—C11—H11A	119.9
C6—C1—H1A	119.8	C10—C11—H11A	119.9
C3—C2—C1	119.8 (2)	C11—C12—C13	122.20 (19)
C3—C2—H2A	120.1	C11—C12—Br1	119.62 (15)

C1—C2—H2A	120.1	C13—C12—Br1	118.16 (16)
C2—C3—C4	120.5 (2)	O2—C13—C12	121.22 (18)
C2—C3—H3A	119.7	O2—C13—C14	121.44 (18)
C4—C3—H3A	119.7	C12—C13—C14	117.32 (19)
C3—C4—C5	119.7 (2)	O3—C14—C15	126.11 (18)
C3—C4—H4A	120.2	O3—C14—C13	112.63 (18)
C5—C4—H4A	120.2	C15—C14—C13	121.26 (18)
C4—C5—C6	120.5 (2)	C14—C15—C10	119.68 (19)
C4—C5—H5A	119.7	C14—C15—H15A	120.2
C6—C5—H5A	119.7	C10—C15—H15A	120.2
C5—C6—C1	119.1 (2)	C9—C16—H16A	109.5
C5—C6—C7	120.85 (18)	C9—C16—H16B	109.5
C1—C6—C7	120.07 (18)	H16A—C16—H16B	109.5
O1—C7—C6	111.27 (17)	C9—C16—H16C	109.5
O1—C7—C9	104.75 (16)	H16A—C16—H16C	109.5
C6—C7—C9	115.34 (19)	H16B—C16—H16C	109.5
O1—C7—H7A	108.4	N1—C17—H17A	109.5
C6—C7—H7A	108.4	N1—C17—H17B	109.5
C9—C7—H7A	108.4	H17A—C17—H17B	109.5
O1—C8—N1	103.74 (17)	N1—C17—H17C	109.5
O1—C8—C10	112.86 (18)	H17A—C17—H17C	109.5
N1—C8—C10	112.88 (17)	H17B—C17—H17C	109.5
O1—C8—H8A	109.1	O3—C18—H18A	109.5
N1—C8—H8A	109.1	O3—C18—H18B	109.5
C10—C8—H8A	109.1	H18A—C18—H18B	109.5
N1—C9—C16	113.81 (18)	O3—C18—H18C	109.5
N1—C9—C7	101.0 (2)	H18A—C18—H18C	109.5
C16—C9—C7	113.98 (18)	H18B—C18—H18C	109.5
C6—C1—C2—C3	-0.5 (5)	C6—C7—C9—N1	149.13 (17)
C1—C2—C3—C4	0.0 (5)	O1—C7—C9—C16	148.90 (18)
C2—C3—C4—C5	0.0 (5)	C6—C7—C9—C16	-88.4 (2)
C3—C4—C5—C6	0.5 (5)	O1—C8—C10—C11	-129.0 (2)
C4—C5—C6—C1	-1.0 (4)	N1—C8—C10—C11	113.7 (2)
C4—C5—C6—C7	178.2 (3)	O1—C8—C10—C15	55.5 (3)
C2—C1—C6—C5	1.0 (4)	N1—C8—C10—C15	-61.8 (3)
C2—C1—C6—C7	-178.2 (2)	C15—C10—C11—C12	1.4 (3)
C8—O1—C7—C6	-124.92 (17)	C8—C10—C11—C12	-174.2 (2)
C8—O1—C7—C9	0.36 (18)	C10—C11—C12—C13	0.0 (3)
C5—C6—C7—O1	50.7 (3)	C10—C11—C12—Br1	-178.39 (17)
C1—C6—C7—O1	-130.1 (2)	C11—C12—C13—O2	178.6 (2)
C5—C6—C7—C9	-68.4 (3)	Br1—C12—C13—O2	-3.0 (3)
C1—C6—C7—C9	110.7 (2)	C11—C12—C13—C14	-2.5 (3)
C7—O1—C8—N1	-27.71 (18)	Br1—C12—C13—C14	175.97 (16)
C7—O1—C8—C10	-150.23 (16)	C18—O3—C14—C15	-10.8 (3)
C17—N1—C8—O1	167.85 (16)	C18—O3—C14—C13	169.59 (19)
C9—N1—C8—O1	44.77 (18)	O2—C13—C14—O3	2.2 (3)
C17—N1—C8—C10	-69.6 (2)	C12—C13—C14—O3	-176.71 (19)

C9—N1—C8—C10	167.27 (17)	O2—C13—C14—C15	−177.4 (2)
C17—N1—C9—C16	71.3 (2)	C12—C13—C14—C15	3.6 (3)
C8—N1—C9—C16	−165.77 (19)	O3—C14—C15—C10	178.0 (2)
C17—N1—C9—C7	−166.13 (17)	C13—C14—C15—C10	−2.3 (3)
C8—N1—C9—C7	−43.20 (18)	C11—C10—C15—C14	−0.2 (3)
O1—C7—C9—N1	26.45 (18)	C8—C10—C15—C14	175.3 (2)

Hydrogen-bond geometry (Å, °)

Cg2 is the centroid of the C1—C6 phenyl ring.

D—H···A	D—H	H···A	D···A	D—H···A
O2—H2···O1 ⁱ	0.85 (1)	2.03 (1)	2.7853 (19)	148 (2)
C15—H15A···O2 ⁱⁱ	0.95	2.46	3.232 (3)	138
C18—H18A···Cg2 ⁱ	0.98	2.96	3.679 (3)	131

Symmetry codes: (i) $-x+1, y+1/2, -z+3/2$; (ii) $-x+1, y-1/2, -z+3/2$.