

Open Research Online

The Open University's repository of research publications and other research outputs

Nitrate-Dependent Iron Oxidation: A Potential Mars Metabolism

Journal Item

How to cite:

Price, Alex; Pearson, Victoria K.; Schwenzer, Susanne P.; Miot, Jennyfer and Olsson-Francis, Karen (2018). Nitrate-Dependent Iron Oxidation: A Potential Mars Metabolism. *Frontiers in Microbiology*, 9(513)

For guidance on citations see [FAQs](#).

© 2018 The Authors

Version: Version of Record

Link(s) to article on publisher's website:
<http://dx.doi.org/doi:10.3389/fmicb.2018.00513>

Copyright and Moral Rights for the articles on this site are retained by the individual authors and/or other copyright owners. For more information on Open Research Online's data [policy](#) on reuse of materials please consult the policies page.

oro.open.ac.uk



Nitrate-Dependent Iron Oxidation: A Potential Mars Metabolism

Alex Price^{1*}, Victoria K. Pearson¹, Susanne P. Schwenzer¹, Jennyfer Miot² and Karen Olsson-Francis¹

¹ Faculty of Science, Technology, Engineering and Mathematics, The Open University, Milton Keynes, United Kingdom,

² CNRS, Institut de Minéralogie, de Physique des Matériaux et de Cosmochimie, Muséum National d'Histoire Naturelle, Université Pierre et Marie Curie – Sorbonne Universités, UMR 7590, Paris, France

This work considers the hypothetical viability of microbial nitrate-dependent Fe²⁺ oxidation (NDFO) for supporting simple life in the context of the early Mars environment. This draws on knowledge built up over several decades of remote and *in situ* observation, as well as recent discoveries that have shaped current understanding of early Mars. Our current understanding is that certain early martian environments fulfill several of the key requirements for microbes with NDFO metabolism. First, abundant Fe²⁺ has been identified on Mars and provides evidence of an accessible electron donor; evidence of anoxia suggests that abiotic Fe²⁺ oxidation by molecular oxygen would not have interfered and competed with microbial iron metabolism in these environments. Second, nitrate, which can be used by some iron oxidizing microorganisms as an electron acceptor, has also been confirmed in modern aeolian and ancient sediment deposits on Mars. In addition to redox substrates, reservoirs of both organic and inorganic carbon are available for biosynthesis, and geochemical evidence suggests that lacustrine systems during the hydrologically active Noachian period (4.1–3.7 Ga) match the circumneutral pH requirements of nitrate-dependent iron-oxidizing microorganisms. As well as potentially acting as a primary producer in early martian lakes and fluvial systems, the light-independent nature of NDFO suggests that such microbes could have persisted in sub-surface aquifers long after the desiccation of the surface, provided that adequate carbon and nitrates sources were prevalent. Traces of NDFO microorganisms may be preserved in the rock record by biomineralization and cellular encrustation in zones of high Fe²⁺ concentrations. These processes could produce morphological biosignatures, preserve distinctive Fe-isotope variation patterns, and enhance preservation of biological organic compounds. Such biosignatures could be detectable by future missions to Mars with appropriate instrumentation.

Keywords: iron, nitrate, Mars, astrobiology, chemolithotrophy, NDFO, nitrate-dependent ferrous iron oxidation, anaerobic

OPEN ACCESS

Edited by:

Virginia P. Edgcomb,
Woods Hole Oceanographic
Institution, United States

Reviewed by:

Trinity L. Hamilton,
University of Minnesota Twin Cities,
United States
Stephen Brian Pointing,
Yale-NUS College, Singapore

*Correspondence:

Alex Price
alex.price@open.ac.uk

Specialty section:

This article was submitted to
Extreme Microbiology,
a section of the journal
Frontiers in Microbiology

Received: 30 August 2017

Accepted: 06 March 2018

Published: 20 March 2018

Citation:

Price A, Pearson VK, Schwenzer SP,
Miot J and Olsson-Francis K (2018)
Nitrate-Dependent Iron Oxidation:
A Potential Mars Metabolism.
Front. Microbiol. 9:513.
doi: 10.3389/fmicb.2018.00513

INTRODUCTION

Mars, the red planet, has inspired the search for extraterrestrial life since the early days of the telescope, and continues to do so with perceptions of its habitability—or even inhabitation—changing with advances in exploration capabilities and knowledge of martian environments from images and data (Filiberto and Schwenzer, 2017). The present-day surface of Mars is cold, dry,

and exposed to ionizing and UV radiation, conditions deemed detrimental to life, but evidence in the geological and geomorphological record of Mars confirms warmer, wetter, and potentially more favorable surface conditions during the Noachian period of early Mars (4.1–3.7 Ga) (Carr and Head, 2010). During this period, evidence for a denser atmosphere and less oxidizing conditions suggests that more hospitable surface environments for life may have prevailed (Carr and Head, 2010; Mangold et al., 2012), including: large-scale fluvial systems (Malin and Edgett, 2003; Irwin et al., 2005; Fassett and Head, 2008; Mangold et al., 2012; Williams et al., 2013), lacustrine environments (Grotzinger et al., 2014; Rampe et al., 2017a), and impact-generated hydrothermal systems (Schwenzer and Kring, 2009; Osinski et al., 2013). Evidence for these environments comes from lake bed sediments, such as those identified at Gale Crater, which the NASA Mars Science Laboratory rover (Curiosity) is investigating in detail (e.g., Grotzinger et al., 2015). Phyllosilicates and other hydrated minerals have also been observed from orbit (Gendrin et al., 2005; Bibring et al., 2006; Chevrier et al., 2007) and from the ground (Squyres et al., 2004; Ehlmann et al., 2011). In light of our developing understanding of Mars as a dynamic planet with a complex history, this review appraises the viability of microbial nitrate-dependent iron oxidation as a candidate metabolism with regard to past and present martian environments.

MARS – GEOLOGICAL BACKGROUND

For a better understanding of the contrast between the detrimental conditions on the surface of present-day Mars and the wetter, more clement past of martian surface environments, two specific potentially habitable environments are discussed here: (1) the ancient lake bed investigated by the Curiosity rover at Gale Crater (Grotzinger et al., 2014, 2015; Palucis et al., 2016) and (2) the impact-generated hydrothermal environment discovered in the rim of Endeavour Crater by the MER Opportunity rover (Squyres et al., 2012; Arvidson et al., 2014; Fox et al., 2016).

The ancient lake bed at Gale Crater is likely to be one of many that formed within impact craters on Mars (Cabrol and Grin, 1999). Conglomerates, cross-bedded sandstones, siltstones, and mudstones have been identified by the Curiosity rover, allowing for a detailed understanding of water flow, standing water conditions, and even temporary periods of desiccation (Vaniman et al., 2013; Williams et al., 2013; Grotzinger et al., 2014, 2015; Palucis et al., 2016; Hurowitz et al., 2017). The mineralogy and geochemistry of Gale Crater sediments suggest that the conditions in this ancient lake were temperate and pH-neutral, suitable for the maintenance of life for most of the time (Grotzinger et al., 2014, 2015), although excursions to, or local areas of, acidic conditions are evidenced by the discovery of jarosite (Rampe et al., 2017a,b). Post-depositional diagenetic and alteration processes, such as the dissolution of primary minerals, the formation of calcium-sulfate veins, cementation, desiccation, or even changes to the chemistry of the incoming sediment load due to external silicic volcanism, will have changed the

environmental conditions multiple times, leading to a complex association of environmental conditions variable in space and time (Bridges et al., 2015; Johnson et al., 2016; Schwenzer et al., 2016; Frydenvang et al., 2017; Nachon et al., 2017; Rampe et al., 2017a; Yen et al., 2017). Further, Gale Crater sediments are reported to contain bioessential elements such as hydrogen, phosphorus, oxygen, and nitrogen, variable iron and sulfur oxidation states as possible energy sources, and perhaps even complex organic molecules at concentrations that could have supported past life (Vaniman et al., 2013; Grotzinger et al., 2014; Stern et al., 2015; Morris et al., 2016; Sutter et al., 2016).

Orbital observations have shown that many craters bear evidence of impact-generated hydrothermal activity (Marzo et al., 2010; Mangold et al., 2012), and ground-based exploration by the MER rover Opportunity revealed an impact-generated hydrothermal system at Endeavour Crater (Squyres et al., 2012; Arvidson et al., 2014; Fox et al., 2016). Characteristic products of such alteration are clay minerals, with the most complete succession of minerals ascribed to impact-generated hydrothermal activity found in the nakhlite meteorites (Changela and Bridges, 2010; Bridges and Schwenzer, 2012; Hicks et al., 2014). While these meteorites have an unknown geological context, and thus the impact-generated nature of the alteration remains an informed guess, the opportunity to investigate the succession of minerals with Earth-based instrumentation adds significant detail to an understanding of the compositional, reduction–oxidation (redox), and pH evolution of such alteration processes. For example, the alteration reactions evident in the nakhlites indicate a change in the redox conditions from Fe^{2+} precipitates to Fe^{3+} precipitates in the course of the formation of the assemblage (Bridges and Schwenzer, 2012; Hicks et al., 2014). Investigating such details is, to date, beyond the capability of rovers and landers, but provides essential information for assessing the habitability of the site during and after the hydrothermal activity.

Active terrestrial hydrothermal systems observed today are linked to active tectonic processes or volcanism, which drive water circulation on present-day Earth; there is no evidence of a sufficiently large or sufficiently young crater in which an active impact-generated hydrothermal system could exist. However, evidence for past hydrothermal systems is observed in the form of hydrothermal mineral veins around many terrestrial craters, e.g., Chicxulub, Manicouagan, Sudbury, and many others (see Pirajno, 2009; Osinski et al., 2013 for reviews). The difference between impact-generated and volcanic hydrothermal systems is the addition of species from degassing magma in the latter system, mainly HCl, H_2HSO_4 , and other volatiles (Pirajno, 2009; Osinski et al., 2013), though fluids in both types of systems dissolve the wall rock and deposit secondary phases as conditions change throughout their lifetime. In both cases, the hydrothermal systems contain abundant bioessential elements (carbon, hydrogen, oxygen, nitrogen, and sulfur) that support diverse microbial communities (Arnold and Sheppard, 1981; Welhan and Craig, 1983; Charlou and Donval, 1993; Wheat et al., 1996; Konn et al., 2009). On Mars, hydrothermal systems caused by large hypervelocity impacts could provide warm water conditions even in periods of cold

climate. With estimated life-times of 150–200k years even for modest craters (100–180 km diameter) the size of Gale, and with cycles of continuous mineral dissolution and precipitation maintaining the availability of redox substrates during that time, impact-generated hydrothermal systems could have provided localized hospitable zones (Abramov and Kring, 2005; Schwenger and Kring, 2009).

These two examples of martian environments (lacustrine and impact-generated hydrothermal systems) demonstrate the diversity of potentially habitable environments (as we understand them today) on ancient Mars. In early surface environments, where the conditions were less inhospitable than the present-day, both phototrophic (solar energy-driven) and chemotrophic (chemical energy-driven) primary producers may have been viable, possibly producing enough organic carbon for the subsequent development of heterotrophy and a complex web of microbial life. As the environment evolved from “warm and wet” to “cold and dry,” life would have likely become limited to the sub-surface environment (Nixon et al., 2012), protected from the adverse surface conditions and, as such, may have become limited to light-independent chemolithotrophic (inorganic chemical energy-driven) metabolisms.

Laboratory-based Mars simulation experiments, using analog regolith or brine, and theoretical modeling have suggested that chemolithotrophic life could persist in the sub-surface martian environment across a wide range of pH, salinity, desiccation, and temperature (Parnell et al., 2004; Amils et al., 2007; Jepsen et al., 2007; Gronstal et al., 2009; Chastain and Kral, 2010; Smith, 2011; Popa et al., 2012; Hoehler and Jørgensen, 2013; Montoya et al., 2013; Summers, 2013; Bauermeister et al., 2014; Oren et al., 2014; King, 2015; Fox-Powell et al., 2016; Schuerger and Nicholson, 2016).

CHEMOLITHOTROPHY ON MARS

Chemolithotrophic microorganisms harvest energy from redox reactions using inorganic substrates that are available in the environment. This metabolic strategy involves the transfer of electrons donated by the inorganic substrate, through the electron transport chain for ATP production, to a final acceptor. Chemolithotrophy is pivotal for biogeochemical cycling on Earth, such as iron, nitrogen, and sulfur cycling, and for rock weathering (Madigan et al., 2009).

The iron-rich nature of Mars raises possibilities regarding the feasibility of iron biogeochemical cycling. Martian crustal geology is dominated by rocks of basaltic composition, which contain abundant FeO in quantities roughly twice those observed in comparable basalts on Earth (McSween et al., 2003, 2009). Though the planet's surface is widely colored by iron oxides, reduced iron, Fe²⁺, exists as little as a few centimeters beneath the surface (Vaniman et al., 2013). Indeed, Fe²⁺-bearing minerals such as olivine [(Mg, Fe²⁺)₂SiO₄] have been detected across wide areas of the martian surface (Hoefen et al., 2003) and large amounts of basaltic glass (amorphous Fe²⁺-containing materials) are contained within martian crustal rocks (Morris et al., 2006a,b; McSween et al., 2009). An active hydrological cycle, combined

with prevailing reducing conditions during the Noachian period, is likely to have facilitated large-scale transport of iron (Figure 1).



On early Earth, iron biogeochemical cycling and the occurrence of iron redox couples were crucial to the biosphere, to provide energy sources and because of the role of iron in many metalloproteins such as cytochromes, nitrogenases, and hydrogenases (Canfield et al., 2006; Hoppert, 2011; Raiswell and Canfield, 2012). Iron can act as either an electron acceptor or donor dependent on its redox state (Miot and Etique, 2016). Iron oxidizing microorganisms have been shown to utilize Fe²⁺ directly after its dissolution from minerals such as olivine (Santelli et al., 2001), and a similar process may have operated within potentially habitable environments on Mars. Conversely, microbial iron reduction commonly utilizes electrons donated from organic substrates, H₂ or S⁰, with oxidized Fe³⁺ as the final electron acceptor (Lovley and Phillips, 1988; Lovley et al., 1989).

A hypothetical ‘loop’ of biologically mediated martian iron cycling (Figure 2) was first proposed by Nealson (1997), which included both iron reduction and also phototrophic iron oxidation (Ehrenreich and Widdel, 1994); the plausibility of iron reduction has been appraised previously (Nixon et al., 2012, 2013; Nixon, 2014). However, Nealson's model has limited applications to present-day Mars because of prohibitive conditions for phototrophic life in surface environments that prevent closure of this ‘loop’ for biogeochemical iron cycling.

Although research suggests that phototrophs may be sufficiently protected inside various micro-habitats within ice, halite, Fe³⁺-rich sediments, and impact-shocked rocks to withstand modern martian UV flux and remain photosynthetically productive (Cockell and Raven, 2004), the effect of desiccation, in combination with UV irradiation, would prevent dispersal and negatively impact viability (Cockell et al., 2005). Additionally, a lack of liquid water at the surface of Mars would be detrimental to life (Martín-Torres et al., 2015). A plausible alternative to a phototrophic iron oxidizer would be a chemolithotrophic iron oxidizer, which can obtain energy from redox reactions involving inorganic substances. This would allow for a light-independent iron cycle, which could have existed at the surface or in the sub-surface of early Mars and even continue today in deep sub-surface groundwaters (Michalski et al., 2013).

BIOTIC IRON OXIDATION

Abiotic Fe²⁺ oxidation occurs as a function of oxidant concentration, pH, temperature, and Fe²⁺ concentration (Ionescu et al., 2015). On Earth, low pH (<4) prevents the abiotic oxidation of Fe²⁺ by atmospheric O₂, allowing biotic oxidation (using oxygen as the electron acceptor) to dominate (Morgan and Lahav, 2007). Evidence from evaporitic palaeoenvironments on Mars suggests historic low pH (<3.5) conditions existed in certain regions (Gendrin et al., 2005; Squyres and Knoll, 2005; Ming et al., 2006), although neutral-alkaline pH-associated clays are also observed in older terrains (Bibring et al., 2006). The transition to more arid conditions is thought to have coincided with a general shift from widespread clay formation to evaporitic sulfate precipitation at the surface (Bibring et al., 2006;



Chevrier et al., 2007), resulting in increasingly acidic brines that may promote this form of biotic iron oxidation (Tosca and McLennan, 2006, 2009). However, given that only trace quantities (1450 ppm) of oxygen exist in the modern martian atmosphere (Mahaffy et al., 2013), aerobic, acidophilic iron oxidation is unlikely at the surface today (Bauermeister et al., 2014).

An alternative to aerobic iron oxidizers is microaerophilic neutrophilic iron oxidizers (NFeOs), which are able to compete with abiotic oxidation at near neutral pH. On Earth, this form of metabolism is largely restricted to oxic-anoxic boundary zones, where chemical oxidation is much slower (Roden et al., 2004). Phylogenetic studies have identified NFeOs in a variety of terrestrial environments including arctic tundra, Icelandic streams, deep-ocean vents, iron-rich soils, and temperate ground waters (Emerson and Moyer, 2002; Edwards et al., 2003; Emerson and Weiss, 2004; Cockell et al., 2011; Hedrich et al., 2011; Emerson et al., 2015). Many NFeOs are psychrophilic (Edwards et al., 2003, 2004), which could be linked to the much lower rate of abiotic iron oxidation at low temperatures (Millero et al., 1987).

On Mars, regions of higher partial pressure of oxygen in the modern sub-surface, relative to the surface, have been proposed as tolerable for microaerophiles today (Fisk and Giovannoni, 1999). King (2015) also argued that aerobic activity could be supported by the oxygen concentrations recorded by the Curiosity rover (Mahaffy et al., 2013); however, aerobic metabolism would be restricted, since oxygen diffusion distances in sediments are often limited to a few millimeters (Revsbech et al., 1980; Reimers et al., 1986; Visscher et al., 1991). Furthermore, there is evidence to suggest that redox stratification, seen in standing water bodies on Earth (Comeau et al., 2012), also occurred in martian lakes such as Gale Crater, resulting in an anoxic bottom layer (Hurowitz et al., 2017). Even assuming an oxygen-rich early martian atmosphere such as that suggested by Tuff et al. (2013), deeper waters, sediments, and the sub-surface would have been largely anoxic. As such, whatever the martian atmospheric oxygen concentration, potential habitats

for anaerobically respiring light-independent chemolithotrophs would have been prevalent on ancient and present-day Mars.

Anaerobic chemotrophic iron oxidation is known to occur in terrestrial anoxic waters and sediments of approximately circumneutral pH (Straub et al., 1996; Benz et al., 1998; Kappler and Straub, 2005; Chakraborty and Picardal, 2013). Data from Curiosity at Gale Crater have shown that the Sheepbed mudstone formation at Yellowknife Bay contains abundant clay minerals, indicating a circumneutral pH environment during sedimentation (Vaniman et al., 2013; Grotzinger et al., 2014; Bridges et al., 2015; Schwenzer et al., 2016). The conditions associated with Gale Crater are not unique and can be inferred for other sites on Mars. For example, circumneutral aqueous alteration during both the Noachian and across the Noachian–Hesperian boundary has been proposed based on orbital data of Jezero crater (Ehlmann et al., 2008, 2009), indicating further environments in which anaerobic iron oxidation may have occurred.

AVAILABILITY OF ELECTRON ACCEPTORS

In the absence of molecular oxygen, chemolithotrophic iron oxidizers would be limited by the availability of alternative electron acceptors, such as perchlorates and nitrates, for metabolic redox reactions (Straub et al., 1996; Benz et al., 1998; Kappler and Straub, 2005; Chakraborty and Picardal, 2013).

Studies at multiple locations on Mars have confirmed the presence of perchlorate (Hecht et al., 2009; Navarro-González et al., 2010; Glavin et al., 2013; Kounaves et al., 2014). Perchlorate-reducing bacteria, some able to grow at 0.4 M ClO_4^- (Oren et al., 2014)—concentrations exceeding those found on Mars (Stern et al., 2017)—have been isolated from terrestrial environments. Many are able to promote Fe^{2+} oxidation when perchlorate or nitrate is provided as an electron acceptor (Bruce et al., 1999; Chaudhuri et al., 2001; Lack et al., 2002), though energy conservation leading to growth is yet to be described in the case of perchlorate reduction coupled to Fe^{2+} oxidation.

Nitrate is thus a more feasible electron acceptor for martian iron oxidation, having been observed as the oxidant in iron-oxidizing metabolisms of growth-phase cultures (Hafenbradl et al., 1996; Straub et al., 1996; Benz et al., 1998; Straub and Buchholz-Cleven, 1998). However, until the recent discovery of nitrates on the surface of Mars (Stern et al., 2015), nitrate reducers have been largely overlooked with regard to Mars astrobiology. The following sections discuss the discovery of nitrates on Mars and the feasibility of nitrate-dependent iron oxidation as a plausible metabolism for now closing the biological iron ‘loop’ on Mars (Figure 2).

NITRATES AND NITROGEN CYCLING ON MARS

The geochemical evidence of nitrates on the surface of Mars comes from *in situ* analysis of mudstone at Gale Crater by


Curiosity (Stern et al., 2015) and from analysis of the EETA79001 and Nakhla martian meteorites (Grady et al., 1995; Kounaves et al., 2014). It has been proposed that these nitrates may have formed through photochemical processing (Smith et al., 2014) of the low abundance molecular nitrogen (1.9%) in the martian atmosphere (Mahaffy et al., 2013), volcanic-induced lightning, or thermal shock from impacts (Stern et al., 2015), and may have resulted in large accumulated quantities of nitrates during the early history of the planet (Manning et al., 2009; Stern et al., 2017) (Figure 3).

Although it is not believed that nitrate deposition currently operates on the martian surface (Stern et al., 2015), interest in the martian nitrogen cycle has been reignited because of recent spacecraft observations of atmospheric nitrogen in the upper atmosphere (Stevens et al., 2015). On Earth, the production of molecular nitrogen is primarily facilitated by microbes through denitrification (Fowler et al., 2013). Biological denitrification on Mars could have contributed to an early nitrogen cycle during the Noachian period, although Mars’ atmosphere (including its primordial atmosphere) has long been suspected to have had a low nitrogen abundance relative to Earth (Fox, 1993). Nevertheless, the presence of nitrates as a plausible electron acceptor expands the range of microbial metabolisms that could be considered potentially viable on Mars. Of particular interest is the coupling of nitrate reduction to iron oxidation, which could exploit the vast martian reservoir of Fe^{2+} ions via nitrate-dependent Fe^{2+} oxidation (NDFO).

NITRATE-DEPENDENT Fe^{2+} OXIDATION (NDFO)

Nitrate-dependent Fe^{2+} oxidation metabolism was identified on Earth two decades ago (Straub et al., 1996), yet the detailed biochemical mechanisms involved are still unresolved (e.g., Carlson et al., 2013). Early studies reported Fe^{2+} oxidation balanced with nitrate reduction in mixed cultures and isolates from anaerobic freshwater, brackish water, and marine sediments (Hafenbradl et al., 1996; Straub et al., 1996; Benz et al., 1998). There are only a few known isolates capable of this metabolism (see Table 1), but this is likely to be an under-representation of the true diversity and prevalence of these organisms (Straub and Buchholz-Cleven, 1998); NDFO may actually be an innate capability of all nitrate reducers (Carlson et al., 2013; Etique et al., 2014). Enzymatic Fe^{2+} oxidation by NDFO has never been proven and a detailed proteomic study of the NDFO species *Acidovorax ebreus* definitively demonstrated that this strain lacks any specific Fe^{2+} oxidoreductase (Carlson et al., 2013). Alternatively, electrons may transit from Fe^{2+} to other periplasmic enzymes (e.g., enzymes from the nitrate reduction chain) and abiotic side reactions between Fe^{2+} and reactive nitrogen species (NO and NO_2^-) produced upon nitrate reduction could also account for Fe^{2+} oxidation (Carlson et al., 2013; Klueglein et al., 2014, 2015).

Nitrate-dependent Fe^{2+} oxidation microorganisms have to balance (a) a potential energy gain from coupled iron oxidation and nitrate reduction and (b) energy consumption to overcome



the toxicity of Fe^{2+} and reactive nitrogen species (Carlson et al., 2012, 2013). Although Fe^{2+} oxidation coupled to nitrate reduction to nitrite provides less energy ($-481.15 \text{ kJ mol}^{-1} \text{ NO}_3^-$) than both organotrophic denitrification ($-556 \text{ kJ mol}^{-1} \text{ NO}_3^-$) and organotrophic nitrate ammonification ($-623 \text{ kJ mol}^{-1} \text{ NO}_3^-$) (Strohm et al., 2007), this reaction is exergonic at circumneutral pH ($-481.15 \text{ kJ mol}^{-1} \text{ NO}_3^-$), and may theoretically provide enough energy to sustain growth under mixotrophic (Muehe et al., 2009; Weber et al., 2009) or autotrophic conditions (Laufer et al., 2016). At the same time, ferruginous conditions stimulate metal efflux pumping and stress response pathways (Carlson et al., 2013) and may thus impair the energetic budget of NDFO.

The terrestrial NDFO microbes currently described in the literature are phylogenetically diverse, including an archaeal species, as well as representatives of the alpha-, beta-, gamma-, and delta-proteobacteria (Hafenbradl et al., 1996; Kappler et al., 2005; Kumaraswamy et al., 2006; Weber et al., 2009; Chakraborty et al., 2011). The isolation of a member of the euryarchaeota capable of NDFO from a submarine vent system (Hafenbradl et al., 1996) is suggestive that NDFO may have been a very early microbial process on Earth, due to the implication of such environments in the earliest evolution of life (Martin et al., 2008). Ilbert and Bonnefoy (2013) postulated that the mechanisms of biological anaerobic iron oxidation have arisen independently several times on Earth in an example of convergent evolution (i.e., similar strategies are adopted by genetically distant species). This widespread phylogeny, evidence from iron palaeochemistry, physiology, and redox protein cofactors involved in these pathways, suggests that NDFO may be the most ancient iron oxidation pathway in terrestrial life (Ilbert and Bonnefoy, 2013). Indeed, NDFO microbes have been implicated, alongside anoxygenic Fe^{2+} -oxidizing phototrophy, in iron cycling and the

production of early banded iron formations prior to the full oxygenation of the atmosphere on Earth (Weber et al., 2006a; Busigny et al., 2013; Ilbert and Bonnefoy, 2013). Thus, NDFO may be relevant to any putative early biosphere on Mars, where the conditions are favorable to this metabolism.

FEASIBILITY OF NDFO ON EARLY MARS

The relevance of NDFO as a plausible metabolism for putative life on Mars had, until recently, been overlooked due to the lack of evidence of nitrogen species on Mars, although the theoretical possibility of NDFO was explored using numerical modeling with hypothetical nitrate sources (Jepsen et al., 2007). The newly found availability of nitrates helps to close the 'loop' of potential chemotrophic iron cycling on Mars (Figure 2), since it could provide a ready source of electron acceptors for NDFO organisms (Figure 4); the concentration of nitrates detected at Gale Crater (Stern et al., 2015) is consistent with predictions of a $5 \times 10^{15} \text{ mol}$ global nitrate reservoir from past impact processing (Manning et al., 2009). It should be noted that the highest nitrate concentrations (1,100 ppm) determined by Curiosity were present in the sedimentary rocks with the least evidence of subsequent alteration, suggesting a period of more active nitrate production during sediment deposition, which was then followed by leaching of some sediments (Stern et al., 2015).

The modern martian atmosphere is 95.9% CO_2 (Mahaffy et al., 2013), and CO_2 is likely to have also formed a major proportion of the denser early Mars atmosphere (Ramirez et al., 2014; Jakosky et al., 2017) (Figure 5). Microbes that can utilize inorganic atmospheric carbon would therefore hold an advantage in the Mars environment. Although a low energy-yielding metabolism, a some species (*Pseudogulbenkiania* sp. strain 2002

TABLE 1 | Examples of microbial species capable of nitrate-dependent iron oxidation.

Isolate	Respiration	e ⁻ Donor	e ⁻ Acceptor	Optimum pH	Optimum temperature (°C)	Metabolism	NDFO Growth	Reference
<i>Thiobacillus denitrificans</i>	Obligate anaerobe	S-species/Fe ²⁺	NO ₃ ⁻	6.90	30	Autotrophic	Unclear	Straub et al., 1996
<i>Pseudogulbenkiania</i> sp. strain 2002	Facultative aerobe	Fe ²⁺	NO ₃ ⁻	6.75–8.00	37	Autotrophic	Yes	Weber et al., 2006b
<i>Paracoccus</i> sp. strain KS1	Facultative aerobe	Organics/S/Fe ²⁺	NO ₃ ⁻	7.00	37	Heterotrophic	No	Kumaraswamy et al., 2006
<i>Acidovorax</i> sp. strain BofFeN1	Facultative anaerobe	Organics/Fe ²⁺	NO ₃ ⁻	6.80	30	Mixotrophic	Unclear	Kappler et al., 2005
<i>Ferroglobus placidus</i>	Obligate anaerobe	Fe ²⁺ /H ₂ /S ²⁻	NO ₃ ⁻	7.00	85	Autotrophic	Yes	Hafenbradl et al., 1996
<i>Azospira</i> sp. strain PS	Facultative anaerobe	Fe ²⁺ /humic acids	NO ₃ ⁻ /ClO ₄ ⁻	7.00	26	Mixotrophic	No	Lack et al., 2002; Byrne-Bailey and Coates, 2012
<i>Acidovorax</i> sp. strain BRG1	Facultative anaerobe	Organics/Fe ²⁺	NO ₃ ⁻	6.70	28–35	Heterotrophic	No	Straub et al., 1996, 2004
<i>Aquabacterium</i> sp. strain BRG2	Facultative anaerobe	Organics/Fe ²⁺	NO ₃ ⁻	6.40–6.70	28	Heterotrophic	Unclear	Straub et al., 1996, 2004
<i>Thermomonas</i> sp. strain BRG3	Facultative anaerobe	Organics/Fe ²⁺	NO ₃ ⁻	6.70	32–35	Heterotrophic	No	Straub et al., 1996, 2004
<i>Klebsiella mobilis</i>	Facultative aerobe	Organics	NO ₃ ⁻	7.00	30	Heterotrophic	No	Etique et al., 2014

and the hyperthermophilic archaeon *Ferroglobus placidus*) have been found to fix carbon autotrophically from CO₂ and other inorganic sources during growth by NDFO (Hafenbradl et al., 1996; Weber et al., 2006b, 2009), providing an alternative carbon assimilatory capability relevant for the early and current Mars environments. Although nitrate reduction can be coupled to anaerobic oxidation of methane (Raghoebarsing et al., 2006; Ettwig et al., 2008), the ability of NDFO strains to use C1 organic compounds as carbon sources has not been investigated. This could prove an important capability when considering the martian environment, given the as yet unexplained detections of methane in the modern atmosphere (Formisano et al., 2004; Webster et al., 2015), and should be investigated further.

Most NDFOs are heterotrophic and require an organic carbon source (Chaudhuri et al., 2001; Kappler et al., 2005; Muehe et al., 2009). Organic carbon has been reported on the martian surface and in martian meteorites (Sephton et al., 2002; Steele et al., 2012; Ming et al., 2014), which may be endogenous (Steele et al., 2012) or have been delivered into the martian crust by meteoritic input (~2.4 × 10⁵ kg/year; Yen et al., 2006) (Figure 5). Sutter et al. (2016) calculated that <1% of the total carbon detected in sedimentary rocks at Gale Crater would have been sufficient to support 10⁵ cells g⁻¹ sediment if present as biologically available organics in the earlier lacustrine environment, and hence could well have sustained heterotrophic NDFOs.

Although today's martian atmosphere is oxidizing, even modest levels of volcanism over the last 3.5 billion years are likely to have produced CO₂ at levels that contributed to periodically reducing conditions (Sholes et al., 2017), favoring NDFO by limiting abiotic iron oxidation. However, there has also been a suggestion that certain locations of the ancient surface environment (>3.5 billion years ago) were, at one point, oxidizing (Lanza et al., 2016). In practical terms, oxidizing atmospheric conditions and potential redox stratified water bodies would not preclude the viability of NDFO, but merely restrict it to anoxic sediment and water regions, as is the case on Earth.

Aside from metabolic requirements, life also needs an environment which falls within other sets of physical parameters that are conducive to life. In contrast to phototrophic iron oxidizers, NDFO could have occurred in near-surface ground waters (Straub et al., 1996), which would have protected the microorganisms even if the surface radiation environment of early Mars was as intense as it is today (Dartnell et al., 2007). In addition, cell encrustation by Fe minerals may have protected them against UV irradiation (Gauger et al., 2016). In the deep sub-surface, neutral-alkaline, Fe²⁺-rich ground waters could have persisted long after the evaporation of most surface bodies (Michalski et al., 2013), greatly extending the period across which NDFO could have been viable, possibly to the present-day.

BIOMINERALIZATION AND PRESERVATION IN THE ROCK RECORD

Under Fe²⁺-rich (>5 mM) conditions, a major limiting factor for the growth of NDFO populations is the progressive




FIGURE 4 | Overview of potential redox substrate sources for nitrate-dependent iron oxidizing microorganisms in the early Mars environment. Nitrates are produced from an early atmospheric nitrogen reservoir by fixation from volcanic lightning (Stern et al., 2015), thermal shock during impacts (Summers and Khare, 2007), and irradiation from solar and cosmic sources (Smith et al., 2014). Reduced iron is released into aqueous environments by mineral dissolution, a process accentuated by hydrothermal activity (Emerson and Moyer, 2002; McSween et al., 2009). A fuller description of abiotic nitrogen fixation pathways is available in Summers et al. (2012).




FIGURE 5 | Summary of the proposed processes in carbon cycling on early Mars. Atmospheric carbon dioxide is sequestered by basalts to form carbonate minerals (Edwards and Ehlmann, 2015). The carbon is then remobilized by hydrothermal fluids and incorporated into simple organic compounds, such as methane, by serpentinization reactions (Chassefière and Leblanc, 2011). Carbon dioxide is gradually lost to space due to erosion of the atmosphere by solar winds. Meteorites are also likely to have delivered an inventory of organic carbon to the surface and sub-surface of Mars (Yen et al., 2006).

encrustation of the periplasm and outer membrane by insoluble Fe^{3+} compounds (Figure 6), resulting in a decline in individual metabolic activity and cell death (Miot et al., 2015). Even the lithoautotrophic *Pseudogulbenkiania* sp. strain 2002

shows evidence of encrustation after batch culture (Klueglein et al., 2014) (Figure 6C). Although the mechanisms remain unexplained, various extracellular Fe^{3+} mineral precipitates also form as by-products of NDFO metabolism, either due to




FIGURE 6 | (A) Transmission electron microscopy (TEM) image of an iron-encrusted cell from an NDFO enrichment culture from the anoxic layer of the ferruginous Lake Pavin, France (Jennyfer Miot, personal communication). **(B)** TEM of BoFeN1 cells fully, partially, and non-encrusted with goethite (from Miot et al., 2015). **(C)** SEM of encrusted *Pseudogulbenkiania* sp. 2002 cells (from Klueglein et al., 2014) (scale bar = 500 nm).

the interaction of released Fe^{3+} ions with dissolved phosphate, sulfate, and carbonate ions, or by oxidation of extracellular Fe^{2+} -bearing minerals (Miot et al., 2009). Persistence of a low proportion of cells that escape encrustation ensures the viability of NDFO microorganisms at the population scale, thus accounting for their occurrence in ferruginous habitats on modern Earth (Miot et al., 2016).

The membrane-associated and extracellular mineral precipitates associated with NDFO metabolism may also present plausible biosignatures that may be detectable by future life detection missions, provided that they would persist over geological time. In particular, periplasmic encrustation leads to mineral shells that entrap protein globules and which display a constant thickness (around 40 nm) (Miot et al., 2011). The nature of the minerals has been shown to be dependent on both the local chemical composition and the pH environment. *Acidovorax* sp. strain BoFeN1, one of the best studied NDFO species, has been found to produce either lepidocrocite [$\gamma\text{-FeO(OH)}$] at pH 7 (Miot et al., 2014b) or a mixture of lepidocrocite and magnetite (Fe_3O_4) at pH 7.6 (Miot et al., 2014a). Likewise, changing the chemical composition of the culture medium at pH 7 results in

the precipitation of either Fe^{3+} phosphates (Miot et al., 2009), goethite [$\alpha\text{-FeO(OH)}$] (Kappler et al., 2005; Schädler et al., 2009), or green rust (mixed $\text{Fe}^{2+}/\text{Fe}^{3+}$ hydroxides) (Pantke et al., 2012).

It is also becoming apparent that encrustation is less likely in environments with low Fe^{2+} concentrations (50–250 μM), i.e., conditions more representative of many terrestrial NDFO sample sites (Chakraborty et al., 2011). Encrustation may occur only when solutions become highly concentrated (millimolar) with Fe^{2+} ions, as may have occurred in hydrothermal and stratified lake settings on early Mars (Hurowitz et al., 2017) or in evaporitic environments during the desiccation of the martian surface (Tosca and McLennan, 2009). Oxide-encrusted cells in both of these contexts could have been deposited and preserved during sedimentation (Figure 4). If deposited and lithified as macroscopic flocs or bands within an otherwise generally reducing sedimentary geological context, these oxidized mineral features may be visible in exposed strata and would serve as prime initial targets for further astrobiological investigation. Alternative mineralization processes such as pyritization (saturation and replacement of biological structures

with iron sulfide) or silicification (saturation and replacement of biological structures with silica) could also contribute to non-specific morphological preservation of microbes in iron and sulfur-rich, predominantly basaltic, early martian environments. Microbial silicification has been observed on Earth *in situ* and *in vivo* around hot springs and under simulated conditions as well as in the fossil record (Toporski et al., 2002; Konhauser et al., 2004) whereas microbial pyritization is recognized only in the context of microfossils (Schieber, 2002; Wacey et al., 2013). Given the ability of microbial communities to thrive in conditions which encourage geologically rapid mineralization of biological material, these processes should not be viewed as prohibitive to microbial life on Noachian Mars, and are beneficial to the search for any traces of early life.

Formation of organo-ferric complexes has also been demonstrated to facilitate the preservation of organic molecules in soils and sediment over geological timescales on Earth (Lalonde et al., 2012), raising the possibility that encrustation of NDFO cells by Fe³⁺-bearing minerals and subsequent complexation may be beneficial to the preservation of organic biosignatures. At the same time, depending on the nature of encrusting minerals and diagenetic (T, P) conditions, Fe minerals may promote the thermal maturation of organic matter and partly erase organic biosignatures (Miot et al., 2017). It may be possible for the Mars Organics Molecule Analyzer (MOMA) mass spectrometer and Raman laser spectrometer (RLS), aboard the ESA ExoMars 2020 rover, to detect biogenic organic molecules in association with Fe³⁺ in iron-rich drill samples and laser targets, respectively (Lopez-Reyes et al., 2013; Arevalo et al., 2015). However, these instruments are not specific enough to distinguish evidence of NDFO microbes from any other potentially biological material encrusted in Fe minerals (e.g., Kish et al., 2016; Mirvaux et al., 2016).

Specific evidence of NDFO metabolism in the geological record on Earth or Mars may, however, come from isotopes. NDFOs have been shown to produce distinctive ⁵⁶Fe/⁵⁴Fe isotope fractionation patterns, discernible from other processes (Kappler et al., 2010). These variations may be detectable in the rock record, for example, in returned samples, using isotope ratio mass spectrometry (Anand et al., 2006; Czaja et al., 2013). The preservation of isotopic anomalies in martian sediments could provide detectable supporting evidence of NDFO on early Mars.

REFERENCES

- Abramov, O., and Kring, D. A. (2005). Impact-induced hydrothermal activity on early Mars. *J. Geophys. Res. Planets* 110:E12S09. doi: 10.1029/2005JE002453
- Amils, R., González-Toril, E., Fernández-Remolar, D., Gómez, F., Aguilera, A., Rodríguez, N., et al. (2007). Extreme environments as Mars terrestrial analogs: the Rio Tinto case. *Planet. Space Sci.* 55, 370–381. doi: 10.1016/j.pss.2006.02.006
- Anand, M., Russell, S., Blackhurst, R., and Grady, M. (2006). Searching for signatures of life on Mars: an Fe-isotope perspective. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 361, 1715–1720. doi: 10.1098/rstb.2006.1899
- Arevalo, R., Brinckerhoff, W., van Amerom, F., Danell, R., Pinnick, V., Xiang, L., et al. (2015). "Design and demonstration of the Mars organic molecule

CONCLUSION

Nitrate-dependent Fe²⁺ oxidation (NDFO) microorganisms oxidase Fe²⁺ compounds while also reducing nitrates under anaerobic, circumneutral conditions. These environments are proposed to have existed on Mars, providing the electron donors and acceptors required for NDFO metabolism. This implies that NDFO is a feasible and logical avenue for investigating hypothetical early martian life.

The discovery of nitrates establishes NDFO as a viable mechanism for hypothetical, biological iron oxidation on present-day Mars. NDFO could help to close a chemotrophic 'loop' of biogeochemical iron cycling on Mars, by providing a potential mechanism for iron oxidation, and allowing chemotrophic iron cycling to occur in both circumneutral ancient surface waters and deep sub-surface waters throughout martian history.

To test the validity of this hypothesis, further research should seek to determine the feasibility of NDFO metabolism under Mars simulation conditions and characterize any associated biomineralization processes. Should the suitability of NDFO to martian environments be supported by the outcomes of these experiments, future life detection missions could be optimized to seek the distinctive mineralized biosignatures of NDFO in the martian rock record.

AUTHOR CONTRIBUTIONS

AP was responsible for writing the manuscript with a large amount of input and revision from KO-F, VP, and SS. JM contributed to the revision, providing experience and expertise in nitrate-dependent iron oxidation and biomineralization processes. The concept for the paper was developed in discussions between AP and KO-F with VP and SS involved from the beginning.

FUNDING

This study was supported by Science and Technology Facilities Council and The Open University. The collaboration was made possible thanks to a Short Term Science Mission (STSM) grant from the EU COST Action ORIGINS TD1308.

analyzer (MOMA) on the ExoMars 2018 rover," in *Proceedings of the 2015 IEEE Aerospace Conference*, Big Sky, MT, 1–11. doi: 10.1109/AERO.2015.7119073

- Arnold, M., and Sheppard, S. M. (1981). East Pacific Rise at latitude 21 N: isotopic composition and origin of the hydrothermal sulphur. *Earth Planet. Sci. Lett.* 56, 148–156. doi: 10.1016/0012-821X(81)90122-9
- Arvidson, R. E., Squyres, S. W., Bell, J. F., Catalano, J. G., Clark, B. C., Crumpler, L. S., et al. (2014). Ancient aqueous environments at Endeavour crater, Mars. *Science* 343:1248097. doi: 10.1126/science.1248097
- Bauermeister, A., Rettberg, P., and Flemming, H.-C. (2014). Growth of the acidophilic iron-sulfur bacterium *Acidithiobacillus ferrooxidans* under Mars-like geochemical conditions. *Planet. Space Sci.* 98, 205–215. doi: 10.1016/j.pss.2013.09.009

- Benz, M., Brune, A., and Schink, B. (1998). Anaerobic and aerobic oxidation of ferrous iron at neutral pH by chemoheterotrophic nitrate-reducing bacteria. *Arch. Microbiol.* 169, 159–165. doi: 10.1007/s002030050555
- Bibring, J.-P., Langevin, Y., Mustard, J. F., Poulet, F., Arvidson, R., Gendrin, A., et al. (2006). Global mineralogical and aqueous Mars history derived from OMEGA/Mars express data. *Science* 312, 400–404. doi: 10.1126/science.1122659
- Bridges, J. C., and Schwenzer, S. P. (2012). The nakhlite hydrothermal brine on Mars. *Earth Planet. Sci. Lett.* 35, 117–123. doi: 10.1016/j.epsl.2012.09.044
- Bridges, J. C., Schwenzer, S. P., Leveille, R., Westall, F., Wiens, R. C., Mangold, N., et al. (2015). Diagenesis and clay mineral formation at Gale Crater, Mars. *J. Geophys. Res. Planets* 120, 1–19. doi: 10.1002/2014JE004757
- Bruce, R. A., Achenbach, L. A., and Coates, J. D. (1999). Reduction of (per) chlorate by a novel organism isolated from paper mill waste. *Environ. Microbiol.* 1, 319–329. doi: 10.1046/j.1462-2920.1999.00042.x
- Busigny, V., Lebeau, O., Ader, M., Krapež, B., and Bekker, A. (2013). Nitrogen cycle in the Late Archean ferruginous ocean. *Chem. Geol.* 362, 115–130. doi: 10.1016/j.chemgeo.2013.06.023
- Byrne-Bailey, K. G., and Coates, J. D. (2012). Complete genome sequence of the anaerobic perchlorate-reducing bacterium *Azospira suillum* strain PS. *J. Bacteriol.* 194, 2767–2768. doi: 10.1128/JB.00124-12
- Cabrol, N. A., and Grin, E. A. (1999). Distribution, classification, and ages of Martian impact crater lakes. *Icarus* 142, 160–172. doi: 10.1006/icar.1999.6191
- Canfield, D. E., Rosing, M. T., and Bjerrum, C. (2006). Early anaerobic metabolisms. *Philos. Trans. R. Soc. Lond. B Biol. Sci.* 361, 1819–1836. doi: 10.1098/rstb.2006.1906
- Carlson, H. K., Clark, I. C., Blazewicz, S. J., Iavarone, A. T., and Coates, J. D. (2013). Fe (II) oxidation is an innate capability of nitrate-reducing bacteria that involves abiotic and biotic reactions. *J. Bacteriol.* 195, 3260–3268. doi: 10.1128/JB.00058-13
- Carlson, H. K., Clark, I. C., Melnyk, R. A., and Coates, J. D. (2012). Toward a mechanistic understanding of anaerobic nitrate-dependent iron oxidation: balancing electron uptake and detoxification. *Front. Microbiol.* 3:57. doi: 10.3389/fmicb.2012.00057
- Carr, M. H., and Head, J. W. (2010). Geologic history of Mars. *Earth Planet. Sci. Lett.* 294, 185–203. doi: 10.1016/j.epsl.2009.06.042
- Chakraborty, A., and Picardal, F. (2013). Neutrophilic, nitrate-dependent, Fe(II) oxidation by a *Dechloromonas* species. *World J. Microbiol. Biotechnol.* 29, 617–623. doi: 10.1007/s11274-012-1217-9
- Chakraborty, A., Roden, E. E., Schieber, J., and Picardal, F. (2011). Enhanced growth of *Acidovorax* sp. Strain 2AN during nitrate-dependent Fe(II) oxidation in batch and continuous-flow systems. *Appl. Environ. Microbiol.* 77, 8548–8556. doi: 10.1128/aem.06214-11
- Changela, H. G., and Bridges, J. C. (2010). Alteration assemblages in the nakhlites: variation with depth on Mars. *Meteorit. Planet. Sci.* 45, 1847–1867. doi: 10.1111/j.1945-5100.2010.01123.x
- Charlou, J. L., and Donval, J. P. (1993). Hydrothermal methane venting between 12° N and 26° N along the Mid-Atlantic Ridge. *J. Geophys. Res. Solid Earth* 98, 9625–9642. doi: 10.1029/92JB02047
- Chassefière, E., and Leblanc, F. (2011). Methane release and the carbon cycle on Mars. *Planet. Space Sci.* 59, 207–217. doi: 10.1016/j.pss.2010.09.004
- Chastain, B. K., and Kral, T. A. (2010). Zero-valent iron on Mars: an alternative energy source for methanogens. *Icarus* 208, 198–201. doi: 10.1016/j.icarus.2010.02.024
- Chaudhuri, S. K., Lack, J. G., and Coates, J. D. (2001). Biogenic magnetite formation through anaerobic biooxidation of Fe (II). *Appl. Environ. Microbiol.* 67, 2844–2848. doi: 10.1128/AEM.67.6.2844-2848.2001
- Chevrier, V., Poulet, F., and Bibring, J.-P. (2007). Early geochemical environment of Mars as determined from thermodynamics of phyllosilicates. *Nature* 448, 60–63. doi: 10.1038/nature05961
- Cockell, C. S., Kelly, L. C., Summers, S., and Marteinson, V. (2011). Following the kinetics: iron-oxidizing microbial mats in cold Icelandic volcanic habitats and their rock-associated carbonaceous signature. *Astrobiology* 11, 679–694. doi: 10.1089/ast.2011.0606
- Cockell, C. S., and Raven, J. A. (2004). Zones of photosynthetic potential on Mars and the early Earth. *Icarus* 169, 300–310. doi: 10.1016/j.icarus.2003.12.024
- Cockell, C. S., Schuerger, A. C., Billi, D., Friedmann, E. I., and Panitz, C. (2005). Effects of a simulated Martian UV flux on the cyanobacterium, *Chroococcidiopsis* sp. 029. *Astrobiology* 5, 127–140. doi: 10.1089/ast.2005.5.127
- Comeau, A. M., Harding, T., Galand, P. E., Vincent, W. F., and Lovejoy, C. (2012). Vertical distribution of microbial communities in a perennially stratified Arctic lake with saline, anoxic bottom waters. *Sci. Rep.* 2:604. doi: 10.1038/srep00604
- Czaja, A. D., Johnson, C. M., Beard, B. L., Roden, E. E., Li, W., and Moorbath, S. (2013). Biological Fe oxidation controlled deposition of banded iron formation in the ca. 3770Ma Isua Supracrustal Belt (West Greenland). *Earth Planet. Sci. Lett.* 363, 192–203. doi: 10.1016/j.epsl.2012.12.025
- Dartnell, L. R., Desorgher, L., Ward, J. M., and Coates, A. J. (2007). Modelling the surface and subsurface Martian radiation environment: implications for astrobiology. *Geophys. Res. Lett.* 34:L02207. doi: 10.1029/2006GL027494
- Edwards, C. S., and Ehlmann, B. L. (2015). Carbon sequestration on Mars. *Geology* 43, 863–866. doi: 10.1130/G36983.1
- Edwards, K. J., Bach, W., McCollom, T. M., and Rogers, D. R. (2004). Neutrophilic iron-oxidizing bacteria in the ocean: their habitats, diversity, and roles in mineral deposition, rock alteration, and biomass production in the deep-sea. *Geomicrobiol. J.* 21, 393–404. doi: 10.1080/01490450490485863
- Edwards, K. J., Rogers, D. R., Wirsén, C. O., and McCollom, T. M. (2003). Isolation and characterization of novel psychrophilic, neutrophilic, Fe-oxidizing, chemolithoautotrophic α - and γ -*Proteobacteria* from the deep sea. *Appl. Environ. Microbiol.* 69, 2906–2913. doi: 10.1128/AEM.69.5.2906-2913.2003
- Ehlmann, B. L., Mustard, J. F., Fassett, C. I., Schon, S. C., Head, J. W. III, Des Marais, D. J., et al. (2008). Clay minerals in delta deposits and organic preservation potential on Mars. *Nat. Geosci.* 1, 355–358. doi: 10.1038/ngeo207
- Ehlmann, B. L., Mustard, J. F., Murchie, S. L., Bibring, J.-P., Meunier, A., Fraeman, A. A., et al. (2011). Subsurface water and clay mineral formation during the early history of Mars. *Nature* 479, 53–60. doi: 10.1038/nature10582
- Ehlmann, B. L., Mustard, J. F., Swayze, G. A., Clark, R. N., Bishop, J. L., Poulet, F., et al. (2009). Identification of hydrated silicate minerals on Mars using MRO-CRISM: geologic context near Nili Fossae and implications for aqueous alteration. *J. Geophys. Res. Planets* 114:E00D08. doi: 10.1029/2009JE003339
- Ehrenreich, A., and Widdel, F. (1994). Anaerobic oxidation of ferrous iron by purple bacteria, a new type of phototrophic metabolism. *Appl. Environ. Microbiol.* 60, 4517–4526.
- Emerson, D., and Moyer, C. L. (2002). Neutrophilic Fe-oxidizing bacteria are abundant at the Loihi Seamount hydrothermal vents and play a major role in Fe oxide deposition. *Appl. Environ. Microbiol.* 68, 3085–3093. doi: 10.1128/AEM.68.6.3085-3093.2002
- Emerson, D., Scott, J. J., Benes, J., and Bowden, W. B. (2015). Microbial iron oxidation in the arctic tundra and its implications for biogeochemical cycling. *Appl. Environ. Microbiol.* 81, 8066–8075. doi: 10.1128/AEM.02832-15
- Emerson, D., and Weiss, J. V. (2004). Bacterial iron oxidation in circumneutral freshwater habitats: findings from the field and the laboratory. *Geomicrobiol. J.* 21, 405–414. doi: 10.1080/01490450490485881
- Etique, M., Jorand, F. P. A., Zegeye, A., Greigoire, B., Despas, C., and Ruby, C. (2014). Abiotic process for Fe (II) oxidation and green rust mineralization driven by a heterotrophic nitrate reducing bacteria (*Klebsiella mobilis*). *Environ. Sci. Technol.* 48, 3742–3751. doi: 10.1021/es403358v
- Ettwig, K. F., Shima, S., Van de Pas-Schoonen, K. T., Kahnt, J., Medema, M. H., Op Den Camp, H. J. M., et al. (2008). Denitrifying bacteria anaerobically oxidize methane in the absence of *Archaea*. *Environ. Microbiol.* 10, 3164–3173. doi: 10.1111/j.1462-2920.2008.01724.x
- Fassett, C. I., and Head, J. W. (2008). The timing of Martian valley network activity: constraints from buffered crater counting. *Icarus* 195, 61–89. doi: 10.1016/j.icarus.2007.12.009
- Filiberto, J., and Schwenzer, S. P. (2017). *Volatiles in the Martian Crust*. Amsterdam: Elsevier.

- Fisk, M. R., and Giovannoni, S. J. (1999). Sources of nutrients and energy for a deep biosphere on Mars. *J. Geophys. Res. Planets* 104, 11805–11815. doi: 10.1029/1999JE900010
- Formisano, V., Atreya, S., Encrenaz, T., Ignatiev, N., and Giuranna, M. (2004). Detection of methane in the atmosphere of Mars. *Science* 306, 1758–1761. doi: 10.1126/science.1101732
- Fowler, D., Coyle, M., Skiba, U., Sutton, M. A., Cape, J. N., Reis, S., et al. (2013). The global nitrogen cycle in the twenty-first century. *Philos. Trans. R. Soc. B Biol. Sci.* 368:20130164. doi: 10.1098/rstb.2013.0164
- Fox, J. L. (1993). The production and escape of nitrogen atoms on Mars. *J. Geophys. Res. Planets* 98, 3297–3310. doi: 10.1029/92JE02289
- Fox, V. K., Arvidson, R. E., Guinness, E. A., McLennan, S. M., Catalano, J. G., Murchie, S. L., et al. (2016). Smectite deposits in Marathon Valley, Endeavour Crater, Mars, identified using CRISM hyperspectral reflectance data. *Geophys. Res. Lett.* 43, 4885–4892. doi: 10.1002/2016GL069108
- Fox-Powell, M. G., Hallsworth, J. E., Cousins, C. R., and Cockell, C. S. (2016). Ionic strength is a barrier to the habitability of Mars. *Astrobiology* 16, 427–442. doi: 10.1089/ast.2015.1432
- Frydenvang, J., Gasda, P. J., Hurowitz, J. A., Grotzinger, J. P., Wiens, R. C., Newsom, H. E., et al. (2017). Diagenetic silica enrichment and late-stage groundwater activity in Gale crater, Mars. *Geophys. Res. Lett.* 44, 4716–4724. doi: 10.1002/2017GL073323
- Gauger, T., Konhauser, K., and Kappler, A. (2016). Protection of nitrate-reducing Fe(II)-oxidizing bacteria from UV radiation by biogenic Fe(III) minerals. *Astrobiology* 16, 301–310. doi: 10.1089/ast.2015.1365
- Gendrin, A., Mangold, N., Bibring, J.-P., Langevin, Y., Gondet, B., Poulet, F., et al. (2005). Sulfates in Martian layered terrains: the OMEGA/Mars express view. *Science* 307, 1587–1591. doi: 10.1126/science.1109087
- Glavin, D. P., Freissinet, C., Miller, K. E., Eigenbrode, J. L., Brunner, A. E., Buch, A., et al. (2013). Evidence for perchlorates and the origin of chlorinated hydrocarbons detected by SAM at the Rocknest aeolian deposit in Gale Crater. *J. Geophys. Res. Planets* 118, 1955–1973. doi: 10.1002/jgre.20144
- Grady, M. M., Wright, I., and Pillinger, C. T. (1995). A search for nitrates in Martian meteorites. *J. Geophys. Res. Planets* 100, 5449–5455. doi: 10.1029/94JE02803
- Gronstal, A., Pearson, V., Kappler, A., Dooris, C., Anand, M., Poitrasson, F., et al. (2009). Laboratory experiments on the weathering of iron meteorites and carbonaceous chondrites by iron-oxidizing bacteria. *Meteorit. Planet. Sci.* 44, 233–247. doi: 10.1111/j.1945-5100.2009.tb00731.x
- Grotzinger, J. P., Gupta, S., Malin, M. C., Rubin, D. M., Schieber, J., Siebach, K., et al. (2015). Deposition, exhumation, and paleoclimate of an ancient lake deposit, Gale crater, Mars. *Science* 350:aac7575. doi: 10.1126/science.aac7575
- Grotzinger, J. P., Sumner, D. Y., Kah, L. C., Stack, K., Gupta, S., Edgar, L., et al. (2014). A habitable fluvio-lacustrine environment at Yellowknife Bay, Gale crater, Mars. *Science* 343:1242777. doi: 10.1126/science.1242777
- Hafenbradl, D., Keller, M., Dirmeier, R., Rachel, R., Roßnagel, P., Burggraf, S., et al. (1996). *Ferroglobus placidus* gen. nov., sp. nov., a novel hyperthermophilic archaeum that oxidizes Fe²⁺ at neutral pH under anoxic conditions. *Arch. Microbiol.* 166, 308–314. doi: 10.1007/s002030050388
- Hecht, M. H., Kounaves, S. P., Quinn, R. C., West, S. J., Young, S. M. M., Ming, D. W., et al. (2009). Detection of perchlorate and the soluble chemistry of Martian soil at the phoenix Lander site. *Science* 325, 64–67. doi: 10.1126/science.1172466
- Hedrich, S., Schlömann, M., and Johnson, D. B. (2011). The iron-oxidizing *Proteobacteria*. *Microbiology* 157, 1551–1564. doi: 10.1099/mic.0.045344-0
- Hicks, L. J., Bridges, J. C., and Gurman, S. (2014). Ferric saponite and serpentine in the nakhlite Martian meteorites. *Geochim. Cosmochim. Acta* 136, 194–210. doi: 10.1016/j.gca.2014.04.010
- Hoefen, T. M., Clark, R. N., Bandfield, J. L., Smith, M. D., Pearl, J. C., and Christensen, P. R. (2003). Discovery of olivine in the Nili Fossae region of Mars. *Science* 302, 627–630. doi: 10.1126/science.1089647
- Hoehler, T. M., and Jørgensen, B. B. (2013). Microbial life under extreme energy limitation. *Nat. Rev. Microbiol.* 11, 83–94. doi: 10.1038/nrmicro2939
- Hoppert, M. (2011). “Metalloenzymes,” in *Encyclopedia of Geobiology*, eds R. Reitner and V. Thiel (Dordrecht: Springer), 558–563. doi: 10.1007/978-1-4020-9212-1_134
- Hurowitz, J. A., Grotzinger, J. P., Fischer, W. W., McLennan, S. M., Milliken, R. E., Stein, N., et al. (2017). Redox stratification of an ancient lake in Gale crater, Mars. *Science* 356:eah6849. doi: 10.1126/science.aah6849
- Ilbert, M., and Bonnefoy, V. (2013). Insight into the evolution of the iron oxidation pathways. *Biochim. Biophys. Acta* 1827, 161–175. doi: 10.1016/j.bbabi.2012.10.001
- Ionescu, D., Heim, C., Polerecky, L., Thiel, V., and De Beer, D. (2015). Biotic and abiotic oxidation and reduction of iron at circumneutral pH are inseparable processes under natural conditions. *Geomicrobiol. J.* 32, 221–230. doi: 10.1080/01490451.2014.887393
- Irwin, R. P., Howard, A. D., Craddock, R. A., and Moore, J. M. (2005). An intense terminal epoch of widespread fluvial activity on early Mars: 2. Increased runoff and paleolake development. *J. Geophys. Res. Planets* 110:E12S15. doi: 10.1029/2005JE002460
- Jakosky, B. M., Slipski, M., Benna, M., Mahaffy, P., Elrod, M., Yelle, R., et al. (2017). Mars’ atmospheric history derived from upper-atmosphere measurements of 38Ar/36Ar. *Science* 355, 1408–1410. doi: 10.1126/science.aai7721
- Jepsen, S. M., Priscu, J. C., Grimm, R. E., and Bullock, M. A. (2007). The potential for lithoautotrophic life on Mars: application to shallow interfacial water environments. *Astrobiology* 7, 342–354. doi: 10.1089/ast.2007.0124
- Johnson, J. R., Bell, J. F., Bender, S., Blaney, D., Cloutis, E., Ehlmann, B., et al. (2016). Constraints on iron sulfate and iron oxide mineralogy from ChemCam visible/near-infrared reflectance spectroscopy of Mt. Sharp basal units, Gale Crater, Mars. *Am. Mineral.* 101, 1501–1514. doi: 10.2138/am-2016-5553
- Kappler, A., Johnson, C., Crosby, H., Beard, B., and Newman, D. (2010). Evidence for equilibrium iron isotope fractionation by nitrate-reducing iron (II)-oxidizing bacteria. *Geochim. Cosmochim. Acta* 74, 2826–2842. doi: 10.1016/j.gca.2010.02.017
- Kappler, A., Schink, B., and Newman, D. K. (2005). Fe (III) mineral formation and cell encrustation by the nitrate-dependent Fe (II)-oxidizer strain BoFeN1. *Geobiology* 3, 235–245. doi: 10.1128/AEM.03277-13
- Kappler, A., and Straub, K. L. (2005). Geomicrobiological cycling of iron. *Rev. Mineral. Geochem.* 59, 85–108. doi: 10.1128/AEM.01151-16
- King, G. M. (2015). Carbon monoxide as a metabolic energy source for extremely halophilic microbes: implications for microbial activity in Mars regolith. *Proc. Natl. Acad. Sci. U.S.A.* 112, 4465–4470. doi: 10.1073/pnas.1424989112
- Kish, A., Miot, J., Lombard, C., Guigner, J.-M., Bernard, S., Zirah, S., et al. (2016). Preservation of archaeal surface layer structure during mineralization. *Sci. Rep.* 6:26152. doi: 10.1038/srep26152
- Klueglein, N., Picardal, F., Zedda, M., Zwiener, C., and Kappler, A. (2015). Oxidation of Fe (II)-EDTA by nitrite and by two nitrate-reducing Fe (II)-oxidizing *Acidovorax* strains. *Geobiology* 13, 198–207. doi: 10.1111/gbi.12125
- Klueglein, N., Zeitvogel, F., Stierhof, Y.-D., Floetenmeyer, M., Konhauser, K. O., Kappler, A., et al. (2014). Potential role of nitrite for abiotic Fe (II) oxidation and cell encrustation during nitrate reduction by denitrifying bacteria. *Appl. Environ. Microbiol.* 80, 1051–1061. doi: 10.1128/AEM.03277-13
- Konhauser, K. O., Jones, B., Phoenix, V. R., Ferris, G., and Renaut, R. W. (2004). The microbial role in hot spring silicification. *Ambio* 33, 552–558. doi: 10.1579/0044-7447-33.8.552
- Konn, C., Charlou, J.-L., Donval, J.-P., Holm, N., Dehairs, F., and Bouillon, S. (2009). Hydrocarbons and oxidized organic compounds in hydrothermal fluids from Rainbow and Lost City ultramafic-hosted vents. *Chem. Geol.* 258, 299–314. doi: 10.1016/j.chemgeo.2008.10.034
- Kounaves, S. P., Carrier, B. L., O’Neil, G. D., Stroble, S. T., and Claire, M. W. (2014). Evidence of Martian perchlorate, chlorate, and nitrate in Mars meteorite EETA79001: implications for oxidants and organics. *Icarus* 229, 206–213. doi: 10.1016/j.icarus.2013.11.012
- Kumaraswamy, R., Sjölema, K., Kuenen, G., van Loosdrecht, M., and Muyzer, G. (2006). Nitrate-dependent [Fe(II)EDTA]²⁻ oxidation by *Paracoccus ferrooxidans* sp. nov., isolated from a denitrifying bioreactor. *Syst. Appl. Microbiol.* 29, 276–286. doi: 10.1016/j.syapm.2005.08.001
- Lack, J. G., Chaudhuri, S. K., Chakraborty, R., Achenbach, L. A., and Coates, J. D. (2002). Anaerobic biooxidation of Fe(II) by *Dechlorosoma suillum*. *Microb. Ecol.* 43, 424–431. doi: 10.1007/s00248-001-1061-1
- Lalonde, K., Mucci, A., Ouellet, A., and Gélinas, Y. (2012). Preservation of organic matter in sediments promoted by iron. *Nature* 483, 198–200. doi: 10.1038/nature10855
- Lanza, N. L., Wiens, R. C., Arvidson, R. E., Clark, B. C., Fischer, W. W., Gellert, R., et al. (2016). Oxidation of manganese in an ancient aquifer,

- Kimberley formation, Gale crater, Mars. *Geophys. Res. Lett.* 43, 7398–7407. doi: 10.1002/2016GL069109
- Laufer, K., Røy, H., Jørgensen, B. B., and Kappler, A. (2016). Evidence for the existence of autotrophic nitrate-reducing Fe (II)-oxidizing bacteria in marine coastal sediment. *Appl. Environ. Microbiol.* 82, 6120–6131. doi: 10.1128/AEM.01570-16
- Lopez-Reyes, G., Rull, F., Venegas, G., Westall, F., Foucher, F., Bost, N., et al. (2013). Analysis of the scientific capabilities of the ExoMars Raman Laser Spectrometer instrument. *Eur. J. Mineral.* 25, 721–733. doi: 10.1127/0935-1221/2013/0025-2317
- Lovley, D. R., Baedecker, M. J., Lonergan, D. J., Cozzarelli, I. M., Phillips, E. J., and Siegel, D. I. (1989). Oxidation of aromatic contaminants coupled to microbial iron reduction. *Nature* 339, 297–300. doi: 10.1038/339297a0
- Lovley, D. R., and Phillips, E. J. (1988). Novel mode of microbial energy metabolism: organic carbon oxidation coupled to dissimilatory reduction of iron or manganese. *Appl. Environ. Microbiol.* 54, 1472–1480.
- Madigan, M. T., Martinko, J. M., Dunlap, P. V., and Clark, D. P. (2009). *Brock Biology of Microorganisms*, 12th Edn. San Francisco, CA: Pearson Benjamin Cummings.
- Mahaffy, P. R., Webster, C. R., Atreya, S. K., Franz, H., Wong, M., Conrad, P. G., et al. (2013). Abundance and isotopic composition of gases in the Martian atmosphere from the Curiosity rover. *Science* 341, 263–266. doi: 10.1126/science.1237966
- Malin, M. C., and Edgett, K. S. (2003). Evidence for persistent flow and aqueous sedimentation on early Mars. *Science* 302, 1931–1934. doi: 10.1126/science.1090544
- Mangold, N., Carter, J., Poulet, F., Dehouck, E., Ansan, V., and Loizeau, D. (2012). Late Hesperian aqueous alteration at Majuro crater, Mars. *Planet. Space Sci.* 72, 18–30. doi: 10.1016/j.pss.2012.03.014
- Manning, C. V., Zahnle, K. J., and McKay, C. P. (2009). Impact processing of nitrogen on early Mars. *Icarus* 199, 273–285. doi: 10.1016/j.icarus.2008.10.015
- Martin, W., Baross, J., Kelley, D., and Russell, M. J. (2008). Hydrothermal vents and the origin of life. *Nat. Rev. Microbiol.* 6, 805–814. doi: 10.1038/nrmicro1991
- Martin-Torres, F. J., Zorzano, M.-P., Valentin-Serrano, P., Harri, A.-M., Genzer, M., Kemppinen, O., et al. (2015). Transient liquid water and water activity at Gale crater on Mars. *Nat. Geosci.* 8, 357–361. doi: 10.1038/ngeo2412
- Marzo, G. A., Davila, A. F., Tornabene, L. L., Dohm, J. M., Fairén, A. G., Gross, C., et al. (2010). Evidence for Hesperian impact-induced hydrothermalism on Mars. *Icarus* 208, 667–683. doi: 10.1016/j.icarus.2010.03.013
- McSween, H. Y., Grove, T. L., and Wyatt, M. B. (2003). Constraints on the composition and petrogenesis of the Martian crust. *J. Geophys. Res. Planets* 108:5135. doi: 10.1029/2003JE002175
- McSween, H. Y., Taylor, G. J., and Wyatt, M. B. (2009). Elemental composition of the Martian crust. *Science* 324, 736–739. doi: 10.1126/science.1165871
- Michalski, J. R., Cuadros, J., Niles, P. B., Parnell, J., Rogers, A. D., and Wright, S. P. (2013). Groundwater activity on Mars and implications for a deep biosphere. *Nat. Geosci.* 6, 133–138. doi: 10.1038/ngeo1706
- Millero, F. J., Sotolongo, S., and Izaguirre, M. (1987). The oxidation kinetics of Fe (II) in seawater. *Geochim. Cosmochim. Acta* 51, 793–801. doi: 10.1016/0016-7037(87)90093-7
- Ming, D. W., Archer, P. D., Glavin, D. P., Eigenbrode, J. L., Franz, H. B., Sutter, B., et al. (2014). Volatile and organic compositions of sedimentary rocks in Yellowknife Bay, Gale crater, Mars. *Science* 343:1245267. doi: 10.1126/science.1245267
- Ming, D. W., Mittlefehldt, D. W., Morris, R. V., Golden, D. C., Gellert, R., Yen, A., et al. (2006). Geochemical and mineralogical indicators for aqueous processes in the Columbia Hills of Gusev crater, Mars. *J. Geophys. Res. Planets* 111:E02S12. doi: 10.1029/2005JE002560
- Miot, J., Benzerara, K., Morin, G., Kappler, A., Bernard, S., Obst, M., et al. (2009). Iron biomineralization by anaerobic neutrophilic iron-oxidizing bacteria. *Geochim. Cosmochim. Acta* 73, 696–711. doi: 10.1016/j.gca.2008.10.033
- Miot, J., Bernard, S., Bourreau, M., Guyot, F., and Kish, A. (2017). Experimental maturation of *Archaea* encrusted by Fe-phosphates. *Sci. Rep.* 7:16984. doi: 10.1038/s41598-017-17111-9
- Miot, J., and Etique, M. (2016). “Formation and transformation of iron-bearing minerals by iron(II)-oxidizing and iron(III)-reducing bacteria,” in *Iron Oxides: From Nature to Applications*, ed. D. Faivre (Weinheim: Wiley-VCH), 53–98.
- Miot, J., Jézéquel, D., Benzerara, K., Cordier, L., Rivas-Lamelo, S., Skouri-Panet, F., et al. (2016). Mineralogical diversity in lake Pavin: connections with water column chemistry and biomineralization processes. *Minerals* 6:24. doi: 10.3390/min6020024
- Miot, J., Li, J., Benzerara, K., Sougrati, M. T., Ona-Nguema, G., Bernard, S., et al. (2014a). Formation of single domain magnetite by green rust oxidation promoted by microbial anaerobic nitrate-dependent iron oxidation. *Geochim. Cosmochim. Acta* 139, 327–343. doi: 10.1016/j.gca.2014.04.047
- Miot, J., Maclellan, K., Benzerara, K., and Boisset, N. (2011). Preservation of protein globules and peptidoglycan in the mineralized cell wall of nitrate-reducing, iron(II)-oxidizing bacteria: a cryo-electron microscopy study. *Geobiology* 9, 459–470. doi: 10.1111/j.1472-4669.2011.00298.x
- Miot, J., Recham, N., Larcher, D., Guyot, F., Brest, J., and Tarascon, J.-M. (2014b). Biomineralized α -Fe₂O₃: texture and electrochemical reaction with Li. *Energy Environ. Sci.* 7, 451–460. doi: 10.1039/C3EE41767K
- Miot, J., Remusat, L., Duprat, E., Gonzalez, A., Pont, S., and Poinot, M. (2015). Fe biomineralization mirrors individual metabolic activity in a nitrate-dependent Fe (II)-oxidizer. *Front. Microbiol.* 6:879. doi: 10.3389/fmicb.2015.00879
- Mirvaux, B., Recham, N., Miot, J., Courty, M., Bernard, S., Beyssac, O., et al. (2016). Iron phosphate/bacteria composites as precursors for textured electrode materials with enhanced electrochemical properties. *J. Electrochem. Soc.* 163, A2139–A2148. doi: 10.1149/2.0101610jes
- Montoya, L., Vizioli, C., Rodríguez, N., Rastoll, M. J., Amils, R., and Marin, I. (2013). Microbial community composition of Tirez lagoon (Spain), a highly sulfated athalassohaline environment. *Aquat. Biosyst.* 9:19. doi: 10.1186/2046-9063-9-19
- Morgan, B., and Lahav, O. (2007). The effect of pH on the kinetics of spontaneous Fe (II) oxidation by O₂ in aqueous solution—basic principles and a simple heuristic description. *Chemosphere* 68, 2080–2084. doi: 10.1016/j.chemosphere.2007.02.015
- Morris, R. V., Klingelhoefer, G., Schröder, C., Rodionov, D. S., Yen, A., Ming, D. W., et al. (2006a). Mössbauer mineralogy of rock, soil, and dust at Gusev crater, Mars: spirit’s journey through weakly altered olive basalt on the plains and pervasively altered basalt in the Columbia Hills. *J. Geophys. Res. Planets* 111:E02S13. doi: 10.1029/2005JE002584
- Morris, R. V., Klingelhoefer, G., Schröder, C., Rodionov, D. S., Yen, A., Ming, D. W., et al. (2006b). Mössbauer mineralogy of rock, soil, and dust at Meridiani Planum, Mars: opportunity’s journey across sulfate-rich outcrop, basaltic sand and dust, and hematite lag deposits. *J. Geophys. Res. Planets* 111:E12S15. doi: 10.1029/2006JE002791
- Morris, R. V., Vaniman, D. T., Blake, D. F., Gellert, R., Chipera, S. J., Rampe, E. B., et al. (2016). Silicic volcanism on Mars evidenced by tridymite in high-SiO₂ sedimentary rock at Gale crater. *Proc. Natl. Acad. Sci. U.S.A.* 113, 7071–7076. doi: 10.1073/pnas.1607098113
- Muehe, E. M., Gerhardt, S., Schink, B., and Kappler, A. (2009). Ecophysiology and the energetic benefit of mixotrophic Fe (II) oxidation by various strains of nitrate-reducing bacteria. *FEMS Microbiol. Ecol.* 70, 335–343. doi: 10.1111/j.1574-6941.2009.00755.x
- Myers, C. R., and Nealon, K. H. (1988). Microbial reduction of manganese oxides: interactions with iron and sulfur. *Geochim. Cosmochim. Acta* 52, 2727–2732. doi: 10.1016/0016-7037(88)90041-5
- Nachon, M., Mangold, N., Forni, O., Kah, L. C., Cousin, A., Wiens, R. C., et al. (2017). Chemistry of diagenetic features analyzed by ChemCam at Pahrump Hills, Gale crater, Mars. *Icarus* 281, 121–136. doi: 10.1016/j.icarus.2016.08.026
- Navarro-González, R., Vargas, E., de La Rosa, J., Raga, A. C., and McKay, C. P. (2010). Reanalysis of the Viking results suggests perchlorate and organics at midlatitudes on Mars. *J. Geophys. Res. Planets* 115:E12010. doi: 10.1029/2010JE003599
- Nealon, K. H. (1997). The limits of life on Earth and searching for life on Mars. *J. Geophys. Res. Planets* 102, 23675–23686. doi: 10.1029/97JE01996
- Nie, N. X., Dauphas, N., and Greenwood, R. C. (2017). Iron and oxygen isotope fractionation during iron UV photo-oxidation: implications for early Earth and Mars. *Earth Planet. Sci. Lett.* 458, 179–191. doi: 10.1016/j.epsl.2016.10.035
- Nixon, S. L. (2014). *Microbial Iron Reduction on Earth and Mars*. Ph.D. thesis: The Open University, Milton Keynes.
- Nixon, S. L., Cockell, C. S., and Tranter, M. (2012). Limitations to a microbial iron cycle on Mars. *Planet. Space Sci.* 72, 116–128. doi: 10.1016/j.pss.2012.04.003

- Nixon, S. L., Cousins, C. R., and Cockell, C. S. (2013). Plausible microbial metabolisms on Mars. *Astron. Geophys.* 54, 1.13–1.16. doi: 10.1093/astrogeo/ats034
- Oren, A., Bardavid, R. E., and Mana, L. (2014). Perchlorate and halophilic prokaryotes: implications for possible halophilic life on Mars. *Extremophiles* 18, 75–80. doi: 10.1007/s00792-013-0594-9
- Osinski, G. R., Tornabene, L. L., Banerjee, N. R., Cockell, C. S., Flemming, R., Izawa, M. R., et al. (2013). Impact-generated hydrothermal systems on Earth and Mars. *Icarus* 224, 347–363. doi: 10.1038/srep03487
- Palucis, M. C., Dietrich, W. E., Williams, R. M., Hayes, A. G., Parker, T., Sumner, D. Y., et al. (2016). Sequence and relative timing of large lakes in Gale crater (Mars) after the formation of Mount Sharp. *J. Geophys. Res. Planets* 121, 472–496. doi: 10.1002/2015JE004905
- Pantke, C., Obst, M., Benzerara, K., Morin, G., Ona-Nguema, G., Dippon, U., et al. (2012). Green rust formation during Fe (II) oxidation by the nitrate-reducing *Acidovorax* sp. strain BoFeN1. *Environ. Sci. Technol.* 46, 1439–1446. doi: 10.1021/es2016457
- Parnell, J., Lee, P., Cockell, C. S., and Osinski, G. R. (2004). Microbial colonization in impact-generated hydrothermal sulphate deposits, Haughton impact structure, and implications for sulphates on Mars. *Int. J. Astrobiol.* 3, 247–256. doi: 10.1017/S14735504004001995
- Pirajno, F. (ed.). (2009). “Hydrothermal processes associated with meteorite impacts,” in *Hydrothermal Processes and Mineral Systems* (Dordrecht: Springer), 1097–1130. doi: 10.1007/978-1-4020-8613-7_11
- Popa, R., Smith, A. R., Popa, R., Boone, J., and Fisk, M. (2012). Olivine-respiring bacteria isolated from the rock-ice interface in a lava-tube cave, a Mars analog environment. *Astrobiology* 12, 9–18. doi: 10.1089/ast.2011.0639
- Raghoebarsing, A. A., Pol, A., van de Pas-Schoonen, K. T., Smolders, A. J. P., Ettwig, K. F., Rijpstra, W. I. C., et al. (2006). A microbial consortium couples anaerobic methane oxidation to denitrification. *Nature* 440, 918–921. doi: 10.1038/nature04617
- Raiswell, R., and Canfield, D. E. (2012). The iron biogeochemical cycle past and present. *Geochem. Perspect.* 1, 1–322. doi: 10.1038/nature20772
- Ramirez, R. M., Koppaparapu, R., Zegger, M. E., Robinson, T. D., Freedman, R., and Kasting, J. F. (2014). Warming early Mars with CO₂ and H₂. *Nat. Geosci.* 7, 59–63. doi: 10.1038/ngeo2000
- Rampe, E. B., Ming, D. W., Blake, D. F., Bristow, T. F., Chipera, S. J., Grotzinger, J. P., et al. (2017a). Mineralogy of an ancient lacustrine mudstone succession from the Murray formation, Gale crater, Mars. *Earth Planet. Sci. Lett.* 471, 172–185. doi: 10.1016/j.epsl.2017.04.021
- Rampe, E. B., Ming, D. W., Grotzinger, J. P., Morris, R. V., Blake, D. F., Vaniman, D. T., et al. (2017b). “Mineral trends in early Hesperian lacustrine mudstone at Gale crater, Mars,” in *Proceedings of the 48th Lunar and Planetary Science Conference*, The Woodlands, TX.
- Reimers, C. E., Fischer, K. M., Merewether, R., Smith, K., and Jahnke, R. A. (1986). Oxygen microprofiles measured in situ in deep ocean sediments. *Nature* 320, 741–744. doi: 10.1038/320741a0
- Revsbech, N. P., Sorensen, J., Blackburn, T. H., and Lomholt, J. P. (1980). Distribution of oxygen in marine sediments measured with microelectrodes. *Limnol. Oceanogr.* 25, 403–411. doi: 10.4319/lo.1980.25.3.0403
- Roden, E. E., Sobolev, D., Glazer, B., and Luther, G. W. (2004). Potential for microscale bacterial Fe redox cycling at the aerobic-anaerobic interface. *Geomicrobiol. J.* 21, 379–391. doi: 10.1080/01490450490485872
- Santelli, C. M., Welch, S. A., Westrich, H. R., and Banfield, J. F. (2001). The effect of Fe-oxidizing bacteria on Fe-silicate mineral dissolution. *Chem. Geol.* 180, 99–115. doi: 10.1016/S0009-2541(01)00308-4
- Schädlar, S., Burkhardt, C., Hegler, F., Straub, K., Miot, J., Benzerara, K., et al. (2009). Formation of cell-iron-mineral aggregates by phototrophic and nitrate-reducing anaerobic Fe (II)-oxidizing bacteria. *Geomicrobiol. J.* 26, 93–103. doi: 10.1111/gbi.12088
- Schieber, J. (2002). Sedimentary pyrite: a window into the microbial past. *Geology* 30, 531–534. doi: 10.1130/0091-7613(2002)030<0531:SPAUIT>2.0.CO;2
- Schuerger, A. C., and Nicholson, W. L. (2016). Twenty species of hypobarophilic bacteria recovered from diverse soils exhibit growth under simulated Martian conditions at 0.7 kPa. *Astrobiology* 16, 964–976. doi: 10.1089/ast.2016.1587
- Schwenzer, S. P., Bridges, J. C., Wiens, R. C., Conrad, P. G., Kelley, S. P., Leveille, R., et al. (2016). Fluids during diagenesis and sulfate vein formation in sediments at Gale crater, Mars. *Meteorit. Planet. Sci.* 51, 2175–2202. doi: 10.1111/maps.12668
- Schwenzer, S. P., and Kring, D. A. (2009). Impact-generated hydrothermal systems capable of forming phyllosilicates on Noachian Mars. *Geology* 37, 1091–1094. doi: 10.1130/G30340A.1
- Sephton, M. A., Wright, I. P., Gilmour, I., de Leeuw, J. W., Grady, M. M., and Pillinger, C. T. (2002). High molecular weight organic matter in Martian meteorites. *Planet. Space Sci.* 50, 711–716. doi: 10.1016/S0032-0633(02)00053-3
- Sholes, S. F., Smith, M. L., Claire, M. W., Zahnle, K. J., and Catling, D. C. (2017). Anoxic atmospheres on Mars driven by volcanism: implications for past environments and life. *Icarus* 290, 46–62. doi: 10.1016/j.icarus.2017.02.022
- Smith, A. R. (2011). *Subsurface Igneous Mineral Microbiology: Iron-Oxidizing Organotrophs on Olivine Surfaces and the Significance of Mineral Heterogeneity in Basalts*. Master thesis, Portland State University, Portland.
- Smith, M. L., Claire, M. W., Catling, D. C., and Zahnle, K. J. (2014). The formation of sulfate, nitrate and perchlorate salts in the Martian atmosphere. *Icarus* 231, 51–64. doi: 10.1016/j.icarus.2013.11.031
- Squyres, S. W., Arvidson, R. E., Bell, J., Calef, F., Clark, B., Cohen, B., et al. (2012). Ancient impact and aqueous processes at Endeavour Crater, Mars. *Science* 336, 570–576. doi: 10.1126/science.1220476
- Squyres, S. W., Grotzinger, J. P., Arvidson, R. E., Bell, J. F., Calvin, W., Christensen, P. R., et al. (2004). In situ evidence for an ancient aqueous environment at Meridiani Planum, Mars. *Science* 306, 1709–1714. doi: 10.1126/science.1104559
- Squyres, S. W., and Knoll, A. H. (2005). Sedimentary rocks at Meridiani Planum: 006Frigin, diagenesis, and implications for life on Mars. *Earth Planet. Sci. Lett.* 240, 1–10. doi: 10.1016/j.epsl.2005.09.038
- Steele, A., McCubbin, F. M., Fries, M., Kater, L., Bockor, N. Z., Fogel, M. L., et al. (2012). A reduced organic carbon component in Martian basalts. *Science* 337, 212–215. doi: 10.1126/science.1220715
- Stern, J. C., Sutter, B., Freissinet, C., Navarro-González, R., McKay, C. P., Archer, P. D., et al. (2015). Evidence for indigenous nitrogen in sedimentary and aeolian deposits from the Curiosity rover investigations at Gale crater, Mars. *Proc. Natl. Acad. Sci. U.S.A.* 112, 4245–4250. doi: 10.1073/pnas.1420932112
- Stern, J. C., Sutter, B., Jackson, W. A., Navarro-González, R., McKay, C. P., Ming, D. W., et al. (2017). The nitrate/(per)chlorate relationship on Mars. *Geophys. Res. Lett.* 44, 2643–2651. doi: 10.1002/2016GL072199
- Stevens, M. H., Evans, J., Schneider, N. M., Stewart, A. I. F., Deighan, J., Jain, S. K., et al. (2015). New observations of molecular nitrogen in the Martian upper atmosphere by IUVS on MAVEN. *Geophys. Res. Lett.* 42, 9050–9056. doi: 10.1002/2015GL065319
- Straub, K. L., Benz, M., Schink, B., and Widdel, F. (1996). Anaerobic, nitrate-dependent microbial oxidation of ferrous iron. *Appl. Environ. Microbiol.* 62, 1458–1460.
- Straub, K. L., and Buchholz-Cleven, B. E. (1998). Enumeration and detection of anaerobic ferrous iron-oxidizing, nitrate-reducing bacteria from diverse European sediments. *Appl. Environ. Microbiol.* 64, 4846–4856.
- Straub, K. L., Schönhuber, W. A., Buchholz-Cleven, B. E., and Schink, B. (2004). Diversity of ferrous iron-oxidizing, nitrate-reducing bacteria and their involvement in oxygen-independent iron cycling. *Geomicrobiol. J.* 21, 371–378. doi: 10.1080/01490450490485854
- Strohm, T. O., Griffin, B., Zumft, W. G., and Schink, B. (2007). Growth yields in bacterial denitrification and nitrate ammonification. *Appl. Environ. Microbiol.* 73, 1420–1424. doi: 10.1128/AEM.02508-06
- Summers, D. P., Basa, R. C., Khare, B., and Rodoni, D. (2012). Abiotic nitrogen fixation on terrestrial planets: reduction of NO to ammonia by FeS. *Astrobiology* 12, 107–114. doi: 10.1089/ast.2011.0646
- Summers, D. P., and Khare, B. (2007). Nitrogen fixation on early Mars and other terrestrial planets: experimental demonstration of abiotic fixation reactions to nitrite and nitrate. *Astrobiology* 7, 333–341. doi: 10.1089/ast.2006.0032
- Summers, S. (2013). *The Bacterial Ecology and Function from a Sub-surface Critical Zone*. Ph.D. thesis, The Open University, Milton Keynes.
- Sutter, B., Eigenbrode, J. L., Steele, A., McAdam, A., Ming, D. W., Archer, D. Jr., et al. (2016). “The sample analysis at Mars (SAM) detections of CO₂ and CO in sedimentary material from Gale crater, Mars: implications for the presence of organic carbon and microbial habitability on Mars,” in *Proceedings of the AGU Fall Meeting*, New Orleans, LA.
- Toporski, J. K., Steele, A., Westall, F., Thomas-Keppta, K. L., and McKay, D. S. (2002). The simulated silicification of bacteria—new clues to the

- modes and timing of bacterial preservation and implications for the search for extraterrestrial microfossils. *Astrobiology* 2, 1–26. doi: 10.1089/153110702753621312
- Tosca, N. J., and McLennan, S. M. (2006). Chemical divides and evaporite assemblages on Mars. *Earth Planet. Sci. Lett.* 241, 21–31. doi: 10.1016/j.epsl.2005.10.021
- Tosca, N. J., and McLennan, S. M. (2009). Experimental constraints on the evaporation of partially oxidized acid-sulfate waters at the Martian surface. *Geochim. Cosmochim. Acta* 73, 1205–1222. doi: 10.1016/j.gca.2008.11.015
- Tuff, J., Wade, J., and Wood, B. (2013). Volcanism on Mars controlled by early oxidation of the upper mantle. *Nature* 498, 342–345. doi: 10.1038/nature12225
- Vaniman, D. T., Bish, D. L., Ming, D. W., Bristow, T. F., Morris, R. V., Blake, D. F., et al. (2013). Mineralogy of a mudstone at Yellowknife Bay, Gale crater, Mars. *Science* 343:1243480. doi: 10.1126/science.1243480
- Visscher, P. T., Beukema, J., and van Gemerden, H. (1991). In situ characterization of sediments: measurements of oxygen and sulfide profiles with a novel combined needle electrode. *Limnol. Oceanogr.* 36, 1476–1480. doi: 10.4319/lo.1991.36.7.1476
- Wacey, D., McLoughlin, N., Kilburn, M. R., Saunders, M., Cliff, J. B., Kong, C., et al. (2013). Nanoscale analysis of pyritized microfossils reveals differential heterotrophic consumption in the 1.9-Ga Gunflint chert. *Proc. Natl. Acad. Sci. U.S.A.* 110, 8020–8024. doi: 10.1073/pnas.1221965110
- Weber, K. A., Achenbach, L. A., and Coates, J. D. (2006a). Microorganisms pumping iron: anaerobic microbial iron oxidation and reduction. *Nat. Rev. Microbiol.* 4, 752–764.
- Weber, K. A., Hedrick, D. B., Peacock, A. D., Thrash, J. C., White, D. C., Achenbach, L. A., et al. (2009). Physiological and taxonomic description of the novel autotrophic, metal oxidizing bacterium, *Pseudogulbenkiania* sp. strain 2002. *Appl. Microbiol. Biotechnol.* 83, 555–565. doi: 10.1007/s00253-009-1934-7
- Weber, K. A., Pollock, J., Cole, K. A., O'Connor, S. M., Achenbach, L. A., and Coates, J. D. (2006b). Anaerobic nitrate-dependent iron (II) bio-oxidation by a novel lithoautotrophic betaproteobacterium, strain 2002. *Appl. Environ. Microbiol.* 72, 686–694.
- Webster, C. R., Mahaffy, P. R., Atreya, S. K., Flesch, G. J., Mischna, M. A., Meslin, P.-Y., et al. (2015). Mars methane detection and variability at Gale crater. *Science* 347, 415–417. doi: 10.1126/science.1261713
- Welhan, J. A., and Craig, H. (1983). “Methane, hydrogen and helium in hydrothermal fluids at 21 N on the East Pacific Rise,” in *Hydrothermal Processes at Seafloor Spreading Centers*, eds P. A. Rona, K. Boström L. Laubier, and K. L. Smith (Berlin: Springer), 391–409. doi: 10.1007/978-1-4899-0402-7_17
- Wheat, C. G., Feely, R. A., and Mottl, M. J. (1996). Phosphate removal by oceanic hydrothermal processes: an update of the phosphorus budget in the oceans. *Geochim. Cosmochim. Acta* 60, 3593–3608. doi: 10.1016/0016-7037(96)00189-5
- Williams, R. M., Grotzinger, J. P., Dietrich, W. E., Gupta, S., Sumner, D. Y., Wiens, R. C., et al. (2013). Martian fluvial conglomerates at Gale crater. *Science* 340, 1068–1072. doi: 10.1126/science.1237317
- Yen, A. S., Ming, D. W., Vaniman, D. T., Gellert, R., Blake, D. F., Morris, R. V., et al. (2017). Multiple stages of aqueous alteration along fractures in mudstone and sandstone strata in Gale Crater, Mars. *Earth Planet. Sci. Lett.* 471, 186–198. doi: 10.1016/j.epsl.2017.04.033
- Yen, A. S., Mittlefehldt, D. W., McLennan, S. M., Gellert, R., Bell, J., McSween, H. Y., et al. (2006). Nickel on Mars: constraints on meteoritic material at the surface. *J. Geophys. Res. Planets* 111:E12S11. doi: 10.1029/2006JE002797

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2018 Price, Pearson, Schwenzer, Miot and Olsson-Francis. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.