



The University of Bradford Institutional Repository

<http://bradscholars.brad.ac.uk>

This work is made available online in accordance with publisher policies. Please refer to the repository record for this item and our Policy Document available from the repository home page for further information.

To see the final version of this work please visit the publisher's website. Access to the published online version may require a subscription.

Link to publisher's version: <https://doi.org/10.3389/fmicb.2015.01226>

Citation: Liu Y, Chan K and Chang C (2015) Modulation of host biology by *Pseudomonas aeruginosa* quorum sensing signal molecules: messengers or traitors. *Frontiers in Microbiology*. 6:1226.

Copyright statement: © 2015 Liu, Chan and Chang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.



Modulation of Host Biology by *Pseudomonas aeruginosa* Quorum Sensing Signal Molecules: Messengers or Traitors

Yi-Chia Liu¹, Kok-Gan Chan² and Chien-Yi Chang^{3,4*}

¹ Division of Molecular Microbiology, School of Life Sciences, University of Dundee, Dundee, UK, ² Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia, ³ Centre for Bacterial Cell Biology, Medical School, Newcastle University, Newcastle upon Tyne, UK, ⁴ Interdisciplinary Computing and Complex BioSystems (ICOS) Research Group, School of Computing Science, Newcastle University, Newcastle upon Tyne, UK

OPEN ACCESS

Edited by:

Katharina F. Kubatzky,
University Hospital Heidelberg,
Germany

Reviewed by:

Rodolfo García-Contreras,
Instituto Nacional de Cardiología,
Mexico

Gertrud Maria Hänsch,
University of Heidelberg, Germany

*Correspondence:

Chien-Yi Chang
chien-yi.chang@ncl.ac.uk

Specialty section:

This article was submitted to
Microbial Immunology,
a section of the journal
Frontiers in Microbiology

Received: 09 September 2015

Accepted: 20 October 2015

Published: 09 November 2015

Citation:

Liu Y-C, Chan K-G and Chang C-Y
(2015) Modulation of Host Biology by
Pseudomonas aeruginosa Quorum
Sensing Signal Molecules:
Messengers or Traitors.
Front. Microbiol. 6:1226.
doi: 10.3389/fmicb.2015.01226

Bacterial cells sense their population density and respond accordingly by producing various signal molecules to the surrounding environments thereby trigger a plethora of gene expression. This regulatory pathway is termed quorum sensing (QS). Plenty of bacterial virulence factors are controlled by QS or QS-mediated regulatory systems and QS signal molecules (QSSMs) play crucial roles in bacterial signaling transduction. Moreover, bacterial QSSMs were shown to interfere with host cell signaling and modulate host immune responses. QSSMs not only regulate the expression of bacterial virulence factors but themselves act in the modulation of host biology that can be potential therapeutic targets.

Keywords: Quorum sensing, *N*-acyl homoserine lactones, *Pseudomonas* quinolone signal, *Pseudomonas aeruginosa*, immunomodulation

INTRODUCTION

Quorum sensing (QS) is coined to describe the phenomenon of an intercellular co-operative behavior of bacteria used to coordinate the activities of individual cells. Diffusible QS signal molecules (QSSMs) play crucial roles in signal transduction of which, when QSSMs reach a threshold concentration, can coordinate multiple gene expression and a change in the behavior of bacterial population through the activation of sensor regulatory proteins (Fuqua et al., 1994; Miller and Bassler, 2001; Williams and Cámara, 2009). Bacteria produce a broad-range of signal molecules. Different types of QSSMs have been identified and characterized (LaSarre and Federle, 2013). Besides prokaryote, bacterial QSSMs also affect the settlement and germination of eukaryotic seaweed zoospores (Joint et al., 2002; Twigg et al., 2013). In several pathogenic bacteria, QS control their virulence determinants and contribute to bacterial pathogenesis. Due to the fact that the population density-dependent regulatory systems used by many bacterial pathogens are not essential for survival under most conditions, the disruption/interference of QS is considered an alternative approach to attenuate bacterial virulence in infections (LaSarre and Federle, 2013). However, this point has recently been argued that the resistance mechanisms against QS inhibitors could be possible and have been identified (Defoirdt et al., 2010, 2013; García-Contreras et al., 2013, 2015a).

Pseudomonas aeruginosa is an ubiquitous Gram-negative bacterium with remarkably large and complex genome and is capable of adapting to versatile environments. In human cystic fibrosis (CF) lungs where *P. aeruginosa* has evolved the ability to form biofilms which are difficult to be eradicated by antibiotics (Heeb et al., 2011; Winsor et al., 2011). QS is responsible for the regulation of a large number of genes, for instance, around 10% of genes in the genome of *P. aeruginosa* are regulated by QS (Williams and Cámara, 2009). Here we review recent advances of *P. aeruginosa* QSSMs focusing on their roles in interference with host cells (Table 1) and the development of novel compounds that counteract the QSSMs activities.

N-ACYL HOMOSERINE LACTONES (AHLs) AND THEIR MODULATIONS IN HOST CELLS

Gram-negative bacteria, like *Aliivibrio fischeri* (previous *Vibrio fischeri*; Urbanczyk et al., 2007), have a conserved QS system with two central components, the LuxR-type and LuxI-type proteins, which serve as the signal receptor and signal synthase, respectively. LuxI catalyzes the synthesis of signaling molecules called *N*-acyl homoserine lactones (AHLs). When an AHLs concentration of 10 nM is reached, AHLs interact with LuxR and form a complex which promotes the expression of target genes, *luxICDABE* for bioluminescence production and also the LuxI production (Kaplan and Greenberg, 1985). This forms a positive loop to produce more signal molecules (Fuqua et al., 1994; Cámara et al., 2002; Fuqua and Greenberg, 2002). The *N*-acyl homoserine lactone consists of a homoserine lactone ring from *S*-adenosylmethionine (SAM) and acyl chain from acyl acyl-carrier-protein (acyl-ACP) linked by an amide bond (Parsek et al., 1999). Based on the acyl-ACP binding site, different LuxI homologs produce different AHLs with various acyl side chains (Watson et al., 2002; Gould et al., 2004). A broad range of AHLs is produced in Gram-negative bacteria and AHL-QS systems control various bacterial behaviors (LaSarre and Federle, 2013). In *A. fischeri* *N*-(3-oxohexanoyl) homoserine lactone (3-oxo-C6-HSL) is produced for controlling bioluminescence production. In *P. aeruginosa* two AHL synthases, RhII and LasI, produce a wide spectrum of AHLs including *N*-butanoyl-homoserine lactone (C4-HSL), *N*-hexanoyl-homoserine lactone (C6-HSL) by RhII and *N*-(3-oxooctanoyl)-homoserine lactone (3-oxo-C8-HSL), *N*-(3-oxodecanoyl)-homoserine lactone (3-oxo-C10-HSL), *N*-(3-oxododecanoyl)-homoserine lactone (3-oxo-C12-HSL) and *N*-(3-oxotetradecanoyl)-homoserine lactone (3-oxo-C14-HSL) by LasI (Ortori et al., 2011). An unusual *N*-(3-oxohexadecanoyl)-homoserine lactone (3-oxo-C16-HSL) secreted by an environmental *Pseudomonas* sp. from a diseased Tilapia fish suggests that 3-oxo-C16-HSL may contribute to the pathogenesis (Chang et al., 2012).

The abundant concentration of 3-oxo-C12-HSL in the culture of *P. aeruginosa* prompted investigations for its role in the pathogenesis with a mechanism potentially distinct from other pathogens. Indeed, 3-oxo-C12-HSL was found to activate

mammalian cells through a mechanism independent of the toll-like receptor (TLR) pathways (Kravchenko et al., 2006). 3-oxo-C12-HSL was shown to activate pro-inflammatory responses in human epithelial and fibroblast cells through the induction the transcriptional factor, nuclear factor kappa-light-chain-enhancer of activated B cells (NF- κ B) via the phosphorylation of ERK/MARK (Smith et al., 2001). However, this molecule selectively disrupts the NF- κ B signaling pathway in activated macrophages (Kravchenko et al., 2008). Studies indicated that 3-oxo-C12-HSL not only induces apoptosis in haematopoietic cells but is cytotoxic to non-haematopoietic cells including airway epithelial cells, endothelial cells, fibroblasts, and mesenchymal stem cells (Tateda et al., 2003; Shiner et al., 2006; Schwarzer et al., 2010, 2012; Grabiner et al., 2014; Holban et al., 2014). *P. aeruginosa* 3-oxo-C12-HSL also impairs the epithelial barrier integrity through the alternations of calcium signaling and phosphorylation status of junctional proteins in the intestinal epithelial cells (Vikström et al., 2009, 2010).

In addition to its cytotoxicity, the role of 3-oxo-C12-HSL in immunomodulation has been intensively investigated (Table 1). Ritchie et al. (2005) reported that 3-oxo-C12-HSL inhibits the differentiation of Th1 and Th2 cells. Human polymorphonuclear neutrophils (PMNs) are attracted by 3-oxo-C12-HSL and increasingly express the adhesion proteins CD11b/CD18 and the immunoglobulin receptors CD16 and CD64 (Zimmermann et al., 2006; Wagner et al., 2007). The downregulation of the immune responses by 3-oxo-C12-HSL was demonstrated in human monocytes and murine macrophage-like cells in the presence of lipopolysaccharides (LPS) that 3-oxo-C12-HSL inhibits the production of pro-inflammatory cytokine tumor necrosis factor α (TNF- α) but promotes the production of anti-inflammatory cytokine interleukin-10 (IL-10; Hooi et al., 2004; Glucksam-Galnoy et al., 2013). Grabiner et al. (2014) noticed that despite the increasing transcriptional expression of the murine interleukin 8 (IL-8) homologs KC and interleukin 6 (IL-6) in murine embryonic fibroblasts (MEFs), KC and IL-6 protein secretion were inhibited by the treatment of 3-oxo-C12-HSL. It was shown that 3-oxo-C12-HSL acts upon the activation of endoplasmic reticulum (ER) stress transducer protein kinase RNA-like ER kinase (PERK) leading to the inhibition of protein synthesis. However, PERK is independent of 3-oxo-C12-HSL induced apoptosis indicating that 3-oxo-C12-HSL interferes with host cell biological activities through different mechanisms (Grabiner et al., 2014). Recent advances on the interactions between 3-oxo-C12-HSL and various types of host cells are highlighted in the review (Holm and Vikström, 2014).

Several host targets of 3-oxo-C12-HSL have been identified (Figure 1). In murine fibroblasts and human lung epithelial cells peroxisome proliferator-activated receptor beta/delta (PPAR β/δ) and PPAR γ may be the 3-oxo-C12-HSL receptors for pro-inflammatory responses (Jahoor et al., 2008; Cooley et al., 2010). 3-oxo-C12-HSL interacts and co-localizes with the IQ-motif-containing GTPase-activating protein IQGAP1 in human intestinal epithelial cells that causes the alternation of cell migration in a Rac1 and Cdc42- dependent manner (Karlsson et al., 2012). MEFs in lack of a transcriptional factor X-box binding protein 1 transcription factor (XBP1s) are

TABLE 1 | Bacterial quorum sensing molecules and their roles in pathogenesis/immunomodulation.

| Quorum sensing (QS) signal molecules (QSSMs) | Mechanism of virulence | Effect concentrations | Reference |
|--|--|---|--|
| 3-oxo-C12-HSL | <p>Apoptosis and cytotoxicity</p> <p>3-oxo-C12-HSL is cytotoxic to murine bone-marrow derived macrophages, neutrophils and monocytic cell lines.</p> <p>3-oxo-C12-HSL induces apoptosis in murine fibroblasts and human vascular endothelial cells (HUVEC)</p> <p>3-oxo-C12-HSL triggers intrinsic apoptotic pathway in airway epithelial cells including depolarization of mitochondrial membrane potential, release of cytochrome C and activation of caspases 3/7, 8, and 9. 3-oxo-C12-HSL-mediated apoptosis is independent of the presence of CFTR in airway epithelial cells</p> <p>3-oxo-C12-HSL promotes human mesenchymal stem cells (MSCs) apoptosis</p> <p>Disruption of barrier integrity</p> <p><i>Pseudomonas aeruginosa</i> 3O-C12-HSL causes the loss of epithelial barrier function via calcium signaling and further alteration in the phosphorylation status of junction proteins</p> <p>Immunomodulation and/or signaling</p> <p>3-oxo-C12-HSL promotes the expression and production of IL-8 in human epithelial and fibroblast cells through the induction of NF-κB via the phosphorylation of ERK/MARK</p> <p>3-oxo-C12-HSL inhibits ConA-activated PBMCs proliferation and IL-2 secretion</p> <p>3-oxo-C12-HSL inhibits the proliferation of anti-CD3/anti-CD28 antibody activated T cells</p> <p>3-oxo-C12-HSL inhibits the differentiation of Th1 and Th2 cells</p> <p>3-oxo-C12-HSL increases the cytosolic calcium levels and calcium release through inositol triphosphate (IP3) receptors in the ER.</p> <p>3-oxo-C12-HSL promotes neutrophil chemotaxis, phagocytosis and up-regulates the expression of CD11b/CD18 and CD16/CD64 receptors</p> <p>3-oxo-C12-HSL selectively disrupts NF-κB signaling but not TLR-dependent pathways in activated macrophages</p> <p>3-oxo-C12-HSL binds to PPARγ ligand binding domain</p> <p>3-oxo-C12-HSL increases the secretion of IL-1β in human MSCs</p> <p>3-oxo-C12-HSL activates NF-κB p65 by preventing the re-synthesis of IκB, increases transcription of KC and IL-6 but inhibits secretion of KC and IL-6 by MEFs. 3-oxo-C12-HSL activates PERK and inhibits protein synthesis</p> | <p>12–50 μM</p> <p>100 μM</p> <p>> 10 μM</p> <p>50 μM</p> <p>20–200 and 10 μM with slower response</p> <p>100 μM</p> <p>^aIC₅₀: 18.24 μM</p> <p>IC₅₀: 44.47 μM</p> <p>5 μM</p> <p>1 mM</p> <p>100 μM</p> <p>50 μM</p> <p>25–50 μM</p> <p>50 μM</p> <p>50 or 100 μM</p> | <p>Tateda et al., 2003</p> <p>Shiner et al., 2006</p> <p>Schwarzer et al., 2010</p> <p>Holban et al., 2014</p> <p>Vikström et al., 2009, 2010</p> <p>Smith et al., 2001</p> <p>Hool et al., 2004</p> <p>Hool et al., 2004</p> <p>Ritchie et al., 2005</p> <p>Shiner et al., 2006</p> <p>Zimmermann et al., 2006;</p> <p>Wagner et al., 2007</p> <p>Kravchenko et al., 2006, 2008</p> <p>Jahoor et al., 2008; Cooley et al., 2010</p> <p>Holban et al., 2014</p> <p>Grabiner et al., 2014</p> <p>Williams and Cámara, 2009</p> <p>Diggle et al., 2007</p> <p>Allesen-Holm et al., 2006</p> <p>Häussler and Becker, 2008</p> |
| Alkylquinolones | <p>Change of bacterial behaviors</p> <p>Bacterial autolysis</p> <p>Iron chelation</p> <p>exDNA release</p> <p>Oxidative functions</p> | <p>Spent culture supernatant 15 μg on filter discs</p> <p>50 μM</p> <p>Genetic and phenotype study</p> <p>> 100 μM <i>in vitro</i></p> | <p>(Continued)</p> |

TABLE 1 | Continued


| Quorum sensing (QS) signal molecules (QSSMs) | Mechanism of virulence | Effect concentrations | Reference |
|--|--|--|--|
| | Apoptosis | | |
| | HHQ promotes human MSCs apoptosis | 50 μ M | Holban et al., 2014 |
| | Immunomodulation and/or signaling | | |
| | <i>Pseudomonas</i> quinolone signal (PQS) inhibits the proliferation of ConA-activated PBMCs | IC ₅₀ : 0.90 μ M | Hooi et al., 2004 |
| | PQS affects IL-2 secretion of ConA-stimulated PBMCs | IC ₅₀ : 2.03 μ M | Hooi et al., 2004 |
| | PQS promotes TNF- α production in LPS-treated monocytes | >25 μ M | Hooi et al., 2004 |
| | Inhibition of IL-12 production in dendritic cells resulting in reduction of T-cell proliferation | IC ₅₀ : 17.2 μ M | Skindersoe et al., 2009 |
| | Inhibition of NF- κ B and HIP-1 α pathways in murine epithelial cells and murine macrophages | Bacterial culture supernatants approx. PQS in PA14, 15 μ M approx. PQS in <i>pqsL</i> , 45 μ M | Kirm et al., 2010; Legendre et al., 2012 |
| | Stimulation of chemotaxis of neutrophils via the MAPK and p38 signaling pathways | 10–100 μ M | Hänsch et al., 2014 |
| | PQS activates the secretion of IL-6 and HHQ induces IL-10 secretion by human MSCs | 50 μ M | Holban et al., 2014 |

^a50% inhibitory concentration (IC₅₀).

protective from 3-oxo-C12-HSL and C14-HSL (*N*-tetradecanoyl-homoserine lactone) mediated apoptosis indicating that XBP1s is a critical host target in response of AHLs (Valentine et al., 2013). Paraoxonase 2, in response to 3-oxo-C12-HSL through its lactonase activity, leads to apoptosis in human and murine embryonic epithelial cells (Schwarzer et al., 2015). Interestingly, 3-oxo-C12-HSL activates the expression of a taste receptor T2R38 on the surface of primary human sinonasal cells (Lee et al., 2014) and neutrophils (Maurer et al., 2015). This recognition regulates calcium-dependent NO production thereby stimulates the mucociliary clearance and antibacterial effects suggesting an alternative innate immune defense mechanism distinct from the activation by canonical pattern recognition receptors (PRRs; Lee et al., 2012, 2014). Identification of the host compartments targeted by QSSMs could be the milestone for developing effective therapeutic methods against infections.

PATHOGENIC ROLES OF ALKYL-QUINOLONE SIGNALS

Pseudomonas aeruginosa also employs the alkyl-quinolone (AQ)-based QS system and the signal molecule was termed *Pseudomonas* quinolone signal (PQS; Pesci et al., 1999). The study of the AQs began from their intriguing structures similar to antimicrobial quinolones, although AQs were found no antimicrobial activities. Further studies unveiled that among more than 50 alkyl-quinolones found in *P. aeruginosa*, 2-heptyl-3-hydroxy-4-(1H)-quinolone (PQS) and its precursor molecular 2-heptyl-4(1H)-quinolone (HHQ) are major QSSMs that cooperates with the AHL-QS (Xiao et al., 2006; Heeb et al., 2011). Synthesis of PQS depends on the *pqsABCDE* operon. PqsA, the anthranilate co-enzyme A ligase, catalyzes anthranilate that is produced by PhnAB to anthraniloyl-coenzyme A. PqsD mediates the synthesis of 2-aminobenzoylacetate (2-ABA) from anthraniloyl-coenzyme A and malonyl-CoA, decarboxylating coupling of 2-ABA to an octanoate group of octanoic acid that linked to PqsBC to produce HHQ (Dulcey et al., 2013). A recent study suggested PqsE is involved in the HHQ synthesis through hydrolysing the 2-ABA-CoA to form 2-ABA (Drees and Fetzner, 2015). HHQ can be transformed to PQS by the mono-oxygenase PqsH encoded by *pqsH* located elsewhere on the chromosome (Pesci et al., 1999; Diggle et al., 2006). PqsR, also known as MvfR, is a LysR-type transcriptional regulator, with a conserved N-terminal DNA-binding helix-turn-helix and a C-terminal co-inducer-binding domain. PqsR activates the transcription of *pqsABCDE* and possibly the *phnAB* operon when binding to PQS or HHQ and triggers the typical QS autoinducing response enhancing AQ biosynthesis (Maddocks and Oyston, 2008; Heeb et al., 2011). PQS has been shown to reach the maximal production at late logarithmic phase (Diggle et al., 2003) and its production is promoted by the availability of the substrate anthranilate and the presence of aromatic amino acids (Palmer et al., 2005). AQ- and AHL-QS in *P. aeruginosa* are hierarchical and involved in the regulation of multiple virulence factors including rhamnolipids, pyocyanin, elastases, exotoxin A, and alkaline protease (Xiao



et al., 2006; Dubern and Diggle, 2008; Nadal Jimenez et al., 2012).

Pseudomonas quinolone signal is considered a multifunctional molecule. PQS is involved in bacterial cell autolysis at high population densities in nutrient deprived conditions (Williams and Cámara, 2009). PQS also has iron-chelating properties that contribute to iron transport and facilitates siderophore-mediated iron delivery (Diggle et al., 2007). It has been demonstrated that there is far less extracellular DNA (exDNA) released by a *pqsA* mutant than its wild-type counterpart either in planktonic or biofilm cultures (Allesen-Holm et al., 2006). PQS has dual pro- and anti-oxidative functions for developing different levels of tolerance in *P. aeruginosa* cells to environmental stress (Häussler and Becker, 2008). This may shape the whole population structure, increase the fitness in hostile environments and lead to the development of resistance to host immune systems (García-Contreras et al., 2015a,b).

A new QS molecule, 2-(2-hydroxyphenyl)-thiazole-4-carbaldehyde (IQS) encoded by the *ambBCDE* operon was discovered recently (Lee et al., 2013). IQS is induced when *P. aeruginosa* is exposed to a phosphate-deprived environment. Under this unfavorable environment, expression of IQS overcomes the *las*-led QS circuit and promotes the expression of virulence factors. This finding may partially explain how *P. aeruginosa* clinical isolates persist in CF respiratory infections in the absence of a functional *las* system. Despite the fact that the AHL-QS of *P. aeruginosa* have been shown to play central roles

in the regulation of virulence and immune modulation *in vitro*, this situation could differ *in vivo*. Around 50% of strains isolated from lungs of late stage CF patients are deficient in *lasR* function (Winstanley and Fothergill, 2009). Moreover, abolishing the whole AHL-QS by generating a quadruple mutation of *rhlIR* and *lasIR* exerted comparable infectivity to the wild-type strain in a mouse lung infection model (Lazenby et al., 2013), suggesting that the AHL-QS may not be required for full pathogenesis *in vivo* and other regulatory mechanism could be involved.

The contribution of the AQ-QS system to *P. aeruginosa* virulence was firstly described by Cao et al. (2001). PqsR positively regulates the expression of *phnAB* operon and the production of elastase, 3-oxo-C12-HSL and PQS that promotes the production of numerous virulence determinants. The *pqsR* mutant was attenuated up to 320-fold in the *Arabidopsis* plant infection model and caused a 65% reduction of mortality in a murine burn wound model (Cao et al., 2001). The interaction of PqsR with the AHL-QS was investigated by other groups showed that the effect of *pqsR* deficiency on pathogenesis is independent from LasRI/RhlRI (Déziel et al., 2004; Dubern and Diggle, 2008). Mutations in the multidrug efflux pump, such as *mexI* and *opmD* led to the inhibition of PQS production and the attenuation of *P. aeruginosa* in rat and plant infection models. Provision of exogenous AQS to these mutants restored the virulence on plants (Aendekerck, 2005). Rampioni et al. (2010) found that both *pqsA* and *pqsE* mutants in PAO1 were attenuated in plant, nematode and mouse burn wound infection models. In an acute urinary tract infection model, PQS molecules were

present in the renal and bladder tissue of mice infected with wild-type *P. aeruginosa* but absent in the mice with PQS mutants infections (Bala et al., 2014). Wild-type *P. aeruginosa* caused more severe inflammation and tissue destruction and greater levels of inflammatory cytokines TNF- α , IL-6, and IL-10 at the site of infection in mice infected with wild-type strain than with PQS mutants. The virulence of PQS mutants can be restored by the addition of exogenous PQS molecules (Bala et al., 2014). These findings indicate that PQS participates in the pathogenesis of *P. aeruginosa*.

Pseudomonas quinolone signal has been identified in sputum, bronchoalveolar lavage fluid (BAL) and mucopurulent fluid from distal airways of end-stage CF lungs removed for transplant and at different stages from asymptotic early stage to late progression, suggesting a potential role of PQS in coordinating virulence factors during the course of infections (Collier et al., 2002; Guina et al., 2003). A study involving 60 CF patients with chronic *P. aeruginosa* infection indicated that the AQs were detectable in the sputum, plasma and urine and the concentrations of molecules are positively correlated to the *P. aeruginosa* bacterial cell density. 2-nonyl-4-hydroxy-quinoline (NHQ) in plasma was suggested to be the biomarker for *P. aeruginosa* infection in CF lungs (Barr et al., 2015). An *in vitro* transcriptomic study investigating the physiology of *P. aeruginosa* grown in CF sputum revealed that the genes associated with PQS metabolism, such as those coding for the aromatic amino acid aminotransferase, 4-hydroxyphenylpyruvate dioxygenase (*hpd*) and *pqsABCDE*, were expressed 10-fold greater than the expression when *P. aeruginosa* was cultured in media containing glucose alone as the carbon source (Palmer et al., 2005).

Pseudomonas aeruginosa AQ molecules have been implicated in the immuno-modulation on host cells. PQS was shown to modulate cell proliferation, the production of interleukin-2 (IL-2) and TNF- α in mitogen-stimulated human peripheral blood mononuclear cells (PBMCs; Hooi et al., 2004). PQS inhibited the production of IL-12 by LPS-stimulated bone marrow-derived dendritic cells which led to reduced T-cell proliferation (Skindersoe et al., 2009). Additionally AQ extracts derived from *P. aeruginosa* PA14 supernatants down-regulated host innate immune responses via inhibition of the NF- κ B and hypoxia-inducible factor 1 alpha (HIF-1 α) pathways in murine macrophages and cells obtained from BAL (Kim et al., 2010; Legendre et al., 2012). A recent study addressed the importance of timing in neutrophil infiltration in relation to the role of PQS in interference with neutrophil chemotaxis. Low levels of PQS stimulated the chemotaxis of neutrophils via the MAPK and p38 signaling pathways, whereas high levels of PQS, most likely produced by biofilm-like *P. aeruginosa*, did not interfere with neutrophils phagocytic capability and viability (Hänsch et al., 2014). Massive neutrophil accumulation is commonly seen in CF airways and high levels of neutrophil elastase correlate with poor pulmonary functions (Downey et al., 2008; Gifford and Chalmers, 2014). PQS may thus provide *P. aeruginosa* with another strategy for bacterial survival via the interference in multiple aspects of host biological activities.

THE DEVELOPMENT OF INHIBITORS AND VACCINES AGAINST QSSMS

Due to the fact that QSSMs have been implicated in the involvement of pathogenesis, the search for inhibitors and the development of vaccines that antagonize QSSMs are currently intensively investigated. Chang et al. (2014) suggested a strategy to screen novel anti-QS compounds from plant extraction that potentially could tackle the QS-mediated infections. The antibody 3-oxo-C12-HSL-BSA conjugate was also shown to alleviate the inflammatory responses by *P. aeruginosa* infections in an acute murine lung infection model (Miyairi et al., 2006). In a burn wound infection model, mice immunized with the vaccine 3-oxo-C12-HSL-r-PcrV conjugate before *P. aeruginosa* infection had higher survival than those without immunization (Golpasha et al., 2015). A high-throughput screening approach based on the inhibition of C12-mediated host responses identified triazolo[4,3-a]quinolines as 3-oxo-C12-HSL inhibitors with nanomolar potency that restore NF- κ B activity in 3-oxo-C12-HSL treated cell lines and shown protective using an *in vivo* dermal infection model (Valentine et al., 2014). Since anthranilate (AA) being the precursor of AQs, halogenated AA analogs were found to inhibit the AQ biosynthesis and down-regulate the expression of PqsR controlled genes. Treatment with AA analogs prior to *P. aeruginosa* infection increased mice survival and lowered the bacterial dissemination to the organs (Lesic et al., 2007). MvfR-regulon inhibitors that bind QS transcriptional regulator MvfR (PqsR) were not only protective in murine acute and persistent infections against *P. aeruginosa* but also effectively reduced the formation of antibiotic-tolerant persisters (Starkey et al., 2014). These studies suggest the therapeutic potential of inhibitors and vaccines against QSSMs in both acute and chronic infections.

CONCLUSION

Quorum sensing-based bacterial communication links the individual bacterial cells to behave as multicellular organisms by employing signal molecules and to promote its population survival in the environment or hosts. QSSMs also interact with host cell signal pathways and the modulation of immune cell biology. For more than a decade strategies have been proposed from the use of inhibitors of QS for containing chronic infections (Hentzer et al., 2003) to the application of QSSMs for modulating immune responses to bacterial infections (Hancock et al., 2012). Understanding how QSSMs interact with host cells seems the promising land to tackle bacterial infections. Here we discussed recent advances on the interference of QSSMs with mammalian cells, the recently identified receptors on mammalian cells that target QSSMs and the QSSM inhibitors and their mechanisms. However, contradictory results suggested that many unknown mechanisms in complex bacteria-host interactions are remained. Also concern about bacterial resistance to QS inhibitors (García-Contreras et al., 2015a) and unexpected impact of QS inhibitors to environment (Decho et al., 2010) is increasing. Extending our understanding of the multiple roles of QSSMs would be valuable

in the development of new therapeutic strategies against bacterial infections.

AUTHOR CONTRIBUTIONS

Y-CL, K-GC, C-YC wrote the paper. Y-CL made the figure. Y-CL and C-YC made the table.

REFERENCES

- Aendekerk, S. (2005). The MexGHI-OpmD multidrug efflux pump controls growth, antibiotic susceptibility and virulence in *Pseudomonas aeruginosa* via 4-quinolone-dependent cell-to-cell communication. *Microbiology* 151, 1113–1125. doi: 10.1099/mic.0.27631-0
- Allesen-Holm, M., Barken, K. B., Yang, L., Klausen, M., Webb, J. S., Kjelleberg, S., et al. (2006). A characterization of DNA release in *Pseudomonas aeruginosa* cultures and biofilms. *Mol. Microbiol.* 59, 1114–1128. doi: 10.1111/j.1365-2958.2005.05008.x
- Bala, A., Chhibber, S., and Harjai, K. (2014). *Pseudomonas* quinolone signalling system: a component of quorum sensing cascade is a crucial player in the acute urinary tract infection caused by *Pseudomonas aeruginosa*. *Int. J. Med. Microbiol.* 304, 1199–1208. doi: 10.1016/j.ijmm.2014.08.013
- Barr, H. L., Halliday, N., Cámara, M., Barrett, D. A., Williams, P., Forrester, D. L., et al. (2015). *Pseudomonas aeruginosa* quorum sensing molecules correlate with clinical status in cystic fibrosis. *Eur. Respir. J.* 46, 1046–1054. doi: 10.1183/09031936.00225214
- Cámara, M., Williams, P., and Hardman, A. (2002). Controlling infection by tuning in and turning down the volume of bacterial small-talk. *Lancet Infect. Dis.* 2, 667–676. doi: 10.1016/S1473-3099(02)00447-4
- Cao, H., Krishnan, G., Goumnerov, B., Tsongalis, J., Tompkins, R., and Rahme, L. G. (2001). A quorum sensing-associated virulence gene of *Pseudomonas aeruginosa* encodes a LysR-like transcription regulator with a unique self-regulatory mechanism. *Proc. Natl. Acad. Sci. U.S.A.* 98, 14613–14618. doi: 10.1073/pnas.251465298
- Chang, C.-Y., Koh, C.-L., Sam, C.-K., Chan, X.-Y., Yin, W. F., and Chan, K. G. (2012). Unusual long-chain N-acyl homoserine lactone production by and presence of quorum quenching activity in bacterial isolates from diseased tilapia fish. *PLoS ONE* 7:e44034. doi: 10.1371/journal.pone.0044034.t002
- Chang, C.-Y., Krishnan, T., Wang, H., Chen, Y., Yin, W. F., Chong, Y. M., et al. (2014). Non-antibiotic quorum sensing inhibitors acting against N-acyl homoserine lactone synthase as druggable target. *Sci. Rep.* 4:7245. doi: 10.1038/srep07245
- Collier, D. N., Anderson, L., McKnight, S. L., Noah, T. L., Knowles, M., Boucher, R., et al. (2002). A bacterial cell to cell signal in the lungs of cystic fibrosis patients. *FEMS Microbiol. Lett.* 215, 41–46. doi: 10.1111/j.1574-6968.2002.tb11367.x
- Cooley, M. A., Whittall, C., and Rolph, M. S. (2010). *Pseudomonas* signal molecule 3-oxo-C12-homoserine lactone interferes with binding of rosiglitazone to human PPAR γ . *Microbes Infect.* 7, 231–237. doi: 10.1016/j.micinf.2009.12.009
- Decho, A. W., Norman, R. S., and Visscher, P. T. (2010). Quorum sensing in natural environments: emerging views from microbial mats. *Trends Microbiol.* 18, 73–80. doi: 10.1016/j.tim.2009.12.008
- Defoirdt, T., Boon, N., and Bossier, P. (2010). Can bacteria evolve resistance to quorum sensing disruption? *PLoS Pathogens* 6:e1000989. doi: 10.1371/journal.ppat.1000989.t001
- Defoirdt, T., Brackman, G., and Coenye, T. (2013). Quorum sensing inhibitors: how strong is the evidence? *Trends Microbiol.* 21, 619–624. doi: 10.1016/j.tim.2013.09.006
- Déziel, E., Gopalan, S., Tampakaki, A. P., Lépine, F., Padfield, K. E., Saucier, M., et al. (2004). The contribution of MvfR to *Pseudomonas aeruginosa* pathogenesis and quorum sensing circuitry regulation: multiple quorum sensing-regulated genes are modulated without affecting lasRI, rhIRI or the production of N-acyl-L-homoserine lactones. *Mol. Microbiol.* 55, 998–1014. doi: 10.1111/j.1365-2958.2004.04448.x
- Diggle, S. P., Cornelis, P., Williams, P., and Cámara, M. (2006). 4-Quinolone signalling in *Pseudomonas aeruginosa*: old molecules, new perspectives. *Int. J. Med. Microbiol.* 296, 83–91. doi: 10.1016/j.ijmm.2006.01.038
- Diggle, S. P., Matthijs, S., Wright, V. J., Fletcher, M. P., Chhabra, S. R., Lamont, I. L., et al. (2007). The *Pseudomonas aeruginosa* 4-quinolone signal molecules HHQ and PQS play multifunctional roles in quorum sensing and iron entrapment. *Chem. Biol.* 14, 87–96. doi: 10.1016/j.chembiol.2006.11.014
- Diggle, S. P., Winzer, K., Chhabra, S. R., Worrall, K. E., Cámara, M., and Williams, P. (2003). The *Pseudomonas aeruginosa* quinolone signal molecule overcomes the cell density-dependency of the quorum sensing hierarchy, regulates rhl-dependent genes at the onset of stationary phase and can be produced in the absence of LasR. *Mol. Microbiol.* 50, 29–43. doi: 10.1046/j.1365-2958.2003.03672.x
- Downey, D. G., Bell, S. C., and Elborn, J. S. (2008). Neutrophils in cystic fibrosis. *Thorax* 64, 81–88. doi: 10.1136/thx.2007.082388
- Drees, S. L., and Fetzner, S. (2015). PqsE of *Pseudomonas aeruginosa* acts as pathway-specific thioesterase in the biosynthesis of alkylquinolone signaling molecules. *Chem. Biol.* 22, 611–618. doi: 10.1016/j.chembiol.2015.04.012
- Dubern, J.-F., and Diggle, S. P. (2008). Quorum sensing by 2-alkyl-4-quinolones in *Pseudomonas aeruginosa* and other bacterial species. *Mol. Biosyst.* 4, 882–888. doi: 10.1039/b803796p
- Dulcey, C. E., Dekimpe, V., Fauvelle, D.-A., Milot, S., Groleau, M.-C., Doucet, N., et al. (2013). The end of an old hypothesis: the *Pseudomonas* signaling molecules 4-hydroxy-2-alkylquinolines derive from Fatty Acids, not 3-ketofatty acids. *Chem. Biol.* 20, 1481–1491. doi: 10.1016/j.chembiol.2013.09.021
- Fuqua, C., and Greenberg, E. P. (2002). Signalling: listening in on bacteria: acyl-homoserine lactone signalling. *Nat. Rev. Mol. Cell Biol.* 3, 685–695. doi: 10.1038/nrm907
- Fuqua, W. C., Winans, S. C., and Greenberg, E. P. (1994). Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J. Bacteriol.* 176, 269–275.
- García-Contreras, R., Maeda, T., and Wood, T. K. (2013). Resistance to quorum-quenching compounds. *Appl. Environ. Microbiol.* 79, 6840–6846. doi: 10.1128/AEM.02378-13
- García-Contreras, R., Maeda, T., and Wood, T. K. (2015a). Can resistance against quorum-sensing interference be selected? *ISME J.* doi: 10.1038/ismej.2015.84
- García-Contreras, R., Nuñez-López, L., Jasso-Chávez, R., Kwan, B. W., Belmont, J. A., Rangel-Vega, A., et al. (2015b). Quorum sensing enhancement of the stress response promotes resistance to quorum quenching and prevents social cheating. *ISME J.* 9, 115–125. doi: 10.1038/ismej.2014.98
- Gifford, A. M., and Chalmers, J. D. (2014). The role of neutrophils in cystic fibrosis. *Curr. Opin. Hematol.* 21, 16–22. doi: 10.1097/MOH.000000000000009
- Glucksam-Galnoy, Y., Sananes, R., Silberstein, N., Krief, P., Kravchenko, V. V., Meijler, M. M., et al. (2013). The bacterial quorum-sensing signal molecule N-3-oxo-dodecanoyl-L-homoserine lactone reciprocally modulates pro- and anti-inflammatory cytokines in activated macrophages. *J. Immunol.* 191, 337–344. doi: 10.4049/jimmunol.1300368
- Golpasha, I. D., Mousavi, S. F., Owlia, P., Siadat, S. D., and Irani, S. (2015). Immunization with 3-oxododecanoyl-L-homoserine lactone-rPcrV conjugate enhances survival of mice against lethal burn infections caused by *Pseudomonas aeruginosa*. *Bosn. J. Basic Med. Sci.* 15, 15–24. doi: 10.17305/bjbm.2015.292
- Gould, T. A., Schweizer, H. P., and Churchill, M. E. A. (2004). Structure of the *Pseudomonas aeruginosa* acyl-homoserine lactone synthase LasI. *Mol. Microbiol.* 53, 1135–1146. doi: 10.1111/j.1365-2958.2004.04211.x
- Grabner, M. A., Fu, Z., Wu, T., Barry, K. C., Schwarzer, C., and Machen, T. E. (2014). *Pseudomonas aeruginosa* quorum-sensing molecule homoserine lactone

ACKNOWLEDGMENTS

We are grateful for the Open Access Funding from the University of Dundee. This work was also supported by the University of Malaya High Impact Research Grants (UM C/625/1/HIR/MOHE/CHAN/01, A-000001-50001, and UM C/625/1/HIR/MOHE/CHAN/14/1, H-50001-A000027) awarded to K-GC.

- modulates inflammatory signaling through PERK and eIF-2. *J. Immunol.* 193, 1459–1467. doi: 10.4049/jimmunol.1303437
- Guina, T., Purvine, S. O., Yi, E. C., Eng, J., Goodlett, D. R., Aebersold, R., et al. (2003). Quantitative proteomic analysis indicates increased synthesis of a quinolone by *Pseudomonas aeruginosa* isolates from cystic fibrosis airways. *Proc. Natl. Acad. Sci. U.S.A.* 100, 2771–2776. doi: 10.1073/pnas.0435846100
- Hancock, R. E. W., Nijnik, A., and Philpott, D. J. (2012). Modulating immunity as a therapy for bacterial infections. *Nat. Rev. Microbiol.* 10, 243–254. doi: 10.1038/nrmicro2745
- Hänsch, G. M., Prior, B., Brenner-Weiss, G., Obst, U., and Overhage, J. (2014). The *Pseudomonas* quinolone signal (PQS) stimulates chemotaxis of polymorphonuclear neutrophils. *JABFM* 12, 21–26. doi: 10.5301/jabfm.5000204
- Häussler, S., and Becker, T. (2008). The *Pseudomonas* quinolone signal (PQS) balances life and death in *Pseudomonas aeruginosa* populations. *PLoS Pathogens* 4:e1000166. doi: 10.1371/journal.ppat.1000166.s002
- Heeb, S., Fletcher, M. P., Chhabra, S. R., Diggle, S. P., Williams, P., and Cámara, M. (2011). Quinolones: from antibiotics to autoinducers. *FEMS Microbiol. Rev.* 35, 247–274. doi: 10.1111/j.1574-6976.2010.00247.x
- Hentzer, M., Wu, H., Andersen, J. B., Riedel, K., Rasmussen, T. B., Bagge, N., et al. (2003). Attenuation of *Pseudomonas aeruginosa* virulence by quorum sensing inhibitors. *EMBO J.* 22, 3803–3815. doi: 10.1093/emboj/cdg366
- Holban, A.-M., Bleotu, C., Chifiriuc, M. C., Bezirtzoglou, E., and Lazar, V. (2014). Role of *Pseudomonas aeruginosa* quorum sensing (QS) molecules on the viability and cytokine profile of human mesenchymal stem cells. *Virulence* 5, 303–310. doi: 10.4161/viru.27571
- Holm, A., and Vikström, E. (2014). Quorum sensing communication between bacteria and human cells: signals, targets, and functions. *Front. Plant Sci.* 5:309. doi: 10.3389/fpls.2014.00309
- Hooi, D. S. W., Bycroft, B. W., Chhabra, S. R., Williams, P., and Pritchard, D. I. (2004). Differential immune modulatory activity of *Pseudomonas aeruginosa* quorum-sensing signal molecules. *Infect. Immun.* 72, 6463–6470. doi: 10.1128/IAI.72.11.6463-6470.2004
- Jahoor, A., Patel, R., Bryan, A., Do, C., Krier, J., Watters, C., et al. (2008). Peroxisome proliferator-activated receptors mediate host cell proinflammatory responses to *Pseudomonas aeruginosa* autoinducer. *J. Bacteriol.* 190, 4408–4415. doi: 10.1128/JB.01444-07
- Joint, I., Tait, K., Callow, M. E., Callow, J. A., Milton, D., Williams, P., et al. (2002). Cell-to-cell communication across the prokaryote-eukaryote boundary. *Science* 298, 1207. doi: 10.1126/science.1077075
- Kaplan, H. B., and Greenberg, E. P. (1985). Diffusion of autoinducer is involved in regulation of the *Vibrio fischeri* luminescence system. *J. Bacteriol.* 163, 1210–1214.
- Karlsson, T., Turkina, M. V., Yakymenko, O., Magnusson, K.-E., and Vikström, E. (2012). The *Pseudomonas aeruginosa* N-acylhomoserine lactone quorum sensing molecules target IQGAP1 and modulate epithelial cell migration. *PLoS Pathogens* 8:e1002953. doi: 10.1371/journal.ppat.1002953
- Kim, K., Kim, Y. U., Koh, B. H., Hwang, S. S., Kim, S.-H., Lépine, F., et al. (2010). HHQ and PQS, two *Pseudomonas aeruginosa* quorum-sensing molecules, down-regulate the innate immune responses through the nuclear factor- κ B pathway. *Immunology* 129, 578–588. doi: 10.1111/j.1365-2567.2009.03160.x
- Kravchenko, V. V., Kaufmann, G. F., Mathison, J. C., Scott, D. A., Katz, A. Z., Grauer, D. C., et al. (2008). Modulation of gene expression via disruption of NF- κ B signaling by a bacterial small molecule. *Science* 321, 259–263. doi: 10.1126/science.1156499
- Kravchenko, V. V., Kaufmann, G. F., Mathison, J. C., Scott, D. A., Katz, A. Z., Wood, M. R., et al. (2006). N-(3-Oxo-acyl) homoserine lactones signal cell activation through a mechanism distinct from the canonical pathogen-associated molecular pattern recognition receptor pathways. *J. Biol. Chem.* 281, 28822–28830. doi: 10.1074/jbc.M606613200
- LaSarre, B., and Federle, M. J. (2013). Exploiting quorum sensing to confuse bacterial pathogens. *Microbiol. Mol. Biol. Rev.* 77, 73–111. doi: 10.1128/MMBR.00046-12
- Lazenby, J. J., Griffin, P. E., Kyd, J., Whitchurch, C. B., and Cooley, M. A. (2013). A quadruple knockout of lasIR and rhlIR of *Pseudomonas aeruginosa* PAO1 that retains wild-type twitching motility has equivalent infectivity and persistence to PAO1 in a mouse model of lung infection. *PLoS ONE* 8:e60973. doi: 10.1371/journal.pone.0060973
- Lee, J., Wu, J., Deng, Y., Wang, J., Wang, C., Wang, J., et al. (2013). A cell-cell communication signal integrates quorum sensing and stress response. *Nat. Chem. Biol.* 9, 339–343. doi: 10.1038/nchembio.1225
- Lee, R. J., Chen, B., Redding, K. M., Margolskee, R. F., and Cohen, N. A. (2014). Mouse nasal epithelial innate immune responses to *Pseudomonas aeruginosa* quorum-sensing molecules require taste signaling components. *Innate Immun.* 20, 606–617. doi: 10.1177/1753425913503386
- Lee, R. J., Xiong, G., Kofonow, J. M., Chen, B., Lysenko, A., Jiang, P., et al. (2012). T2R38 taste receptor polymorphisms underlie susceptibility to upper respiratory infection. *J. Clin. Invest.* 122, 4145–4159. doi: 10.1172/JCI64240
- Legendre, C., Reen, F. J., Mooij, M. J., McGlacken, G. P., Adams, C., and O’Gara, F. (2012). *Pseudomonas aeruginosa* alkyl quinolones repress hypoxia-inducible factor 1 (HIF-1) signaling through HIF-1 α degradation. *Infect. Immun.* 80, 3985–3992. doi: 10.1128/IAI.00554-12
- Lesic, B., Lépine, F., Déziel, E., Zhang, J., Zhang, Q., Padfield, K., et al. (2007). Inhibitors of pathogen intercellular signals as selective anti-infective compounds. *PLoS Pathogens* 3:e126. doi: 10.1371/journal.ppat.0030126.st003
- Maddocks, S. E., and Oyston, P. C. F. (2008). Structure and function of the LysR-type transcriptional regulator (LTTR) family proteins. *Microbiology* 154, 3609–3623. doi: 10.1099/mic.0.2008/022772-0
- Maurer, S., Wabnitz, G. H., Kahle, N. A., Stegmaier, S., Prior, B., Giese, T., et al. (2015). Tasting *Pseudomonas aeruginosa* biofilms: human neutrophils express the bitter receptor T2R38 as sensor for the quorum sensing molecule N-(3-oxododecanoyl)-L-Homoserine Lactone. *Front. Immunol.* 6:369. doi: 10.3389/fimmu.2015.00369
- Miller, M. B., and Bassler, B. L. (2001). Quorum sensing in bacteria. *Ann. Rev. Microbiol.* 55, 165–199. doi: 10.1146/annurev.micro.55.1.165
- Miyairi, S., Tateda, K., Fuse, E. T., Ueda, C., Saito, H., Takabatake, T., et al. (2006). Immunization with 3-oxododecanoyl-L-homoserine lactone-protein conjugate protects mice from lethal *Pseudomonas aeruginosa* lung infection. *J. Med. Microbiol.* 55, 1381–1387. doi: 10.1099/jmm.0.46658-0
- Nadal Jimenez, P., Koch, G., Thompson, J. A., Xavier, K. B., Cool, R. H., and Quax, W. J. (2012). The multiple signaling systems regulating virulence in *Pseudomonas aeruginosa*. *Microbiol. Mol. Biol. Rev.* 76, 46–65. doi: 10.1128/MMBR.05007-11
- Ortori, C. A., Dubern, J.-F., Chhabra, S. R., Cámara, M., Hardie, K., Williams, P., et al. (2011). Simultaneous quantitative profiling of N-acyl-L-homoserine lactone and 2-alkyl-4(1H)-quinolone families of quorum-sensing signaling molecules using LC-MS/MS. *Anal. Bioanal. Chem.* 399, 839–850. doi: 10.1007/s00216-010-4341-0
- Palmer, K. L., Mashburn, L. M., Singh, P. K., and Whiteley, M. (2005). Cystic fibrosis sputum supports growth and cues key aspects of *Pseudomonas aeruginosa* physiology. *J. Bacteriol.* 187, 5267–5277. doi: 10.1128/JB.187.15.5267-5277.2005
- Parsek, M. R., Val, D. L., Hanzelka, B. L., Cronan, J. E., and Greenberg, E. P. (1999). Acyl homoserine-lactone quorum-sensing signal generation. *Proc. Natl. Acad. Sci. U.S.A.* 96, 4360–4365. doi: 10.1073/pnas.96.8.4360
- Pesci, E. C., Milbank, J. B., Pearson, J. P., McKnight, S., Kende, A. S., Greenberg, E. P., et al. (1999). Quinolone signaling in the cell-to-cell communication system of *Pseudomonas aeruginosa*. *Proc. Natl. Acad. Sci. U.S.A.* 96, 11229–11234. doi: 10.1073/pnas.96.20.11229
- Rampioni, G., Pustelny, C., Fletcher, M. P., Wright, V. J., Bruce, M., Rumbaugh, K. P., et al. (2010). Transcriptomic analysis reveals a global alkyl-quinolone-independent regulatory role for PqsE in facilitating the environmental adaptation of *Pseudomonas aeruginosa* to plant and animal hosts. *Environ. Microbiol.* 12, 1659–1673. doi: 10.1111/j.1462-2920.2010.02214.x
- Ritchie, A. J., Jansson, A., Stallberg, J., Nilsson, P., Lysaght, P., and Cooley, M. A. (2005). The *Pseudomonas aeruginosa* quorum-sensing molecule N-3-(oxododecanoyl)-L-homoserine lactone inhibits T-cell differentiation and cytokine production by a mechanism involving an early step in T-cell activation. *Infect. Immun.* 73, 1648–1655. doi: 10.1128/IAI.73.3.1648-1655.2005
- Schwarzer, C., Fu, Z., Morita, T., Whitt, A. G., Neely, A. M., Li, C., et al. (2015). Paraoxonase 2 serves a proapoptotic function in mouse and human cells in response to the *Pseudomonas aeruginosa* quorum-sensing molecule N-(3-oxododecanoyl)-homoserine lactone. *J. Biol. Chem.* 290, 7247–7258. doi: 10.1074/jbc.M114.620039

- Schwarzer, C., Fu, Z., Patanwala, M., Hum, L., Lopez-Guzman, M., Illek, B., et al. (2012). *Pseudomonas aeruginosa* biofilm-associated homoserine lactone C12 rapidly activates apoptosis in airway epithelia. *Cell. Microbiol.* 14, 698–709. doi: 10.1111/j.1462-5822.2012.01753.x
- Schwarzer, C., Wong, S., Shi, J., Matthes, E., Illek, B., Ianowski, J. P., et al. (2010). *Pseudomonas aeruginosa* homoserine lactone activates store-operated cAMP and cystic fibrosis transmembrane regulator-dependent Cl⁻ secretion by human airway epithelia. *J. Biol. Chem.* 285, 34850–34863. doi: 10.1074/jbc.M110.167668
- Shiner, E. K., Terentyev, D., Bryan, A., Sennoune, S., Martinez-Zaguilan, R., Li, G., et al. (2006). *Pseudomonas aeruginosa* autoinducer modulates host cell responses through calcium signalling. *Cell. Microbiol.* 8, 1601–1610. doi: 10.1111/j.1462-5822.2006.00734.x
- Skindersoe, M. E., Zeuthen, L. H., Brix, S., Fink, L. N., Lazenby, J., Whittall, C., et al. (2009). *Pseudomonas aeruginosa* quorum-sensing signal molecules interfere with dendritic cell-induced T-cell proliferation. *FEMS Immun. Med. Microbiol.* 55, 335–345. doi: 10.1111/j.1574-695X.2008.00533.x
- Smith, R. S., Fedyk, E. R., Springer, T. A., Mukaida, N., Iglewski, B. H., and Phipps, R. P. (2001). IL-8 production in human lung fibroblasts and epithelial cells activated by the *Pseudomonas* autoinducer N-3-oxododecanoyl homoserine lactone is transcriptionally regulated by NF-kappa B and activator protein-2. *J. Immunol.* 167, 366–374. doi: 10.4049/jimmunol.167.1.366
- Starkey, M., Lépine, F., Maura, D., Bandyopadhyaya, A., Lesic, B., He, J., et al. (2014). Identification of anti-virulence compounds that disrupt quorum-sensing regulated acute and persistent pathogenicity. *PLoS Pathogens* 10:e1004321. doi: 10.1371/journal.ppat.1004321.s009
- Tateda, K., Ishii, Y., Horikawa, M., Matsumoto, T., Miyairi, S., Pechere, J. C., et al. (2003). The *Pseudomonas aeruginosa* autoinducer N-3-oxododecanoyl homoserine lactone accelerates apoptosis in macrophages and neutrophils. *Infect. Immun.* 71, 5785–5793. doi: 10.1128/IAI.71.10.5785-5793.2003
- Twigg, M. S., Tait, K., Williams, P., Atkinson, S., and Cámara, M. (2013). Interference with the germination and growth of *Ulva* zoospores by quorum-sensing molecules from *Ulva*-associated epiphytic bacteria. *Environ. Microbiol.* 16, 445–453. doi: 10.1111/1462-2920.12203
- Urbanczyk, H., Ast, J. C., Higgins, M. J., Carson, J., and Dunlap, P. V. (2007). Reclassification of *Vibrio fischeri*, *Vibrio logei*, *Vibrio salmonicida* and *Vibrio wodanis* as *Aliivibrio fischeri* gen. nov., comb. nov., *Aliivibrio logei* comb. nov., *Aliivibrio salmonicida* comb. nov. and *Aliivibrio wodanis* comb. nov. *Int. J. Syst. Evol. Microbiol.* 57, 2823–2829. doi: 10.1099/ijs.0.65081-0
- Valentine, C. D., Anderson, M. O., Papa, F. R., and Haggie, P. M. (2013). X-box binding protein 1 (XBP1s) is a critical determinant of *Pseudomonas aeruginosa* homoserine lactone-mediated apoptosis. *PLoS Pathog.* 9:e1003576. doi: 10.1371/journal.ppat.1003576
- Valentine, C. D., Zhang, H., Phuan, P.-W., Nguyen, J., Verkman, A. S., and Haggie, P. M. (2014). Small molecule screen yields inhibitors of *Pseudomonas* homoserine lactone-induced host responses. *Cell. Microbiol.* 16, 1–14. doi: 10.1111/cmi.12176
- Vikström, E., Bui, L., Konradsson, P., and Magnusson, K.-E. (2009). The junctional integrity of epithelial cells is modulated by *Pseudomonas aeruginosa* quorum sensing molecule through phosphorylation-dependent mechanisms. *Exp. Cell Res.* 315, 313–326. doi: 10.1016/j.yexcr.2008.10.044
- Vikström, E., Bui, L., Konradsson, P., and Magnusson, K.-E. (2010). Role of calcium signalling and phosphorylations in disruption of the epithelial junctions by *Pseudomonas aeruginosa* quorum sensing molecule. *Eur. J. Cell Biol.* 89, 584–597. doi: 10.1016/j.ejcb.2010.03.002
- Wagner, C., Zimmermann, S., Brenner-Weiss, G., Hug, F., Prior, B., Obst, U., et al. (2007). The quorum-sensing molecule N-3-oxododecanoyl homoserine lactone (3OC12-HSL) enhances the host defence by activating human polymorphonuclear neutrophils (PMN). *Anal. Bioanal. Chem.* 387, 481–487. doi: 10.1007/s00216-006-0698-5
- Watson, W. T., Minogue, T. D., Val, D. L., von Bodman, S. B., and Churchill, M. E. A. (2002). Structural basis and specificity of acyl-homoserine lactone signal production in bacterial quorum sensing. *Mol. Cell* 9, 685–694. doi: 10.1016/S1097-2765(02)00480-X
- Williams, P., and Cámara, M. (2009). Quorum sensing and environmental adaptation in *Pseudomonas aeruginosa*: a tale of regulatory networks and multifunctional signal molecules. *Curr. Opin. Microbiol.* 12, 182–191. doi: 10.1016/j.mib.2009.01.005
- Winsor, G. L., Lam, D. K. W., Fleming, L., Lo, R., Whiteside, M. D., Yu, N. Y., et al. (2011). *Pseudomonas* genome database: improved comparative analysis and population genomics capability for *Pseudomonas* genomes. *Nucleic Acids Res.* 39, D596–D600. doi: 10.1093/nar/gkq869
- Winstanley, C., and Fothergill, J. L. (2009). The role of quorum sensing in chronic cystic fibrosis *Pseudomonas aeruginosa* infections. *FEMS Microbiol. Lett.* 290, 1–9. doi: 10.1111/j.1574-6968.2008.01394.x
- Xiao, G., He, J., and Rahme, L. G. (2006). Mutation analysis of the *Pseudomonas aeruginosa* mvfR and pqsABCDE gene promoters demonstrates complex quorum-sensing circuitry. *Microbiology* 152, 1679–1686. doi: 10.1099/mic.0.28605-0
- Zimmermann, S., Wagner, C., Muller, W., Brenner-Weiss, G., Hug, F., Prior, B., et al. (2006). Induction of neutrophil chemotaxis by the quorum-sensing molecule N-(3-oxododecanoyl)-L-homoserine lactone. *Infect. Immun.* 74, 5687–5692. doi: 10.1128/IAI.01940-05

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2015 Liu, Chan and Chang. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) or licensor are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.