



The University of Bradford Institutional Repository

<http://bradscholars.brad.ac.uk>

This work is made available online in accordance with publisher policies. Please refer to the repository record for this item and our Policy Document available from the repository home page for further information.

To see the final version of this work please visit the publisher's website. Access to the published online version may require a subscription.

Link to publisher's version: <http://dx.doi.org/10.1128/genomeA.00100-15>

Citation: Chan K, Tan W, Chang C et al (2015) Genome sequence analysis reveals evidence of quorum-sensing genes present in *Aeromonas hydrophila* strain M062, isolated from freshwater. *Genome Announcements*. 3(2): e00100-15.

Copyright statement: © 2015 Chan et al. This is an open-access article distributed under the terms of the Creative Commons Attribution 3.0 Unported license.

Genome Sequence Analysis Reveals Evidence of Quorum-Sensing Genes Present in *Aeromonas hydrophila* Strain M062, Isolated from Freshwater

Kok-Gan Chan,^a Wen-Si Tan,^a Chien-Yi Chang,^{b,c} Wai-Fong Yin,^a Nina Yusrina Mumahad Yunos^a

Division of Genetics and Molecular Biology, Institute of Biological Sciences, Faculty of Science, University of Malaya, Kuala Lumpur, Malaysia^a; Interdisciplinary Computing and Complex BioSystems (ICOS) Research Group, School of Computing Science, Newcastle University, Newcastle upon Tyne, United Kingdom^b; The Centre for Bacterial Cell Biology, Medical School, Newcastle University, Newcastle upon Tyne, United Kingdom^c

***Aeromonas hydrophila* has emerged worldwide as a human pathogen. Here, we report the draft whole-genome sequence of a freshwater isolate from Malaysia, *A. hydrophila* strain M062, and its *N*-acylhomoserine lactone genes are also reported here.**

Received 26 January 2015 Accepted 2 February 2015 Published 12 March 2015

Citation Chan K-G, Tan W-S, Chang C-Y, Yin W-F, Mumahad Yunos NY. 2015. Genome sequence analysis reveals evidence of quorum-sensing genes present in *Aeromonas hydrophila* strain M062, isolated from freshwater. *Genome Announc* 3(2):e00100-15. doi:10.1128/genomeA.00100-15.

Copyright © 2015 Chan et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](#).

Address correspondence to Kok-Gan Chan, kokgan@um.edu.my.

Quorum sensing (QS) is a term coined by Fuqua and coworkers to define the phenomenon of a bacterial cell-to-cell communication system that controls bacterial physiological processes in a population density-dependent manner (1). This process describes the event whereby an increase in the population density of the bacterial cell is proportional to an increase in the concentration of signal molecule(s) in the extracellular environment (2, 3). *Aeromonas hydrophila* is QS bacterium and is a known pathogen, as it can cause minor skin infections and gastroenteritis in humans (4, 5). In this study, we report the whole-genome sequence and the QS genes of *A. hydrophila* strain M062.

A. hydrophila M062 was isolated from the Sungai Tua waterfall, Selangor, Malaysia. The genomic DNA of *A. hydrophila* M062 was extracted using the MasterPure DNA purification kit (Epicentre, Inc., Madison, WI, USA). A Qubit 2.0 fluorometer (Invitrogen, USA) and NanoDrop spectrophotometer (Thermo Scientific, USA) were used to quantify and qualify the DNA preceding whole-genome shotgun sequencing on an Illumina MiSeq personal sequencer (Illumina, Inc., San Diego, CA, USA) and generated 4,984,734 paired-end reads. Subsequently, the paired-end reads were trimmed and *de novo* assembled with CLC Genomics Workbench version 5.1 (CLC bio, Denmark). Gene prediction was conducted using Prodigal (version 2.60) (6), followed by gene annotation using RAST (7). Next, tRNAs and rRNAs were predicted with tRNAscan-SE version 1.21 (8) and RNAmmer (9), respectively.

From this whole-genome sequencing, a total of 4,974,350 reads were generated. After trimming (1,548,021 quality reads), the reads were assembled into 140 contigs, with an N_{50} of 93,561 and an average coverage of 51.4-fold. The G+C content of the draft genome of *A. hydrophila* strain M062 is 60.5%. An analysis of its genome sequences led to the identification of 4,392 coding DNA sequences (CDSs), 95 tRNAs, two copies of 5S rRNA, and one copy each of 16S rRNA and 23S rRNA.

The annotation result showed that the *luxI* and *luxR* homo-

logues of *A. hydrophila* strain M062 are located at contig 35. With the availability of the whole-genome sequence of *A. hydrophila* strain M062, it may provide a better understanding of the QS system in this strain.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited at DDBJ/EMBL/GenBank under the accession no. [JSXE00000000](#). The version described in this paper is the first version, JSXE01000000.

ACKNOWLEDGMENTS

This work was supported by the University of Malaya via High-Impact Research Grants (UM C/625/1/HIR/MOHE/CHAN/01, no. A-000001-50001), and a UM-MOHE HIR grant (UM C/625/1/HIR/MOHE/CHAN/14/1, no. H-50001-A000027), which were awarded to Kok-Gan Chan.

REFERENCES

1. Fuqua WC, Winans SC, Greenberg EP. 1994. Quorum sensing in bacteria: the LuxR-LuxI family of cell density-responsive transcriptional regulators. *J Bacteriol* 176:269–275.
2. Atkinson S, Williams P. 2009. Quorum sensing and social networking in the microbial world. *J R Soc Interface* 6:959–978. <http://dx.doi.org/10.1098/rsif.2009.0203>.
3. Yunos NY, Tan WS, Koh CL, Sam CK, Mohamad NI, Tan PW, Adrian TG, Yin WF, Chan KG. 2014. *Pseudomonas cremoricolorata* strain nd07 produces *N*-acyl homoserine lactones as quorum sensing molecules. *Sensors (Basel)* 14:11595–11604. <http://dx.doi.org/10.3390/s140711595>.
4. Garde C, Bjarnsholt T, Givskov M, Jakobsen TH, Hentzer M, Clausen A, Sniepen K, Ferkinghoff-Borg J, Sams T. 2010. Quorum sensing regulation in *Aeromonas hydrophila*. *J Mol Biol* 396:849–857. <http://dx.doi.org/10.1016/j.jmb.2010.01.002>.
5. Chan XY, Chua KH, Yin WF, Puthuchery SD, Chan KG. 2014. Whole-genome analysis of *Aeromonas hydrophila* strain 187, exhibiting quorum-sensing activity. *Genome Announc* 2(6):e01360-14. <http://dx.doi.org/10.1128/genomeA.01360-14>.
6. Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
7. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formisano

- K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
8. Lowe TM, Eddy SR. 1997. tRNAscan-SE: a program for improved detection of transfer RNA genes in genomic sequence. *Nucleic Acids Res* 25: 955–964. <http://dx.doi.org/10.1093/nar/25.5.0955>.
9. Lagesen K, Hallin P, Rødland EA, Staerfeldt HH, Rognes T, Ussery DW. 2007. RNAmmer: consistent and rapid annotation of ribosomal RNA genes. *Nucleic Acids Res* 35:3100–3108. <http://dx.doi.org/10.1093/nar/gkm160>.