

CIAT Research Online - Accepted Manuscript

Effect of feeding oat and vetch forages on milk production and quality in smallholder dairy farms in Central Kenya

The International Center for Tropical Agriculture (CIAT) believes that open access contributes to its mission of reducing hunger and poverty, and improving human nutrition in the tropics through research aimed at increasing the eco-efficiency of agriculture.

CIAT is committed to creating and sharing knowledge and information openly and globally. We do this through collaborative research as well as through the open sharing of our data, tools, and publications.

Citation:

Mwendia, Solomon W.; Mwangu, Chris M.; Ng'ang'a, Stanley Karanja; Njenga, David; Notenbaert, An. 2018. Effect of feeding oat and vetch forages on milk production and quality in smallholder dairy farms in Central Kenya. *Tropical Animal Health and Production* 1-7 p.

Publisher's DOI:

<https://doi.org/10.1007/s11250-018-1529-3>

Access through CIAT Research Online:

<http://hdl.handle.net/10568/90961>

Terms:

© 2018. CIAT has provided you with this accepted manuscript in line with CIAT's open access policy and in accordance with the Publisher's policy on self-archiving.



This work is licensed under a [Creative Commons Attribution-NonCommercial 4.0 International License](https://creativecommons.org/licenses/by-nc/4.0/). You may re-use or share this manuscript as long as you acknowledge the authors by citing the version of the record listed above. You may not use this manuscript for commercial purposes.

For more information, please contact CIAT Library at CIAT-Library@cgiar.org.

Tropical Animal Health and Production

Effect of feeding oat and vetch forages on milk production and quality in smallholder dairy farms in Central Kenya

--Manuscript Draft--

Manuscript Number:	TROP-D-17-00871R2	
Full Title:	Effect of feeding oat and vetch forages on milk production and quality in smallholder dairy farms in Central Kenya	
Article Type:	Regular Articles	
Corresponding Author:	Solomon Mwendia Centro Internacional de Agricultura Tropical Nairobi, KENYA	
Corresponding Author Secondary Information:		
Corresponding Author's Institution:	Centro Internacional de Agricultura Tropical	
Corresponding Author's Secondary Institution:		
First Author:	Solomon Mwendia	
First Author Secondary Information:		
Order of Authors:	Solomon Mwendia	
	Chris M. Mwangu, MSc	
	Stanley Nga'ng'a Karanja, PhD	
	David Njenga, MSc	
	An Notenbaert, MSc	
Order of Authors Secondary Information:		
Funding Information:	IFDC (2Scale project)	Not applicable

COMMENTS FOR THE AUTHOR:

Just about ready, I think.

Submit the next version, without highlighting individual changes in the text.

A couple of very minor points:

Line 24

Low livestock productivity happens in many smallholder farms in Africa, attributable to inadequate fodder of good quality (Manaye et al., 2009).

Better as

The low levels of livestock productivity in many smallholder farms in Africa, are largely attributable to inadequate fodder of good quality (Manaye et al., 2009).

Line 48

036° 24' E

36° 24' E

Comment	Revision
<p>Line 24 Low livestock productivity happens in many smallholder farms in Africa, attributable to inadequate fodder of good quality (Manaye et al., 2009).</p>	<p>The low levels of livestock productivity in many smallholder farms in Africa, are largely attributable to inadequate fodder of good quality (Manaye et al., 2009).</p>
<p>Line 48 036° 24' E</p>	<p>36° 24' E</p>

1 **Effect of feeding oat and vetch forages on milk production and quality in smallholder dairy**
2 **farms in Central Kenya**

3 *Solomon W. Mwendia*^{1,4} (<http://orcid.org/0000-0002-3203-8770>), *Chris M. Mwangi*¹, *Ng'ang'a,*
4 *Stanley Karanja*³ (<https://orcid.org/0000-0002-6166-7920>), *David Njenga*², and *An Notenbaert*¹

5 ¹ *International Center for Tropical Agriculture (CIAT), P.O. Box 823-00621 Nairobi, Kenya*

6 ² *International Fertilizer Development Center (IFDC), P.O. Box 30772-00100, Nairobi, Kenya*

7 ³ *International Center for Tropical Agriculture (CIAT), Pan - Africa Bean Research Alliance*

8 *National Agricultural Research Laboratories- Kawanda P.O. Box 6247, Kampala, Uganda*

9 ⁴Corresponding author: mwendia2007@gmail.com

10

11 ***Abstract***

12 Despite the significant livestock contribution to households' nutrition and incomes in many African smallholder farms,
13 milk productivity remains low. Inadequate feeding is the main reason for the underperformance. To contribute towards
14 addressing this, an on-farm feeding trial was undertaken in Ol-joro-Orok Central Kenya. A feed basket using oat
15 (*Avena sativa*) cv Conway and vetch (*Vicia villosa*) was compared to farmers practice. Milk production (kg) and
16 quality parameters, including butterfat, protein, lactose, and density, were monitored, and cost-benefit analysis (CBA)
17 undertaken. Feeding both oat and vetch increased milk production by 21% (morning) and 18%, (evening), equivalent
18 to 1.4 kg/day. Increases (%) in quality were; butter fat (18.2), solid-non-fat (16.5), lactose (16.2) and protein (16.1).
19 Concomitantly, the CBA returned positive results, supporting the hypothesis of economic advantage in using oat and
20 vetch in milk production in the area, and possibly in other similar areas.

21 Keywords: Forage, milk production, cost-benefit

22

23 **Introduction**

24 The low levels of livestock productivity in many smallholder farms in Africa, are largely attributable to inadequate
25 fodder of good quality (Manaye et al., 2009). The increasing human population in Sub-Saharan Africa (SSA), coupled
26 with urbanization and expansion of middle class with disposable income (Cohen, 2006), contribute to a projected
27 increase in demand for livestock products, predisposing dire need to increase livestock productivity. If productivity
28 does not rise at the same rate or more to keep up with the demand, a food crisis is likely, and this may exacerbate the
29 situation of poor human nutrition, already a major concern in SSA (FAO 2017).

30 Amongst constraints, inadequate quantity and quality feed is the main limiting factor to dairy improvement in African,
31 yet improved forages can support and enhance livestock productivity (Yami et al., 2013). Forage cultivation is still
32 low with preference accorded to food crops, whose residues are often used as livestock feed despite their low feeding
33 value (Methu et al., 2001), resulting in poor performance especially milk production. To improve lactation yields,
34 cows require access to enough nutrients and clean water (Lukuyu et al., 2012). In smallholder dairy in eastern Africa,
35 feeding accounts for 55-70% of the costs (Odero-Waitituh 2017). As such, using technologies that can increase milk
36 yields and lower production cost would be preferable. Recent evidence shows it is possible to lower cost of milk
37 production by 4.4% without decreasing the output (Odero-Waitituh, 2017).

38 Most studies on cost-benefit analysis (CBA) of milk are rarely technology specific e.g. Mburu et al., (2007). In
39 addition, important indicators of CBA e.g. Net Present Value (NPV), Internal Rate of Return (IRR) and Pay Back
40 Period (PBP) are not estimated. Therefore, the work entailed here set out to evaluate (a) whether the use of oat and
41 vetch forages has any impact on milk production and quality compared to farmers' practice, and (b) if an economically
42 positive and sound outcome would be realized when feeding oat and vetch.

43 **Materials and methods**

44 *Area of study*

45 The study was done in Ol-joro-Orok sub-county, in central Kenya. The area lies between 0° 09' S and 36° 24' E
46 covering an area of about 359 km² and, about 2359 meters above sea level. The average minimum and maximum
47 temperature for the last 24 years ranges 5-8 °C and 20-23 °C respectively, while annual rainfall over the same period
48 averages 817–977mm (Jaetzold et al., 2006).

49 *Production of oat and vetch for trial*

50 The study is linked to Eldoville Dairies in Ol-joro-Orok with an existing smallholder dairy farmers' network, who
51 supply milk. Interest in increased milk in this area has increased, and in the attempt to close the supply gap Eldoville
52 Dairies processing factory with a daily capacity of about 70,000 litres provided 1.5 acres of land for planting oats and
53 vetch.

54 We used oat (*Avena sativa*) cv Conway and *Vicia villosa* cv purple vetch. Farmers had previously selected Conway
55 as the best bet (Mwendia et al. 2017). Conway seeds from Aberystwyth UK and vetch seeds from Kenya Agricultural
56 and Livestock Research Organization- Ol-joro-Orok) were used. The land was ploughed and harrowed in September
57 2016. An acre was established with Conway and half an acre with vetch on 9th October 2016. Oat was planted in
58 furrows spaced at 15cm apart at 100 kg/ha seed rate, while vetch was in 30 cm apart furrows, at 20kg/ha seed rate. At
59 planting, inorganic NPK fertiliser, 23:23:0, was applied at 50 kg Nha⁻¹ for oat while none was applied for vetch. After
60 establishment, vetch was weeded manually as necessary while oat field was sprayed with broadleaf herbicide –
61 Bellamine 72%.

62 Vetch was harvested at flowering stage and dried under shade, producing 308 kg of hay (equivalent to 1.5 t DM /ha).
63 Oat production was estimated by measuring harvest from three randomly selected 2 m² plots, with a mean of 3.37 kg
64 (fresh) equivalent to 6700 kg ~16.7 t/ha. This translated to about 2.18 t DM/ha at 13% DM content (Mwendia et al.,
65 2017).

66 *Farmer selection and roles*

67 Ten dairy farmers were selected on the criteria i.e. sell milk to Eldoville Dairies; own a cow in early to mid-lactation,
68 and at 2nd or 3rd parity, and willingness to cooperate with data collection from their cows. In a meeting at Eldoville,
69 the feeding trial was explained and roles shared. While farmers were to provide lactating cows and allow data
70 collection, Eldoville Dairies was to do milk analysis and coordinate issuance of test forages to the farmers. Researchers
71 were to provide forage for the trial and collect data.

72 *Feeding protocol and data collection*

73 A Local agricultural extension officer was assigned to collect data from the selected 10 farms on daily basis. Starting
74 4th January 2017, data was collected on milk production and quality for 2 weeks. Amounts of morning and evening
75 milk produced (kg) were recorded. Upon milk delivery to the Eldoville Dairies, a milk sample of ~ 50ml was collected

76 daily for quality analysis (described later). Further, type of feeds fed to the cows was quantified where possible with
77 a spring balance and recorded.

78 After the 2 weeks, feeding was switched to oat and vetch for selected 10 cows and, for 10 consecutive days. A daily
79 ration of 60 kg of wilted fodder oat (7.8 kg DM) and 2kg of vetch hay (1.7 kg DM) thus 9.5 kg DM/day was fed.
80 Where under farmer practice the cows were grazed or supplemented, the type and quantity were maintained under the
81 intervention. As such, the difference was the change of the basal roughage (oat-vetch vs farmer practice). Throughout
82 the trial, cows had access to clean drinking water adlib. Comparison between farmers' practice to oat-vetch was within
83 the animals (University of Reading, 2000) and not between animals. After 10 days of intervention feeding, the farmers
84 resorted to farmer practice, which was further trailed for 2 weeks. However, two farmers who were not cooperative
85 were dropped.

86 *Milk quality analysis*

87 Milk quality was daily analyzed with a Milk Lactoscan (SL Ultrasonic Milk Analyzer, Tamil Nadu, India) throughout
88 the trial. Measurements included butterfat, solids-non-fat (SNF), density, lactose and protein levels.

89 *CBA data and variables measurement*

90 The CBA data was collected from the 8 farmers who fully participated in the feeding trial. Structured questionnaires
91 through face to face interviews were administered, four months after the experiment (April 2017). Details on expenses
92 including; labour, feed, veterinary, maintenance, and milk income during the feeding trials were collected. For the
93 CBA indicators, we treated the cost of feed additives, water, veterinary services and commercial feeds as constant.
94 Fixed costs such as depreciation in the value of the milking cows, shed, machinery and interest cost on capital were
95 ignored. Total milk produced including milk fed to calves and consumed at home was valued at the market price of
96 0.37 USD/litre, the price at which Eldoville pay farmers.

97 *Analytical Model*

98 We adopted cost-benefit approach that comprises of net present value (NPV), internal rate of return (IRR) and payback
99 period (PBP) (Kimenju & De Groote, 2010). An intervention is economically viable if the payback is less than the
100 time taken to recover the initially invested amount. In this study, we calculated payback period using cash flow
101 amounts based on non-discounted dollar amounts.

102 NPV is the difference between present value of cash inflows and the present value of cash outflows. It is determined
103 by applying a discount rate to the identified costs and benefits. With investments, one decides what to do with the
104 money today. An investment is viable if NPV is greater than zero. Benefits flow for farmers after adopting an
105 intervention, over initial practice can be given by:

$$106 \quad NPV = \sum_0^t \frac{(B_t - C_t)}{(1+r)^t}$$

107 Where

108 B_t - Benefits at time t

109 C_t - cost at time t

110 r - discount rate

111 To compute IRR using the formulae, NPV is set to zero and solve for r -discount rate = IRR. Investment is viable if
112 IRR is positive and greater than the market discount rate.

113 *Data analyses*

114 All data were managed in Microsoft Excel spreadsheets. Standard errors were calculated as σ/\sqrt{n} and plots are done
115 in Excel. Where applicable, analysis of variance was done in GenStat (2011) software and means separated by least
116 significant (LSD). CBA data was managed in STATA version 14.1. Measures of central tendency (mean, median and
117 mode) and dispersion (range and standard deviation) were computed using macros in excel 2013. PBP, NPV and IRR
118 were calculated in an online excel based tool (www.cbatool.ciat.cgiar.org) developed by the International Center for
119 Tropical Agriculture (CIAT) to assess the economic viability of climate-smart agricultural (CSA) technologies.

120 **Results**

121 *Farmers' practice*

122 Among the study farms, feeding was on Napier grass and bought hay (~2 USD/bale), and to a lesser extent, fodder oat
123 and sorghum. Maize stovers, bean haulms and weeds were also included. In addition, animals spent at least 3 h/day
124 grazing. There was no estimation of the feed intake from grazing, but fields were overgrazed suggesting minimal
125 benefit. Supplementation with dairy a meal and minerals was adopted in all the farms. While farmer practice fed dry

126 matter ranging 3.3—19.6/day, in some cases higher than the intervention (9.5 DM/day), all the animals increased milk
127 production under intervention. Table 1 summarizes quantities (kg) of feeds offered to the animals.

128 “Insert table 1”

129 *Milk yields and quality*

130 Morning and evening milk production (Table 2), increased by 21 and 18% respectively, under the intervention. The
131 difference between milk production under intervention and farmer practice, divided by milk production under farmer
132 practice multiplied by 100 constituted the percentages. More milk was produced in the morning than in the evenings,
133 with an average increase of ~1.4 kg/day.

134 Over the 42-day trial period, pooled milked yields across the 8 farms, separately for morning and evening production
135 increased steadily (Figure 1) to a peak at day-22, after which there was a drop especially after reverting to farmer
136 practice at day-25. The drop continued steadily to the end of the trial, day-42. At no time did the evening production
137 surpass the morning production, however, the trend was similar.

138 “insert figure 1”

139 Figure 2 summarizes effect on butter fat, lactose, solid-non-fat, density, protein. Except for butterfat, increases in milk
140 when fed on oat-vetch compared to farmers’ practice were not significant. However, were highly significant for butter
141 fat, lactose, solid-non-fat and protein when computed on the net increase in daily milk production (Table 2). Absolute
142 percentage (%), increases in the weights were in the order; butterfat (18.2), protein (16.1), lactose (16.2) and solid-
143 non-fat (16.5) Table 2.

144 “Insert figure 2”

145 “Insert table 2”

146 *Fodder production costs*

147 Table 3 shows the cost of production for the common feeds in the farms. The costs for all the inputs were based on
148 farmer’s land size and scaled accordingly to one acre. Largely inputs include; fertilizer/manure, seeds, pesticides,
149 herbicides, and labor. Vetch had the highest cost of production and Conway-oat the lowest, per acre. On production,

150 Napier grass yields more per acre. Costs of producing maize stovers, other crop residues and weeds were not available,
151 but farmers assigned values assuming buying or selling them from an acre.

152 “Table 3”

153 The CBA indicators revealed that it is economically viable to adopt oat and vetch. The benefit of adopting oat and
154 vetch over 4 months would generate an NPV of \$22. The money invested through inputs and labor is recovered after
155 65 days while IRR of 15% indicates that money invested in producing forage crops will increase by 15%.

156 “Insert table 4”

157 **Discussion**

158 While the focus of the study was to compare milk quality and production under farmers’ practice with improved
159 feeding, understanding what constituted the farmers’ practice was also important. Some of the feeds used are of poor
160 quality. In particular, maize stovers contain <3% crude protein, in addition to poor digestibility (Methu et al., 2001)
161 compared to 13–18% considered appropriate (Lukuyu et al., 2012). Napier grass is neither of superior quality with
162 crude protein ranging 8 –10% (Tessema et al., 2010). However, maize stovers and Napier grass are used substantially
163 in the study area to support livestock production and may be contributing to the low productivity. Although farmers
164 grazed animals (Table 1), visually, the overgrazed paddocks suggested minimal nutritional animal benefit. Poor
165 feeding limits the production potential and negates gains in livestock breed improvement. As such, most likely the
166 farmers in the study were not fully exploiting their cross-bred animals.

167 Feeding oat and vetch compared to the farmers practice increased milk production and quality (Table 2). Although
168 improved milk production is influenced by several aspects, including animal breeding, health, and feeding; feeding
169 has immediate results as shown here. If a large number of farmers embraced improved feeding, increasing milk
170 production is possible, and contribute to addressing present and future demands. Studies about feeding rarely look at
171 the economic potential. However, positive economic benefits in this study, are likely to provide the impetus for
172 adoption. Rising demand for animal products linked to population growth is likely to provide market-pull that could
173 favorably catalyze adoption (Kimenye & McEwan, 2014).

174

175 Milk is a raw material for other products e.g. butter, whey, and cheese which require high-quality milk. To increase
176 butter and cheese at processing, high milk butter fat and protein contents respectively, are paramount (Rønholt et al.,
177 2013; Wedholm et al. 2006). Elsewhere raw milk prices are pegged on the milk quality (Jesse and Cropp 2004), and
178 likely to attract improved feeding. Eldoville Dairies is involved in butter and cheese production, and envisage paying
179 milk based on quality (A. Waithaka Pers. Comm.) to encourage dairy keepers improve feeding.

180 To realize adoption at a scale of such promising forage technologies, awareness creation and functioning forage seed
181 systems need to happen concurrently. Lack of forage seeds is a major bottleneck curtailing adoption of improved
182 forage technologies (Negassa et al., 2016). In Kenya, forage seed system is limited, with few options from the private
183 sector (Mwendia et al., 2016). Governments should facilitate certification of proven forage technologies without
184 lengthy institutional requirements.

185 In conclusion, our results show feeding improved forages has the potential to increase milk production and quality.
186 Unimproved feeding suggests that farmers are not exploiting the productivity potential of their animals. However,
187 economic benefit would most likely drive farmers to improve feeding. Awareness creation is vital coupled with
188 strengthening the forage seed systems.

189 **Acknowledgements**

190 We sincerely thank the farmers who allowed data collection from their animals and responded to interviews. Gratitude
191 to Mr. B.M. Kimani from Ministry of Agriculture who assisted with daily data collection, Eldoville Dairies, especially
192 Mr. Andrew Waithaka who facilitated oat and vetch growing, Mr. Mutembei who supervised the forage fields and
193 Mr. Julius Njuguna who measured the milk quality. Lastly, we thank the International Fertilizer Development Center
194 (IFDC) for financing the study.

195 **Compliance with ethical standards**

196 Farmers gave informed consent to data collection from their farms and animals.

197 **Conflict of interest**

198 None

199 **References**

200 References

201 Ayoade, J. A. ; Makhambera, P. E. ; Bodzalekani, M. Z., 1983. Evaluation of crop residues as
202 feeds for goats. Part 1. Voluntary intakes, digestibility and nitrogen utilization of groundnut
203 and bean haulms. South Afr. J. Anim. Sci., 13 (1): 12-13

204 Cohen B. 2006. Urbanization in developing countries: Current trends, future projections, and key
205 challenges for sustainability Technology in Society 28, 63–80
206 doi:10.1016/j.techsoc.2005.10.005

207 Food and Agriculture Organization (FAO) 2017. The future of food and agriculture- Trends and
208 challenges. Rome.

209 Genstat 2011. GenStat statistical software, version 14 for windows. VSN International
210 Ltd, Hertfordshire, UK.

211 Gietema B. 2005. Modern dairy farming in warm climate zones. © Agromisa Foundation,
212 Wageningen, ISBN: 90 5285 016 1/016 X

213 International livestock research institute (ILRI) 2016. Feed assessment tool.
214 <https://www.ilri.org/feast>

215 Jaetzold, R., Schimidt, H., Hornetz, B. and Shisanya, C. 2006. Farm Management Handbook of
216 Kenya Vol. II. Natural Conditions and Farm Management Information, 2nd edn. Nairobi,
217 Kenya: Ministry of Agriculture.

218 Jesse, E., and Cropp, B. 2004. Basic milk pricing concepts for Dairy Farmers. University of
219 Wisconsin Extension A3379 retrieved from
220 <http://future.aae.wisc.edu/publications/a3379.pdf> on 21st August 2017

221 Kimenju, S. C., & De Groote, H. 2010. Economic analysis of alternative maize storage
222 technologies in Kenya (pp. 19–23). Presented at the Joint 3rd African Association of
223 Agricultural Economists (AAAE) and 48th Agricultural Economists Association of South
224 Africa (AEASA) Conference, Cape Town, South Africa.

225 Kimenye, L., & McEwan, M. 2014. Scaling up, Dissemination and Adoption of Agricultural
226 Technologies using Innovation Platforms-Lessons from Eastern and Central Africa.
227 Entebbe: ASARECA.

228 Lukuyu, B., Gachuri, C. K., Lukuyu, M. N., Lusweti, C., Mwendia, S. 2012. Feeding dairy cattle
229 in East Africa. East Africa Dairy Development Project, 1–112. Retrieved from
230 [https://scholar.google.com/scholar?cluster=8864360012852157201&hl=en&as_sdt=2005](https://scholar.google.com/scholar?cluster=8864360012852157201&hl=en&as_sdt=2005&sciodt=0,5)
231 [https://scholar.google.com/scholar?cluster=8864360012852157201&hl=en&as_sdt=2005](https://scholar.google.com/scholar?cluster=8864360012852157201&hl=en&as_sdt=2005&sciodt=0,5)
&sciodt=0,5

232 Manaye, T., Tolera, A., & Zewdu, T. 2009. Feed intake, digestibility and body weight gain of
233 sheep fed Napier grass mixed with different levels of Sesbania sesban. *Livestock Science*,
234 122(1), 24–29.

235 Mburu, L., Gitu, K., & Wakhungu, J. 2007. A cost-benefit analysis of smallholder dairy cattle
236 enterprises in different agro-ecological zones in Kenya highlands. *Livestock Research for*
237 *Rural Development*, 19(7), 2007.

238 Methu, J., Owen, E., Abate, A., & Tanner, J. 2001. Botanical and nutritional composition of maize
239 stover, intakes and feed selection by dairy cattle. *Livestock Production Science*, 71(2), 87–
240 96.

241 Mwendia, S., Notenbaert, A.M.O., & Paul, B. 2016. Forage seed systems in Kenya
242 <http://hdl.handle.net/10568/72588><http://hdl.handle.net/10568/72588>.

243 Mwendia S, Maass, B., Njenga D., Nyakundi F. & Notenbaert A. (2017). Evaluating oat cultivars
244 for dairy forage production in the central Kenyan highlands. *African Journal of Range &*
245 *Forage Science* <http://www.tandfonline.com/doi/pdf/10.2989/10220119.2017.1358214>

246 Negassa, A., Shapiro, B., Kidane, T., Abdena, A., & Hanson, J. 2016. Ex-ante assessment of
247 demand for improved forage seed and planting materials among smallholder farmers in
248 Ethiopia: A contingent valuation analysis.

249 Odero-Waitituh JA. 2017. Smallholder dairy production in Kenya; a review. *Livestock Research*
250 *for Rural Development*. 29, Article #139 accessed on 3rd July, 2017 from
251 <http://www.lrrd.org/lrrd29/7/atiw29139.html>

252 Rønholt, S., Mortensen, K., & Knudsen, J. C. 2013. The Effective Factors on the Structure of
253 Butter and Other Milk Fat- Based Products. *Comprehensive Reviews in Food Science and*
254 *Food Safety*, 12(5), 468–482.


255 Tessema, Z., Mihret, J., & Solomon, M. 2010. Effect of defoliation frequency and cutting height
256 on growth, dry- matter yield and nutritive value of Napier grass (*Pennisetum purpureum*
257 (L.) Schumach). *Grass and Forage Science*, 65(4), 421–430.

258 University of Reading. 2000. One Animal per Farm? University of Reading Statistical Services
259 Centre, (March). Retrieved from
260 <http://www.reading.ac.uk/ssc/resources/OneAnimalPerFarm.pdf>
261 <http://www.reading.ac.uk/ssc/resources/OneAnimalPerFarm.pdf>

262 Wedholm, A., Larsen, L. B., Lindmark-Månsson, H., Karlsson, A. H., & Andrén, A. 2006. Effect
263 of protein composition on the cheese-making properties of milk from individual dairy
264 cows. *Journal of Dairy Science*, 89(9), 3296–3305.

265 Yami, M., Begna, B., & Teklewold, T. 2013. Enhancing the productivity of livestock production
266 in highland of Ethiopia: Implication for improved on- farm feeding strategies and
267 utilization. *International Journal of Livestock Production*, 4(8), 113–127.


268



1

2 Figure 1. Mean morning and evening milk production (kg) over 6 weeks' experimental period at Ol-joro-Orok, in
3 Nyandarua county in Kenya. The 10-day period between the arrows depict intervention feeding.

4



1

2 Figure 2. Means (\pm se) for milk quality attributes measured under farmer practice or intervention for (a) fat content

3 (b) protein (c) lactose (d) density and solid non-fat (e) at Ol-joro-Orok, in Nyandarua county in Kenya in January

4 2017.

1 Table 1. Daily feeds and fresh forages offered, under farmers' practice during the study in January-February 2017.

Farm	Average /day (kg)											Estimated Total dry matter intake /day	ME (MJ)	CP (g)
	NG (20)	MS (90)	Weeds (30)	Hay (87)	BH (53)	FS (23)	FO (18)	DM (90)	Bran (90)	GZ (hrs.)	MN (g)			
1	21.9	4.7	-	4.7	-	-	-	1.2	1.1	5.6	120	14.8	132.1	1219.8
2	2.6	-	2.7	-	1.6	-	-	1.2	-	7.0	adlib	3.3	36.9	361.0
3	3.7	-	2.8	4.7	0.9	-	2.0	1.4	-	-	100	7.8	77.8	729.6
4	1.1	2.0	13.1	14.5	-	-	-	1.2	-	-	80	19.6	192	1421.4
5	5.3	2.8	3.9	-	-	1.9	-	1.2	-	5.0	80	6.3	65.5	418.7
6	-	5.4	2.5	-	-	-	-	1.2	-	6.1	80	6.7	66.1	248.5
7	6.1	3.0	-	-	0.5	-	-	0.6	-	3.0	80	4.8	46.3	261.3
8	1.5	5.2	8.3	-	3.5	-	-	1.3	-	7.8	adlib	10.5	109.4	574.8

2 *NG (Napier grass); MS (Maize stovers); BH (bean haulms); FS (Fodder Sorghum); FO (Fodder Oat); DM (Dairy*

3 *meal); GZ (Grazing); MN (minerals); - (indicates not applicable). Values in brackets denote (%) dry matter content*

4 *adapted from Ayoade et al. 1983 and Gietema 2005. ILRI, 2016*

1
2 Table 2. Farmers' average milk production (kg) under farmer practice (FP) and intervention (IN) with associated
3 quality attributes (g) during the trial period in Ol-joro-Orok, in Nyandarua county in Kenya.

Farm	Treatments	milk production (kg)		BF (g)	Lactose (g)	Protein (g)	SNF (g)
		Morning	Evening				
1	FP	4.5	3.9	229.5	345.1	230.2	671.9
	IN	5.7	5.3	219.3	477.8	318.4	929.0
2	FP	2.6	2.1	222.6	220.0	146.9	430.6
	IN	2.9	2.6	223.4	225.4	150.6	441.1
3	FP	4.4	4.0	250.6	335.4	223.7	655.4
	IN	4.7	4.3	310.2	356.4	241.7	706.7
4	FP	5.5	5.1	319.9	422.9	281.9	840.3
	IN	6.5	5.5	436.9	475.1	321.7	937.7
5	FP	3.6	2.7	261.0	251.4	167.8	489.2
	IN	4.6	3.3	299.1	326.3	217.8	635.2
6	FP	2.8	2.2	173.4	198.8	132.8	376.5
	IN	3.3	2.8	203.5	234.2	147.3	458.7
7	FP	3.7	3.2	230.2	270.6	182.6	531.9
	IN	4.8	3.7	291.2	307.1	204.9	603.4
8	FP	2.7	2.5	141.1	194.4	129.6	383.1
	IN	3.3	2.7	183.5	224.6	149.7	441.2
<i>LSD</i>		0.5***	0.4***	52.7***	34.2***	24.1***	71.9***
<i>All Farms</i>	Farmer practice	3.7	3.2	230.2	284.3	189.9	556.0
	Intervention	4.5	3.8	272.2	330.3	220.4	648.0
<i>LSD</i>		0.3***	0.3**	24.6***	26.5***	18.1***	53.0***

4 *Degree of freedom (df) 209. BF- butterfat; SNF-Solids-Non-Fat; **P < 0.01; ***P < 0.001*

5

1 Table 3: Cost (in US\$) of producing main fodder crops per acre and value of crop residues and weeds used as livestock feed

Cost	Oat (Conway)	Vetch	Napier Grass	Local oats	Maize Stover	Irish potatoes residues	Beans haulms	Weeds
Inputs								
Vegetative Materials (Cuttings/Splits)	0	0	33.95	0	-	-	-	-
Seeds	38.8	38.8	0	43.65	-	-	-	-
Fertilizer (DAP)	38.8	0	0	29.1	-	-	-	-
Fertilizer (CAN)	0	0	0	27.16	-	-	-	-
Organic Manure	0	0	58.2	0	-	-	-	-
Herbicide (Round up)	7.76	0	0	0	-	-	-	-
Omex (Foliar feed- oats)	2.43	0	0	0	-	-	-	-
Bellamine (Herbicide broad leaf)	5.82	0	0	0	-	-	-	-
Orus (control rust in oat)	12.61	0	0	0	-	-	-	-
Labour								
Ploughing and Harrowing	38.8	38.8	38.8	38.8	-	-	-	-
Planting and fertilizer/manure application	29.1	19.4	38.8	29.1	-	-	-	-
Manual weeding	0	38.8	29.1	0	-	-	-	-
Spraying herbicides and pesticides	4.85	0	0	0	-	-	-	-
Harvesting and Transportation	83.42	102.82	14.55	33.95	-	-	-	-
Total cost of production per acre	83.42	228.92	213.4	201.76	-	-	-	-
Production potential (Kg/acre)	7769.97	623.22	19600	2000				
Value per acre	-	-	-	-	19.4	17.46	24.25	17.46

2 Source: Authors Survey, 2017; - indicates not applicable

3

1 Table 4: Profitability of oats and vetch acre/cow/season (Values in US\$)

Attribute	Farmer Practice	Intervention
<i>Costs</i>		
Cost of inputs	134.93	201.08
Cost of labour	38.80	26.68
Total Costs	173.73	237.46
<i>Benefits</i>		
Revenue – Sale of Milk	331.74	419.84
Profit of milk/acre/cow	158.01	192.08
<i>CBA Indicator</i>		
NPV	22	
IRR	15%	
PBP	65 days	

2

3