Growing soybean and maize in rotation

in partnership with IAR, SG2000, ADPs, NEARLS, and DGDC–Belgium

Growing soybean and maize in rotation

Year 1: Grow soybean Year 2: Grow maize on the same plot

Some benefits:


- Soybean brings nitrogen from air into soil as a free source of urea.
- You can obtain good maize yield after soybean with less inorganic fertilizer application due to soil improvement by soybean.
- Growing soybean before maize reduces *striga* infestation in your maize farm.
- Growing soybean and maize ensures food security and diversification (good animal feed, high quality food per family).




Year 1: Grow soybean




Buy seed of improved soybean variety with good haulm and grain yield from a reliable seed company. 40–50 kg Soybean seeds per hectare is required. **2** Apply fertilizers: For a farm of a hectare size, broadcast six bags of SSP before ridging.






3 Make ridges 50 cm apart either by hand, animal traction or by tractor.


Plant after a good rain. Make shallow grooves (1–2 cm deep) on ridges and drill soybean seed (1 seed every 5 cm) and cover lightly with soil.






S Scare birds during the first two weeks after planting.


Control weed manually or with herbicides recommended for soybean by extension agent.






When crop is infested by insects, spray with recommended insecticides.

B Harvest by cutting at ground level with sickle when leaves turn yellow. Do not harvest by uprooting the plants.





Thresh soybean on the field. Spread the haulm on the field or feed them to livestock and return the manure to the field.




Year 2: Maize on the same plot




• Buy improved maize seeds from a reliable seed company.


Break old ridges and make new ridges75 cm apart.





Plant after a good rain. Use a stick to make planting hole 5 cm deep at every 25 cm on the ridge. Plant one seed per hole and cover lightly with soil. Control weed by spraying recommended herbicide before maize germination or manually after germination.





UREA
UREA

Apply 3 bags of NPK when maize is at emergence (7 days after planting). Use a stick to make a hole at 5 cm away from plants. Place the fertilizer in the hole using a coca-cola cap (1 cap full) and cover the hole.

Apply 3 bags of Urea when plants are at the knee height. Use a stick to make a hole at 5 cm away from plants. Place the fertilizer in the hole using a coco-cola cap (1 cap full) and cover the hole. **6** Bend cobs downward at maturity to hasten drying and to minimize spoilage.





Harvest by dehusking and removing cobs immediately from the field.

B Dry cobs properly before storage in a dry place.

