

1 **Blanket application rates for synthetic grain protectants across agro-climatic zones: Do**  
2 **they work? Evidence from field efficacy trials using sorghum grain**

3 Macdonald Mubayiwa<sup>1</sup>, Brighton M. Mvumi<sup>1</sup>, Tanya E. Stathers<sup>2</sup>, Shaw Mlambo and Tinashe  
4 Nyabako<sup>1</sup>

5 <sup>1</sup>Department of Soil Science and Agricultural Engineering, University of Zimbabwe. P. O. Box  
6 MP 167 Mt Pleasant, Harare

7 <sup>2</sup>Natural Resources Institute, University of Greenwich, Central Avenue, Chatham Maritime,  
8 Kent, ME4 4TB, United Kingdom

9 Corresponding author: [mvumibm@hotmail.com](mailto:mvumibm@hotmail.com); [mvumibm@agric.uz.ac.zw](mailto:mvumibm@agric.uz.ac.zw)

---

10

11 **Abstract**

12 Many smallholder farmers in sub-Saharan Africa rely on synthetic pesticides for protecting  
13 stored-grain. Recommendations on use of these grain protectants are typically based on  
14 “blanket” application rates which are fixed rates that are not varied according to grain type, pest  
15 range or agro-climatic regions. There are numerous anecdotal reports of storage pesticide failure  
16 or reduced efficacy from farmers. Might rising global temperatures be a contributory factor?  
17 Smallholder farmers are responding by over-applying pesticides, increasing the application  
18 frequency or switching to non-recommended pesticides; leading to a pesticide treadmill. Trials to  
19 determine the efficacy and persistence of five commercially-available synthetic pesticides  
20 applied at manufacturer’s recommended rates on stored sorghum grain under contrasting climatic  
21 conditions were conducted in Mbire (mean temperatures of 32 – 42 °C and 30 – 50 % rh) and  
22 Harare (18 – 32 °C; 42 – 75 % rh) districts in Zimbabwe. Grain samples were collected at 8-

23 week intervals throughout a 10 month period in the 2014/15 and 2015/16 storage seasons. The  
24 samples were analyzed for insect grain damage, weight loss, total number of storage insects by  
25 species) and grain moisture content. Results showed significant differences in the performance of  
26 treatments ( $p < 0.001$ ). Grain damage was consistently higher in Harare than in Mbire. *Tribolium*  
27 *castaneum* was the dominant pest in Mbire, while *Sitotroga cerealella* and *Sitophilus oryzae*  
28 were dominant in Harare. *Tribolium castaneum* populations were high in the Shumba Super  
29 dust<sup>®</sup> (fenitrothion 1% + deltamethrin 0.13%) treatment in Mbire, while *S. cerealella* was  
30 dominant in Super guard<sup>®</sup> (pirimiphos-methyl 1.6% + permethrin 0.4%) and Actellic Gold dust<sup>®</sup>  
31 (pirimiphos-methyl 1.6% + thiamethoxam 0.36%) treated grain in Harare. Grain moisture  
32 content varied with ambient conditions, and was high in treatments with high insect pest levels.  
33 The results show that differences in climatic conditions influence insect pest species dynamics  
34 and response to pesticide treatments. Storage pesticides are not equally effective across different  
35 climatic conditions; thus more context-specific application recommendations are required.

36 **Key words:** synthetic pesticides, pesticide degradation, pesticide tolerance, sorghum grain  
37 storage, climate change and variability, storage pest dynamics

---

38 **1. Introduction**

39 Improvements in the food and nutrition security status of many sub-Saharan African (SSA)  
40 countries through enhanced crop production are being hampered by a rapidly growing human  
41 population and the effects of climate change and variability (Kaminski and Christiaensen, 2014).  
42 Rising temperatures and less predictable rainfall patterns and amounts are already occurring in  
43 SSA (Niang et al., 2014). Temperatures are projected to increase by up to 4 °C by the end of the  
44 century, depending on the development pathway chosen (IPCC, 2014; Serdeczny et al., 2016).  
45 Due to this and inadequate and more intermittent rains during the growing season, the success of  
46 dryland crop production is getting more unpredictable. The importance of efficient postharvest  
47 grain management to protect whatever is harvested against loss due to storage insect pest damage  
48 is becoming ever more important (Stathers et al., 2013; Vassilakos et al., 2015). To guard against  
49 storage pest damage, many smallholder farmers in SSA rely on applying synthetic pesticides  
50 composed of organophosphates and synthetic pyrethroids (Arthur, 1996; Stathers et al., 2002,  
51 Vassilakos, 2015). However, rising temperatures may promote increased degradation of these  
52 synthetic pesticides (Ismail et al., 2012) and may also favour development of many grain storage  
53 insect pests (Gornall et al., 2010) and affect their distribution and biology (Palikhe, 2007) such as  
54 shortening of life cycles. Shortened life cycles can increase the chances of insects developing  
55 resistance to pesticides as the insects more quickly adapt to treatments (Musolin and Saulich,  
56 2012, Velázquez-Fernández et al., 2012).

57 In many areas characterized by inadequate rainfall and high temperatures in SSA, small grains  
58 such as sorghum are staples and therefore widely grown as a coping and resilience strategy to  
59 these climatic conditions. However, the postharvest losses that occur reduce the amount of grain  
60 available for human consumption. Informed estimates suggest annual sorghum postharvest

61 weight losses of 12.1 % for SSA (APHLIS, 2014). Most of these farm-level postharvest losses  
62 are due to poor postharvest handling and insect pest attack (World Bank, 2011) and the latter  
63 necessitates effective control strategies, such as synthetic pesticides.

64 However, over-reliance on a narrow range of synthetic pesticides makes the development of pest  
65 resistance inevitable (Hagstrum and Subramanyam, 2006). Although most of the evidence is  
66 anecdotal, farmers across SSA frequently report storage pesticide failure or reduced efficacy (De  
67 Groote et al., 2013; Mlambo et al., 2017), and tend to respond by increasing the pesticide  
68 application rates, using non-recommended pesticides and/or increasing application frequencies to  
69 effect kill, which increases safety risks for users, consumers and the environment, and can trigger  
70 a pesticide treadmill effect.

71 The application of synthetic pesticides on stored grain is based on “blanket” or generalised  
72 application rates (Pretty, 2012). There are fixed rates that are not varied according to grain type,  
73 pest range or agro-climatic regions (temperature, relative humidity), despite the often wide  
74 variability in these factors under field conditions. This is also irrespective of possible  
75 implications on pesticide stability and dominant insect pest species in the different physical  
76 environments. These synthetic grain protectants are supposed to be applied just once at the start  
77 of a 6 to 12-month storage period. The use of “blanket” application rates, make manufacturers’  
78 recommendations easier for agricultural extension agents to extend, and are simpler to  
79 implement for farmers. However, this practice could eventually render the pesticides ineffective;  
80 a potential drawback not always understood by stakeholders. The objective of the current study  
81 was to determine the suitability of “blanket” application rates across different agro-climatic  
82 regions, a topic which has not previously been well-investigated under field conditions.

83 Much of the research undertaken to investigate insect response to synthetic pesticides has  
84 focussed on acute effects of adult mortality under rigidly controlled experimental conditions in  
85 the laboratory. There is, therefore, limited information on the effect on insect fecundity, pesticide  
86 persistence, or the effects on mixed populations of insects *in vivo*. The current studies were on  
87 sorghum, a small-grain grown in some of the more marginal agro-climatic zones; areas which are  
88 likely to get even warmer in the future. Therefore, information generated in this study is  
89 important in terms of deepening understanding of crop postharvest protection and food security  
90 in already highly vulnerable situations. The study, therefore, also sought to determine the  
91 diversity of storage insect species found on sorghum under the prevailing ambient conditions in  
92 two contrasting agro-climatic zones


93

## 94 **2. Materials and methods**

### 95 *2.1 Description of trial sites*

96 On-station researcher-managed trials to determine the persistence and effectiveness of synthetic  
97 grain protectants under contrasting agro-climatic conditions were set up at Mahuwe Rural  
98 Service Centre in Mbire district (20° 43' S; 30° 34' E), Zimbabwe, and at the Department of Crop  
99 Science, University of Zimbabwe in Harare (17° 47' S; 31° 03' E), Zimbabwe, during the  
100 2014/15 and 2015/16 storage seasons. Mbire district is located in the Zambezi valley, in northern  
101 Zimbabwe, about 283 km from the capital city, Harare (Fig. 1). The district is characterized by  
102 high annual temperature ranges of 32 – 42 °C, low rainfall of less than 450 mm per annum, and  
103 low mean annual relative humidity of 30 – 50 %. Harare, is located in central Zimbabwe, and

104 receives warm to high temperatures ranging between 18 – 30 °C, a high mean annual rainfall  
105 range of 900 – 1000 mm, and has a mean annual relative humidity of 42 – 75 %.



106  
107 Figure 1: Map of Zimbabwe showing the trial sites: Harare and Mbire district

108  
109 *2.2 Experimental layout*

110 Five commercially available synthetic grain protectant pesticides commonly used in Zimbabwe  
111 and an untreated control treatment were evaluated in this study (Table 1). In response to  
112 numerous anecdotal reports from farmers that locally-purchased grain protectants had lower  
113 efficacy than those bought at agro-dealers in urban centres, two Shumba Super dust<sup>®</sup> pesticide  
114 products were evaluated in the first season; one was purchased from a registered agro-dealer in  
115 Harare and the other bought from a local agro-dealer in Mbire. The Mbire local Shumba Super  
116 dust was not applied in Harare. However, as no significant differences between the locally-  
117 bought and Harare-bought pesticides were found in the first season trials, the Mbire locally  
118 purchased Shumba Super treatment was replaced by a newly introduced synthetic pesticide dust,  
119 Actellic Gold dust<sup>®</sup> during the second season.

120

121 Sorghum grain weighing 450 kg was thoroughly mixed to ensure it was as homogenous as  
122 possible before dividing it into lots of 75kgs of threshed grain per treatment. The 75 kgs of the  
123 threshed grain was sub-divided into three 25 kg lots before being separately admixed with  
124 pesticides at the application rates recommended on their labels (Table 1). During the 2014/15  
125 season, SC Sila variety of sorghum was used, while in the 2015/16 storage season, a 1:1 mixture  
126 of SC Sila and Macia varieties of sorghum grain was used. These smallholder farmers commonly  
127 change and mix the crop varieties they store as grain for food. Therefore grain protectants need  
128 to be effective on a range of varieties and mixtures.

129

130 After admixing, each treatment replicate was loaded into new polypropylene bags, which were  
131 then labelled and placed on brick dunnage to avoid direct contact with the floor and therefore  
132 reduce the chances of moisture accumulation in the stored grain. The trials were conducted over  
133 a 40-week period (~10 months) during each storage season. The treatments were laid out in a  
134 completely randomized design with three replicates of each treatment per site. Temperature and  
135 relative humidity data were collected using Easylog data loggers (Model EL-USB-1,  
136 Whiteparish, Wiltshire, SP5 2SJ, United Kingdom) installed 1.5 m above the ground in the  
137 storage rooms. The trials were housed in brick-walled rooms, with ceilings, and iron sheet roofs  
138 at both sites.

139

### 140 2.3 *Sampling, sample analysis and measurements*

141 Composite samples of 500 g were collected from each treatment at each sampling timepoint  
142 using a Seedburo Bag Trier spear (No. 76 13", Nickel Plated steel, 1½" outside diameter at the

143 large end, 7 3/8" long with top slot of 1 1/4" tapering down to 1/4"). The 500 g (approximately 18  
144 000 grains) sample was used for analysis of grain moisture content, grain damage, storage insect  
145 numbers present by species, and grain weight loss. The grain samples were collected by probing  
146 from at least five equidistant points around the bag, ensuring that grain was collected from the  
147 top, middle and bottom sections of each bag. Sampling was done every eight weeks until trial  
148 termination at 40 weeks. Insect counts per species were expressed per kilogram of grain; % grain  
149 damage was calculated as a proportion of the total number of grains in the sub-sample; and grain  
150 weight loss was determined using the count-and-weigh method (Boxall, 1986). Grain moisture  
151 content was measured using a pre-calibrated Dickey-John digital moisture meter (M3G™ model;  
152 Dickey-John Corporation, Minneapolis, USA).



153 Table 1: Grain protectant treatments used in the trial, and their active ingredients

Trade name	Active ingredients	Application rate (g / 25kg grain)*	Pesticide groups combined in product			Harare district		Mbire district	
			Organophosphates	Pyrethroids	Neonicotinoids	2014/15 season	2015/16 season	2014/15 season	2015/16 season
1. Super guard dust <sup>®</sup>	pirimiphos-methyl 1.6% + permethrin 0.4%	13.9	•	•		✓	✓	✓	✓
2. Chikwapuro <sup>®</sup>	pirimiphos-methyl 2.5% + deltamethrin 0.1%	10.0	•	•		✓	✓	✓	✓
3. Harare Shumba super dust <sup>®</sup>	fenitrothion 1% + deltamethrin 0.13%	12.5	•	•		✓	✓	✓	✓
5. Mbire Shumba super dust <sup>®</sup>	fenitrothion 1% + deltamethrin 0.13%	12.5	•	•		x	x	✓	x
6. Actellic gold dust <sup>®</sup>	pirimiphos-methyl 1.6% + thiamethoxam 0.36%	12.5	•		•	x	✓	x	✓
7. Ngwena yedura <sup>®</sup>	pirimiphos-methyl 2.5% + deltamethrin 0.2%	10.0	•	•		✓	✓	✓	✓
8. Untreated control	N/A					✓	✓	✓	✓

154 \* For pesticides, manufacturer's label application rates were used. Shumba super dust<sup>®</sup> obtained from Mbire agro-  
 155 dealers for use in the 2014/15 storage season was replaced by Actellic gold dust<sup>®</sup> during the 2015/16 storage  
 156 season; x – indicates absence of the treatment in that specific trial

157  
158

159 2.4 Data analysis and data presentation

160 The percentage number of damaged grains, and the percentage weight loss data were analysed in  
 161 Genstat version 14, using a repeated measures analysis of variance (rANOVA) as the samples  
 162 were collected from the same experimental unit throughout the trial. Initially all data were  
 163 subjected to the Shapiro-Wilk test for normality. Data on percentage number of damaged grains

164 and percentage weight loss which failed the normality test were transformed using the square  
165 root transformation (Bartlett et al., 1936). Where the ANOVA showed significant differences,  
166 means were further separated using Fisher's protected LSD at 5 % probability.

167 Data on total *Sitotroga cerealella* (Olivier) (Lepidoptera: Gelechiidae) populations in Harare  
168 were presented separately in tables derived from ANOVA means because of their abundance  
169 which masked other insect species. Data on total counts of *S. cerealella* did not meet the  
170 assumptions of analysis of variance and were  $\log_{10}(x + 1)$  transformed, with  $x$  representing the  
171 recorded original mean (De Muth, 2014).

172

### 173 3. Results

#### 174 3.1 Pest dynamics


175 The dominant insect pest species in the Harare trial during the 2014/15 storage season was  
176 *S. cerealella*, which developed extensively in grain treated with Super guard dust and in  
177 untreated grain (Table 2 compared to Fig. 2). The highest populations of the moth were recorded  
178 in July 2015 (at 32 weeks storage) and the total populations decreased thereafter. However, the  
179 protectants Chikwapuro and Ngwena yedura managed to suppress the development of the moth  
180 throughout the 40 week storage period. High numbers of *Sitophilus oryzae* L. (Coleoptera:  
181 Curculionidae) developed in the untreated control grain in the Harare trial during the 2014/15  
182 storage season (Fig 2).

183

184 Table 2: Mean total *Sitotroga cerealella* counts ( $\pm$  SEM) per kilogram of sorghum grain in the different storage protectant treatments  
 185 during the 2014/15 storage season in Harare (n = 3)


Treatment	Storage duration (weeks)					
	0	8	16	24	32	40
Untreated control	7.24 $\pm$ 2.07	9.11 $\pm$ 2.55a	556.08 $\pm$ 148.39c	1382.84 $\pm$ 68.62c	6163.77 $\pm$ 2831.63b	1395.86 $\pm$ 119.03b
Super guard dust	10.35 $\pm$ 1.94	207.5 $\pm$ 17.04b	251.35 $\pm$ 19.26c	1289.02 $\pm$ 90.4c	9818.79 $\pm$ 1126.41b	1627.91 $\pm$ 201.77b
Ngwena yedura	5.07 $\pm$ 4.28	6.75 $\pm$ 1.76a	7.17 $\pm$ 0.07b	4.06 $\pm$ 1.45a	8.14 $\pm$ 2.31a	3.41 $\pm$ 1.01a
Chikwapuro	6.97 $\pm$ 5.12	3.68 $\pm$ 1.01a	0.0 $\pm$ 0.00a	8.01 $\pm$ 2.05ab	12.24 $\pm$ 6.42a	1.83 $\pm$ 0.99a
Shumba Super dust (Harare)	6.85 $\pm$ 2.68	24.76 $\pm$ 22.36a	24.67 $\pm$ 21.54b	61.07 $\pm$ 53.48b	93.28 $\pm$ 81.35a	246.14 $\pm$ 240.95a
<b>p-value</b>	<b>0.79</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>
<b>F<sub>4,10</sub></b>	<b>0.42</b>	<b>48.59</b>	<b>12.7</b>	<b>164.37</b>	<b>11.11</b>	<b>27.82</b>

186 \*Means in the same column followed by different alphabetical letters are significantly different using Fisher's protected 5 % LSD test




187  
 188 Figure 2: Comparison of mean total number of adult insects by species (excluding *Sitotroga*  
 189 *cerealella*) per kilogram of sorghum grain in different grain protectant treatments in Harare  
 190 during the 2014/15 storage season (n = 3)

191  
 192 Although *S. cerealella* and *S. oryzae* were dominant in Harare, they failed to establish in the  
 193 Mbire trial during the two seasons, except during the relatively warmer months, between April  
 194 and June where temperatures of 28 – 30 °C were experienced. *Tribolium castaneum* (Herbst)  
 195 (Coleoptera: Tenebrionidae) and *Rhyzopertha dominica* (Fabricius) (Coleoptera: Bostrichidae)  
 196 were the dominant storage pests in Mbire during the 2014/15 storage season (Fig. 3). *Sitotroga*  
 197 *cerealella* populations were recorded between January and May 2015 (week 8 to 24) (end of  
 198 summer season to the onset winter).



199  
 200 Figure 3: Comparison of the mean total number of adult insects by species per kilogram of  
 201 sorghum grain in Mbire during the 2014/15 storage season (n = 3)

202  
 203 Higher initial pest populations were recorded in the Harare trial during the 2015/16 than the  
 204 2014/15 storage season. In the Harare 2015/16 trial, there was a decline in the number of insect  
 205 pests recorded in all the pesticide-treatments after the baseline samples were collected. *Sitotroga*  
 206 *cerealella* and *S. oryzae* were the dominant pest species in Harare during the 2015/16 season,  
 207 with *S. cerealella* managing to develop in Super guard treated grain while the other pesticide  
 208 treatments suppressed it (Table 3). Numbers of *S. oryzae* increased in the untreated control (Fig.  
 209 4), but failed to increase in the synthetic pesticide treatments. Low populations of *T. castaneum*  
 210 were recorded in the untreated grain and in Shumba Super dust.



212

213 Figure 4: Comparison of the mean total number of adult insects by species in different grain

214 protectant treatments in Harare during the 2015/16 storage season (n = 3)

215

216


217 Table 3: Comparison of the mean total adult *Sitotroga cerealella* counts ( $\pm$  SEM) in different grain protectant treatment in Harare  
 218 during the 2015/16 storage period (n = 3)

<b>Treatment</b>	<b>0 weeks</b>	<b>8 weeks</b>	<b>16 weeks</b>	<b>24 weeks</b>	<b>32 weeks</b>	<b>40 weeks</b>
Untreated control	5.82 $\pm$ 1.32	5.46 $\pm$ 1.67	34.04 $\pm$ 15.12ab	34.04 $\pm$ 15.12ab	41.38 $\pm$ 9.62bc	135.84 $\pm$ 61.22abc
Super guard dust	3.47 $\pm$ 1.35	17.46 $\pm$ 8.23	119.05 $\pm$ 41.24c	119.05 $\pm$ 41.24c	199.09 $\pm$ 163.87c	1060.51 $\pm$ 633.64c
Ngwena yedura	5.86 $\pm$ 1.41	17.67 $\pm$ 10.16	6.86 $\pm$ 4.84a	6.86 $\pm$ 4.84a	30.14 $\pm$ 11.86abc	34.00 $\pm$ 14.26a
Chikwapuro	3.50 $\pm$ 1.84	3.86 $\pm$ 2.57	40.54 $\pm$ 10.38b	40.54 $\pm$ 10.38b	8.70 $\pm$ 2.84a	40.77 $\pm$ 21.48a
Shumba Super dust (Harare)	9.45 $\pm$ 0.76	21.23 $\pm$ 1.54	30.26 $\pm$ 4.92ab	30.26 $\pm$ 4.92ab	19.03 $\pm$ 6.84ab	22.97 $\pm$ 6.14a
Actellic Gold dust	7.88 $\pm$ 1.49	26.37 $\pm$ 18.49	52.58 $\pm$ 16.49bc	61.83 $\pm$ 7.47bc	105.47 $\pm$ 29.17c	174.19 $\pm$ 20.68bc
<b>p-value</b>	<b>0.07</b>	<b>0.28</b>	<b>0.01</b>	<b>0.01</b>	<b>0.02</b>	<b>&lt;0.01</b>
<b>F<sub>5,12</sub></b>	<b>2.27</b>	<b>0.85</b>	<b>3.72</b>	<b>4.19</b>	<b>1.14</b>	<b>2.41</b>

219 \*Means in the same column followed by different alphabetical letters are significantly different using Fisher's protected 5 % LSD test

220 In the 2015/16 Mbire trial, the dominant insect pest species were *S. oryzae* and *T. castaneum*  
 221 (Fig. 5). *Tribolium castaneum* build-up was high in the untreated and the Shumba Super dust  
 222 treated grain. The pest also developed in lower numbers in all the other pesticide treatments from  
 223 week 24 onwards (Fig. 5).

224



225  
 226 Figure 5: Comparison of the mean total adult insect counts per kilogram of stored sorghum grain  
 227 in different grain protectant treatments in Mbire district during the 2015/16 storage season (n =3)

228

229 **3.2 Grain damage**

230 Grain damage was high in the untreated and the Super guard dust treated grain in Harare in  
 231 2014/15 (Table 4). While damage was suppressed in the Chikwapuro, Ngwena yedura and  
 232 Shumba Super dust treatments throughout the storage season. From 8 weeks storage onwards,  
 233 significant differences in the percentage number of damaged grains were noted between



234 treatments ( $F_{4, 10} = 46.06$ ;  $p < 0.01$ ), and the differences became wide at the 40 weeks ( $F_{4, 10} =$   
235  $32.34$ ;  $p < 0.01$ ). In Mbire, significant differences ( $F_{5, 12} = 6.47$ ;  $p < 0.004$ ) in grain damage  
236 between grain protectant treatments were recorded from 16 weeks of storage (Table 4), with an  
237 increase in grain damage in the two Shumba Super dust treatments, Super guard dust and  
238 untreated control. However, the overall changes in grain damage were low in Mbire district as  
239 compared to Harare during the 2014/15 storage season.

240

241 The level of grain damage was significantly higher ( $p < 0.001$ ) in the untreated (36.5 %), Actellic  
242 Gold dust (19.0 %) and Super guard dust (11.9%) treated grain than in the other treatments in  
243 Harare by the end of the 2015/16 storage season (Table 5). Significant differences between  
244 treatments were recorded from 8 weeks storage onwards in Harare, whereas in Mbire they only  
245 occurred at 24 and 32 weeks storage. Grain damage levels were higher in Harare than Mbire  
246 district during the 2015/16 storage season (Table 5). In Harare, the pesticides Ngwena yedura,  
247 Chikwapuro and Shumba Super dust suppressed insect grain damage throughout the 2015/16  
248 storage season, whilst Chikwapuro, Actellic Gold dust and Super guard dust were effective in  
249 Mbire although grain damage levels were low in all treatments ( $< 10\%$ ) for up to 32 weeks  
250 (Table 5).

### 251 3.3 *Grain weight loss*

252 Grain weight loss was lower in Harare (0.42 %) than in Mbire (1.28 %) at trial set-up in 2014/15.  
253 From 16 weeks of storage onwards, significant differences in grain weight loss were recorded  
254 between the different protectant treatments ( $F_{5, 10} = 5.31$ ;  $p = 0.015$ ). In the Harare trial, grain  
255 weight loss was significantly higher in the untreated and Super guard dust treated grain while in

256 Mbire district, higher grain weight loss was recorded in the untreated control and the two  
257 Shumba Super dust pesticide treatments (Table 6). Overall, grain weight loss was higher in  
258 Harare than in Mbire district during the 2014/15 storage season.

259 During the 2015/16 storage season, significant differences ( $F_{5, 17} = 17.13$ ;  $p < .001$ ) in grain  
260 weight loss were recorded from week 24 weeks until trial termination in Harare, but only at 40  
261 week's storage in Mbire district ( $F_{5, 17} = 6.35$ ;  $p = 0.004$ ) (Table 7). Weight loss was significantly  
262 higher in the untreated grain than in the treated grain from 24 weeks' storage in Harare although  
263 by 32 weeks' storage weight loss in the Super guard dust and Actellic Gold dust was also  
264 significantly higher than in the grain treated with the other protectants in Harare. In Mbire,  
265 significantly higher grain weight loss was only recorded in the untreated control and Shumba  
266 Super dust treatments after 40 weeks' storage (Table 7).

267

#### 268 3.4 Grain moisture content and storage room temperatures

269 Mean temperatures in storage rooms in Harare were lower (22 – 24 °C) than those in Mbire  
270 district (29 – 34 °C) (Figs. 6a and b) during the 2014/15 storage season. In Harare, grain  
271 moisture content fluctuated in all treatments, with highest peaks occurring in the untreated and  
272 Super guard dust treated grain. However, in Mbire district, grain moisture content increased from  
273 week 8 (November) and reached a peak in week 16 ( $13.5 \pm 0.08$ ). An increase in temperatures  
274 within the storage rooms was linked to an increase in grain moisture content at both sites (Figs  
275 6a and b).

276 The room temperatures within the storage rooms in Harare were lower than those of Mbire  
277 district during the 2015/16 storage season. The mean temperature ranges within the storage

278 rooms in Harare were between 20 and 26 °C, and dropped slightly from week 24 until the end of  
279 the season. Grain moisture content in the Harare trial increased from 9 to 12 % between week 8  
280 and 16 of storage (Fig. 7a). However, untreated control and Super guard treated grain had higher  
281 moisture contents at week 40. In the Mbire trial, high mean temperatures (27 – 33 °C) were  
282 recorded in the storage room (Fig. 7b) and low grain moisture content of (9 – 10 %) ranges were  
283 recorded. A drop in grain moisture content at week 8 in Mbire district coincided with an increase  
284 in mean temperatures during the same period (Fig. 7b).

285 Table 4: Comparison of mean percentage number of damaged grains in different grain protectant treatments in Harare and Mbire  
 286 district during the 2014/15 storage season (n = 3)

Treat- ment	Harare						Mbire					
	0 weeks	8 weeks	16 weeks	24 weeks	32 weeks	40 weeks	0 weeks	8 weeks	16 weeks	24 weeks	32 weeks	40 weeks
1	1.92±0.15	7.36±0.30c	10.33±2.09c	24.95±0.03b	19.97±2.75b	35.59±6.27b	7.22±0.88	6.67±0.92	12.18±0.83cd	12.31±0.43	16.75±0.83c	17.83±1.50c
2	2.54±0.15	4.99±0.50b	6.20±0.30b	23.23±1.10b	30.33±2.41c	41.46±4.00b	6.63±0.41	6.07±0.75	9.49±1.01ab	10.70±0.46	10.70±0.78a	12.78±0.54ab
3	2.16±0.26	2.43±0.15a	2.53±0.23a	2.54±0.07a	2.69±0.23a	2.75±0.47a	7.06±0.89	4.88±0.49	10.13±0.73abc	10.45±0.49	10.93±1.73a	11.65±0.45a
4	2.10±0.24	2.92±0.22a	2.74±0.23a	3.18±0.41a	3.05±0.36a	2.30±0.36a	6.99±0.10	6.33±0.40	8.88±0.29a	9.56±0.21	11.12±0.14a	11.01±0.11a
5	2.21±0.14	3.25±0.15a	4.70±1.34ab	4.57±1.50a	8.96±5.53a	5.13±0.37a	7.37±0.73	7.44±0.63	13.30±0.39d	12.05±1.49	11.51±0.57a	15.88±0.81bc
6	*	*	*	*	*	*	8.21±0.79	6.94±0.78	11.43±0.44bcd	10.79±0.08	14.15±0.62b	14.04±0.70ab
<b>P-</b>	<b>0.31</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>0.71</b>	<b>0.22</b>	<b>&lt;0.01</b>	<b>0.12</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>
<b>value</b>												
<b>F<sub>5,12</sub></b>	<b>1.38</b>	<b>46.06</b>	<b>10.17</b>	<b>177.18</b>	<b>16.21</b>	<b>32.34</b>	<b>0.59</b>	<b>1.65</b>	<b>6.47</b>	<b>2.19</b>	<b>15.38</b>	<b>5.51</b>

287 1 = Untreated control ; 2 = Super guard dust; 3 = Ngwena yedura ; 4 = Chikwapuro; 5 = Shumba Super dust (Harare); 6 = Shumba Super dust (Mbire)

288 All means in the same column followed by a different letter are significantly different from one another at 5% LSD

289

290 Table 5: Comparison of mean percentage number of damaged grains ( $\pm$ SEM) in different grain protectant treatments in Harare and  
 291 Mbire district during the 2015/16 storage season (n=3).

Treat- ment	Harare						Mbire					
	0 weeks	8 weeks	16 weeks	24 weeks	32 weeks	40 weeks	0 weeks	8 weeks	16 weeks	24 weeks	32 weeks	40 weeks
1	4.10 $\pm$ 0.01	6.97 $\pm$ 0.41b	18.41 $\pm$ 1.11c	25.13 $\pm$ 2.29c	31.28 $\pm$ 1.49c	36.49 $\pm$ 1.15c	4.60 $\pm$ 0.60	4.98 $\pm$ 0.93	4.40 $\pm$ 0.58	6.62 $\pm$ 0.74b	8.71 $\pm$ 1.19c	10.25 $\pm$ 2.10
2	4.81 $\pm$ 0.31	5.29 $\pm$ 0.51a	5.03 $\pm$ 0.60a	6.50 $\pm$ 0.43ab	10.70 $\pm$ 0.77b	11.92 $\pm$ 0.97ab	4.20 $\pm$ 0.23	4.25 $\pm$ 0.23	4.85 $\pm$ 0.57	4.27 $\pm$ 0.28a	4.80 $\pm$ 0.47ab	5.8 $\pm$ 0.52
3	4.58 $\pm$ 0.07	5.90 $\pm$ 0.30ab	4.72 $\pm$ 0.32a	4.85 $\pm$ 0.37a	4.50 $\pm$ 0.31a	4.99 $\pm$ 0.30a	4.11 $\pm$ 0.45	4.26 $\pm$ 0.41	4.21 $\pm$ 0.24	3.9 $\pm$ 0.14a	6.80 $\pm$ 1.21bc	9.29 $\pm$ 1.93
4	4.27 $\pm$ 0.06	4.76 $\pm$ 0.62a	5.30 $\pm$ 0.26a	5.88 $\pm$ 0.08ab	5.07 $\pm$ 0.32a	5.44 $\pm$ 0.38a	3.48 $\pm$ 0.49	3.62 $\pm$ 0.51	3.44 $\pm$ 0.11	3.72 $\pm$ 1.00a	5.04 $\pm$ 0.64ab	7.65 $\pm$ 1.58
5	4.37 $\pm$ 0.21	6.63 $\pm$ 0.10b	5.61 $\pm$ 0.66ab	5.99 $\pm$ 0.57ab	5.42 $\pm$ 0.14a	6.38 $\pm$ 0.41a	5.02 $\pm$ 1.10	4.93 $\pm$ 1.05	4.37 $\pm$ 0.08	6.69 $\pm$ 0.50b	8.16 $\pm$ 0.30c	8.65 $\pm$ 0.11
6	4.34 $\pm$ 0.20	5.04 $\pm$ 0.32a	8.19 $\pm$ 1.53b	9.67 $\pm$ 2.64b	11.19 $\pm$ 2.6b	18.96 $\pm$ 6.31b	3.94 $\pm$ 0.79	3.93 $\pm$ 0.79	3.24 $\pm$ 0.30	3.51 $\pm$ 0.12a	4.31 $\pm$ 0.13a	4.62 $\pm$ 0.15
<b>P- value</b>	<b>0.17</b>	<b>0.02</b>	<b>&lt;0 .01</b>	<b>&lt;0 .01</b>	<b>&lt;0 .01</b>	<b>&lt;0 .01</b>	<b>0.69</b>	<b>0.73</b>	<b>0.69</b>	<b>&lt;0.01</b>	<b>0.01</b>	<b>0.12</b>
<b>F<sub>5,12</sub></b>	<b>1.89</b>	<b>4.56</b>	<b>37.03</b>	<b>27.86</b>	<b>53.46</b>	<b>21.06</b>	<b>0.62</b>	<b>0.56</b>	<b>2.76</b>	<b>6.73</b>	<b>5.22</b>	<b>2.23</b>

292 1 = Untreated control ; 2 = Super guard dust; 3 = Ngwena yedura ; 4 = Chikwapuro; 5 = Shumba Super dust (Harare); 6 = Actellic Gold dust

293 Means in the same column followed by different alphabetical letters are significantly different using Fisher's protected 5 % LSD test

294

295

296 Table 6: Comparison of mean % grain weight loss ( $\pm$ SEM) in different grain protectant treatments in Harare and Mbire during the  
 297 2014/15 storage season (n = 3)

Treatment	Harare				Mbire			
	16 weeks	24 weeks	32 weeks	40 weeks	16 weeks	24 weeks	32 weeks	40 weeks
Untreated control	2.34 $\pm$ 0.46b	6.1 $\pm$ 0.26b	4.4 $\pm$ 0.46bc	8.39 $\pm$ 0.49b	3.1 $\pm$ 0.45b	3.18 $\pm$ 0.09d	4.19 $\pm$ 0.21c	3.81 $\pm$ 0.2c
Super guard dust	1.44 $\pm$ 0.12ab	4.98 $\pm$ 1.22b	6.74 $\pm$ 0.41c	9.79 $\pm$ 0.41b	1.63 $\pm$ 0.4a	2.13 $\pm$ 0.06ab	1.98 $\pm$ 0.11a	2.16 $\pm$ 0.15a
Ngwena yedura	0.61 $\pm$ 0.05a	0.59 $\pm$ 0.06a	0.46 $\pm$ 0.05a	0.5 $\pm$ 0.05a	2.01 $\pm$ 0.16a	2.08 $\pm$ 0.14ab	2.16 $\pm$ 0.22a	2.03 $\pm$ 0.17a
Chikwapuro	0.62 $\pm$ 0.04a	0.34 $\pm$ 1.22a	0.52 $\pm$ 0.41a	0.57 $\pm$ 0.1a	1.79 $\pm$ 0.46a	1.69 $\pm$ 0.27a	2.23 $\pm$ 0.09a	1.95 $\pm$ 0.05a
Shumba Super dust (Harare)	1.2 $\pm$ 0.5a	1.27 $\pm$ 0.46a	2.79 $\pm$ 1.97ab	2.14 $\pm$ 1.97a	3.58 $\pm$ 0.28b	2.58 $\pm$ 0.24bc	2.86 $\pm$ 0.12b	3.31 $\pm$ 0.36bc
Shumba Super dust (Mbire)	*	*	*	*	3.15 $\pm$ 0.14b	2.77 $\pm$ 0.19cd	3.81 $\pm$ 0.08c	3.2 $\pm$ 0.1b
<b>p-value</b>	<b>0.02</b>	<b>&lt;0 .01</b>	<b>&lt;0.01</b>	<b>&lt;0 .01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>
<b>F<sub>5,12</sub></b>	<b>5.31</b>	<b>20.16</b>	<b>8.27</b>	<b>19.85</b>	<b>8.28</b>	<b>8.75</b>	<b>38.11</b>	<b>15.99</b>

298 *Figures presented in the table are original mean losses per treatment, and were compared per column. Figures within a column*  
 299 *followed by a different letter are significantly different from one another. (\*) indicates treatment excluded*

300

301

302



303

304 Table 7: Mean % grain weight loss ( $\pm$ SEM) in Mbire and Harare during the 2015/16 storage season (n = 3)

Treatment	Harare				Mbire			
	16 weeks	24 weeks	32 weeks	40 weeks	16 weeks	24 weeks	32 weeks	40 weeks
Untreated control	1.58 $\pm$ 0.58	5.97 $\pm$ 0.14b	6.58 $\pm$ 1.61b	8.14 $\pm$ 1.94c	0.4 $\pm$ 0.13	1.79 $\pm$ 0.32	1.72 $\pm$ 0.32	2.43 $\pm$ 0.41c
Super guard dust	1.01 $\pm$ 0.1	1.69 $\pm$ 0.33a	4.49 $\pm$ 0.76b	5.8 $\pm$ 0.48bc	1.17 $\pm$ 0.52	1.64 $\pm$ 0.35	0.34 $\pm$ 0.11	1.25 $\pm$ 0.08ab
Ngwena yedura	0.51 $\pm$ 0.1	0.5 $\pm$ 0.1a	0.58 $\pm$ 0.12a	0.85 $\pm$ 0.08a	0.95 $\pm$ 0.32	0.74 $\pm$ 0.11	1.11 $\pm$ 0.55	0.96 $\pm$ 0.32a
Chikwapuro	1.13 $\pm$ 0.57	1.35 $\pm$ 0.74a	0.59 $\pm$ 0.2a	0.97 $\pm$ 0.13a	0.71 $\pm$ 0.23	1.36 $\pm$ 0.68	0.67 $\pm$ 0.29	0.62 $\pm$ 0.11a
Shumba Super dust (Harare)	0.68 $\pm$ 0.11	0.62 $\pm$ 0.19a	1.36 $\pm$ 0.52a	1.17 $\pm$ 0.11a	0.97 $\pm$ 0.01	1.87 $\pm$ 0.28	1.17 $\pm$ 0.16	1.78 $\pm$ 0.14bc
Actellic Gold dust	0.57 $\pm$ 0.18	1.6 $\pm$ 0.84a	5.51 $\pm$ 1.29b	3.35 $\pm$ 1.06ab	0.62 $\pm$ 0.11	0.88 $\pm$ 0.02	0.64 $\pm$ 0.03	0.97 $\pm$ 0.27a
<b>p-value</b>	<b>0.3</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>&lt;0.01</b>	<b>0.47</b>	<b>0.20</b>	<b>0.11</b>	<b>&lt;0.01</b>
<b>F<sub>5,12</sub></b>	<b>1.38</b>	<b>17.13</b>	<b>14.92</b>	<b>10.56</b>	<b>0.97</b>	<b>1.75</b>	<b>2.29</b>	<b>6.35</b>

305 *Figures presented in the table are original mean losses per treatment, and were compared per column. Figures within a column followed by a different letter are*  
306 *significantly different from one another*

307



308

309 Figure 6: Comparison of the mean % grain moisture content ( $\pm$ SEM) in different grain protectant treatments and mean store room temperature in  
 310 Harare (a) and Mbire (b) during the 2014/15 storage season ( $n = 3$ ). In some cases, error bars are not visible due to little variation between the  
 311 treatment replications.

312

313



314

315



316

317



318

319 Figure 7: Comparison of the mean percentage grain moisture content ( $\pm$  SEM) in different grain protectant treatments and mean storage room  
320 temperatures in Harare (a) and Mbire (b) during the 2015/16 storage season ( $n = 3$ ). In some cases, error bars are not visible due to little variation  
321 between the treatment replications.

322

## 323 Discussion

324 The climatic conditions at the two trial sites influenced which insect pest species became  
325 dominant. The warm mean temperatures ( $23.6 \pm 0.8$  °C) in Harare during the 2014/15 storage  
326 season, with mean temperature range of 22 – 24 °C, favoured the development of *S. oryzae* and  
327 *S. cerealella* which develop well between temperatures of 18 – 30 °C (Mason and McDonough,  
328 2012; Akter et al., 2013). The high mean seasonal ambient store temperatures ( $31.8 \pm 1.4$  °C) in  
329 Mbire in the 2014/15 storage season, with a mean temperature range of 27 – 34 °C, favoured the  
330 development of pests such as *T. castaneum* and *R. dominica* which are tolerant to higher  
331 temperatures (Baldassari et al., 2004). Mean diurnal temperatures at the Mbire site ranged  
332 between 26.3 – 36.1 °C, during the 2014/15 storage season. High temperatures above 35 °C  
333 cause high mortality of many insect species due to disruption of their nervous and endocrine  
334 systems (Subramanyam and Hagstrum, 1991; Neven, 2000; Fields et al., 2015), and will have  
335 suppressed *S. cerealella* and *S. oryzae* population development in the Mbire trial. When  
336 temperatures rise beyond an individual storage insect's optimum conditions, they will enter into  
337 diapause (Bell, 2014), which stops their development and some of their metabolic processes, but  
338 leads to death if the conditions persist (Fleurat-Lessard and Dupuis, 2010).

339 Higher populations of *S. cerealella* in Super guard treated grain than in untreated control grain  
340 suggest the possibility of high tolerance levels to this pesticide by the pest. The pest, if tolerant,  
341 may manage to develop particularly well where there is a lack of interspecific competition with  
342 other species (Makundi et al., 2010). Such increased fecundity of one pesticide-tolerant species  
343 was also observed in trials involving the larger grain borer, *Prostephanus truncatus* on maize  
344 grain (Chigoverah and Mvumi, 2016). Differential responses to pesticide treatments by different  
345 species of storage insect pests has been reported by various researchers. Selected examples

346 include: spinosad-treated wheat grain where *S. oryzae* was less susceptible than *R. dominica*  
347 (Athanassiou et al., 2009); diatomaceous earths (DEs) treatment where different strains of *T.*  
348 *castaneum* responded differently (Rigaux et al., 2001) and malathion treatment where major  
349 storage insect pests showed differences (Arthur, 1996). In field stored sorghum trials Stathers et  
350 al. (2002) observed that *R. dominica* was less susceptible than *S. oryzae* to DEs.

351 The occurrence of high *S. cerealella* numbers in Actellic Gold dust treated grain in Harare is also  
352 attributed to high levels of tolerance to the pesticide by this pest, which managed to develop in  
353 treated grain as from 8 weeks of storage. Both Actellic Gold dust and Super guard dust have low  
354 levels of pirimiphos-methyl (1.6 %) compared to other pesticides which share the same active  
355 ingredient. Ngwena yedura and Chikwapuro have higher levels of pirimiphos-methyl (2.6 %) in  
356 addition to deltamethrin, and these pesticides displayed better levels of control of *S. cerealella*.  
357 Elimination of natural enemies could also be the result of increased build-up of *S. cerealella* and  
358 *T. castaneum* in Super guard dust in Harare and Shumba Super dust in Mbire, respectively. This  
359 was demonstrated in earlier research on *S. oryzae* and *A. calandreae* where the latter was more  
360 susceptible than the former to a range of active ingredients (Baker and Weaver, 1993). Similarly,  
361 Stathers et al. (2002) attributed the dominance of *R. dominica* in DE-treated sorghum grain to  
362 elimination of its natural enemies compared to untreated grain, coupled with the lower efficacy  
363 of DEs against bostrichid beetles.

364 *Sitotroga cerealella* is known to use behavioral avoidance mechanisms to escape treatments,  
365 avoiding unfavorable patches within the grain, then re-infesting the grain when pesticide levels  
366 become sub-lethal (Oppert et al., 2010; Trematerra, 2015). Storage insects which are able to  
367 detect chemical treatments and avoid them have better chances of survival (Bell, 2014). The high  
368 mobility of this *S. cerealella* increases its ability to escape soon after treatment (Trematerra,

369 2015), reducing its exposure to pesticides and providing it with a chance to recover (Rumbos et  
370 al., 2016). High susceptibility to pirimiphos-methyl has been reported previously for *S. oryzae*,  
371 *T. confusum* and *T. castaneum* (Rumbos et al., 2016) and was also found in the current  
372 experiment where all the pesticides containing this active ingredient suppressed these pest  
373 species.

374 Massive build-up of *T. castaneum* in Shumba Super dust treated sorghum grain in Mbire from  
375 two months of storage onwards indicates the possibility that the pest has developed resistance to  
376 the grain protectant which has been in use for a long period of time (at least 15 years) in  
377 Zimbabwe. The initial grain damage level of around 7 % in Mbire district during the 2014/15  
378 season, and approximately 4 % during the 2015/16 storage season would have been sufficient to  
379 support the development of *T. castaneum* which is a secondary pest. The point at which  
380 *T. castaneum* development commenced in treated grain can also mark a point whereby the  
381 pesticide's active compounds are degraded below potent levels (Afridi et al., 2001) due to  
382 predominantly high temperatures in the area. The quick and high rate of infestation by  
383 *T. castaneum* in Shumba Super dust treated grain is problematic, as in SSA smallholder farmers  
384 usually only treat the grain they intend to store for periods of more than 4 months (Stathers et al.,  
385 2013). Hence, this pesticide would fail to give the desirable protection. *Tribolium castaneum*  
386 build-up was low under the relatively low temperatures in Harare in all treatments. This could  
387 suggest that the pesticides were persistent for longer periods under cooler environments as  
388 compared to higher temperature environments, or that the temperatures were below the optimum  
389 development temperatures for this pest, or that the *T. castaneum* strain in Harare is not resistant  
390 to Shumba Super dust. Pesticide active ingredients have variable persistence periods in different  
391 grain ecosystems (Afridi et al., 2001).

392 An increase in resistance to pesticides, particularly to organophosphates, has previously been  
393 reported for *R. dominica*, *T. castaneum* and *Sitophilus* spp. in Brazil (Lorini and Galley, 1999)  
394 and some parts of Australia (Collins et al., 2017). Some *R. dominica* strains resistant to  
395 deltamethrin were found in Taiwan and the resistance was attributed to selection pressure (Chen  
396 and Chen, 2013). Pesticide resistance in insect pests is achieved through the death of  
397 homozygous susceptible individuals within the treated grain, leaving homozygous resistant and  
398 heterozygous susceptible individuals (Hagstrum and Subramanyam, 2006) which may require  
399 increased dosages or different control treatments to effect kill. As a result of ineffective control,  
400 farmers often resort to increased application rates and/or frequencies with some resorting to  
401 using non-recommended pesticides e.g. the use of carbaryl on stored grain, which is normally  
402 used for controlling cotton pests in Mbire district.

403 The results of this experiment challenge the rationale behind blanket pesticide application  
404 recommendations across different climatic regions. Pesticides generally degrade faster under  
405 high temperatures (Katagi, 2004) due to increased photodegradation, and given the same  
406 application rates greater efficacy is expected in small grains than in large grains due to the  
407 smaller intergranular spaces which do not allow insects to so easily escape from areas in the  
408 grain bulk where pesticide dust concentrations are high (Athanasios et al., 2003, Rumbos et al.,  
409 2016). However, other factors such as the texture of the grain surface also play a critical role in  
410 pesticide adherence and retention (Kabir et al., 2013; Kavallieratos et al., 2005) as the pesticides  
411 adhere more on rough grains.

412 The significant increase in grain damage levels at four months' storage in Mbire, largely during  
413 the 2014/15 storage season, could reflect a decrease in pesticide efficacy to below potent levels,  
414 enabling pest re-infestation (Sallam, 2008). Chikwapuro, Ngwena yedura and Shumba Super

415 dust suppressed grain damage and subsequent weight loss throughout the entire storage season in  
416 Harare. This could be due to high persistence of these pesticides under the relatively low  
417 temperature environments which prevailed in Harare. The superior performance of Shumba  
418 Super dust against sorghum pests in Harare as opposed to in Mbire shows the possible effects of  
419 differences in climatic conditions and dominant storage insect pest species on pesticide  
420 effectiveness. The failure of Shumba Super dust pesticides in suppressing grain damage in Mbire  
421 was also recorded against maize pests in the same district. An increase in percentage grain  
422 damage in Actellic Gold dust and Super guard dust and the development of *S. cerealella* in these  
423 treatments during the 2015/16 storage season in both Mbire and Harare indicates a high level of  
424 insecticide tolerance by this insect pest.

425 Low levels of grain damage in Mbire can be attributed to the high temperatures which  
426 suppressed many insect pests, with the exception of *T. castaneum* (Fleurat-Lessard and Dupuis,  
427 2010; Wilches et al., 2016). When temperatures go beyond the optimum range of most of the  
428 pest species capable of damaging whole grains, lower grain damage is likely to occur (Fleurat-  
429 Lessard and Dupuis, 2010). High temperature-related insect mortality and suppression in Mbire  
430 district can explain why grain damage was low for a longer period of time in all treatments in  
431 Mbire (16 weeks), as compared to Harare (8 weeks). However, beyond 32 weeks storage in  
432 Mbire all the pesticides tested failed to prevent grain damage increases, indicating a loss of  
433 potency due to the high temperatures (Palikhe, 2007, Ismail, 2012).

434 Temperatures were much higher (up to 39 °C) and grain moisture content was lower (9 - 10 %  
435 mc) in the store room in Mbire district than in Harare during the 2014/15 and 2015/16 storage  
436 seasons. The rise in grain moisture content in Super guard dust and untreated control in Harare  
437 may have been linked to high infestations of storage insects. Temperature and relative humidity

438 (local environmental conditions) play a critical role in influencing grain moisture content, pest  
439 development (Musolin and Saulich, 2012; Bendito and Twomlow, 2015), and influencing  
440 pesticide persistence and success (Katagi, 2004; Ismail et al., 2012; Bell, 2014). If the pesticides  
441 degrade beyond potent levels, re-infestation will occur (Afridi et al., 2001). The high  
442 temperatures ( $> 30\text{ }^{\circ}\text{C}$ ) experienced in Mbire and low grain moisture content of  $\sim 9\%$  can be  
443 catastrophic for most storage insects, with the exception of species such as *T. castaneum* and the  
444 bostrichids which can tolerate high temperatures (Fields et al., 2015). Most storage insects favour  
445 minimum relative humidity of  $50\%$  for optimum survival (Bell, 2014), and their optimum  
446 temperatures for survival and development are narrow, and vary from species to species. For  
447 every  $2\text{ }^{\circ}\text{C}$  increase in temperature from the minimum developmental temperature of any pest,  
448 the rate of development doubles until optimum temperatures are reached (Subramanyam *et al.*,  
449 1991; Sharma and Prabhakar, 2014). The low grain moisture content in samples collected from  
450 Mbire district is likely to have led to suppressed insect development and grain damage levels in  
451 that location, particularly during the second season. Most storage insects favour high moisture  
452 content when temperatures are above their optimum, as was found by Bell (2014) in  
453 *S. granarius*. Increasing grain moisture in non-effective pesticide treatments can be attributed to  
454 insect respiration and presence of immature stages of insects within the grains.

455 The high temperatures, as experienced in Mbire district and the general rising regional  
456 temperatures in SSA, may cause the emergence of new pest species in new sites and/ or  
457 extinction of some storage insect species (Karuppaiah and Sujayanad, 2012) shorter pest  
458 lifecycles, more rapid build-up of pests in stored commodities and in the rate of field to store  
459 contamination, in addition to causing accelerated pesticide breakdown (Stathers et al., 2013).  
460 Warming temperatures expand insect overwintering areas and volatinism (Karuppaiah and

461 Sujayanad, 2012; Sharma and Prabhakar, 2014) and may modify insect behaviour and dispersal  
462 (Bell, 2014). This may require more frequent pesticide treatment (Sharma and Prabhakar, 2014)  
463 Under accelerated pest development, and increased pesticide degradation due to higher  
464 temperatures resulting from global warming, farmers may resort to increased pesticide  
465 applications to effect control (Gatto et al., 2014; Delcour et al., 2015). It is therefore, imperative  
466 that climate-smart agricultural investments across SSA build the understanding of smallholder  
467 farmers and their service providers, about these likely impacts and the many postharvest  
468 adaptation strategies which can be adopted (Stathers et al., 2013; Mvumi and Stathers, 2014), to  
469 help avoid situations of increased losses of stored grains and/or increased pesticide use and the  
470 associated food safety risks.

471 The findings of this study show that climatic conditions may influence the success or failure of  
472 grain protectant dusts. The recommended application rate of Super guard dust failed to suppress  
473 a build-up of *S. cerealella* in Harare, while Shumba Super dust was ineffective against  
474 *T. castaneum* at both sites. The dominant pest species plays a major role in determining pesticide  
475 efficacy. Further investigations are required to examine the exact nature of the pesticide failure  
476 to control *S. cerealella* and *T. castaneum* and how this is mediated by temperature and how  
477 natural enemies such as predators and parasitoids come into play.

478 The development and levels of damage caused by these insect pests, alongside the potency of the  
479 pesticide, are also influenced by the prevailing climatic conditions. *Tribolium castaneum*  
480 dominated the pest complex in the hotter climate (Mbire), while *S. cerealella* and *S. oryzae*  
481 dominated in the cooler environment (Harare). Chikwapuro and Ngwena yedura were effective  
482 at both sites, while Shumba Super dust and Actellic Gold dust were only effective in Harare and  
483 Mbire, respectively. This study suggests the current blanket application rate recommendations



484 for grain protectants are insufficient, and there is a need to generate more context-specific  
485 pesticide recommendations based on both the expected dominant pest species, and the expected  
486 climatic conditions.

#### 487 **Acknowledgements**

488 The authors are grateful to the European Union (EU) for funding the project “*Supporting*  
489 *smallholder farmers in southern Africa to better manage climate-related risks to crop production*  
490 *and post-harvest handling*” through the contribution agreement DCI-Food/2012/304-807, under  
491 the leadership of the Food and Agriculture Organization (FAO) of the United Nations. National  
492 extension services in Mbire district are appreciated for their active engagement and contributions  
493 in day-to-day management of the experiment. Gratitude is also extended to Dr Susan  
494 Richardson-Kageler (Department of Crop Science, University of Zimbabwe) for assistance with  
495 statistical analysis. The Department of Crop Science is acknowledged for providing space where  
496 the experiment was conducted. Mention of a trademark or proprietary product does not constitute  
497 a guarantee or warranty of the product by the University of Zimbabwe or Natural Resources  
498 Institute and does not imply its approval to the exclusion of other products that may also be  
499 suitable.

500

#### 501 **References**

502 Afridi, I. A., Parveen, Z. & Masud, S. Z. 2001. Stability of organophosphate and pyrethroid  
503 pesticides on wheat in storage. *Journal of Stored Products Research*, 37, 199-204.

504 Akter, T., Jahan, M. & Bhuiyan, M. 2013. Biology of the angoumois grain moth, *Sitotroga*  
505 *Cerealella* (Oliver) on stored rice grain in laboratory condition. *Journal of the Asiatic*  
506 *Society of Bangladesh, Science*, 39, 61-67.

507 APHLIS, 2014. Estimated Postharvest Losses 2003 – 2015. Available at  
508 [http://www.aphlis.net/?form=losses\\_estimates](http://www.aphlis.net/?form=losses_estimates) (accessed on 14-11-2016).

509 Arthur, F. H. 1996. Grain protectants: Current status and prospects for the future. *Journal of*  
510 *Stored Products Research*, 32(4), 293-302

511 Athanassiou, C., Kavallieratos, N., Tsaganou, F., Vayias, B., Dimizas, C. & Buchelos, C. T.  
512 2003. Effect of grain type on the insecticidal efficacy of SilicoSec against *Sitophilus*  
513 *oryzae* (L.)(Coleoptera: Curculionidae). *Crop Protection*, 22, 1141-1147.

514 Athanassiou, C.G., Arthur, F.H. & Throne, J.E., 2009. Efficacy of spinosad in layer-treated  
515 wheat against five stored-product insect species. *Journal of Stored Products Research*,  
516 45, 236–240.

517 Baldassari, N., Berluti, A., Martini, A. & Baronio, P. 2004. Analysis of the sensitivity of  
518 different stages of *Rhyzopertha dominica* and *Tribolium castaneum* to diatomaceous  
519 earth. *Bulletin of Insectology*, 57, 95-102.

520 Baker, J.E. & Weaver, D.K., 1993. Resistance in field strains of the parasitoid *Anisopteromalus*  
521 *calandrae* (Hymenoptera: Pteromalidae) and its host, *Sitophilus oryzae* (Coleoptera:  
522 Curculionidae), to malathion, chlorpyrifos-methyl, and pirimiphos-methyl. *Biological*  
523 *control*, 3(3), 233-242.

524 Bartlett, A. M. S., Supplement, S. & Society, S., 1936. The square root transformation in analysis  
525 of variance. *Supplement for the Royal Statistical Society*, 3, 68-78

526 Bell, C.H., 2014. A review of insect responses to variations encountered in the managed storage  
527 environment. *Journal of Stored Product Research* 59, 260–274.

528 Bendito, A. & Twomlow, S. 2015. Promoting climate smart approaches to post-harvest  
529 challenges in Rwanda. *International Journal of Agricultural Sustainability*, 13, 222-239.

530 Boxall, R. 1986. A critical review of the methodology for assessing farm-level grain losses after  
531 harvest (G191). Tropical Development and Research Institute. London, Britain

532 Chen, C. Y. & Chen, M. E. 2013. Susceptibility of field populations of the lesser grain borer,  
533 *Rhyzopertha dominica* (F.), to deltamethrin and spinosad on paddy rice in Taiwan.  
534 *Journal of Stored Products Research*, 55, 124-127.

535 Chigoverah, A. A. & Mvumi, B. M. 2016. Efficacy of metal silos and hermetic bags against  
536 stored-maize insect pests under simulated smallholder farmer conditions. *Journal of*  
537 *Stored Products Research*, 69, 179-189.

538 Collins, P. J., Falk, M. G., Nayak, M. K., Emery, R. N. & Holloway, J. C. 2017. Monitoring  
539 resistance to phosphine in the lesser grain borer, *Rhyzopertha dominica*, in Australia: A  
540 national analysis of trends, storage types and geography in relation to resistance  
541 detections. *Journal of Stored Products Research*, 70, 25-36.

542 Costa, S. 2014. Reducing food losses in sub-Saharan Africa (improving post-harvest  
543 management and storage technologies of smallholder farmers.)—an ‘action  
544 research’ evaluation trial from Uganda and Burkina Faso. *UN World Food Programme*,  
545 *Kampala, Uganda*.

546 De Groote, H., Kimenju, S. C., Likhayo, P., Kanampiu, F., Tefera, T. & Hellin, J., 2013.  
547 Effectiveness of hermetic systems in controlling maize storage pests in Kenya. *Journal of*  
548 *Stored Products Research* 53, 27–36.

549 Delcour, I., Spanoghe, P. & Uyttendaele, M., 2015. Literature review: Impact of climate change  
550 on pesticide use. *Food Research International*, 68, 7–15.

551 De Muth, J. E. 2014. *Basic Statistics And Pharmaceutical Statistical Applications*, 3<sup>rd</sup> edition.  
552 CRC Press, New York.

553 Fields, P., Subramanyam, B. & Hulasare, R. 2015. Extreme temperatures. In D. W. Hagstrum,  
554 T. W. Phillips, & G. W. Cuperus (Eds.), *Stored Product Protection* (pp. 179-190).  
555 Manhattan, KS: Kansas State University.

556 Fleurat-Lessard, F. & Dupuis, S. A. 2010. Comparative analysis of upper thermal tolerance and  
557 CO<sub>2</sub> production rate during heat shock in two different European strains of *Sitophilus*  
558 *zeamais* (Coleoptera: Curculionidae). *Journal of Stored Products Research*, 46, 20-27.

559 Gatto, M.P., Cabella, R. & Gherardi, M., 2014. Climate change: the potential impact on  
560 occupational exposure to pesticides. *Annali dell'Istituto Superiore di Sanità*, 52, 374–385.

561 Gornall, J., Betts, R., Burke, E., Clark, R., Camp, J., Willett, K. & Wiltshire, A. 2010.  
562 Implications of climate change for agricultural productivity in the early twenty-first  
563 century. *Philosophical Transactions of the Royal Society of London B: Biological*  
564 *Sciences*, 365, 2973-2989.

565 Hagstrum, D. W. & Subramanyam, B. 2006. *Fundamentals of Stored-Product Entomology*,  
566 American Association of Cereal Chemists, Inc (AACCC). St. Paul, Minnesota, USA

567 IPCC, 2014: Climate Change 2014: Synthesis Report. *Contribution of Working Groups I, II and*  
568 *III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*  
569 *[Core Writing Team, R.K. Pachauri and L.A. Meyer (eds.)]*. IPCC, Geneva, Switzerland,  
570 151 pp.

571 Ismail, B., Mazlinda, M. & Zuriati, Z. 2012. Effects of temperature, soil moisture content and  
572 soil type on the degradation of cypermethrin in two types of Malaysian agricultural soils.  
573 *World Applied Sciences Journal*, 17, 428-432.

574 Kaminski, J. & Christiaensen, L. 2014. Post-harvest loss in sub-Saharan Africa—what do  
575 farmers say? *Global Food Security*, 3, 149-158.

576 Karuppaiah, V., Sujayanad, G.K., 2012. Impact of climate change on population dynamics of  
577 insect pests. *World Journal of Agricultural Science* 8, 240–246

578 Katagi, T. 2004. Photodegradation of pesticides on plant and soil surfaces. *Reviews of*  
579 *Environmental Contamination And Toxicology*. Springer.

580 Lorini, I. & Galley, D. 1999. Deltamethrin resistance in *Rhyzopertha dominica* (F.)(Coleoptera:  
581 Bostrichidae), a pest of stored grain in Brazil. *Journal of Stored Products Research*, 35,  
582 37-45.

583 Makundi, R.H., Swila, N.N., Misangu, R.N., Reuben, S.W., Mwatawala, M., Sikira, A., Kilonzo,  
584 B.S., Lyimo, H., Massawe, A.W. & Ishengoma, C., 2010. Dynamics of infestation and  
585 losses of stored maize due to the larger grain borer (*Prostephanus truncatus* Horn) and  
586 maize weevils (*Sitophilus zeamais* Motschulsky). *Archives of Phytopathology and Plant*  
587 *Protection*, 43 (14), 1346-1355.

588 Mason, L. J. & Mcdonough, M. 2012. Biology, behavior, and ecology of stored grain and  
589 legume insects. *Stored product protection*, 7. Kansas State University.

590 Mlambo, S., Mvumi, B. M., Stathers, T., Mubayiwa, M. & Nyabako, T., 2017. Field efficacy of  
591 hermetic and other maize grain storage options under smallholder farmer management.  
592 *Crop Protection*, 98, 198-210

593 Mvumi, B.M. & Stathers, T.E. 2014. Food security challenges in Sub-Saharan Africa: The  
594 potential contribution of postharvest skills, science and technology in closing the gap.  
595 pp. 32-43 In Arthur, F.H, Kengkanpanich, R., Chayaprasert, W. & Suthisut, D. (Eds.),  
596 Proceedings of the 11<sup>th</sup> International Working Conference on Stored Product Protection.  
597 24-28 November 2014, Chiang Mai, Thailand.

598 Musolin, D. & Saulich, A. K. 2012. Responses of insects to the current climate changes: from  
599 physiology and behavior to range shifts. *Entomological Review*, 92, 715-740.

600 Neven, L. 2000, Physiological response of insects to heat. *Postharvest Biology and Technology*,  
601 21, 103-111

602 Niang, I., Ruppel O. C., Abdrabo M. A., Essel A., Lennard C., Padgham J. & Urquhart P. 2014.  
603 In: Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional  
604 Aspects. Contribution of Working Group II to the Fifth Assessment Report of the  
605 Intergovernmental Panel on Climate Change [Barros, V.R., Field C. B., Dokken D.J.,  
606 Mastrandrea M.D., Mach K.J., Bilir T.E., Chatterjee M., Ebi K.L., Estrada Y.O., Genova  
607 R.C., Girma B., Kissel E.S., Levy A.N., MacCracken S., Mastrandrea P.R., and White  
608 L.L. (eds.)]. *Cambridge University Press, Cambridge, United Kingdom and New York,*  
609 *NY, USA, 1199-1265.*

610 Oppert, B., Ellis, R. T. & Babcock, J. 2010. Effects of Cry1F and Cry34Ab1/35Ab1 on storage  
611 pests. *Journal of Stored Products Research*, 46, 143-148.

612 Palikhe, B. R. 2007. Relationship between pesticide use and climate change for crops. *Journal of*  
613 *Agriculture and Environment*, 8, 83-91.

614 Pretty, J. N. 2012. *The pesticide detox: towards a more sustainable agriculture*, Earthscan.

615 Rigaux, M., Haubruge, E. & Fields, P. G. 2001. Mechanisms for tolerance to diatomaceous earth  
616 between strains of *Tribolium castaneum*. *Entomologia Experimentalis et Applicata*, 101:  
617 33-39

618 Rumbos, C. I., Dutton, A. C. & Athanassiou, C. G. 2016. Insecticidal efficacy of two pirimiphos-  
619 methyl formulations for the control of three stored-product beetle species: Effect of  
620 commodity. *Crop Protection*, 80, 94-100.

621 Sallam, M. N. 2008. Insect damage: damage on post-harvest. *AGSI/FAO: INPhO*. Available via  
622 <http://www.fao.org/inpho/content/compend/text/ch02-01.htm>. Accessed, 22.08.2017

623 Serdeczny, O., Adams, S., Baarsch, F., Coumou, D., Robinson, A., Hare, W., Schaeffer, M.,  
624 Perrette, M. & Reinhardt, J. 2016. Climate change impacts in Sub-Saharan Africa: from  
625 physical changes to their social repercussions. *Regional Environmental Change*, 15 (8):  
626 1-16

627 Sharma, H.C. & Prabhakar, C.S., 2014. Impact of climate change on pest management and food  
628 security. *Integrated Pest Management*, 2, 23–36.

629 Stathers, T., Lamboll, R. & Mvumi, B. M. 2013. Postharvest agriculture in changing climates: its  
630 importance to African smallholder farmers. *Food Security*, 5, 361-392.

631 Stathers, T., Mvumi, B. & Golob, P. 2002. Field assessment of the efficacy and persistence of  
632 diatomaceous earths in protecting stored grain on small-scale farms in Zimbabwe. *Crop*  
633 *Protection*, 21, 1033-1048.

634 Subramanyam, B. & Hagstrum, D. 1991. Quantitative analysis of temperature, relative humidity,  
635 and diet influencing development of the larger grain borer, *Prostephanus truncatus*  
636 (Horn)(Coleoptera: Bostrichidae). *International Journal of Pest Management*, 37, 195-  
637 202.

- 638 Trematerra, P. 2015. Adult dispersal of *Sitotroga cerealella* in a conventional small-farm in  
639 Southern Italy. *Bulletin of Insectology*, 68, 111-118.
- 640 Vassilakos, T., Athanassiou, C. & Tsiropoulos, N. 2015. Persistence and efficacy of spinetoram  
641 against three major stored grain beetle on wheat. *Crop Protection*, 69, 44-51.
- 642 Velázquez-Fernández, J. B., Martínez-Rizo, A. B., Domínguez-Ojeda, D. & Ramírez-Sandoval,  
643 M. 2012. *Biodegradation And Bioremediation Of Organic Pesticides*, INTECH Open  
644 Access Publisher,12, 1-21. Rijeka, Croatia
- 645 Wilches, D., Laird, R. A., Floate, K. D. & Fields, P. 2016. A review of diapause and tolerance to  
646 extreme temperatures in dermestids (Coleoptera). *Journal of Stored Products Research*,  
647 68, 50-62.
- 648 World Bank, 2011. Missing Food :The Case of postharvest Grain Losses in Sub-Saharan African.  
649 world Bank 60371–AFR, 116. doi:Report No. 60371-AFR



## Highlights

- The efficacy of the synthetic pesticides varied across the different agro-climatic sites
- *Sitotroga cerealella* and *Sitophilus oryzae* were dominant in the cooler climatic area
- *Tribolium castaneum* was the dominant pest in the hotter climatic area
- *Tribolium castaneum* displayed tolerance to fenitrothion 1% + deltamethrin 0.13% combination