



## LJMU Research Online

**Boddy, LM, Noonan, RJ, Kim, Y, Rowlands, A, Welk, G, Knowles, ZR and Fairclough, S**

**Comparability of children's sedentary time estimates derived from wrist worn GENEActiv and hip worn ActiGraph accelerometer thresholds**

<http://researchonline.ljmu.ac.uk/8357/>

### Article

**Citation** (please note it is advisable to refer to the publisher's version if you intend to cite from this work)

**Boddy, LM, Noonan, RJ, Kim, Y, Rowlands, A, Welk, G, Knowles, ZR and Fairclough, S (2018) Comparability of children's sedentary time estimates derived from wrist worn GENEActiv and hip worn ActiGraph accelerometer thresholds. Journal of Science and Medicine in Sport. 21 (10). pp. 1045-**

LJMU has developed [LJMU Research Online](http://researchonline.ljmu.ac.uk) for users to access the research output of the University more effectively. Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Users may download and/or print one copy of any article(s) in LJMU Research Online to facilitate their private study or for non-commercial research. You may not engage in further distribution of the material or use it for any profit-making activities or any commercial gain.

The version presented here may differ from the published version or from the version of the record. Please see the repository URL above for details on accessing the published version and note that access may require a subscription.

For more information please contact [researchonline@ljmu.ac.uk](mailto:researchonline@ljmu.ac.uk)

<http://researchonline.ljmu.ac.uk/>

1 **Title page**

2 Title: Comparability of children's sedentary time estimates derived from wrist worn GENEActiv and  
3 hip worn ActiGraph accelerometer thresholds

4 Lynne M. Boddy<sup>1</sup>, Robert J. Noonan<sup>1</sup>, Youngwon Kim<sup>2</sup>, Alex V. Rowlands<sup>3,4,5</sup> Greg J. Welk<sup>6</sup> Zoe R.  
5 Knowles<sup>1</sup>, Stuart J. Fairclough<sup>7,8</sup>

6 <sup>1</sup>Physical Activity Exchange, Research Institute for Sport and Exercise Sciences, Liverpool John Moores  
7 University, Liverpool, UK

8 <sup>2</sup> MRC Epidemiology Unit, University of Cambridge School of Clinical Medicine, Cambridge  
9 Biomedical Campus. Cambridge, UK

10 <sup>3</sup>Diabetes Research Centre, University of Leicester, Leicester General Hospital, Leicester, UK;

11 <sup>4</sup>NIHR Leicester Biomedical Research Centre, UK;

12 <sup>5</sup>Alliance for Research in Exercise, Nutrition and Activity (ARENA), Sansom Institute for Health  
13 Research, Division of Health Sciences, University of South Australia, Adelaide, Australia.

14 <sup>6</sup>Department of Kinesiology, College of Human Sciences, Iowa State University, Ames, IA

15 <sup>7</sup>Department of Sport and Physical Activity, Edge Hill University, Ormskirk, UK

16 <sup>8</sup>Department of Physical Education and Sport Sciences, University of Limerick, Limerick, Ireland

17 Corresponding author details: Dr Lynne M. Boddy  
18 The Physical Activity Exchange,  
19 Research Institute for Sport & Exercise Sciences,  
20 Liverpool John Moores University,  
21 62 Great Crosshall Street,  
22 Liverpool,  
23 L3 2AT  
24 UK  
25 Email: [L.M.Boddy@ljmu.ac.uk](mailto:L.M.Boddy@ljmu.ac.uk)  
26 Tel: [+44 \(0\)151 231 4275](tel:+44(0)1512314275)

27 **Word count (excluding abstract and references): 2894**

28

29

30

31

32

33 Title: Comparability of children's sedentary time estimates derived from wrist worn GENEActiv and  
34 hip worn ActiGraph accelerometer thresholds  
35

36 **Abstract:**

37 Objectives: to examine the comparability of children's free-living sedentary time (ST) derived from raw  
38 acceleration thresholds for wrist mounted GENEActiv accelerometer data, with ST estimated using the  
39 waist mounted ActiGraph 100 count·min<sup>-1</sup> threshold.

40 Design: Secondary data analysis

41 Method: 108 10-11-year-old children (n=43 boys) from Liverpool, UK wore one ActiGraph GT3X+  
42 and one GENEActiv accelerometer on their right hip and left wrist, respectively for seven days. Signal  
43 vector magnitude (SVM; mg) was calculated using the ENMO approach for GENEActiv data. ST was  
44 estimated from hip-worn ActiGraph data, applying the widely used 100 count·min<sup>-1</sup> threshold. ROC  
45 analysis using 10-fold hold-out cross-validation was conducted to establish a wrist-worn GENEActiv  
46 threshold comparable to the hip ActiGraph 100 count·min<sup>-1</sup> threshold. GENEActiv data were also  
47 classified using three empirical **wrist** thresholds and equivalence testing was completed.

48 Results: Analysis indicated that a GENEActiv SVM value of 51mg demonstrated fair to moderate  
49 agreement (Kappa: 0.32-0.41) with the 100 count·min<sup>-1</sup> threshold. **However**, the generated and empirical  
50 thresholds for GENEActiv devices were not significantly equivalent to ActiGraph 100 count·min<sup>-1</sup>.  
51 GENEActiv data classified using the 35.6 mg threshold intended for ActiGraph devices generated  
52 significantly equivalent ST estimates as the ActiGraph 100 count·min<sup>-1</sup>.

53 Conclusions: The newly generated and empirical GENEActiv **wrist** thresholds do not provide equivalent  
54 estimates of ST to the ActiGraph 100 count·min<sup>-1</sup> approach. More investigation is required to assess the  
55 validity of applying ActiGraph cutpoints to GENEActiv data. Future studies are needed to examine the  
56 backward compatibility of ST data and to produce a robust method of classifying SVM-derived ST.

57 Keywords: children, physical activity, inactivity, accelerometry, measurement

## 58 **Introduction**

59 Sedentary behaviour is increasingly viewed as an important health risk factor in children <sup>1</sup>, and the  
60 detrimental effects of reallocating PA time to sedentary behaviours have been established <sup>2</sup>. Sedentary  
61 behaviour is defined as ‘any waking behaviour characterized by an energy expenditure  $\leq 1.5$  METS  
62 while in a sitting, reclining or lying posture’ <sup>3</sup> however for children the recommended upper boundary  
63 of energy expenditure is  $\leq 2$  METs or  $\leq 1.5$  child-METs <sup>4</sup>. It is common for researchers to assess  
64 sedentary time (ST) which is commonly defined as the time spent below the threshold of proprietary  
65 accelerometer counts representing light physical activity, rather than focussing on sedentary behaviour  
66 per se.

67 Accelerometers have been used for several years to quantify children’s ST, but heterogeneous data  
68 processing and researcher decisions related to for example, device location, wear time criteria, and  
69 choice of thresholds, often mean that study methods lack consistency and comparability. The advent of  
70 newer accelerometer devices capable of raw acceleration data collection removes the reliance on  
71 proprietary counts and allows researchers more autonomy when examining data, whilst producing  
72 estimates of acceleration that in theory should be comparable between devices <sup>5</sup>. Therefore, devices that  
73 produce raw acceleration data for researchers to use, such as the GENEActiv and ActiGraph GT3X+  
74 offer an opportunity to increase comparability between studies aiming to estimate ST using  
75 accelerometers.

76 Raw acceleration data from GENEActiv and ActiGraph accelerometers are increasingly being processed  
77 in the open source R package GGIR (<http://cran.r-project.org>). GGIR auto-calibrates the data using local  
78 gravity as a reference <sup>6</sup>, detects sustained abnormally high values and generates the average magnitude  
79 of dynamic acceleration (termed the Euclidean Norm Minus One (ENMO))<sup>5, 7-9</sup>. Recently, the ENMO  
80 metric has been used to estimate ST and physical activity in both children and adults<sup>9-11</sup>, but significant  
81 differences have been reported for ST and PA estimated from counts and from raw acceleration signals.  
82 Authors have attributed these differences to the various intensity thresholds used to classify acceleration  
83 data across the reduction approaches and differences in wear-site <sup>11</sup>, but they may also be due to the  
84 inherent differences between the proprietary counts and raw acceleration data. One recent study,

85 conducted in children, provided a method of calibrating raw acceleration data from wrist-worn monitors  
86 to counts based hip-worn physical activity estimates in an effort to harmonise data <sup>9</sup>. The study classified  
87 raw accelerations using a range of ENMO thresholds for wrist-worn monitors and aligned these to  
88 counts-based thresholds for hip-worn monitors, demonstrating that incremental thresholds enable simple  
89 group level comparisons to past estimates of physical activity derived from hip-worn accelerometer  
90 counts cutpoints. For traditional accelerometer counts-based protocols using hip-worn ActiGraphs,  
91 studies have widely adopted 100 vertical axis count·min<sup>-1</sup> as the upper threshold for ST in children <sup>12</sup>.  
92 To date, the comparability of wrist-worn GENEActiv ENMO ST estimates to those generated using the  
93 ActiGraph 100 vertical axis count·min<sup>-1</sup> method is unknown. Studies have utilised the ENMO regression  
94 equation published by Hildebrand et al. <sup>8</sup> which was generated using a laboratory protocol to classify  
95 ST, however, these thresholds have not been cross-validated for classifying ST or examined in  
96 comparison with other methods. More recent studies <sup>13</sup> utilised the Hildebrand et al. <sup>8</sup> laboratory  
97 protocol to general thresholds then examined the agreement between ST and activPAL (which was  
98 considered as a criterion reference standard measure) using free-living data. The thresholds  
99 demonstrated low specificity, overestimating sedentary time in comparison to the activPAL. The  
100 equivalence of wrist worn data classified using these approaches to the 100 count·min<sup>-1</sup> standard is  
101 unknown. Therefore, researchers wishing to represent raw accelerations through ENMO cannot compare  
102 ST to previous counts-based research, and so a pragmatic solution to classifying ST is required.

103 The aims of this secondary data analysis were to examine the comparability of children's free-living ST  
104 derived using the ENMO metric for wrist mounted GENEActiv accelerometer data, with ST estimated  
105 using the waist mounted ActiGraph 100 count·min<sup>-1</sup> threshold. This aim was addressed by examining,  
106 [1] if comparable ST estimates could be attained from wrist-mounted GENEActiv raw acceleration data  
107 anchored to the widely adopted 100 count·min<sup>-1</sup> uniaxial hip-mounted ActiGraph ST threshold, and [2]  
108 the equivalence of ST estimates between the newly generated threshold, those published by Hildebrand  
109 et al. <sup>8,13</sup> and the 100 count·min<sup>-1</sup> uniaxial hip-mounted ActiGraph ST threshold.

110

111 METHODS

112 This is a secondary data analysis of data generated by a previous study <sup>7</sup>. After gaining University ethics  
113 approval, informed parental consent, and participant assent 108 10-11-year-old children (n=43 boys)  
114 were involved in this study. Data collection took place on school sites from January to May 2014. Stature  
115 and body mass were assessed to the nearest 0.1cm using a portable stadiometer (Leicester Height  
116 Measure, Seca, Birmingham, UK) and nearest 0.1kg (Seca, Birmingham, UK) respectively using  
117 standard techniques <sup>14</sup>. Body mass index (BMI), was calculated for each participant.

118 Sedentary time was assessed using two tri-axial accelerometers, one worn on the non-dominant wrist  
119 (GENEActiv; Activinsights, Cambs, UK) and one worn on the right hip (ActiGraph GT3X+; ActiGraph,  
120 Pensacola, FL). Both monitors were initialised using the same computer to record at a frequency of 100  
121 Hz, and participants were asked to wear the monitors at all times for 7 consecutive days except when  
122 sleeping and engaging in water based activities (e.g., bathing, swimming).

123 ActiGraph monitors were analysed using ActiLife v 6.11.4 software (ActiGraph, Pensacola, FL).  
124 Twenty minutes of consecutive zero counts (1 minute spike tolerance) defined non-wear time, and these  
125 periods were subtracted from daily wear time <sup>15</sup>. Sedentary time was coded as  $\leq 100$  count $\cdot$ min<sup>-1</sup> <sup>12</sup>. Valid  
126 days were defined as  $\geq 540$  min for a weekday <sup>16</sup> and  $\geq 480$  min for weekend days <sup>17</sup>. For each participant  
127 the valid weekday and weekend day with the longest wear time were selected and retained for analysis.  
128 For participants with no valid weekend data, the valid weekday only with the longest wear time was  
129 included within analysis. After establishing daily wear time, data for the included days were converted  
130 to 1-s epoch csv output files for further analysis.

131 GENEActiv data were downloaded using GENEActiv v 2.2 software (Activinsights, Cambs, UK) and  
132 saved as binary files. These were then processed in R (<http://cran.r-project.org>) using the GGIR package  
133 (version 1.1-4). To correct for sensor calibration error autocalibration was completed <sup>6</sup>. GGIR processing  
134 produced files in csv format. Each csv file contained the ENMO-derived average magnitude of dynamic  
135 acceleration values expressed in average mg <sup>18</sup>. GENEActiv csv files corresponding to the selected  
136 ActiGraph weekday and/or weekend days were taken forward to the next stage of analysis.

137 ActiGraph and GENEActiv time stamped data were synched, resulting in one csv file for each  
138 participant containing date- and time-stamped ActiGraph and GENEActiv data in 1 s epochs. Non-wear  
139 times were removed from each merged file according to the ActiLife wear time details generated for  
140 each participant's ActiGraph data. For the ROC analysis each participant's ActiGraph and GENEActiv  
141 data were then summed into 1 min epochs to allow data scoring using the ActiGraph vertical axis 100  
142 count·min<sup>-1</sup> as the reference value for sedentary time<sup>12</sup>. These data were then stacked into one csv file  
143 to create a dataset including all participants (n = 108, 43 boys).

144 To establish GENEActiv classification criteria anchored to the ActiGraph 100 count·min<sup>-1</sup> ST threshold,  
145 ROC analysis was performed on the whole sample, which represented 126,999 minutes of monitor wear  
146 time. Threshold values were cross-validated using 10-fold hold-out groups stratified by sex<sup>19</sup>, whereby  
147 separate cross-validation analyses were conducted with a randomly selected hold-out group for each  
148 iteration (11 participants [6 girls and 5 boys] per analysis cycle)<sup>20</sup>. Therefore, each ROC analysis was  
149 completed with 97 participants with 11 excluded to enable cross-validation. For each hold-out group  
150 2x2 contingency tables were used to check classification agreement based on the GENEActiv  
151 classifications generated from each cross-validation ROC analysis. Computed sensitivity and specificity,  
152 Cohen's kappa coefficients, and percentage agreement between classifications were assessed.

153  
154 After generating the classification threshold, ST data were scored using 1 minute epochs. Data were  
155 classified for each participant using the newly generated GENEActiv threshold, ActiGraph 100  
156 count·min<sup>-1</sup>. Additionally GENEActiv ST was scored using the solved regression equation published by  
157 Hildebrand et al<sup>8</sup>, where ST was defined as  $\leq 1.5$  child-METS<sup>4</sup>, resulting in a threshold of 22.6 mg.  
158 GENEActiv ST was also scored using the 56.3 mg GENEActiv and 35.6 mg ActiGraph thresholds from  
159 the Hildebrand et al. 2016 study<sup>13</sup>. The ActiGraph threshold was included as theoretically using the raw  
160 data methods should allow the application of the threshold to the GENEActiv device. Pairwise  
161 equivalence testing was completed between all combinations of the thresholds. For this study a 95%  
162 equivalence test was performed to examine whether the 90% confidence intervals for mean ST for each  
163 classification method completely fell within the proposed equivalence zone ( $\pm 10\%$  of the mean of ST)



164 defined by the other classification method, representing statistically significant equivalence.  
165 Equivalence testing has been increasingly used in recent PA research where differences testing is not  
166 appropriate <sup>11, 21-24</sup>. Difference testing provides information on whether two methods are statistically  
167 different, where in this context it is more useful to know whether two methods are statistically equivalent  
168 at the group level, thus providing similar estimates. Analyses were conducted using IBM SPSS Statistics  
169 v.22 (IBM, Armonk, NY) and Microsoft Excel 2010 (Microsoft, Redmond, WA) and R for Windows  
170 (<http://cran.r-project.org>).

171

172

### 173 Results

174 Mean anthropometric data, weekend and weekday accelerometer wear times and the number of days  
175 included within analysis for boys and girls are displayed in Table 1.

176

177 [TABLE 1 ABOUT HERE]

178

179 The ROC curve for the whole cohort (N = 108) indicated that a GENEActiv threshold of 51 mg  
180 (sensitivity = 81.2%, specificity = 57.4%, AUC 0.760, 95% CI = 0.758, 0.763) provided the most  
181 accurate classification of ST. The ROC generated cutpoints, sensitivity and specificity, agreement, and  
182 Kappa values for each hold-out analysis for ST can be viewed in supplementary material A. The hold-  
183 out analysis found that the ST ENMO threshold performed significantly better than random  
184 classification, with agreement ranging from 64.7-69.7% and Kappa values ranging from 0.32-0.41 (fair  
185 to moderate agreement <sup>25</sup>). The mean GENEActiv ST cutpoint generated was 51 mg, corresponding with  
186 the whole group threshold, therefore 51 mg was used for subsequent equivalence analysis.

187

188 Figure 1 displays the results of the equivalence testing using ActiGraph count·min<sup>-1</sup> as the reference  
189 threshold. Mean time spent in ST for each classification is displayed in supplementary file B. None of  
190 the 90% CIs for the newly generated GENEActiv 51mg (630.6-666.7 min), Hildebrand 2014 22.6 mg

191 (323.2-362 min) or Hildebrand 2016 GENEActiv 56.3mg (673.5-711.1 min) were completely included  
192 within the zone of equivalence for the ActiGraph 100 count·min<sup>-1</sup> (443.2-541.6 min), suggesting no  
193 statistically significant equivalence between the cut-points compared and the ActiGraph 100 count·min<sup>-1</sup>,  
194 on average. The Hildebrand ActiGraph 2016 35.6mg threshold, applied to GENEActiv data yielded  
195 90% CIs (492.9-527.5) that fell within the zone of equivalence, so is considered statistically equivalent  
196 to the GENEActiv, on average. The newly generated GENEActiv threshold ST estimates were, on  
197 average, significantly equivalent to the 2016 Hildebrand GENEActiv threshold, with the 90% CIs for  
198 the Hildebrand 2016 GENEActiv threshold of 56.3mg falling within the zone of equivalence for the  
199 threshold generated by our study (689.4-695.1 min, zone of equivalence 583.8-713.5 min). No other  
200 combinations exhibited statistically significant equivalence.

201

202 [FIGURE 1 ABOUT HERE]

203

204

205 **Discussion**

206 The aims of this secondary data analysis were to examine the comparability of children's free-living  
207 sedentary time (ST) derived from raw acceleration thresholds for wrist mounted GENEActiv  
208 accelerometer data, with ST estimated using the waist mounted ActiGraph 100 count·min<sup>-1</sup> threshold. A  
209 GENEActiv wrist ST threshold of 51 mg was generated which demonstrated fair to moderate agreement  
210 between the cross-validation and whole samples. The fact that the free-living data reflected a typical  
211 range of sedentary activities undertaken by children gave it a high degree of ecological validity.  
212 Irrespective of this, ST estimated using the 51 mg was not equivalent to the ActiGraph 100 count·min<sup>-1</sup>  
213 threshold and therefore is not an acceptable value to use to generate ST estimates from GENEActiv  
214 wrist accelerations that are compatible with estimates from waist-worn ActiGraphs. However, when  
215 applied to the GENEActiv data, the Hildebrand 35.6 mg ActiGraph wrist acceleration threshold  
216 produced significantly equivalent estimates of ST as the waist ActiGraph 100 count·min<sup>-1</sup> suggesting  
217 that this threshold could potentially be applied to GENEActiv data to provide comparable estimates of  
218 ST. Whether this provides an accurate estimate of ST when compared to criterion reference methods

219 such as activPAL warrants further investigation, however this was not the purpose of the analysis  
220 conducted.

221  
222 Field-based approaches to generating acceptable ST thresholds may be desirable because of their  
223 greater ecological validity, and because they may reduce the risk of misclassification associated with  
224 laboratory-derived thresholds being used in the field<sup>26</sup>. However, our findings suggest that the current  
225 thresholds used to classify ST using ENMO do not produce comparable estimates to those reported  
226 when using the standard 100 count·min<sup>-1</sup> approach. The challenges of estimating ST from wrist  
227 accelerometry are becoming more established<sup>27</sup>. Accelerometers are predominantly designed to  
228 measure movement rather than postural allocations. Accelerations from hip- and wrist-worn  
229 accelerometers are highly correlated in children during ST and physical activities of moderate through  
230 to vigorous intensities<sup>28</sup>. However, correlations are weaker during stationary light intensity physical  
231 activity which can involve a combination of sitting and standing activities, as well as transitions  
232 between the two<sup>28</sup>. Sitting and standing often encompass a combination of sedentary time and time in  
233 light intensity physical activity, whereby a high degree of hip and wrist acceleration decoupling occurs.  
234 For example, an individual may be sitting but gesturing with their hands, or standing and throwing a  
235 ball, both of which involve movements that a hip monitor may not detect but that could be detected by  
236 a wrist mounted device. This lack of consistency between hip and wrist accelerations during some  
237 sedentary and light intensity activities provides some explanation of the moderate levels of agreement  
238 observed in the cross-validation analyses, and the lack of equivalence with the hip 100 count·min<sup>-1</sup>  
239 threshold in particular.

240  
241 The accuracy of classifying ST is not explored in this study, we simply looked at the comparability of  
242 the GENEActiv thresholds to the standard ActiGraph vertical axis 100 count·min<sup>-1</sup> threshold. Whether  
243 the standard approach provides a more or less accurate estimate of ST is not examined and warrants  
244 further evaluation. To examine the accuracy of ST thresholds within a field-based protocol, a criterion  
245 measure, such as an inclinometer is needed. Theoretically this would increase participant burden through

246 the need to wear two devices, increase the cost of undertaking the research and data would still not allow  
247 for cross-comparisons between previous counts based studies. An alternative approach, that negates the  
248 need for additional devices, is to use accelerometers to examine assumed postural changes relative to  
249 arm elevation and wrist orientation (i.e., the Sedentary Sphere approach <sup>27</sup>). Recent evidence suggests  
250 that the Sedentary Sphere method provides comparable estimates of ST in adults when compared to the  
251 activPAL <sup>29</sup>, however, this method has not been validated in children, and so further work is required to  
252 examine its utility of this method in this population.

253  
254 The Hildebrand 22.6 mg ST threshold is based on GENEActiv wrist ENMO values, but was generated  
255 using VO<sub>2</sub> data rather than ActiGraph counts as in the current study. This may explain why the  
256 thresholds were not equivalent. In addition, the laboratory protocol used by Hildebrand et al.<sup>8, 13</sup> only  
257 included lying watching TV and sitting using a computer as sedentary activities <sup>8</sup>. Whilst such activities  
258 are common among children they do not reflect the wide range of free-living sedentary behaviours that  
259 the children involved in this study were likely to have engaged in. Further, the Hildebrand et al. (2014)  
260 22.6 mg estimated ST threshold was calculated from a regression equation anchored to energy  
261 expenditure. As sedentary behaviours are characterised by posture and low energy expenditure,  
262 determining sedentary time using energy expenditure alone without posture classification may be a less  
263 accurate approach than using criterion measures such as inclinometers or direct observation <sup>27</sup>.  
264 Hildebrand et al.'s 2016 GENEActiv 56.5mg wrist threshold was similar to our 51mg threshold, though  
265 the former demonstrated low specificity, overestimating sedentary time in comparison to the activPAL  
266 when examining free-living data, which may be due to the limited number of sedentary stations  
267 included in the original laboratory protocol.

268  
269 There are a number of limitations to this study. Our study was conducted in one geographical area of  
270 the UK and as such the results may not be representative of other populations. To classify GENEActiv  
271 data against the 100 count·min<sup>-1</sup> criterion, we used a 1-minute epoch setting. Though this would likely  
272 result in the inability to detect movement at higher intensities, as sedentary behaviour is characterised

273 by a lack of movement the 1-minute epoch setting would have less impact upon the ST estimates  
274 generated. We did not use a criterion reference standard device such as activPAL within this study. This  
275 was by design, as the primary aim was to examine the comparability of simple accelerometer estimates  
276 rather than investigate the accuracy of the measurement of ST. Future studies should aim to utilise the  
277 activPAL and other reference methods to develop and validate ST thresholds for use in children.

278

### 279 **Conclusions**

280 Despite displaying fair to moderate agreement, the generated GENEActiv ST threshold does not provide  
281 an equivalent estimate of ST to the hip mounted ActiGraph 100 count·min<sup>-1</sup> approach. Furthermore, ST  
282 data generated using Hildebrand thresholds were not equivalent to the 100 count·min<sup>-1</sup> method. Future  
283 studies are needed to examine the backwards compatibility of ST data and to produce a robust method  
284 of classifying ENMO-derived ST.

285

### 286 **Practical implications**

- 287 • Estimates of children's sedentary time generated from GENEActiv wrist ENMO and ActiGraph  
288 100 count·min<sup>-1</sup> are not comparable.
- 289 • Researchers should not compare data generated using the two different methods.
- 290 • Future studies are required to provide methods of data harmonization and to establish valid and  
291 reliable sedentary time thresholds for children.

292

293 **References**

294

- 295 1. Cliff DP, Hesketh KD, Vella SA, et al. Objectively measured sedentary behaviour and health and  
 296 development in children and adolescents: systematic review and meta-analysis. *Obesity*  
 297 *Reviews*. 2016; 17(4):330-344.
- 298 2. Fairclough SJ, Dumuid D, Taylor S, et al. Fitness, fatness and the reallocation of time between  
 299 children's daily movement behaviours: an analysis of compositional data. *Int J Behav Nutr Phys*  
 300 *Act*. 2017; 14(1):64.
- 301 3. Tremblay MS, Aubert S, Barnes JD, et al. Sedentary behavior network (SBRN)- Terminology  
 302 Consensus Project process and outcome. *International Journal of Behavioural Nutrition and*  
 303 *Physical Activity*. 2017; 14(75).
- 304 4. Saint-Maurice PF, Kim Y, Welk GJ, Gaesser GA. Kids are not little adults: what MET threshold  
 305 captures sedentary behavior in children? *European Journal of Applied Physiology*. 2016;  
 306 116(1):29-38.
- 307 5. Rowlands AV, Yates T, Davies M, Khunti K, Edwardson CL. Raw Accelerometer Data Analysis  
 308 with GGIR R-package: Does Accelerometer Brand Matter? *Med Sci Sports Exerc*. 2016.
- 309 6. van Hees VT, Fang Z, Langford J, et al. Autocalibration of accelerometer data for free-living  
 310 physical activity assessment using local gravity and temperature: an evaluation on four  
 311 continents. *Journal of applied physiology*. 2014; 117(7):738-744.
- 312 7. Fairclough SJ, Noonan R, Rowlands AV, VANH, Knowles Z, Boddy LM. Wear Compliance and  
 313 Activity in Children Wearing Wrist- and Hip-Mounted Accelerometers. *Med Sci Sports Exerc*.  
 314 2016; 48(2):245-253.
- 315 8. Hildebrand M, VANH, Hansen BH, Ekelund U. Age group comparability of raw accelerometer  
 316 output from wrist- and hip-worn monitors. *Med Sci Sports Exerc*. 2014; 46(9):1816-1824.
- 317 9. Rowlands AV, Cliff DP, Fairclough SJ, et al. Moving Forward with Backward Compatibility:  
 318 Translating Wrist Accelerometer Data. *Med Sci Sports Exerc*. 2016; 48(11):2142-2149.
- 319 10. Noonan RJ, Boddy LM, Kim Y, Knowles ZR, Fairclough SJ. Comparison of children's free-living  
 320 physical activity derived from wrist and hip raw accelerations during the segmented week. *J*  
 321 *Sports Sci*. 2017; 35(21):2067-2072.
- 322 11. Kim Y, Hibbing P, Saint-Maurice PF, et al. Surveillance of Youth Physical Activity and Sedentary  
 323 Behavior With Wrist Accelerometry. *Am J Prev Med*. 2017; 52(6):872-879.
- 324 12. Trost SG, Loprinzi PD, Moore R, Pfeiffer KA. Comparison of accelerometer cut points for  
 325 predicting activity intensity in youth. *Med Sci Sports Exerc*. 2011; 43(7):1360-1368.
- 326 13. Hildebrand M, Hansen BH, van Hees VT, Ekelund U. Evaluation of raw acceleration sedentary  
 327 thresholds in children and adults. *Scand J Med Sci Sports*. 2016.
- 328 14. Lohman T, Roche AF, Martorell R. *Anthropometric standardization reference manual*,  
 329 Champaign, Illinois, Human Kinetics; 1988.
- 330 15. Catellier DJ, Hannan PJ, Murray DM, et al. Imputation of missing data when measuring activity  
 331 by accelerometry. *Med Sci Sports Exerc*. 2005; 37(Suppl 11):S555-S562.
- 332 16. Rich C, Geraci M, Griffiths L, Sera F, Dezateux C, Cortina-Borja M. Quality control methods in  
 333 accelerometer data processing: defining minimum wear time. *PLoS one*. 2013; 8(6):e67206.
- 334 17. Rowlands AV, Pilgrim EL, Eston RG. Patterns of habitual activity across weekdays and weekend  
 335 days in 9-11-year-old children. *Prev Med*. 2008; 46(4):317-324.
- 336 18. van Hees VT, Gorzelniak L, Dean Leon EC, et al. Separating movement and gravity components  
 337 in an acceleration signal and implications for the assessment of human daily physical activity.  
 338 *PLoS One*. 2013; 8(4):e61691.
- 339 19. Welk GJ, McClain J, Ainsworth BE. Protocols for evaluating equivalency of accelerometry-based  
 340 activity monitors. *Med Sci Sports Exerc*. 2012; 44(1 Suppl 1):S39-49.
- 341 20. Staudenmayer J, Zhu W, Catellier DJ. Statistical considerations in the analysis of accelerometry-  
 342 based activity monitor data. *Med Sci Sports Exerc*. 2012; 44(1 Suppl 1):S61-67.

- 343 21. Kim Y, Welk GJ. Criterion Validity of Competing Accelerometry-Based Activity Monitoring  
344 Devices. *Med Sci Sports Exerc.* 2015; 47(11):2456-2463.
- 345 22. Lee JM, Kim Y, Welk GJ. Validity of consumer-based physical activity monitors. *Med Sci Sports*  
346 *Exerc.* 2014; 46(9):1840-1848.
- 347 23. Welk GJ, Kim Y, Stanfill B, et al. Validity of 24-h physical activity recall: physical activity  
348 measurement survey. *Med Sci Sports Exerc.* 2014; 46(10):2014-2024.
- 349 24. Dixon PM, Saint-Maurice PF, Kim Y, Hibbing P, Bai Y, Welk GJ. A primer on the use of  
350 equivalence testing for evaluating measurement agreement. *Medicine and Science in Sports*  
351 *and Exercise.* 2017; Ahead of Print.
- 352 25. Altman DG. *Practical Statistics for Medical Research.*, London, Chapman and Hall; 1991.
- 353 26. Freedson PS, John D. Comment on "Estimating Activity and Sedentary Behavior from an  
354 Accelerometer on the Hip and Wrist". *Medicine & Science in Sports & Exercise.* 2013; 45(5):962-  
355 963 910.1249/MSS.1240b1013e31827f31024d.
- 356 27. Rowlands AV, Olds TS, Hillsdon M, et al. Assessing sedentary behavior with the GENEActiv:  
357 introducing the sedentary sphere. *Med Sci Sports Exerc.* 2014; 46(6):1235-1247.
- 358 28. Rowlands AV, Rennie K, Kozarski R, et al. Children's physical activity assessed with wrist- and  
359 hip-worn accelerometers. *Med Sci Sports Exerc.* 2014; 46(12):2308-2316.
- 360 29. Pavey TG, Gomersall SR, Clark BK, Brown WJ. The validity of the GENEActiv wrist-worn  
361 accelerometer for measuring adult sedentary time in free living. *Journal of science and*  
362 *medicine in sport / Sports Medicine Australia.*  
363 2015:<http://dx.doi.org/10.1016/j.jsams.2015.1004.1007>.

364

365

366 **Table legend**367 Table 1. Mean (SD) anthropometric, wear time and number of days included within analysis for boys368 and girls

369

	Boys N = 43		Girls N = 65	
	Mean or	SD	Mean or	SD
	Frequency		Frequency	
Age (y)	10.03	0.35	10.04	0.31
Height (cm)	139.49	7.89	137.97	7.37
Body mass (kg)	35.64	8.24	34.23	8.60
BMI (kg·m <sup>2</sup> )	18.15	3.00	17.78	3.18
ActiGraph weekday wear (min·day <sup>-1</sup> )	739.88	115.55	738.75	100.35
ActiGraph weekend day wear (min·day <sup>-1</sup> )	631.83	110.82	661.50	108.28
ActiGraph valid weekdays included	41	N/A	64	N/A
ActiGraph valid weekend days included	30	N/A	46	N/A
Total valid included days	71	N/A	110	N/A


370

371



372 **Figure legend**

373 Figure 1. ActiGraph 100 count·min<sup>-1</sup> zone of equivalence (dotted lines) and 90% confidence intervals  
 374 for the GENEActiv sedentary time data



375