

Manipulating Type-I and Type-II Dirac Polaritons in Cavity-Embedded Honeycomb Metasurfaces

Charlie-Ray Mann,¹ Thomas J. Sturges,¹ Guillaume Weick,² William L. Barnes,¹ and Eros Mariani¹

¹*EPSRC Centre for Doctoral Training in Metamaterials (XM²),*

Department of Physics and Astronomy, University of Exeter, Exeter, EX4 4QL, United Kingdom.

²*Université de Strasbourg, CNRS, Institut de Physique et Chimie des Matériaux de Strasbourg, UMR 7504, F-67000 Strasbourg, France.*

Pseudorelativistic Dirac quasiparticles have emerged in a plethora of artificial graphene systems that mimic the underlying honeycomb symmetry of graphene. However, it is notoriously difficult to manipulate their properties without modifying the lattice structure. Here we theoretically investigate polaritons supported by honeycomb metasurfaces and, despite the trivial nature of the resonant elements, we unveil rich Dirac physics stemming from a non-trivial winding in the light-matter interaction. A new kind of type-II Dirac point emerges which simultaneously exists with its conventional type-I counterpart. By modifying only the photonic environment via an enclosing cavity, one can manipulate the location of the type-II Dirac points, leading to distinct polariton phases. This enables one to alter the fundamental properties of the emergent Dirac polaritons while preserving the lattice structure—a unique scenario which has no analog in real or artificial graphene systems so far. Exploiting the photonic environment will thus give rise to unexplored Dirac physics at the subwavelength scale.

The groundbreaking discovery of monolayer graphene [1] has inspired an extensive quest to emulate massless Dirac quasiparticles in a myriad of distinct artificial graphene systems [2–11], ranging from ultracold atoms in optical lattices [3] to evanescently coupled photonic waveguide arrays [4]. Owing to their honeycomb symmetry, linear band-degeneracies manifest in the quasiparticle spectrum which we call conventional Dirac points (CDPs). These belong to the ubiquitous type-I class of two-dimensional (2D) Dirac points that are characterized by Dirac cones with closed isofrequency contours. As a result, the corresponding quasiparticles are described by the rather exotic 2D massless Dirac Hamiltonian [12], in stark contrast to the more conventional Schrödinger equation. Thus, these emergent Dirac quasiparticles offer fundamental insight into pseudorelativistic phenomena such as the iconic Klein paradox, corresponding to the perfect transmission through arbitrarily large potential barriers [13]. The latter is responsible for the suppression of backscattering and antilocalization of Dirac quasiparticles, which are highly desirable properties for efficient quasiparticle propagation in novel devices.

Since the existence of type-I CDPs is intrinsically linked to the honeycomb structure, the fundamental properties of the massless Dirac quasiparticles are notoriously robust and difficult to manipulate. However, by exploiting meticulous control over the lattice structure, artificial graphene systems have enabled the exploration of Dirac quasiparticles in new regimes that are difficult, if not impossible to achieve in graphene itself [14–19]. An archetypal example which has attracted considerable interest is the paradigm of strain-engineering, where it has been shown that lattice anisotropy can induce the merging and annihilation of type-I CDPs [3, 14–16, 20–23], and that aperiodicity can generate large pseudomagnetic fields [17, 24].

This concept of Hamiltonian engineering, and the hunt for exotic quasiparticles, has recently entered the realm of polaritonics [25–30]. The true potential of polaritons lies in their hybrid nature, where their light and matter constituents can be manipulated independently, thereby providing additional tunable degrees of freedom. As an example, recent work has shown the tantalizing prospect of engineering novel topological polaritons by introducing a winding coupling between ordinary photons and excitons [27]. Shortly thereafter, these non-trivial interactions were exploited to gap out photonic Dirac points as an alternative route to topological polaritons [29].

In this work we exploit the hybrid nature of polaritons in a different setting, namely metamaterials. We explore Dirac physics in an entirely new scenario, shifting the focus from the lattice structure and its deformations to the effect of manipulating the surrounding photonic environment. In particular, we theoretically study the polaritons supported by crystalline metasurfaces consisting of a honeycomb array of resonant, dipolar meta-atoms. Such a structure is imminently realizable across the electromagnetic spectrum from arrays of plasmonic nanoparticles to microwave helical resonators. Despite the elementary nature of these metasurfaces, we show that complex




Figure 1. Schematic of a cavity-embedded honeycomb metasurface. The honeycomb array of meta-atoms is composed of two inequivalent (A and B) hexagonal sublattices—defined by lattice vectors $\mathbf{a}_1 = a(-\frac{\sqrt{3}}{2}, \frac{3}{2})$ and $\mathbf{a}_2 = a(\frac{\sqrt{3}}{2}, \frac{3}{2})$ —which are connected by nearest-neighbor vectors $\mathbf{e}_1 = a(0, -1)$, $\mathbf{e}_2 = a(\frac{\sqrt{3}}{2}, \frac{1}{2})$ and $\mathbf{e}_3 = a(-\frac{\sqrt{3}}{2}, \frac{1}{2})$, where a is the subwavelength nearest-neighbor separation. Each meta-atom is modelled as an electric dipole, oriented normal to the plane of the lattice. The honeycomb metasurface is embedded inside a photonic cavity of height L , which is composed of two perfectly conducting metallic plates, enabling one to modify the photonic environment while preserving the lattice structure. This general model can be realized across the electromagnetic spectrum, from arrays of plasmonic nanorods to microwave helical resonators.

non-local interactions, mediated by the electromagnetic environment, give rise to rich and unique Dirac physics at the subwavelength scale. In particular, we unveil the emergence of a new kind of type-II Dirac point, which simultaneously exists with the type-I CDPs. In contrast to the latter, the former are characterized by critically tilted Dirac cones with open, hyperbolic isofrequency contours.

Recently, the discovery of type-II Dirac/Weyl semimetals has sparked a growing interest in the prospect of type-II Dirac/Weyl points since the corresponding ‘Lorentz-violating’ Dirac quasiparticles exhibit markedly different properties from their type-I counterparts [31–35]. After their realization, the pursuit for electromagnetic analogs commenced [36–42]. However, just like type-I CDPs, the existence of type-II Dirac/Weyl points is usually tied to judiciously engineered lattice structures, and are thus impossible to manipulate without modifying the structure [36–42].

In stark contrast, here the existence of the type-II Dirac points is intrinsically linked to the hybrid nature of the polaritons, emerging from a non-trivial winding in the light-matter interaction. Crucially, the combination of this unique physical origin and the truly 2D nature of the metasurface results in an unprecedented tunability of emergent Dirac polaritons by modifying only the surrounding photonic environment. Specifically, we show that by embedding the honeycomb metasurface inside a planar photonic cavity and simply changing the cavity height, one can induce novel phase

transitions, including the multi-merging of type-I and type-II Dirac points and the annihilation of type-II Dirac points. This striking tunability results in distinct polariton phases, despite the preserved lattice structure. In particular, we unveil a morphing between a linear and a parabolic spectrum accompanied by a change in topological Berry phase, and an unprecedented environment-induced inversion of chirality, all of which have no analog in graphene or artificial graphene systems studied thus far. Therefore, this unique paradigm of exploiting the photonic environment will give rise to novel Dirac-related phenomena at the subwavelength scale, such as anomalous Klein tunnelling, negative refraction and pseudo-magnetic Landau levels, which can all be tuned via the photonic environment alone.

RESULTS

Hamiltonian Formulation

While metamaterials have traditionally been described in terms of macroscopic effective properties [36, 39, 43], the importance of crystallinity is becoming increasingly apparent [44]. Therefore, to capture the essential physics related to complex non-local effects that arise from strong multiple-scattering [45], here we study the properties of the cavity-embedded honeycomb metasurface by means of a microscopic Hamiltonian formalism. This allows us to clearly identify the distinct physical origins of the type-I and type-II Dirac points.

The full polariton Hamiltonian of this system reads $H_{\text{pol}} = H_{\text{mat}} + H_{\text{ph}} + H_{\text{int}}$, where the interaction Hamiltonian H_{int} couples the matter and photonic subspaces whose free dynamics are governed by H_{mat} and H_{ph} , respectively. We employ the Coulomb gauge, where the instantaneous Coulomb interaction between the meta-atoms is incorporated within the matter Hamiltonian H_{mat} [46]. The effects of the dynamic photonic environment, described by the transverse vector potential, is included through the principle of minimal-coupling [46].

We model each meta-atom by a single dynamical degree of freedom describing the electric-dipole moment associated with its (non-degenerate) fundamental eigenmode with resonant frequency ω_0 . These meta-atoms are then oriented such that their dipole moments point normal to the plane of the lattice. Furthermore, we consider subwavelength nearest-neighbor separation a such that the light cone intersects the Brillouin zone edge above ω_0 , ensuring the existence of evanescently bound, subwavelength polaritons. The strength of the Coulomb dipole-dipole interaction between neighboring meta-atoms is parametrized by Ω . Finally, the metasurface is embedded at the center of a planar photonic cavity of height L , as depicted in Fig. 1, where the cavity walls are assumed to be lossless and perfectly conducting metallic plates.

Emergence of Type-I Dirac Points

The matter Hamiltonian within the nearest-neighbor approximation reads

$$H_{\text{mat}} = \hbar\tilde{\omega}_0 \sum_{\mathbf{q}} (a_{\mathbf{q}}^\dagger a_{\mathbf{q}} + b_{\mathbf{q}}^\dagger b_{\mathbf{q}}) + \hbar\tilde{\Omega} \sum_{\mathbf{q}} (f_{\mathbf{q}} b_{\mathbf{q}}^\dagger a_{\mathbf{q}} + \text{H.c.}), \quad (1)$$

where, for brevity, we have not presented the non-resonant terms (see Methods for derivation). In equation (1), $\tilde{\omega}_0$ is the renormalized resonant frequency and $\tilde{\Omega}$ is the renormalized Coulombic interaction strength due to the cavity-induced image dipoles (see Methods for their dependence on the cavity height). The bosonic operators $a_{\mathbf{q}}^\dagger$ and $b_{\mathbf{q}}^\dagger$ create quanta of the quasistatic collective-dipole modes that extend across the A and B sublattices, respectively, with wavevector \mathbf{q} in the first Brillouin zone (see Fig. 2a). Finally, the function $f_{\mathbf{q}} = \sum_{j=1}^3 \exp(i\mathbf{q} \cdot \mathbf{e}_j)$ encodes the honeycomb geometry of the lattice with nearest-neighbor vectors \mathbf{e}_j (see Fig. 1).

We diagonalize the matter Hamiltonian (1) as $H_{\text{mat}} = \sum_{\tau=\pm} \sum_{\mathbf{q}} \hbar\omega_{\mathbf{q}\tau}^{\text{mat}} \beta_{\mathbf{q}\tau}^\dagger \beta_{\mathbf{q}\tau}$ where the bosonic operators $\beta_{\mathbf{q}\tau}^\dagger = \psi_{\mathbf{q}}^\dagger |\psi_{\mathbf{q}\tau}\rangle$ create quasistatic collective-dipole normal modes with dispersion $\omega_{\mathbf{q}\tau}^{\text{mat}} = \tilde{\omega}_0 + \tau\tilde{\Omega}|f_{\mathbf{q}}|$. Here, τ indexes the upper ($\tau = +1$) and lower ($\tau = -1$) bands and $\psi_{\mathbf{q}}^\dagger = (a_{\mathbf{q}}^\dagger, b_{\mathbf{q}}^\dagger)$ is a spinor creation operator. The spinors $|\psi_{\mathbf{q}\tau}\rangle = (1, \tau e^{i\varphi_{\mathbf{q}}})^T / \sqrt{2}$ describe an emergent pseudospin degree of freedom where the two components encode the relative amplitude and phase of the dipolar oscillations on the two inequivalent A and B sublattices, respectively, with $\varphi_{\mathbf{q}} = \arg(f_{\mathbf{q}})$. These spinors can be represented by a pseudospin vector on the Bloch sphere which reads $\mathbf{S}_{\mathbf{q}\tau} = \tau(\cos \varphi_{\mathbf{q}}, \sin \varphi_{\mathbf{q}}, 0)$.

At the high symmetry K and K' points (see Fig. 2a) the sublattices decouple with no well-defined relative phase (i.e., $f_{\mathbf{q}} = 0$), giving rise to two inequivalent CDPs located at $\pm\mathbf{K} = \pm \left(\frac{4\pi}{3\sqrt{3}a}, 0 \right)$ as observed in Fig. 2b. As expected from the symmetry of the metasurface, the existence of CDPs is robust against long-range Coulomb interactions as shown in Supplementary Note 1. In fact, for small cavity heights, the image dipoles quench long-range Coulomb interactions and the nearest-neighbor approximation becomes increasingly accurate as shown in Supplementary Fig. 1. The CDPs correspond to vortices in the pseudospin vector field $\mathbf{S}_{\mathbf{q}\tau}$, which give rise to topological singularities in the Berry curvature [47]. Therefore, CDPs are sources of quantized Berry flux $w\pi$, where $w = \pm 1$ is the topological charge of the Dirac point corresponding to the winding number of the vortex.

To quadratic order in $\mathbf{k} = \mathbf{q} - \mathbf{K}$ ($ka \ll 1$), the effective matter Hamiltonian near the K point is $H_{\mathbf{K}}^{\text{eff}} = \sum_{\mathbf{k}} \psi_{\mathbf{k}}^\dagger \mathcal{H}_{\mathbf{K},\mathbf{k}}^{\text{eff}} \psi_{\mathbf{k}}$, with spinor creation operator $\psi_{\mathbf{k}}^\dagger = (a_{\mathbf{k}}^\dagger, b_{\mathbf{k}}^\dagger)$ and Bloch Hamiltonian

$$\mathcal{H}_{\mathbf{K},\mathbf{k}}^{\text{eff}} = \hbar\tilde{\omega}_0 \mathbb{1}_2 - \hbar\tilde{v} \boldsymbol{\sigma} \cdot \mathbf{k} + \hbar\tilde{t} (\boldsymbol{\sigma}^* \cdot \mathbf{k})^{\circ 2}. \quad (2)$$

Here, $\mathbb{1}_2$ is the 2×2 identity matrix, $\boldsymbol{\sigma} = (\sigma_x, \sigma_y)$ and $\boldsymbol{\sigma}^* = (\sigma_x, -\sigma_y)$ are vectors of Pauli matrices, and $\circ 2$ represents the Hadamard (element-wise) square. Note that the




Figure 2. **Evolution of the polariton dispersion as the cavity height is reduced.** **a**, First Brillouin zone defined by primitive reciprocal lattice vectors $\mathbf{b}_1 = \frac{2\pi}{3a}(-\sqrt{3}, -1)$ and $\mathbf{b}_2 = \frac{2\pi}{3a}(\sqrt{3}, -1)$. **b**, Quasistatic dispersion of the collective-dipole normal modes, where the upper band corresponds to a bright, symmetric dipole configuration ($\uparrow\uparrow$) and the lower band corresponds to a dark, antisymmetric dipole configuration ($\uparrow\downarrow$). The light-cone (shaded region) is bounded by the linear dispersion of the TEM mode. Due to the non-trivial winding in the light-matter interaction (see Fig. 3), the band crossings are expected to result in large (band crossings ‘1’ and ‘2’) or small (band crossings ‘3’ and ‘4’) direction-dependent anticrossings in the polariton spectrum. **c-e**, Polariton dispersion obtained from the polariton Hamiltonian H_{pol} (solid black lines) and the two-band Hamiltonian H_{mat} (orange dashed lines), for **c** subcritical ($L = 5a$), **d** critical ($L = L_c = 1.75a$), and **e** supercritical ($L = a$) cavity heights, respectively. While type-I CDPs with an isotropic Dirac cone (see inset of **c**) exist even in the quasistatic dispersion (see **b**), new type-II SDPs with a critically tilted Dirac cone (see inset in **c**) emerge due to the vanishing light-matter interaction for the dark quasistatic band along the $\Gamma - K(K')$ directions (see Fig. 3). At the critical cavity height L_c , three type-II SDPs merge with the type-I CDPs (see Fig. 5) resulting in a quadratic band-degeneracy at $K(K')$ (see inset in **d**). After criticality, the type-II SDPs annihilate one another and the massless Dirac cone re-emerges at the type-I CDPs (see inset in **e**) accompanied by an inversion of chirality (see Fig. 5). Plots obtained with parameters $\omega_{\mathbf{K}0}^{\text{ph}} = 2.5\omega_0$ and $\Omega = 0.01\omega_0$.

image dipoles do not qualitatively affect the physics, but simply lead to a renormalization of the group velocity $\tilde{v} = 3\tilde{\Omega}a/2$ and trigonal warping parameter $\tilde{t} = 3\tilde{\Omega}a^2/8$. Apart from a global energy shift, equation (2) is equivalent to a 2D massless Dirac Hamiltonian to leading order in \mathbf{k} , with an isotropic Dirac cone spectrum $\omega_{\mathbf{k}\tau}^{\text{mat}} = \tilde{\omega}_0 + \tau\tilde{v}|\mathbf{k}|$ that is characterized by closed isofrequency contours. Therefore, as expected from the honeycomb symmetry, the CDP belongs to the type-I class of 2D Dirac points, and the corresponding spinors $|\psi_{\mathbf{k}\tau}\rangle = (1, -\tau e^{i\theta_{\mathbf{k}}})^T/\sqrt{2}$, where $\theta_{\mathbf{k}} = \arctan(k_y/k_x)$, represent massless Dirac collective-dipoles with a topological Berry phase of π . The effective Hamiltonian near the K' point is given by $\mathcal{H}_{K',\mathbf{k}}^{\text{eff}} = (\mathcal{H}_{K,-\mathbf{k}}^{\text{eff}})^*$, where the CDP is a source of $-\pi$ Berry flux as required by time-reversal symmetry.

Hybridization with the Photonic Environment

Given the subwavelength nearest-neighbor separation, it is tempting to assert that the near-field Coulomb interactions in

H_{mat} capture the essential physics. In fact, we will show that this quasistatic description misses the profound influence of the surrounding photonic environment, which has a remarkably non-trivial effect on the Berry curvature and, therefore, on the corresponding nature of the polaritons.

Crucially, the metallic cavity supports a fundamental transverse electromagnetic (TEM) mode whose polarization (parallel to the dipole moments) and linear dispersion (see Fig. 2b) are independent of the cavity height. For brevity, in what follows we do not present the contributions from the other cavity modes since the essential physics emerges from the interaction with the fundamental TEM mode (see Methods for full expressions). In fact, the higher order cavity modes become increasingly negligible for smaller cavities as they are progressively detuned from the dipole resonances.

The effects of the photonic environment are encoded in the free photonic Hamiltonian

$$H_{\text{ph}} = \hbar \sum_{\mathbf{q}^n} \omega_{\mathbf{q}^n}^{\text{ph}} c_{\mathbf{q}^n}^\dagger c_{\mathbf{q}^n} \quad (3)$$

and in the light-matter interaction Hamiltonian $H_{\text{int}} = H_{\text{int}}^{(1)} + H_{\text{int}}^{(2)}$, with

$$H_{\text{int}}^{(1)} = \hbar \sum_{\mathbf{q}n} i\xi_{\mathbf{q}n} \phi_n^* \left(a_{\mathbf{q}}^\dagger c_{\mathbf{q}n} + a_{\mathbf{q}}^\dagger c_{-\mathbf{q}n}^\dagger \right) + \hbar \sum_{\mathbf{q}n} i\xi_{\mathbf{q}n} \phi_n \left(b_{\mathbf{q}}^\dagger c_{\mathbf{q}n} + b_{\mathbf{q}}^\dagger c_{-\mathbf{q}n}^\dagger \right) + \text{H.c.} \quad (4)$$

and

$$H_{\text{int}}^{(2)} = \hbar \sum_{\mathbf{q}n n'} \frac{2\xi_{\mathbf{q}n} \xi_{\mathbf{q}n'}}{\omega_0} \text{Re} \left(\phi_n \phi_{n'}^* \right) \left(c_{\mathbf{q}n}^\dagger c_{\mathbf{q}n'} + c_{\mathbf{q}n}^\dagger c_{-\mathbf{q}n'}^\dagger \right) + \text{H.c.}, \quad (5)$$

where $\xi_{\mathbf{q}n} \propto L^{-1/2}$ parametrizes the strength of the light-matter interaction (see Methods for analytical expression). The bosonic operator $c_{\mathbf{q}n}^\dagger$ creates a TEM photon with wavevector \mathbf{q} in the first Brillouin zone and dispersion $\omega_{\mathbf{q}n}^{\text{ph}} = c|\mathbf{q} - \mathbf{G}_n|$, where n indexes the set of reciprocal lattice vectors \mathbf{G}_n . The complex phase factors $\phi_n = \exp(i\mathbf{a}\mathbf{G}_n \cdot \hat{\mathbf{y}})$ are associated with Umklapp processes that arise due to the discrete, in-plane translational symmetry of the metasurface, and must be retained as they are critical for maintaining the point-group symmetry of the polariton Hamiltonian.

We diagonalize H_{pol} using a generalized Hopfield-Bogoliubov transformation [48] (see Methods for details), and in Fig. 2c-e we present the resulting polariton dispersion for different cavity heights. Also, in Supplementary Fig. 2 we present the full polariton dispersion that includes long-range Coulomb interactions. For small cavity heights, the full polariton dispersion is almost indistinguishable from that obtained in the nearest-neighbor approximation, and therefore one can conclude that long-range Coulomb interactions do not qualitatively affect the physics presented here. It is important to stress that our general model captures the essential physics that will emerge in a variety of different experimental setups. To show this, in Supplementary Fig. 3 and Supplementary Fig. 4 we present the polariton dispersions obtained from full-wave numerical simulations of a honeycomb array of plasmonic nanorods and microwave helical resonators, respectively. These different physical realizations indeed show the same evolution of the polariton spectrum as shown in Fig. 2c-e and Supplementary Fig. 2.

Emergence of Type-II Dirac Points

Given the elementary nature of the individual resonant elements, it is tempting to assume that nothing peculiar could emerge from the ordinary dipole-dipole interactions between the meta-atoms that are mediated by the electromagnetic field. However, by expressing the interaction Hamiltonian (4) in terms of the $\beta_{\mathbf{q}\tau}$ and $\beta_{\mathbf{q}\tau}^\dagger$ operators that diagonalize the matter




Figure 3. Non-trivial winding in the light-matter interaction. **a**, Dependence of the magnitude of the light-matter coupling constant $|\Lambda_{\mathbf{q}0\tau}| \propto |e^{i\varphi_{\mathbf{q}}} + \tau|$ on the direction of \mathbf{q} from $\Gamma - M$ to $\Gamma - K(K')$ (see inset), for the upper (red line) and lower (blue line) quasistatic bands. Plots obtained with $|\mathbf{q}| = |\mathbf{K}|/2$. **b-e**, Schematics of the bright ($\uparrow\uparrow$) and dark ($\uparrow\downarrow$) configurations of the two sublattices interacting with the photonic mode, indicated by the field profile. Panels **b** and **d** represent the crossings labeled ‘1’ and ‘3’ in Fig. 2 along the $\Gamma - K(K')$ directions, respectively, while panels **c** and **e** represent the crossings labelled ‘2’ and ‘4’ along the $\Gamma - M$ directions, respectively. Crucially, the light-matter interaction strength for the dark mode vanishes ($|\Lambda_{\mathbf{q}0-}| = 0$) along the $\Gamma - K(K')$ directions due to the complete destructive interference between the two sublattices (see **d**), leading to the emergence of six inequivalent type-II Dirac points in the polariton spectrum.

Hamiltonian,

$$H_{\text{int}}^{(1)} = \hbar \sum_{\tau=\pm} \sum_{\mathbf{q}n} i\Lambda_{\mathbf{q}n\tau} \left(\beta_{\mathbf{q}\tau}^\dagger c_{\mathbf{q}n} + \beta_{\mathbf{q}\tau}^\dagger c_{-\mathbf{q}n}^\dagger \right) + \text{H.c.}, \quad (6)$$

we find that complex non-local interactions, which arise from strong multiple scattering in the bipartite structure, result in a non-trivial winding of the light-matter coupling as a function of wavevector direction

$$\Lambda_{\mathbf{q}n\tau} \propto \xi_{\mathbf{q}n} \left(\phi_n^* e^{i\varphi_{\mathbf{q}}} + \tau \phi_n \right). \quad (7)$$

Naively, one may expect all of the band crossings in Fig. 2b to be avoided as a result of the hybridization between the collective-dipole and photonic modes, as it is a characteristic feature of polaritonic systems [48]. Indeed, this is the case for the crossings with the upper quasistatic band where $\Lambda_{\mathbf{q}0+} \propto (e^{i\varphi_{\mathbf{q}}} + 1)$ (see red line in Fig. 3a) due to the constructive interference between the sublattices of this bright ($\uparrow\uparrow$) configuration (see Fig. 3b-c). This results in a large anticrossing for all wavevector directions, as observed in Fig. 2c. In stark contrast, for the lower quasistatic band the coupling constant is significantly reduced $\Lambda_{\mathbf{q}0-} \propto (e^{i\varphi_{\mathbf{q}}} - 1)$ (see blue line in Fig. 3a) due to the destructive interference between the sublattices of this dark ($\uparrow\downarrow$) configuration (see Fig. 3e). Consequently, this results in a small anticrossing for a general wavevector direction.

Crucially, however, the light-matter interaction for the lower quasistatic band completely vanishes ($\Lambda_{\mathbf{q}0-} = 0$) along the high-symmetry $\Gamma - K(K')$ directions, where $\varphi_{\mathbf{q}} = 0$, due to the complete destructive interference between the two sublattices (see Fig. 3d). As a result, along these high-symmetry directions the crossings are protected, leading to six inequivalent Dirac points emerging in the polariton spectrum—we call these satellite Dirac points (SDPs) to distinguish them from the CDPs. As we will see below, these SDPs belong to the type-II class of 2D Dirac points where the dispersion takes the form of a critically tilted Dirac cone (see inset of Fig. 2c), characterized by open, hyperbolic isofrequency contours.

Effective Hamiltonian in the Matter Subspace

To explore the nature of the polaritons in the vicinity of the different Dirac points, we first neglect non-resonant terms in the matter Hamiltonian and perform a unitary Schrieffer-Wolff transformation [49] on H_{pol} to integrate out the photonic degrees of freedom (see Methods for details). Finally, we extract the two-band Hamiltonian in the matter sublattice space

$$\begin{aligned} \bar{H}_{\text{mat}} = H_{\text{mat}} - 2\hbar \sum_{\mathbf{q}n} \frac{\xi_{\mathbf{q}n}^2 \omega_{\mathbf{q}n}^{\text{ph}}}{(\omega_{\mathbf{q}n}^{\text{ph}})^2 - \tilde{\omega}_0^2} \\ (a_{\mathbf{q}}^\dagger a_{\mathbf{q}} + b_{\mathbf{q}}^\dagger b_{\mathbf{q}} + \phi_n^2 b_{\mathbf{q}}^\dagger a_{\mathbf{q}} + \phi_n^{*2} a_{\mathbf{q}}^\dagger b_{\mathbf{q}}). \end{aligned} \quad (8)$$

Diagonalizing the two-band Hamiltonian (8) leads to an effective dispersion (see Methods) which provides an excellent description of the polaritons as indicated by the orange dashed lines in Fig. 2c-e. Finally, we expand the two-band Hamiltonian (8) up to quadratic order in \mathbf{k} and obtain the effective Hamiltonian near the K point $\bar{H}_{\text{K}}^{\text{eff}} = \sum_{\mathbf{k}} \psi_{\mathbf{k}}^\dagger \bar{\mathcal{H}}_{\text{K},\mathbf{k}}^{\text{eff}} \psi_{\mathbf{k}}$ (see Supplementary Note 2 for derivation) with Bloch Hamiltonian

$$\bar{H}_{\text{K},\mathbf{k}}^{\text{eff}} = \hbar\tilde{\omega}_0 \mathbb{1}_2 - \hbar\tilde{v} \boldsymbol{\sigma} \cdot \mathbf{k} + \hbar\tilde{t} (\boldsymbol{\sigma}^* \cdot \mathbf{k})^{\circ 2} - \hbar D |\mathbf{k}|^2 \mathbb{1}_2. \quad (9)$$

Similarly, the effective Hamiltonian near the K' point is given by $\bar{\mathcal{H}}_{\text{K}',\mathbf{k}}^{\text{eff}} = (\bar{\mathcal{H}}_{\text{K},-\mathbf{k}}^{\text{eff}})^*$. In equation (9), the resonant frequency $\tilde{\omega}_0$, group velocity \tilde{v} , and trigonal warping parameter \tilde{t} , now encode non-trivial contributions from the hybridization with the photonic environment. There is also an additional wavevector-dependent diagonal term parametrized by D , which breaks the symmetry between the upper and lower polariton bands. The dependence of these parameters on the cavity height is shown in Fig. 4 (see Methods for analytical expressions). To leading order in \mathbf{k} , one can observe that the effective Hamiltonian (9) near the CDP is equivalent to a 2D massless Dirac Hamiltonian. Therefore, the polariton CDPs remain in the type-I class and are robust against the coupling with the photonic environment—this is not surprising given that their physical origin is intrinsically linked to the lattice structure alone, which is preserved here.

To elucidate the nature of the SDPs, we expand the effective Hamiltonian (9) near one of the SDPs located at $\mathbf{K}_{\text{S}} =$




Figure 4. **Tunable parameters in the effective Hamiltonian.** Dependence of the parameters in the effective polariton Hamiltonian (9) on the inverse cavity height. The blue dashed line shows the variation of the group velocity \bar{v} which changes sign at the critical cavity height L_c , leading to the inversion of chirality. The orange dot-dashed line shows the variation of the trigonal warping parameter \bar{t} which becomes dominant close to criticality. These parameters have been normalized to $v = 3\Omega a/2$ and $t = 3\Omega a^2/8$ which are the group velocity and trigonal warping parameters, respectively, in the absence of image dipoles and light-matter interactions. The orange dot-dashed line in the inset shows the variation of the CDP frequency $\tilde{\omega}_0$, while the blue dashed line in the inset shows the variation of the wavevector-dependent diagonal term D . Plots obtained with $\omega_{\text{K}0}^{\text{ph}} = 2.5\omega_0$ and $\Omega = 0.01\omega_0$.

$(\bar{v}/\bar{t}, 0)$ and obtain

$$\bar{\mathcal{H}}_{\text{K}_S,\mathbf{k}'}^{\text{eff}} = \hbar \left(\tilde{\omega}_0 - \frac{D\bar{v}^2}{\bar{t}^2} - \frac{2D\bar{v}}{\bar{t}} k'_x \right) \mathbb{1}_2 + \hbar \boldsymbol{\sigma}^* \cdot \bar{\mathbf{v}} \cdot \mathbf{k}', \quad (10)$$

where \mathbf{k}' measures the deviation from \mathbf{K}_{S} and $\bar{\mathbf{v}} = \bar{v} \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}$ is the velocity tensor. Apart from a global energy shift, the effective Hamiltonian (10) near the SDP takes the form of a generalized 2D massless Dirac Hamiltonian $\mathcal{H}_{\mathbf{k}} = \sum_{i=x,y} \hbar u_i k_i \mathbb{1}_2 + \sum_{i=x,y} \hbar v_i k_i \sigma_i$. If the parameters u_i and v_i satisfy the condition $u_x^2/v_x^2 + u_y^2/v_y^2 < 1$, then the Dirac cone becomes tilted and anisotropic [50] but still belongs to the type-I class with closed isofrequency contours. However, the condition $u_x^2/v_x^2 + u_y^2/v_y^2 > 1$ defines a distinct type-II class of 2D Dirac points, giving rise to a critically tilted Dirac cone with open, hyperbolic isofrequency contours. Hence, the distinct type-I and type-II classes are related via a Lifshitz transition in the topology of the isofrequency contours. Indeed, since we have $u_y = 0$ and $u_x^2/v_x^2 = 4D^2/\bar{t}^2 > 1$, the SDPs belong to the type-II class of 2D Dirac points. Furthermore, since the Hamiltonian (10) is expressed in terms of $\boldsymbol{\sigma}^*$, the pseudospin winds in the opposite direction around the SDPs as compared to the CDP, and therefore the SDPs located along the $\Gamma - K$ directions are sources of $-\pi$ Berry flux. As required by time-reversal symmetry, the SDPs located along the $\Gamma - K'$ directions are sources of π Berry flux (opposite to

the CDP located at the K' point) To the authors knowledge, this is the first realization of type-II Dirac points in a truly 2D system with subwavelength dimension in the transverse direction.

Manipulation of Type-I and Type-II Dirac Points

We have demonstrated the simultaneous existence of type-I and type-II Dirac points in the 2D polariton spectrum. Since the latter are intrinsically linked to the hybridization between the light and matter degrees of freedom, one can manipulate their location within the Brillouin zone by simply modifying the light-matter interaction via the cavity height. As a result, the polariton spectrum evolves into qualitatively distinct phases as highlighted in Fig. 2c-e. To elucidate the differences between these phases, we study the spinor eigenstates (see Methods) of the two-band Hamiltonian (8). In Fig. 5a-c we plot the pseudospin vector field near the K point for different cavity heights and schematically depict the location of the Dirac points, along with their associated Berry flux. Finally, in Fig. 5d-f we illustrate the corresponding effective polariton spectrum to leading order in \mathbf{k} . Note that similar analysis can be performed for the K' point.

In the subcritical phase ($L > L_c$), three type-II SDPs are located along the $\Gamma - K$ directions, each with $-\pi$ Berry flux surrounding a type-I CDP with π Berry flux (see Fig. 5a). To leading order in \mathbf{k} , the polariton spectrum disperses linearly about the type-I CDPs (see Fig. 2c) forming an isotropic Dirac cone with a group velocity \bar{v} that is tunable via the cavity height (see Fig. 4). Here, the effective Hamiltonian (9) is equivalent to a 2D massless Dirac Hamiltonian with spinor eigenstates $|\psi_{\mathbf{k}\tau}\rangle = (1, -\tau e^{i\theta_{\mathbf{k}}})^T / \sqrt{2}$. These represent massless Dirac polaritons with chirality $\langle \psi_{\mathbf{k}\tau} | \boldsymbol{\sigma} \cdot \hat{\mathbf{k}} | \psi_{\mathbf{k}\tau} \rangle = -\tau$, resulting in a pseudospin that winds once around the CDP and a topological Berry phase of π (see Fig. 5d).

At the critical cavity height ($L = L_c$), the group velocity of the massless Dirac polaritons vanishes $\bar{v}(L_c) = 0$ (see Fig. 4) as the type-II SDPs merge with the type-I CDP, forming a quadratic band-degeneracy (see Fig. 2d) with combined -2π Berry flux (see Fig. 5b). The leading order term in the effective Hamiltonian (9) is now quadratic in \mathbf{k} with corresponding spinor eigenstates $|\psi_{\mathbf{k}\tau}\rangle = (1, -\tau e^{-i2\theta_{\mathbf{k}}})^T / \sqrt{2}$. Therefore, during this critical merging transition, the massless Dirac polaritons morph into massive chiral polaritons with qualitatively distinct physical properties. These include a parabolic spectrum and chirality $\langle \psi_{\mathbf{k}\tau} | (\boldsymbol{\sigma}^* \cdot \mathbf{k})^2 | \psi_{\mathbf{k}\tau} \rangle = -\tau$, resulting in a pseudospin that winds twice as fast compared to the subcritical phase and a topological Berry phase of -2π (see Fig. 5e).

Since the point-group symmetry is preserved, the type-II SDPs do not annihilate the type-I CDP, but they re-emerge and continue to migrate along the $K - M$ directions as the cavity height is reduced past criticality ($L < L_c$) (see inset of Fig. 5c). After a small decrease in cavity height, these SDPs annihilate with other SDPs that migrate along the opposite di-

rection and have opposite Berry flux. This topological transition leaves only the type-I CDP remaining in the spectrum with π Berry flux (see Fig. 2e and Fig. 5c).

In this supercritical phase, we recover the linear dispersion near the type-I CDP to leading order in \mathbf{k} (see Fig. 2e), and the effective Hamiltonian (9) is equivalent to a 2D massless Dirac Hamiltonian with corresponding spinor eigenstates $|\psi_{\mathbf{k}\tau}\rangle = (1, \tau e^{i\theta_{\mathbf{k}}})^T / \sqrt{2}$. Remarkably, massless Dirac polaritons thus re-emerge past criticality with an unprecedented environment-induced inversion of chirality $\langle \psi_{\mathbf{k}\tau} | \boldsymbol{\sigma} \cdot \hat{\mathbf{k}} | \psi_{\mathbf{k}\tau} \rangle = \tau$ (see Fig. 5f). Physically, this corresponds to a π -rotation in the relative phase between the dipole oscillations on the two sublattices, which is also accompanied by a π -rotation in the isofrequency domains (compare Fig. 5a and Fig. 5c). We emphasize that it is the chirality of massless Dirac fermions that is responsible for most of the remarkable properties of monolayer graphene, including the Klein tunnelling phenomenon [13]. Consequently, this novel environment-induced inversion of chirality could give rise to unconventional phenomena such as anomalous Klein transport. For example, near the K point, the right-propagating polaritons correspond to an antisymmetric dipole configuration $|\psi_{\mathbf{k}\tau}\rangle = (1, -1)^T / \sqrt{2}$ in the subcritical phase and to a symmetric configuration $|\psi_{\mathbf{k}\tau}\rangle = (1, 1)^T / \sqrt{2}$ in the supercritical phase. Thus, due to the orthogonality between these two eigenstates, the inversion of chirality removes the channel responsible for the perfect transmission in the conventional Klein tunnelling effect [13] (see Fig. 5d and Fig. 5f). Such a scenario could be realized in a simple setup characterized by two neighboring regions with different cavity heights.

As a side remark, we note that the polariton spectrum near criticality bears some resemblance with the low-energy spectrum of bilayer graphene with its central type-I Dirac point and three type-I SDPs [51, 52]. However, given the type-II nature of the polariton SDPs, the topology of the polariton isofrequency contours are markedly different from that of the bilayer spectrum. This is further highlighted at criticality where the polariton bands have the same curvature, which is in stark contrast to the electronic bands in bilayer graphene.

We also note that recent works explored the possibility to manipulate the (3+1) type-I Dirac points in bilayer graphene through the application of lattice deformations [53–56], leading to the merging and annihilation of CDPs and SDPs. In addition, a multi-merging transition of all (3+1) type-I Dirac points has been proposed theoretically within tight-binding models involving the artificial tuning of third-nearest-neighbour hopping amplitudes in a graphene-like honeycomb structure [57–60]. However, these proposals have no physical realization so far. In stark contrast, the imminently-realizable metasurfaces in our work enable the exploration of rich Dirac phases with ease by simply modifying the photonic environment via an enclosing cavity.

As a final remark, we briefly comment on how one might probe these novel Dirac phases. Given that the Dirac points exist in a polaritonic excitation spectrum, one must drive the system with photons at the required frequency in order to




Figure 5. **Merging of type-I and type-II Dirac points with chirality inversion.** **a-c**, Pseudospin vector field and isofrequency contours for the upper polariton band near the K point for **a** subcritical ($L = 2.3a$), **b** critical ($L = L_c = 1.78a$), and **c** supercritical ($L = 1.4a$) cavity heights, respectively, as obtained from \vec{H}_{mat} . The Dirac points (corresponding to vortices) are depicted by orange circles along with their associated Berry flux. Before criticality, three type-II SDPs ($-\pi$ Berry flux) are driven towards the type-I CDP (π Berry flux) along the $\Gamma - K$ directions as the cavity height is reduced (see inset in **a**). At the critical cavity height L_c , they merge together forming a quadratic band-degeneracy with combined Berry flux of -2π (see **b**). After criticality, the type-II SDPs re-emerge and are driven past the type-I CDP along the $K - M$ directions (see inset in **c**). After a small decrease in cavity height, these SDPs annihilate other SDPs that migrate along the opposite direction and have opposite Berry flux, leaving only the type-I CDP remaining in the spectrum with π Berry flux (see **c**). **d-f**, Effective polariton spectrum near the K point to leading order in \mathbf{k} . The colors of the bands correspond to the chirality of the Dirac polaritons as defined in the main text, where the orange and blue bands indicate a chirality of $+1$ and -1 , respectively. The spinor eigenstates, represented by pseudospin vectors (gold arrows), describe **d** massless Dirac polaritons with linear dispersion and Berry phase π , **e** massive chiral polaritons with parabolic dispersion and Berry phase -2π , and **f** massless Dirac polaritons with linear dispersion and Berry phase π , but with inverted chirality. All pseudospin and contour plots are obtained with $\omega_{\mathbf{K}0}^{\text{ph}} = 2.5\omega_0$ and $\Omega = 0.01\omega_0$.

probe them. Both classes of Dirac points lie outside of the light-line and thus one must overcome the momentum mismatch with photons. The specific experimental technique that one would employ will depend on the nature of the metasurface and the corresponding frequency regime. For example, techniques for plasmonic systems have traditionally involved coupling via evanescent waves with prisms, gratings, and local scatterers [61], or more recent techniques such as non-linear wave-mixing [62]. In contrast, realizations in the microwave regime can be probed using point-like antenna sources and detectors [39]. In fact, microwave metamaterials are proving to be a versatile platform for exploring Dirac/Weyl

physics, since one can directly probe the field distributions using near-field scanning techniques [39], and thus one could directly probe the environment-induced chirality inversion.

DISCUSSION

To conclude, we have revealed rich and unique Dirac physics that emerges even in the most elementary honeycomb metasurfaces. In particular, we have shown that the polariton spectrum simultaneously exhibits type-I and type-II Dirac points, where the latter emerge from a non-trivial winding in the light-

matter interaction. We would like to emphasize that it is this unique physical origin together with the truly 2D nature of the metasurface that enables one to qualitatively modify the fundamental properties of the emergent Dirac polaritons by manipulating the surrounding photonic environment. This stands in stark contrast to conventional artificial graphene systems where the fundamental properties are dictated by the lattice structure alone. Therefore, exploiting the rich tunability of the polariton spectrum with the environment offers a new paradigm that opens a variety of opportunities to explore novel Dirac-related physics at the subwavelength scale.

For example, one can simultaneously probe the dynamics of type-I and type-II Dirac quasiparticles, where the latter are predicted to exhibit intriguing anomalous refraction behaviour [40]. Furthermore, the environment-induced redshift of the CDP frequency $\tilde{\omega}_0$ (see Fig. 4) will allow the investigation of polaritonic Klein tunnelling through interfaces separating regions with different cavity heights. Consequently, negative refraction can be induced by simple variations in cavity height, which could be exploited in novel schemes for guiding and manipulating light at the subwavelength scale, including polaritonic Veselago lensing [63, 64]. Moreover, the tunable group velocity will also enable the exploration of velocity barriers for the unprecedented guiding and localization of massless Dirac quasiparticles [65, 66], which is extremely difficult to achieve in real graphene. Furthermore, one could also combine the effects of the environment with inhomogeneous strain deformations, giving rise to unique pseudomagnetic-related effects, including the intriguing ability to induce a pseudo-Landau level spectrum for polaritons that can be qualitatively tuned via the cavity height. Finally, the ability to controllably invert the chirality of Dirac polaritons opens new perspectives for anomalous pseudorelativistic transport through interfaces separating regions in distinct polaritonic phases.

METHODS

Derivation of the polaritonic Hamiltonian

The cavity-embedded metasurface is composed of a honeycomb array of identical meta-atoms located at $\mathbf{R}_A = \mathbf{R} + a\hat{y} + \frac{L}{2}\hat{z}$ and $\mathbf{R}_B = \mathbf{R} - a\hat{y} + \frac{L}{2}\hat{z}$ on the A and B sublattices, respectively. Here, $\mathbf{R} = l_1\mathbf{a}_1 + l_2\mathbf{a}_2$ is an in-plane lattice translation vector with primitive vectors \mathbf{a}_1 and \mathbf{a}_2 (see Fig. 1) and integers l_1 and l_2 . Each meta-atom is modelled by a single dynamical degree of freedom h (with dimensions of length), where the electric-dipole moment associated with its fundamental eigenmode is $\mathbf{p} = -Qh\hat{z}$, with effective charge Q . The Coulomb potential energy between two dipole moments \mathbf{p} and \mathbf{p}' located at generic positions \mathbf{r} and \mathbf{r}' , respectively, is given by

$$V_{\text{Coul}} = \frac{\mathbf{p} \cdot \mathbf{p}' - 3(\mathbf{p} \cdot \hat{\mathbf{n}})(\mathbf{p}' \cdot \hat{\mathbf{n}})}{4\pi\epsilon_0|\mathbf{r} - \mathbf{r}'|^3}, \quad (11)$$

where $\hat{\mathbf{n}} = (\mathbf{r} - \mathbf{r}')/|\mathbf{r} - \mathbf{r}'|$ and ϵ_0 is the vacuum permittivity.

The presence of the perfectly conducting metallic plates, placed at $z = 0$ and $z = L$, modifies the boundary conditions on the scalar potential and, therefore, the Coulomb interaction between the meta-atoms. Using the method of images to ensure the vanishing of the scalar potential at the cavity walls [67], we introduce an infinite series of image dipoles located outside the cavity at positions $\mathbf{R}_s + lL\hat{z}$, where $s = A, B$ indexes the two sublattices and l is a non-zero integer. Noting that the Coulomb potential energy between a real and image dipole is 1/2 of that given by equation (11) [68], the matter

Hamiltonian within the nearest-neighbor approximation reads

$$\begin{aligned} H_{\text{mat}} = & \sum_{s=A,B} \sum_{\mathbf{R}_s} \left(\frac{\Pi_{\mathbf{R}_s}^2}{2M} + \frac{M}{2} \omega_0^2 h_{\mathbf{R}_s}^2 \right) \\ & + \frac{Q^2}{4\pi\epsilon_0 a^3} \sum_{\mathbf{R}_B} \sum_{j=1}^3 h_{\mathbf{R}_B} h_{\mathbf{R}_B + \mathbf{e}_j} \\ & - \frac{Q^2}{8\pi\epsilon_0 a^3} \sum_{s=A,B} \sum_{\mathbf{R}_s} \sum_{l=-\infty}^{+\infty} 2 \left| \frac{a}{lL} \right|^3 h_{\mathbf{R}_s}^2 \\ & - \frac{Q^2}{8\pi\epsilon_0 a^3} \sum_{\mathbf{R}_B} \sum_{j=1}^3 \sum_{l=-\infty}^{+\infty} \frac{2 \left| \frac{lL}{a} \right|^2 - 1}{\left(1 + \left| \frac{lL}{a} \right|^2 \right)^{\frac{5}{2}}} h_{\mathbf{R}_B} h_{\mathbf{R}_B + \mathbf{e}_j} \\ & - \frac{Q^2}{8\pi\epsilon_0 a^3} \sum_{\mathbf{R}_A} \sum_{j=1}^3 \sum_{l=-\infty}^{+\infty} \frac{2 \left| \frac{lL}{a} \right|^2 - 1}{\left(1 + \left| \frac{lL}{a} \right|^2 \right)^{\frac{5}{2}}} h_{\mathbf{R}_A} h_{\mathbf{R}_A - \mathbf{e}_j}, \end{aligned} \quad (12)$$

where the primed summations exclude the $l = 0$ term. Here, $\Pi_{\mathbf{R}_s}$ is the conjugate momentum to the dynamical coordinate $h_{\mathbf{R}_s}$ corresponding to the meta-atom located at \mathbf{R}_s , and M is an effective mass. Next we introduce the bosonic operators

$$a_{\mathbf{R}_A} = \sqrt{\frac{M\omega_0}{2\hbar}} h_{\mathbf{R}_A} + i\sqrt{\frac{1}{2\hbar M\omega_0}} \Pi_{\mathbf{R}_A} \quad (13)$$

and

$$b_{\mathbf{R}_B} = \sqrt{\frac{M\omega_0}{2\hbar}} h_{\mathbf{R}_B} + i\sqrt{\frac{1}{2\hbar M\omega_0}} \Pi_{\mathbf{R}_B} \quad (14)$$

that annihilate quanta of the fundamental eigenmode on the meta-atom located at \mathbf{R}_A and \mathbf{R}_B , respectively, and satisfy the commutation relations $[a_{\mathbf{R}}, a_{\mathbf{R}'}^\dagger] = \delta_{\mathbf{R}\mathbf{R}'}$, $[b_{\mathbf{R}}, b_{\mathbf{R}'}^\dagger] = \delta_{\mathbf{R}\mathbf{R}'}$, and $[a_{\mathbf{R}}, b_{\mathbf{R}'}^\dagger] = 0$. In terms of these operators, the matter Hamiltonian (12) reads

$$\begin{aligned} H_{\text{mat}} = & \hbar\omega_0 \sum_{\mathbf{R}_A} a_{\mathbf{R}_A}^\dagger a_{\mathbf{R}_A} + \hbar\omega_0 \sum_{\mathbf{R}_B} b_{\mathbf{R}_B}^\dagger b_{\mathbf{R}_B} \\ & + \hbar\Omega(1 - \mathcal{I}) \sum_{\mathbf{R}_B} \sum_{j=1}^3 \left[b_{\mathbf{R}_B}^\dagger \left(a_{\mathbf{R}_B + \mathbf{e}_j} + a_{\mathbf{R}_B + \mathbf{e}_j}^\dagger \right) + \text{H.c.} \right] \\ & - \frac{\hbar}{2} \Omega S \sum_{\mathbf{R}_A} \left[a_{\mathbf{R}_A}^\dagger \left(a_{\mathbf{R}_A} + a_{\mathbf{R}_A}^\dagger \right) + \text{H.c.} \right] \\ & - \frac{\hbar}{2} \Omega S \sum_{\mathbf{R}_B} \left[b_{\mathbf{R}_B}^\dagger \left(b_{\mathbf{R}_B} + b_{\mathbf{R}_B}^\dagger \right) + \text{H.c.} \right] \end{aligned} \quad (15)$$

where $\Omega = Q^2/8\pi\epsilon_0 M\omega_0 a^3$ parametrizes the strength of the nearest-neighbor Coulomb interaction, and the parameters

$$S = 4 \sum_{l=1}^{\infty} \left(\frac{a}{lL} \right)^3, \quad \mathcal{I} = 2 \sum_{l=1}^{\infty} \frac{2 \left(\frac{lL}{a} \right)^2 - 1}{\left[1 + \left(\frac{lL}{a} \right)^2 \right]^{\frac{5}{2}}}, \quad (16)$$

encode renormalizations due to the cavity-induced image dipoles. We apply Born-von Kármán boundary conditions over a lattice with $\mathcal{N} \gg 1$ unit cells and introduce the Fourier transform of the bosonic operators $a_{\mathbf{R}_A} = \mathcal{N}^{-1/2} \sum_{\mathbf{q}} a_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{R}_A}$ and $b_{\mathbf{R}_B} = \mathcal{N}^{-1/2} \sum_{\mathbf{q}} b_{\mathbf{q}} e^{i\mathbf{q} \cdot \mathbf{R}_B}$, which transforms the matter Hamiltonian (15) into the local and block-diagonal form

$$\begin{aligned} H_{\text{mat}} = & \hbar \sum_{\mathbf{q}} (\omega_0 - \Omega S) \left(a_{\mathbf{q}}^\dagger a_{\mathbf{q}} + b_{\mathbf{q}}^\dagger b_{\mathbf{q}} \right) \\ & + \hbar \sum_{\mathbf{q}} \Omega(1 - \mathcal{I}) \left[f_{\mathbf{q}} b_{\mathbf{q}}^\dagger \left(a_{\mathbf{q}} + a_{-\mathbf{q}}^\dagger \right) + \text{H.c.} \right] \\ & - \hbar \sum_{\mathbf{q}} \frac{1}{2} \Omega S \left(a_{\mathbf{q}}^\dagger a_{-\mathbf{q}} + b_{\mathbf{q}}^\dagger b_{-\mathbf{q}} + \text{H.c.} \right). \end{aligned} \quad (17)$$

In the main text we do not present the non-resonant terms (e.g. $b_{\mathbf{q}}^\dagger a_{-\mathbf{q}}^\dagger$), leading to equation (1) where $\tilde{\omega}_0 = \omega_0 - \Omega S$ and $\tilde{\Omega} = \Omega(1 - \mathcal{I})$.

In the Coulomb gauge, the light-matter interaction is described by the minimal-coupling Hamiltonian [46] which, within the dipole approximation, reads

$$H_{\text{int}} = \underbrace{\frac{Q}{M} \sum_{s=A,B} \sum_{\mathbf{R}_s} \Pi_{\mathbf{R}_s} A_z(\mathbf{R}_s)}_{H_{\text{int}}^{(1)}} + \underbrace{\frac{Q^2}{2M} \sum_{s=A,B} \sum_{\mathbf{R}_s} A_z^2(\mathbf{R}_s)}_{H_{\text{int}}^{(2)}}, \quad (18)$$

where we have used $\Pi_{\mathbf{R}} = \Pi_{\mathbf{R}} \hat{\mathbf{z}}$. The vector potential can be decomposed into transverse electric (TE) and transverse magnetic (TM) modes of the cavity. However, the photons corresponding to the TE modes have an in-plane polarization, and therefore only TM modes contribute to the z -component of the vector potential

$$A_z(\mathbf{r}, z) = \sum_{\mathbf{q}mn} \sqrt{\frac{\hbar}{\epsilon_0 N_m \mathcal{A} L \omega_{\mathbf{q}mn}^{\text{ph}}}} \frac{|\mathbf{q} - \mathbf{G}_n|}{|\mathbf{q} - \mathbf{G}_n + \frac{m\pi}{L} \hat{\mathbf{z}}|} \cos\left(\frac{m\pi}{L} z\right) \left[c_{\mathbf{q}mn}^\dagger e^{i(\mathbf{q} - \mathbf{G}_n) \cdot \mathbf{r}} + c_{\mathbf{q}mn}^\dagger e^{-i(\mathbf{q} - \mathbf{G}_n) \cdot \mathbf{r}} \right], \quad (19)$$

where $\mathcal{A} = 3\sqrt{3}a^2/2$ is the area of a unit cell and $N_m = 1 + \delta_{m0}$. The bosonic operator $c_{\mathbf{q}mn}^\dagger$ creates a TM photon with wavevector \mathbf{q} in the first Brillouin zone and dispersion $\omega_{\mathbf{q}mn}^{\text{ph}} = c|\mathbf{q} - \mathbf{G}_n + \hat{\mathbf{z}}m\pi/L|$. Here, $\mathbf{G}_n = n_1 \mathbf{b}_1 + n_2 \mathbf{b}_2$ is a reciprocal lattice vector with primitive vectors \mathbf{b}_1 and \mathbf{b}_2 , where n indexes the set of ordered pairs of integers (n_1, n_2) , and m is a non-negative integer indexing the different TM cavity modes. Only TM photons with even m couple to the dipoles due to the parity selection rule at the centre of the cavity.

Substituting the vector potential (19) into equation (18) we obtain the light-matter interaction Hamiltonian

$$H_{\text{int}}^{(1)} = \hbar \sum_{\mathbf{q}mn} i\xi_{\mathbf{q}mn} \phi_n^* \left(a_{\mathbf{q}}^\dagger c_{\mathbf{q}mn} + a_{\mathbf{q}}^\dagger c_{-\mathbf{q}mn}^\dagger \right) + \hbar \sum_{\mathbf{q}mn} i\xi_{\mathbf{q}mn} \phi_n \left(b_{\mathbf{q}}^\dagger c_{\mathbf{q}mn} + b_{\mathbf{q}}^\dagger c_{-\mathbf{q}mn}^\dagger \right) + \text{H.c.} \quad (20)$$

and

$$H_{\text{int}}^{(2)} = \hbar \sum_{\mathbf{q}mm'n'n'} \frac{2\xi_{\mathbf{q}mn} \xi_{\mathbf{q}m'n'}}{\omega_0} \text{Re} \left(\phi_n \phi_{n'}^* \right) \left(c_{\mathbf{q}mn}^\dagger c_{\mathbf{q}m'n'} + c_{\mathbf{q}mn}^\dagger c_{-\mathbf{q}m'n'}^\dagger \right) + \text{H.c.} \quad (21)$$

The strength of the light-matter interaction is parametrized by

$$\xi_{\mathbf{q}mn} = \omega_0 \mathcal{F}(\omega_{\mathbf{q}mn}^{\text{ph}}) \frac{\omega_{\mathbf{q}0n}^{\text{ph}}}{\omega_{\mathbf{q}mn}^{\text{ph}}} \left(\frac{8\pi}{3\sqrt{3}N_m} \frac{a}{L} \frac{\Omega}{\omega_{\mathbf{q}mn}^{\text{ph}}} \right)^{\frac{1}{2}}, \quad (22)$$

where, to take into account the finite size of the meta-atoms, we have introduced a phenomenological function $\mathcal{F}(\omega_{\mathbf{q}mn}^{\text{ph}})$ that provides a smooth cut-off for the interaction with short-wavelength photonic modes where the dipole approximation breaks down. We choose the phenomenological cut-off function to be of the Fermi-Dirac distribution form

$$\mathcal{F}(\omega_{\mathbf{q}mn}^{\text{ph}}) = \frac{1}{1 + e^{2(\omega_{\mathbf{q}mn}^{\text{ph}} - 3\omega_0)/\omega_0}}, \quad (23)$$

which is smooth enough to avoid spurious artifacts appearing in the polariton spectrum. Finally, the free photonic Hamiltonian of the cavity reads

$$H_{\text{ph}} = \hbar \sum_{\mathbf{q}mn} \omega_{\mathbf{q}mn}^{\text{ph}} c_{\mathbf{q}mn}^\dagger c_{\mathbf{q}mn}. \quad (24)$$

In equations (3), (4), (5) in the main text we only present the contribution from the TEM mode ($m = 0$), dropping the corresponding index. In Supplementary Note 1 we discuss the effect of the higher order ($m \neq 0$) TM cavity modes for larger cavities.

Hopfield-Bogoliubov diagonalization

The polariton Hamiltonian $H_{\text{pol}} = H_{\text{mat}} + H_{\text{ph}} + H_{\text{int}}$, where H_{mat} is given by equation (17), H_{ph} by equation (24), and H_{int} by equations (20) and (21), can be recast into matrix form as $H_{\text{pol}} = \frac{1}{2} \sum_{\mathbf{q}} \Psi_{\mathbf{q}}^\dagger \mathcal{H}_{\mathbf{q}}^{\text{pol}} \Psi_{\mathbf{q}}$ where $\Psi_{\mathbf{q}}^\dagger = (\psi_{\mathbf{q}}^\dagger, C_{\mathbf{q}}^\dagger, \psi_{-\mathbf{q}}^T, C_{-\mathbf{q}}^T)$. Here, $\psi_{\mathbf{q}}^\dagger = (a_{\mathbf{q}}^\dagger, b_{\mathbf{q}}^\dagger)$ is the spinor creation operator in the matter sublattice space and

$C_{\mathbf{q}}^\dagger = (c_{\mathbf{q}1}^\dagger, c_{\mathbf{q}2}^\dagger, \dots, c_{\mathbf{q}p}^\dagger, \dots, c_{\mathbf{q}N}^\dagger)$ is the vector of TM photon creation operators, where p indexes the set of ordered triplets of integers (n_1, n_2, m) , and N is the total number of photonic operators considered. The Hermitian $[2(N+2)] \times [2(N+2)]$ matrix $\mathcal{H}_{\mathbf{q}}^{\text{pol}}$ can be written in block form as

$$\mathcal{H}_{\mathbf{q}}^{\text{pol}} = \begin{pmatrix} \mathcal{H}_{\mathbf{q}}^+ & \mathcal{H}_{\mathbf{q}}^- - \mathcal{W}_{\mathbf{q}} \\ (\mathcal{H}_{\mathbf{q}}^- - \mathcal{W}_{\mathbf{q}})^\dagger & (\mathcal{H}_{\mathbf{q}}^+)^* \end{pmatrix}, \quad (25)$$

where

$$\mathcal{W}_{\mathbf{q}} = \hbar \text{Diag} \left(\omega_0, \omega_0, \omega_{\mathbf{q}1}^{\text{ph}}, \omega_{\mathbf{q}2}^{\text{ph}}, \dots, \omega_{\mathbf{q}p}^{\text{ph}}, \dots, \omega_{\mathbf{q}N}^{\text{ph}} \right) \quad (26)$$

is the $(N+2) \times (N+2)$ diagonal matrix of resonant frequencies of the free oscillators. The $(N+2) \times (N+2)$ block matrices $\mathcal{H}_{\mathbf{q}}^\pm$ can be sub-divided into block matrices

$$\mathcal{H}_{\mathbf{q}}^\pm = \begin{pmatrix} \mathcal{H}_{\mathbf{q}}^{\text{mat}} & \mathcal{H}_{\mathbf{q}}^{\text{int}} \\ \pm (\mathcal{H}_{\mathbf{q}}^{\text{int}})^\dagger & \mathcal{H}_{\mathbf{q}}^{\text{ph}} \end{pmatrix}, \quad (27)$$

where the upper-diagonal block

$$\mathcal{H}_{\mathbf{q}}^{\text{mat}} = \hbar \begin{pmatrix} \tilde{\omega}_0 & \tilde{\Omega} f_{\mathbf{q}}^* \\ \tilde{\Omega} f_{\mathbf{q}} & \tilde{\omega}_0 \end{pmatrix} \quad (28)$$

is the 2×2 matrix in the matter subspace, and the lower-diagonal block $\mathcal{H}_{\mathbf{q}}^{\text{ph}}$ is the $N \times N$ matrix in the photonic subspace with components

$$(\mathcal{H}_{\mathbf{q}}^{\text{ph}})_{pp'} = \hbar \omega_{\mathbf{q}p}^{\text{ph}} \delta_{pp'} + 4\hbar \frac{\xi_{\mathbf{q}p} \xi_{\mathbf{q}p'}}{\omega_0} \text{Re} \left\{ \phi_p \phi_{p'}^* \right\}. \quad (29)$$

Finally, the off-diagonal block $\mathcal{H}_{\mathbf{q}}^{\text{int}}$ in equation (27) is the $2 \times N$ interaction matrix, where the p^{th} column reads

$$\left(\mathcal{H}_{\mathbf{q}}^{\text{int}} \right)_p = \hbar \begin{pmatrix} i\xi_{\mathbf{q}p} \phi_p^* \\ i\xi_{\mathbf{q}p} \phi_p \end{pmatrix}. \quad (30)$$

The polariton Hamiltonian H_{pol} is diagonalized by a generalized Hopfield-Bogoliubov transformation [48] $\Psi_{\mathbf{q}} = T_{\mathbf{q}} X_{\mathbf{q}}$, where $X_{\mathbf{q}}^\dagger = (\chi_{\mathbf{q}}^\dagger, \chi_{-\mathbf{q}}^T)$ and $\chi_{\mathbf{q}}^\dagger = (\gamma_{\mathbf{q}1}^\dagger, \gamma_{\mathbf{q}2}^\dagger, \dots, \gamma_{\mathbf{q}p}^\dagger, \dots, \gamma_{\mathbf{q}N+2}^\dagger)$. To ensure the invariance of the bosonic commutation relations for the transformed operators, $T_{\mathbf{q}}$ must be a $[2(N+2)] \times [2(N+2)]$ paraunitary matrix [69] that satisfies $T_{\mathbf{q}} \eta_z T_{\mathbf{q}}^\dagger = T_{\mathbf{q}}^\dagger \eta_z T_{\mathbf{q}} = \eta_z$, where $\eta_z = \sigma_z \otimes \mathbb{1}_2$ and σ_z is the Pauli matrix. The transformed bosonic operators $\gamma_{\mathbf{q}\nu}^\dagger = \Psi_{\mathbf{q}}^\dagger \eta_z |\Psi_{\mathbf{q}\nu}\rangle$ and $\gamma_{\mathbf{q}\nu} = \langle \Psi_{\mathbf{q}\nu} | \eta_z \Psi_{\mathbf{q}}$ that diagonalize the polariton Hamiltonian as

$$H_{\text{pol}} = \sum_{\mathbf{q}\nu} \hbar \omega_{\mathbf{q}\nu}^{\text{pol}} \gamma_{\mathbf{q}\nu}^\dagger \gamma_{\mathbf{q}\nu}, \quad (31)$$

create and annihilate polaritons in the ν^{th} band, respectively. The polariton dispersion $\omega_{\mathbf{q}\nu}^{\text{pol}}$ (black solid lines in Fig. 2c-e) and the corresponding linearly-independent eigenvectors $|\Psi_{\mathbf{q}\nu}\rangle$ (first two columns of $T_{\mathbf{q}}$) are determined from the positive eigenvalue solutions to the non-Hermitian eigenvalue equation $\eta_z \mathcal{H}_{\mathbf{q}}^{\text{pol}} |\Psi_{\mathbf{q}\nu}\rangle = \hbar \omega_{\mathbf{q}\nu}^{\text{pol}} |\Psi_{\mathbf{q}\nu}\rangle$.

Schrieffer-Wolff transformation

To obtain an effective two-band Hamiltonian in the matter sublattice space, we neglect non-resonant terms in the matter Hamiltonian (since $\Omega/\omega_0 \ll 1$ for practical realizations of the metasurface), but not in the light-matter interaction Hamiltonian since the photons are not resonant with the collective-dipoles near the corners of the Brillouin zone (see Fig. 2b). Next, we perform a unitary transformation

$$\bar{H}_{\text{pol}} = e^S H_{\text{pol}} e^{-S} = H_{\text{pol}} + [S, H_{\text{pol}}] + \frac{1}{2} [S, [S, H_{\text{pol}}]] + \dots \quad (32)$$

and impose the Schrieffer-Wolff condition [49]

$$[S, H_{\text{mat}} + H_{\text{ph}}] = -H_{\text{int}}^{(1)} \quad (33)$$

which eliminates the light-matter interaction to first order in $\xi_{\mathbf{q}mn}$. From equation (33) the particular form of the anti-Hermitian operator S reads

$$S = - \sum_{\mathbf{q}mn} \frac{i\xi_{\mathbf{q}mn}}{\omega_{\mathbf{q}mn}^{\text{ph}} - \tilde{\omega}_0} (\phi_n^* a_{\mathbf{q}}^\dagger + \phi_n b_{\mathbf{q}}^\dagger) c_{\mathbf{q}mn} + \sum_{\mathbf{q}mn} \frac{i\xi_{\mathbf{q}mn}}{\omega_{\mathbf{q}mn}^{\text{ph}} + \tilde{\omega}_0} (\phi_n^* a_{\mathbf{q}}^\dagger + \phi_n b_{\mathbf{q}}^\dagger) c_{-\mathbf{q}mn}^\dagger - \text{H.c.} \quad (34)$$

where we have used the approximation $|\omega_{\mathbf{q}mn}^{\text{ph}} \pm \omega_0| \gg \Omega|f_{\mathbf{q}}|$ that is valid near the K and K' points. Retaining leading-order terms in $\xi_{\mathbf{q}mn}$, the transformed polariton Hamiltonian (32) reads

$$\tilde{H}_{\text{pol}} \simeq H_{\text{mat}} + \underbrace{\frac{1}{2} [S, H_{\text{int}}^{(1)}]}_{\tilde{H}_{\text{mat}}} + \underbrace{H_{\text{ph}} + H_{\text{int}}^{(2)}}_{\tilde{H}_{\text{ph}}}, \quad (35)$$

where the matter and photonic subspaces are decoupled to quadratic order in $\xi_{\mathbf{q}mn}$. Calculating the commutator in equation (35) and extracting the Hamiltonian within the matter sublattice space we obtain the two-band Hamiltonian

$$\tilde{H}_{\text{mat}} = H_{\text{mat}} - 2\hbar \sum_{\mathbf{q}mn} \frac{\xi_{\mathbf{q}mn}^2 \omega_{\mathbf{q}mn}^{\text{ph}}}{(\omega_{\mathbf{q}mn}^{\text{ph}})^2 - \tilde{\omega}_0^2} (a_{\mathbf{q}}^\dagger a_{\mathbf{q}} + b_{\mathbf{q}}^\dagger b_{\mathbf{q}} + \phi_n^2 b_{\mathbf{q}}^\dagger a_{\mathbf{q}} + \phi_n^{*2} a_{\mathbf{q}}^\dagger b_{\mathbf{q}}). \quad (36)$$

In equation (8) in the main text we only present the contribution from the TEM mode ($m = 0$), dropping the corresponding index. We can recast the Hamiltonian (36) into matrix form as $\tilde{H}_{\text{mat}} = \sum_{\mathbf{q}} \psi_{\mathbf{q}}^\dagger \tilde{\mathcal{H}}_{\mathbf{q}}^{\text{mat}} \psi_{\mathbf{q}}$, with Bloch Hamiltonian

$$\tilde{\mathcal{H}}_{\mathbf{q}}^{\text{mat}} = \hbar \begin{pmatrix} W_{\mathbf{q}} & F_{\mathbf{q}}^* \\ F_{\mathbf{q}} & W_{\mathbf{q}} \end{pmatrix}. \quad (37)$$

Here $W_{\mathbf{q}} = \tilde{\omega}_0 - \Omega \sum_{mn} \Delta_{\mathbf{q}mn}$ and $F_{\mathbf{q}} = \tilde{\Omega} f_{\mathbf{q}} - \Omega \sum_{mn} \Delta_{\mathbf{q}mn} \phi_n^2$ with

$$\Delta_{\mathbf{q}mn} = \frac{16\pi}{3\sqrt{3}N_m} \frac{a}{L} \frac{\omega_0^2}{(\omega_{\mathbf{q}mn}^{\text{ph}})^2 - \tilde{\omega}_0^2} \left(\frac{\omega_{\mathbf{q}0n}^{\text{ph}}}{\omega_{\mathbf{q}mn}^{\text{ph}}} \right)^2 \mathcal{F}^2(\omega_{\mathbf{q}mn}^{\text{ph}}). \quad (38)$$

Diagonalizing \tilde{H}_{mat} leads to the two-band dispersion $\tilde{\omega}_{\mathbf{q}\tau}^{\text{mat}} = W_{\mathbf{q}} + \tau|F_{\mathbf{q}}|$, which is indicated by the orange-dashed lines in Fig. 2c-e. The corresponding spinor eigenstates $|\psi_{\mathbf{q}\tau}\rangle = (1, \tau e^{i\varphi_{\mathbf{q}}})^T / \sqrt{2}$, where $\varphi_{\mathbf{q}} = \arg(F_{\mathbf{q}})$, can be represented by the pseudospin vector $\mathbf{S}_{\mathbf{q}\tau} = \tau(\cos \varphi_{\mathbf{q}}, \sin \varphi_{\mathbf{q}}, 0)$ from which we obtain the pseudospin vector field plots in Fig 5a-c.

Expansion of the effective two-band Hamiltonian

Near the K point, the function $\Delta_{\mathbf{q}mn}$, given by equation (38), expands as

$$\begin{aligned} \Delta_{\mathbf{k}mn} &\simeq \Delta_{\mathbf{K}mn}^{(0)} - a^2 \Delta_{\mathbf{K}mn}^{(1)} [(\mathbf{K} - \mathbf{G}_n)_x k_x + (\mathbf{K} - \mathbf{G}_n)_y k_y] \\ &+ \frac{1}{2} [-a^2 \Delta_{\mathbf{K}mn}^{(1)} + a^4 \Delta_{\mathbf{K}mn}^{(2)} (\mathbf{K} - \mathbf{G}_n)_x^2] k_x^2 \\ &+ \frac{1}{2} [-a^2 \Delta_{\mathbf{K}mn}^{(1)} + a^4 \Delta_{\mathbf{K}mn}^{(2)} (\mathbf{K} - \mathbf{G}_n)_y^2] k_y^2 \\ &+ a^4 \Delta_{\mathbf{K}mn}^{(2)} (\mathbf{K} - \mathbf{G}_n)_x (\mathbf{K} - \mathbf{G}_n)_y k_x k_y \end{aligned} \quad (39)$$

to quadratic order in \mathbf{k} , where the real parameters $\Delta_{\mathbf{K}mn}^{(v)}$ ($v = 0, 1, 2$) depend only on the photon frequencies $\omega_{\mathbf{K}mn}^{\text{ph}}$ at the K point. Collecting the contributions from the degenerate photons (see Supplementary Note 2 for details) we obtain the effective Hamiltonian (9), where parameters are given by

$$\frac{\tilde{\omega}_0}{\omega_0} = 1 - \frac{\Omega}{\omega_0} S - \frac{\Omega}{\omega_0} \sum_{mn} \Delta_{\mathbf{K}mn}^{(0)}, \quad (40)$$

$$\frac{\tilde{v}}{v} = 1 - \mathcal{I} - \frac{4\pi}{27} \sum_{mn} A_n \Delta_{\mathbf{K}mn}^{(1)}, \quad (41)$$

$$\frac{\tilde{t}}{t} = 1 - \mathcal{I} - \frac{8\pi^2}{81} \sum_{mn} B_n \Delta_{\mathbf{K}mn}^{(2)}, \quad (42)$$

and

$$\frac{D}{\omega_0 a^2} = \frac{\Omega}{\omega_0} \sum_{mn} \left(\frac{4\pi^2}{27} C_n \Delta_{\mathbf{K}mn}^{(2)} - \frac{1}{2} \Delta_{\mathbf{K}mn}^{(1)} \right), \quad (43)$$

with

$$A_n = \frac{\sqrt{3}}{2} (2 - 3n_1) \cos \left[\frac{4\pi}{3} (n_1 + n_2) \right] + \frac{1}{2} (6n_2 - 3n_1) \sin \left[\frac{4\pi}{3} (n_1 + n_2) \right], \quad (44)$$

$$B_n = (3n_1^2 - 6n_2^2 - 6n_1 + 6n_1 n_2 + 2) \cos \left[\frac{4\pi}{3} (n_1 + n_2) \right] + \sqrt{3} (2n_1 - 4n_2 + 6n_1 n_2 - 3n_1^2) \sin \left[\frac{4\pi}{3} (n_1 + n_2) \right], \quad (45)$$

and

$$C_n = 1 + 3n_1 (n_1 - 1) + 3n_2 (n_2 - n_1). \quad (46)$$

For brevity, we retain only the dominant ($m = 0$) TEM contribution for the plots in Fig. 4, where the coefficients $\Delta_{\mathbf{K}0n}^{(v)}$ in equation (39) are given by

$$\Delta_{\mathbf{K}0n}^{(0)} = 2 \left(\frac{4\pi}{3\sqrt{3}} \right) \left(\frac{a}{L} \right) \left[\frac{\omega_0^2}{(\omega_{\mathbf{K}0n}^{\text{ph}})^2 - \tilde{\omega}_0^2} \right] \mathcal{F}^2(\omega_{\mathbf{K}0n}^{\text{ph}}), \quad (47)$$

$$\begin{aligned} \Delta_{\mathbf{K}0n}^{(1)} &= 4 \left(\frac{4\pi}{3\sqrt{3}} \right)^{-1} \left(\frac{a}{L} \right) \left(\frac{\omega_{\mathbf{K}00}^{\text{ph}}}{\omega_0} \right)^2 \left[\frac{\omega_0^2}{(\omega_{\mathbf{K}0n}^{\text{ph}})^2 - \tilde{\omega}_0^2} \right]^2 \\ &\left\{ \mathcal{F}^2(\omega_{\mathbf{K}0n}^{\text{ph}}) - \frac{\partial [\mathcal{F}^2(\omega_{\mathbf{K}0n}^{\text{ph}})]}{\partial \omega_{\mathbf{K}0n}^{\text{ph}}} \frac{(\omega_{\mathbf{K}0n}^{\text{ph}})^2 - \tilde{\omega}_0^2}{2\omega_{\mathbf{K}0n}^{\text{ph}}} \right\}, \end{aligned} \quad (48)$$

and

$$\begin{aligned} \Delta_{\mathbf{K}0n}^{(2)} &= 16 \left(\frac{4\pi}{3\sqrt{3}} \right)^{-3} \left(\frac{a}{L} \right) \left(\frac{\omega_{\mathbf{K}00}^{\text{ph}}}{\omega_0} \right)^4 \left[\frac{\omega_0^2}{(\omega_{\mathbf{K}0n}^{\text{ph}})^2 - \tilde{\omega}_0^2} \right]^3 \\ &\left\{ \mathcal{F}^2(\omega_{\mathbf{K}0n}^{\text{ph}}) - \frac{\partial [\mathcal{F}^2(\omega_{\mathbf{K}0n}^{\text{ph}})]}{\partial \omega_{\mathbf{K}0n}^{\text{ph}}} \frac{[5(\omega_{\mathbf{K}0n}^{\text{ph}})^2 - \tilde{\omega}_0^2] [(\omega_{\mathbf{K}0n}^{\text{ph}})^2 - \tilde{\omega}_0^2]}{8(\omega_{\mathbf{K}0n}^{\text{ph}})^3} \right. \\ &\left. + \frac{\partial^2 [\mathcal{F}^2(\omega_{\mathbf{K}0n}^{\text{ph}})]}{(\partial \omega_{\mathbf{K}0n}^{\text{ph}})^2} \frac{[(\omega_{\mathbf{K}0n}^{\text{ph}})^2 - \tilde{\omega}_0^2]^2}{8(\omega_{\mathbf{K}0n}^{\text{ph}})^2} \right\}. \end{aligned} \quad (49)$$

Data Availability

All relevant data are available from the corresponding authors upon reasonable request.

- [1] Novoselov, K. S. et al. Electric field effect in atomically thin carbon films. *Science* **306**, 666-669 (2004).
- [2] Polini, M., Guinea, F., Lewenstein, M., Manoharan, H. C. & Pellegrini, V. Artificial honeycomb lattices for electrons, atoms and photons. *Nat. Nanotechnol.* **8**, 625-633 (2013).
- [3] Zhu, S.-L., Wang, B. & Duan, L.-M. Simulation and detection of Dirac fermions with cold atoms in an optical lattice. *Phys. Rev. Lett.* **98**, 260402 (2007).
- [4] Peleg, O. et al. Conical diffraction and gap solitons in honeycomb photonic lattices. *Phys. Rev. Lett.* **98**, 103901 (2007).
- [5] Han, D., Lai, Y., Zi, J., Zhang, Z. Q. & Chan, C. T. Dirac spectra and edge states in honeycomb plasmonic lattices. *Phys. Rev. Lett.* **102**, 123904 (2009).

- [6] Gibertini, M. et al. Engineering artificial graphene in a two-dimensional electron gas. *Phys. Rev. B* **79**, 241406(R) (2009).
- [7] Torrent, D. & Sánchez-Dehesa, J. Acoustic analogue of graphene: Observation of Dirac cones in acoustic surface waves. *Phys. Rev. Lett.* **108**, 174301 (2012).
- [8] Gomes, K. K., Mar, W., Ko, W., Guinea, F. & Manoharan, H. C. Designer Dirac fermions and topological phases in molecular graphene. *Nature* **483**, 306-310 (2012).
- [9] Bellec, M., Kuhl, U., Montambaux, G. & Mortessagne, F. Tight-binding couplings in microwave artificial graphene. *Phys. Rev. B* **88**, 115437 (2013).
- [10] Weick, G., Woollacott, C., Barnes, W. L., Hess, O. & Mariani, E. Dirac-like plasmons in honeycomb lattices of metallic nanoparticles. *Phys. Rev. Lett.* **110**, 106801 (2013).
- [11] Yu, S. et al. Surface phononic graphene. *Nat. Mater.* **15**, 1243?1247 (2016).
- [12] Castro Neto, A. H., Peres, N. M. R., Novoselov, K. S., Geim, A. K. & Guinea, F. The electronic properties of graphene. *Rev. Mod. Phys.* **81**, 109-162 (2009).
- [13] Katsnelson, M. I., Novoselov, K. S. & Geim, A. K. Chiral tunnelling and the Klein paradox in graphene. *Nat. Phys.* **2**, 620-625 (2006).
- [14] Tarruell, L., Greif, D., Uehlinger, T., Jotzu, G. & Esslinger, T. Creating, moving and merging Dirac points with a Fermi gas in a tunable honeycomb lattice. *Nature* **483**, 302-305 (2012).
- [15] Bellec, M., Kuhl, U., Montambaux, G. & Mortessagne, F. Topological transition of Dirac points in a microwave experiment. *Phys. Rev. Lett.* **110**, 033902 (2013).
- [16] Rechtsman, M. C. et al. Topological creation and destruction of edge states in photonic graphene. *Phys. Rev. Lett.* **111**, 103901 (2013).
- [17] Rechtsman, M. C. et al. Strain-induced pseudomagnetic field and photonic Landau levels in dielectric structures. *Nat. Photonics* **7**, 153-158 (2013).
- [18] Rechtsman, M. C. et al. Photonic Floquet topological insulators. *Nature* **496**, 196-200 (2013).
- [19] Ni, X., Purtseladze, D., Smirnova, D. A., Slobzhanyuk, A. & Alù, A. Spin and valley polarized one-way Klein tunneling in photonic topological insulators. Preprint at <https://arxiv.org/abs/1707.05802> (2017).
- [20] Hasegawa, Y., Konno, R., Nakano, H. & Kohmoto, M. Zero modes of tight-binding electrons on the honeycomb lattice. *Phys. Rev. B* **74**, 033413 (2006).
- [21] Wunsch, B., Guinea, F. & Sols, F. Dirac-point engineering and topological phase transitions in honeycomb optical lattices. *New J. Phys.* **10**, 103027 (2008).
- [22] Pereira, V. M., Castro Neto, A. H. & Peres, N. M. R. Tight-binding approach to uniaxial strain in graphene. *Phys. Rev. B* **80**, 045401 (2009).
- [23] Montambaux, G., Piéchon, F., Fuchs, J. N. & Goerbig, M. O. Merging of Dirac points in a two-dimensional crystal. *Phys. Rev. B* **80**, 153412 (2009).
- [24] Guinea, F., Katsnelson, M. I. & Geim, A. K. Energy gaps and a zero-field quantum Hall effect in graphene by strain engineering. *Nat. Phys.* **6**, 30-33 (2009).
- [25] Jacqmin, T. et al. Direct observation of Dirac cones and a flatband in a honeycomb lattice for polaritons. *Phys. Rev. Lett.* **112**, 116402 (2014).
- [26] Nalitimov, A. V., Solnyshkov, D. D. & Malpuech, G. Polariton Z topological insulator. *Phys. Rev. Lett.* **114**, 116401 (2015).
- [27] Karzig, T., Bardyn, C. E., Lindner, N. H. & Refael, G. Topological polaritons. *Phys. Rev. X* **5**, 031001 (2015).
- [28] Bardyn, C. E., Karzig, T., Refael, G. & Liew, T. C. H. Topological polaritons and excitons in garden-variety systems. *Phys. Rev. B* **91**, 161413(R) (2015).
- [29] Yi, K. & Karzig, T. Topological polaritons from photonic Dirac cones coupled to excitons in a magnetic field. *Phys. Rev. B* **93**, 104303 (2016).
- [30] Yuen-Zhou, J. et al. Plexciton Dirac points and topological modes. *Nat. Commun.* **7**, 11783 (2016).
- [31] Soluyanov, A. A. et al. Type-II Weyl semimetals. *Nature* **527**, 495?498 (2015).
- [32] Deng, K. et al. Experimental observation of topological Fermi arcs in type-II Weyl semimetal MoTe₂. *Nat. Phys.* **12**, 1105?1110 (2016).
- [33] Huang, L. et al. Spectroscopic evidence for a type II Weyl semimetallic state in MoTe₂. *Nat. Mater.* **15**, 1155-1160 (2016).
- [34] Huang, H., Zhou, S. & Duan, W. Type-II Dirac fermions in the PtSe₂ class of transition metal dichalcogenides. *Phys. Rev. B* **94**, 121117(R) (2016).
- [35] Yan, M. et al. Lorentz-violating type-II Dirac fermions in transition metal dichalcogenide PtTe₂. *Nat. Commun.* **8**, 257 (2017).
- [36] Xiao, M., Lin, Q. & Fan, S. Hyperbolic Weyl point in reciprocal chiral metamaterials. *Phys. Rev. Lett.* **117**, 057401 (2016).
- [37] Chen, W.-J., Xiao, M. & Chan, C. T. Photonic crystals possessing multiple Weyl points and the experimental observation of robust surface states. *Nat. Commun.* **7**, 13038 (2016).
- [38] Noh, J. et al. Experimental observation of optical Weyl points and Fermi arc-like surface states. *Nat. Phys.* **13**, 611?618 (2017).
- [39] Yang, B. et al. Direct observation of topological surface-state arcs in photonic metamaterials. *Nat. Commun.* **8**, 97 (2017).
- [40] Wang, H.-X., Chen, Y., Hang, Z. H., Kee, H.-Y. & Jiang, J.-H. Type-II Dirac photons. *npj Quantum Mater.* **2**, 54 (2017).
- [41] Lin, J. Y., Hu, N. C., Chen, Y. J., Lee, C. H. & Zhang, X. Line nodes, Dirac points, and Lifshitz transition in two-dimensional nonsymmorphic photonic crystals. *Phys. Rev. B* **96**, 075438 (2017).
- [42] Pyrialakos, G. G., Nye, N. S., Kantartzis, N. V & Christodoulides, D. N. Emergence of type-II Dirac points in graphynelike photonic lattices. *Phys. Rev. Lett.* **119**, 113901 (2017).
- [43] Alù, A. First-principles homogenization theory for periodic metamaterials. *Phys. Rev. B* **84**, 075153 (2011).
- [44] Yves, S. et al. Crystalline metamaterials for topological properties at subwavelength scales. *Nat. Commun.* **8**, 16023 (2017).
- [45] Kaina, N., Lemoult, F., Fink, M. & Lerosey, G. Negative refractive index and acoustic superlens from multiple scattering in single negative metamaterials. *Nature* **525**, 77?81 (2015).
- [46] Craig, D. P. & Thirunamachandran, T. *Molecular Quantum Electrodynamics: An Introduction to Radiation-Molecule Interactions*. (Academic Press, London, 1984).
- [47] Fuchs, J. N., Piéchon, F., Goerbig, M. O. & Montambaux, G. Topological Berry phase and semiclassical quantization of cyclotron orbits for two dimensional electrons in coupled band models. *Eur. Phys. J. B* **77**, 351-362 (2010).
- [48] Hopfield, J. J. Theory of the contribution of excitons to the complex dielectric constant of crystals. *Phys. Rev. Lett.* **112**, 1555-1567 (1958).
- [49] Schrieffer, J. R. & Wolff, P. A. Relation between the Anderson and Kondo Hamiltonians. *Phys. Rev.* **149**, 491-492 (1966).
- [50] Goerbig, M. O., Fuchs, J. N., Montambaux, G. & Piéchon, F. Tilted anisotropic Dirac cones in quinoid-type graphene and $\alpha - (\text{BEDT-TTF})_2\text{I}_3$. *Phys. Rev. B* **78**, 045415 (2008).
- [51] McCann, E. & Fal'ko, V. I. Landau-level degeneracy and quantum Hall effect in a graphite bilayer. *Phys. Rev. Lett.* **96**, 086805 (2006).
- [52] McCann, E. & Koshino, M. The electronic properties of bilayer graphene. *Reports Prog. Phys.* **76**, 056503 (2013).
- [53] Mucha-Kruczyński, M., Aleiner, I. L. & Fal'ko, V. I. Strained bilayer graphene: Band structure topology and Landau level spectrum. *Phys. Rev. B* **84**, 041404(R) (2011).
- [54] Mariani, E., Pearce, A. J. & Von Oppen, F. Fictitious gauge fields in bilayer graphene. *Phys. Rev. B* **86**, 165448 (2012).
- [55] Verberck, B., Partoens, B., Peeters, F. M. & Trauzettel, B. Strain-induced band gaps in bilayer graphene. *Phys. Rev. B* **85**, 125403 (2012).
- [56] Pearce, A. J., Cavaliere, F. & Mariani, E. Conductance and shot noise in strained bilayer graphene. *J. Phys. Condens. Matter* **25**, 375301 (2013).
- [57] Bena, C. & Simon, L. Dirac point metamorphosis from third-neighbor couplings in graphene and related materials. *Phys. Rev. B* **83**, 115404 (2011).
- [58] Montambaux, G. An equivalence between monolayer and bilayer honeycomb lattices. *Eur. Phys. J. B* **85**, 375 (2012).
- [59] Sticlet, D. & Piéchon, F. Distant-neighbor hopping in graphene and Haldane models. *Phys. Rev. B* **87**, 115402 (2013).
- [60] Bhattacharya, U., Hutchinson, J. & Dutta, A. Quenching in Chern insulators with satellite Dirac points: The fate of edge states. *Phys. Rev. B* **95**, 144304 (2017).
- [61] Maier, S. A. *Plasmonics: Fundamentals and Applications* (Springer-Verlag, Berlin, 2007).
- [62] Renger, J., Quidant, R., Van Hulst, N., Palomba, S. & Novotny, L. Free-space excitation of propagating surface plasmon polaritons by nonlinear

- four-wave mixing. *Phys. Rev. Lett.* **103**, 266802 (2009).
- [63] Cheianov, V. V, Fal'ko, V. & Altshuler, B. L. The focusing of electron flow and a Veselago lens in graphene p-n junctions. *Science* **315**, 1252-1255 (2007).
- [64] Lee, G.-H., Park, G. & Lee, H. Observation of negative refraction of Dirac fermions in graphene. *Nat. Phys.* **11**, 925-929 (2015).
- [65] Raoux, A. et al. Velocity-modulation control of electron-wave propagation in graphene. *Phys. Rev. B* **81**, 073407 (2010).
- [66] Downing, C. A. & Portnoi, M. E. Localization of massless Dirac particles via spatial modulations of the Fermi velocity. *J. Phys. Condens. Matter* **29**, 315301 (2017).
- [67] Jackson, J. D. *Classical Electrodynamics*, 3rd ed. (Wiley, New York, 1999).
- [68] Power, E. A. & Thirunamachandran, T. Quantum electrodynamics in a cavity. *Phys. Rev. A* **25**, 2473-2484 (1982).
- [69] Colpa, J. H. P. Diagonalization of the quadratic boson Hamiltonian. *Physica A* **93**, 327-353 (1978).

ACKNOWLEDGMENTS

We would like to thank Ian Hooper for useful discussions and help with the numerical simulations. C.-R.M. and T.J.S. acknowledge financial support from the Engineering and Physical Sciences Research Council (EPSRC) of the United Kingdom. In addition, C.-R.M. acknowledges the EPSRC Centre for Doctoral Training in Metamaterials (Grant No. EP/L015331/1) and QinetiQ for additional funding. G.W. acknowledges financial support from Agence Nationale de la Recherche (Project ANR-14-CE26-0005 Q-

MetaMat) and the Centre National de la Recherche Scientifique through the Projet International de Coopération Scientifique program (Contract No. 6384 APAG). W.L.B. acknowledges the financial support from EPSRC (Grant No. EP/K041150/1) and EU ERC project Photmat (ERC-2016-ADG-742222). E.M. acknowledges financial support from the Leverhulme Trust (Research Project Grant RPG-2015-101), and the Royal Society (International Exchange Grant No. IE140367, Newton Mobility Grants NI160073, and Theo Murphy Award TM160190).

AUTHOR CONTRIBUTIONS

C.-R.M. performed the theoretical calculations and wrote the manuscript with E.M.; T.J.S. contributed to the theoretical calculations; E.M. and G.W. conceived the project; and E.M. and W.L.B. supervised the project. All authors commented on the manuscript.


ADDITIONAL INFORMATION

Supplementary information accompanies this work. Correspondence and requests for materials should be addressed to C.-R.M. (email: cm433@exeter.ac.uk) or to E.M. (email: E.Mariani@exeter.ac.uk)


COMPETING FINANCIAL INTERESTS

The authors declare no competing financial interests.


Supplementary Information: Manipulating Type-I and Type-II Dirac Polaritons in Cavity-Embedded Honeycomb Metasurfaces




Supplementary Figure 1. **Quasistatic dispersion beyond the nearest-neighbor approximation.** **a**, Quasistatic dispersion within the nearest-neighbor approximation for cavity heights of $L = 1.3a$ (blue dashed line) and $L = a$ (orange dot-dashed line). For comparison we also have plotted the quasistatic dispersion without a cavity (black solid line). One observes that the effect of the image dipoles is to renormalize the resonant frequency and the Coulomb interaction strength, resulting in the quasistatic dispersion shifting down in frequency and reducing in bandwidth. Panel **b** is the same as **a** but with the full quasistatic dispersion (S8) that includes Coulomb interactions beyond nearest-neighbors until convergence. Long-range Coulomb interactions introduce an asymmetry between the upper and lower quasistatic bands as shown in **b**, but the CDPs remain robust at the K and K' points. As the cavity height is reduced, the image dipoles quench long-range Coulomb interactions, which become negligible for cavity heights on the order of the nearest-neighbor separation. All plots obtained with $\Omega = 0.01\omega_0$.



Supplementary Figure 2. **Evolution of the polariton dispersion beyond the nearest-neighbor approximation.** Full polariton dispersion that includes long-range Coulomb interactions (solid black lines) compared to the polariton dispersion obtained within the nearest-neighbor approximation (orange dashed lines) for **a** very large ($L = 15a$), **b** sub-critical ($L = 5a$), **c** critical ($L = 1.75a$), and **d** super-critical ($L = a$) cavity heights, respectively. Higher-order cavity modes can be observed for very large cavities as seen in **a**. Along the $\Gamma - K(K')$ directions the band crossings between the photon dispersion and the lower quasistatic band are protected, which gives rise to many additional linear-band degeneracies within the light-cone. As the cavity height is decreased, these additional Dirac points are driven towards the Γ point (opposite direction to the SDPs emerging from the TEM mode) where they annihilate one another. Comparing panels **a** and **b**, one can see that the dispersion near the CDPs does not noticeably change, and qualitative changes outside the light-cone only occur for small cavity heights where the higher-order cavity modes become increasingly detuned from the dipole resonances (see panels **b-d**). For small cavity heights, the nearest-neighbor approximation becomes increasingly accurate and indistinguishable from the full polariton dispersion (see **d** for example). Plots obtained with parameters $\omega_{\mathbf{K}0}^{\text{ph}} = 2.5\omega_0$ and $\Omega = 0.01\omega_0$.



Supplementary Figure 3. **Numerical simulation of the polariton dispersion for a cavity-embedded honeycomb array of plasmonic nanorods.** We consider ellipsoidal nanorods (prolate spheroids) with semiminor axis and semimajor axis of lengths 10 nm and 20 nm, respectively, and model them using the simple Drude model permittivity with plasma frequency $\omega_p = 1.32 \times 10^{16} \text{ rad s}^{-1}$ and nonradiative damping frequency $\gamma = 6.9 \times 10^{13} \text{ s}^{-1}$. The plasmonic nanorods are arranged in a honeycomb array with nearest-neighbor separation $a = 45 \text{ nm}$. We apply perfect electric conductor boundary conditions for the cavity walls and periodic boundary conditions surrounding the unit cell (see inset on panel **c**). **a-c**, Numerical simulations of the polariton dispersion for the lowest two bands, calculated using the eigenmode solver in COMSOL Multiphysics, for cavity heights of **a** $L = 3a$, **b** $L = 1.7a$, and **c** $L = 1.3a$, respectively. The orange dots represent numerical solutions and the dashed line is a guide for the eye. The frequencies are normalized with respect to $\omega_0 = 5.3 \times 10^{15} \text{ rad s}^{-1}$.



Supplementary Figure 4. **Numerical simulation of the polariton dispersion for a cavity-embedded honeycomb array of microwave helical resonators.** We consider microwave helical resonators with minor radius 2 mm, major radius 6 mm, pitch 6 mm, and 5 turns (see inset on panel **c**). We model the helices and cavity walls as perfect electric conductors, which is a good approximation at microwave frequencies, and we apply periodic boundary conditions surrounding the unit cell. The helices are arranged in a honeycomb array with nearest-neighbor separation $a = 45 \text{ mm}$. **a-c**, Numerical simulations of the polariton dispersion for the lowest two bands, calculated using the eigenmode solver in COMSOL Multiphysics, for cavity heights of **a** $L = 3.33a$, **b** $L = 1.56a$, and **c** $L = 1.11a$, respectively. The orange dots represent numerical solutions and the dashed line is a guide for the eye. The frequencies are normalized with respect to $\omega_0 = 6.2 \times 10^9 \text{ rad s}^{-1}$.

SUPPLEMENTARY NOTE 1: BEYOND THE NEAREST-NEIGHBOR APPROXIMATION

Including all Coulomb interactions beyond the nearest-neighbor approximation leads to the full matter Hamiltonian within the dipole approximation

$$H_{\text{mat}} = \hbar\omega_0 \sum_{\mathbf{q}} (a_{\mathbf{q}}^\dagger a_{\mathbf{q}} + b_{\mathbf{q}}^\dagger b_{\mathbf{q}}) + \hbar\Omega \sum_{\mathbf{q}} \left[f_{\mathbf{q}}^{\text{AB}} b_{\mathbf{q}}^\dagger (a_{\mathbf{q}} + a_{-\mathbf{q}}^\dagger) + \text{H.c.} \right] \\ + \frac{\hbar\Omega}{2} \sum_{\mathbf{q}} \left[f_{\mathbf{q}}^{\text{AA}} a_{\mathbf{q}}^\dagger (a_{\mathbf{q}} + a_{-\mathbf{q}}^\dagger) + \text{H.c.} \right] + \frac{\hbar\Omega}{2} \sum_{\mathbf{q}} \left[f_{\mathbf{q}}^{\text{BB}} b_{\mathbf{q}}^\dagger (b_{\mathbf{q}} + b_{-\mathbf{q}}^\dagger) + \text{H.c.} \right], \quad (\text{S1})$$

where

$$f_{\mathbf{q}}^{\text{AA}} = f_{\mathbf{q}}^{\text{BB}} = -\mathcal{S} + \sum_{\mathbf{R} \neq 0} \left(\frac{a}{|\mathbf{R}|} \right)^3 (1 - \mathcal{I}_{\mathbf{R}}^{\text{AA}}) e^{i\mathbf{q} \cdot \mathbf{R}} \quad (\text{S2})$$

and

$$f_{\mathbf{q}}^{\text{AB}} = \sum_{\mathbf{R}} \left(\frac{a}{|\mathbf{R} + \mathbf{e}_1|} \right)^3 (1 - \mathcal{I}_{\mathbf{R}}^{\text{AB}}) e^{i\mathbf{q} \cdot (\mathbf{R} + \mathbf{e}_1)}. \quad (\text{S3})$$

In equations (S2) and (S3), the parameters

$$\mathcal{I}_{\mathbf{R}}^{\text{AA}} = 2 \sum_{l=1}^{\infty} \frac{2 \left(\frac{lL}{|\mathbf{R}|} \right)^2 - 1}{\left[1 + \left(\frac{lL}{|\mathbf{R}|} \right)^2 \right]^{\frac{5}{2}}}, \quad \mathcal{I}_{\mathbf{R}}^{\text{AB}} = 2 \sum_{l=1}^{\infty} \frac{2 \left(\frac{lL}{|\mathbf{R} + \mathbf{e}_1|} \right)^2 - 1}{\left[1 + \left(\frac{lL}{|\mathbf{R} + \mathbf{e}_1|} \right)^2 \right]^{\frac{5}{2}}} \quad (\text{S4})$$

encode the renormalizations due to the interaction with cavity-induced image dipoles of neighboring meta-atoms, and \mathcal{S} , given by equation (16) in the Methods, encodes renormalizations due to interactions with their self-images. We can recast the matter Hamiltonian (S1) into matrix form as $H_{\text{mat}} = \frac{1}{2} \sum_{\mathbf{q}} \Psi_{\mathbf{q}}^\dagger \mathcal{H}_{\mathbf{q}}^{\text{mat}} \Psi_{\mathbf{q}}$, where $\Psi_{\mathbf{q}}^\dagger = (\psi_{\mathbf{q}}^\dagger, \psi_{-\mathbf{q}}^{\text{T}})$ and $\psi_{\mathbf{q}}^\dagger = (a_{\mathbf{q}}^\dagger, b_{\mathbf{q}}^\dagger)$. The Hermitian 4×4 matrix $\mathcal{H}_{\mathbf{q}}^{\text{mat}}$ can be written in block form as

$$\mathcal{H}_{\mathbf{q}}^{\text{mat}} = \begin{pmatrix} \mathcal{H}_{\mathbf{q}}^{\text{mat}} & \mathcal{H}_{\mathbf{q}}^{\text{mat}} - \hbar\omega_0 \mathbb{1}_2 \\ \mathcal{H}_{\mathbf{q}}^{\text{mat}} - \hbar\omega_0 \mathbb{1}_2 & \mathcal{H}_{\mathbf{q}}^{\text{mat}} \end{pmatrix}, \quad (\text{S5})$$

with

$$\mathcal{H}_{\mathbf{q}}^{\text{mat}} = \hbar \begin{pmatrix} \omega_0 + \Omega \text{Re}(f_{\mathbf{q}}^{\text{AA}}) & \Omega (f_{\mathbf{q}}^{\text{AB}})^* \\ \Omega f_{\mathbf{q}}^{\text{AB}} & \omega_0 + \Omega \text{Re}(f_{\mathbf{q}}^{\text{BB}}) \end{pmatrix}. \quad (\text{S6})$$

We diagonalize H_{mat} via a Bogoliubov transformation $\Psi_{\mathbf{q}} = T_{\mathbf{q}} X_{\mathbf{q}}$, where $X_{\mathbf{q}}^\dagger = (\chi_{\mathbf{q}}^\dagger, \chi_{-\mathbf{q}}^{\text{T}})$ and $\chi_{\mathbf{q}}^\dagger = (\beta_{\mathbf{q}+}^\dagger, \beta_{\mathbf{q}-}^\dagger)$, with $T_{\mathbf{q}}$ being a paraunitary matrix (see Methods). The bosonic operators $\beta_{\mathbf{q}\tau}^\dagger = (\cosh \vartheta_{\mathbf{q}\tau} \psi_{\mathbf{q}}^\dagger - \sinh \vartheta_{\mathbf{q}\tau} \psi_{-\mathbf{q}}^{\text{T}}) |\psi_{\mathbf{q}\tau}\rangle$ diagonalize the matter Hamiltonian (S1) as

$$H_{\text{mat}} = \sum_{\tau=\pm} \sum_{\mathbf{q}} \hbar\omega_{\mathbf{q}\tau}^{\text{mat}} \beta_{\mathbf{q}\tau}^\dagger \beta_{\mathbf{q}\tau}, \quad (\text{S7})$$

with quasistatic dispersion

$$\omega_{\mathbf{q}\tau}^{\text{mat}} = \sqrt{\omega_0 + 2\Omega \Re(f_{\mathbf{q}}^{\text{AA}}) + 2\tau\Omega |f_{\mathbf{q}}^{\text{AB}}|}. \quad (\text{S8})$$

As described in the main text, the spinors $|\psi_{\mathbf{q}\tau}\rangle = (1, \tau e^{i\varphi_{\mathbf{q}}})^{\text{T}} / \sqrt{2}$ describe a pseudospin degree of freedom where the two components encode the relative amplitude and phase of the dipolar oscillations on the two inequivalent A and B sublattices, respectively, where $\varphi_{\mathbf{q}} = \arg(f_{\mathbf{q}}^{\text{AB}})$. The coefficients $\cosh \vartheta_{\mathbf{q}\tau} = (\omega_0 + \omega_{\mathbf{q}\tau}^{\text{mat}}) / 2\sqrt{\omega_0 \omega_{\mathbf{q}\tau}^{\text{mat}}}$ and $\sinh \vartheta_{\mathbf{q}\tau} = (\omega_0 - \omega_{\mathbf{q}\tau}^{\text{mat}}) / 2\sqrt{\omega_0 \omega_{\mathbf{q}\tau}^{\text{mat}}}$ describe the contributions of the resonant and non-resonant terms, respectively. Neglecting non-resonant terms we obtain $H_{\text{mat}} = \sum_{\mathbf{q}} \psi_{\mathbf{q}}^\dagger \mathcal{H}_{\mathbf{q}}^{\text{mat}} \psi_{\mathbf{q}}$ which can be diagonalized as $H_{\text{mat}} = \sum_{\tau=\pm} \sum_{\mathbf{q}} \hbar\omega_{\mathbf{q}\tau}^{\text{mat}} \beta_{\mathbf{q}\tau}^\dagger \beta_{\mathbf{q}\tau}$, where the operators $\beta_{\mathbf{q}\tau}^\dagger = \psi_{\mathbf{q}}^\dagger |\psi_{\mathbf{q}\tau}\rangle$ and the corresponding dispersion $\omega_{\mathbf{q}\tau}^{\text{mat}} = \omega_0 + \Omega \Re(f_{\mathbf{q}}^{\text{AA}}) + \tau\Omega |f_{\mathbf{q}}^{\text{AB}}|$ take a simpler form.

In Supplementary Fig. 1a we plot the quasistatic dispersion within the nearest-neighbor approximation for different cavity heights and compare it to the full quasistatic dispersion in Fig. 1b, which is given by equation (S8). Indeed, due to the honeycomb symmetry, one observes that the type-I CDPs remain pinned at the \mathbf{K} and \mathbf{K}' points. To prove this robustness against long-range Coulomb interactions, one must show that the function $f_{\mathbf{q}}^{\text{AB}}$, given by equation (S3), vanishes at the corners of the Brillouin zone, i.e., one must show that the phase factors associated with the interactions with dipoles residing at the same separation distance and different sublattices sum to zero. To this aim, we define separation vectors $\mathbf{A}_{l_1 l_2} = l_1 \mathbf{a}_1 + l_2 \mathbf{a}_2 + \mathbf{e}_3$ that connect meta-atoms residing on different sublattices, with primitive lattice vectors $\mathbf{a}_1 = a(\sqrt{3}, 0)$ and $\mathbf{a}_2 = a(-\frac{\sqrt{3}}{2}, \frac{3}{2})$, and integers l_1 and l_2 . We can then find six separation vectors with the same magnitude by exploiting the three-fold rotational symmetry $\mathbf{A}_{l_1 l_2}^{j+} = O_j \mathbf{A}_{l_1 l_2}$, where the operator O_j rotates a vector by the angle $2\pi j/3$ ($j = 0, 1, 2$), and the mirror symmetry about the y -axis $\mathbf{A}_{l_1 l_2}^{j-} = M_x \mathbf{A}_{l_1 l_2}^{j+}$, where the operator M_x inverts the sign of the x component. Explicitly, the six separation vectors (not necessarily all different) read

$$\begin{aligned}\mathbf{A}_{l_1 l_2}^{0\pm} &= \pm a \left[\frac{\sqrt{3}}{2}(2l_1 - l_2 - 1) \right] \hat{\mathbf{x}} + a \left[\frac{1}{2}(3l_2 + 1) \right] \hat{\mathbf{y}} \\ \mathbf{A}_{l_1 l_2}^{1\pm} &= \pm a \left[\frac{\sqrt{3}}{2}(-l_1 - l_2) \right] \hat{\mathbf{x}} + a \left[\frac{1}{2}(3l_1 - 3l_2 - 1) \right] \hat{\mathbf{y}} \\ \mathbf{A}_{l_1 l_2}^{2\pm} &= \pm a \left[\frac{\sqrt{3}}{2}(-l_1 + 2l_2 + 1) \right] \hat{\mathbf{x}} + a \left[\frac{1}{2}(-3l_1 + 1) \right] \hat{\mathbf{y}}\end{aligned}\tag{S9}$$

and evaluating equation (S3) at \mathbf{K} one indeed finds $f_{\mathbf{K}}^{\text{AB}} = 0$.

Without a cavity, the $f_{\mathbf{q}}^{\text{AA}}$ term, given by equation (S2), introduces strong asymmetry between the upper and lower quasistatic bands. Since the lower band corresponds to a dark ($\uparrow\downarrow$) configuration of the two sublattices, the dispersion converges rapidly with increasing number of interactions taken into account as long-range Coulomb interactions quickly average to zero. The most prominent change is seen in the upper band where the group velocity increases towards the Γ point, which is in contrast to the dispersion within the nearest-neighbor approximation. This is because the upper quasistatic band corresponds to a bright ($\uparrow\uparrow$) configuration of the two sublattices and therefore long-range Coulomb interactions do not average to zero, especially near the Γ point. However, as the cavity height is reduced, the image dipoles quench long-range Coulomb interactions, which eventually become negligible for cavity heights on the order of the nearest-neighbor separation.

The polariton dispersion plots presented in the main text (Fig. 2c-e) are within the nearest-neighbor approximation. In Supplementary Fig. 2a-d we present the polariton dispersion including all Coulomb interactions beyond nearest-neighbors until convergence, i.e., using equation (S6) instead of equation (28) in the polariton Hamiltonian (see Methods). As expected from the quasistatic dispersion plots in Supplementary Fig. 1b, long-range Coulomb interactions do not qualitatively affect the physics near the \mathbf{K} and \mathbf{K}' points. In fact, for small cavity heights, the nearest-neighbor approximation becomes increasingly accurate and indistinguishable from the full polariton dispersion (see Supplementary Fig. 2d for example).

As described in the main text, six inequivalent type-II SDPs emerge in the polariton spectrum due to the protected crossing with the TEM mode along the $\Gamma - \mathbf{K}(\mathbf{K}')$ directions. Protected crossings with higher-order cavity modes ($m \neq 0$) result in a large number of additional type-II Dirac points being generated within the light-cone for large cavity heights (see Supplementary Fig. 2a). However, as the cavity height is decreased, these additional Dirac points migrate towards the Γ point (opposite direction to the SDPs), where they eventually merge with and annihilate one another. Comparing Supplementary Fig. 2a and Supplementary Fig. 2b, one observes that the polariton dispersion near the CDPs does not noticeably change, and qualitative changes outside the light-cone only occur at small cavity heights where the higher-order cavity modes become increasingly detuned with the dipole resonances.

SUPPLEMENTARY NOTE 2

In the Methods section we performed a unitary Schrieffer-Wolff transformation on the polariton Hamiltonian and obtained a two-band Hamiltonian in the matter sublattice space given by equation (36). In this section, we seek an effective Hamiltonian describing the polaritons near the corners of the Brillouin zone by expanding the two-band Bloch Hamiltonian (37) near the \mathbf{K} point (see Methods). To quadratic order in $\mathbf{k} = \mathbf{q} - \mathbf{K}$, the diagonal matrix elements expand as $W_{\mathbf{q}} \simeq W_{\mathbf{K},\mathbf{k}}^{(0)} + W_{\mathbf{K},\mathbf{k}}^{(1)} + W_{\mathbf{K},\mathbf{k}}^{(2)}$ and the off-diagonal elements as $F_{\mathbf{q}} \simeq F_{\mathbf{K},\mathbf{k}}^{(0)} + F_{\mathbf{K},\mathbf{k}}^{(1)} + F_{\mathbf{K},\mathbf{k}}^{(2)}$, where the superscript index denotes the corresponding order in \mathbf{k} . To determine the real and imaginary components in the expansion, we must sum the phase contributions from the degenerate photons at the \mathbf{K} point. To this aim, we define reciprocal lattice vectors $\mathbf{G}_n = n_1 \mathbf{b}_1 + n_2 \mathbf{b}_2$ with primitive vectors $\mathbf{b}_1 =$

$\frac{2\pi}{3a}(\sqrt{3}, -1)$ and $\mathbf{b}_2 = \frac{4\pi}{3a}(0, 1)$. In analogy with the analysis in Supplementary Note 1, we can find the reciprocal lattice vectors associated with six degenerate photonic modes at K by exploiting the three-fold rotational symmetry of the reciprocal lattice $\mathbf{G}_n^{j+} = \mathbf{K} - O_j(\mathbf{K} - \mathbf{G}_n)$, where the operator O_j rotates a vector by the angle $2\pi j/3$ ($j = 0, 1, 2$), and the mirror symmetry about the $k_y = 0$ line $\mathbf{G}_n^{j-} = M_y \mathbf{G}_n^{j+}$, where the operator M_y inverts the sign of the y -component. Explicitly, the six reciprocal lattice vectors (not necessarily all different) read

$$\begin{aligned}\mathbf{G}_n^{0\mu} &= \mathbf{K} + \left[\frac{2\pi}{3\sqrt{3}a}(3n_1 - 2) \right] \hat{\mathbf{x}} + \mu \left[\frac{2\pi}{3a}(2n_2 - n_1) \right] \hat{\mathbf{y}}, \\ \mathbf{G}_n^{1\mu} &= \mathbf{K} + \left[\frac{2\pi}{3\sqrt{3}a}(1 - 3n_2) \right] \hat{\mathbf{x}} + \mu \left[\frac{2\pi}{3a}(2n_1 - n_2 - 1) \right] \hat{\mathbf{y}}, \\ \mathbf{G}_n^{2\mu} &= \mathbf{K} + \left[\frac{2\pi}{3\sqrt{3}a}(1 + 3n_2 - 3n_1) \right] \hat{\mathbf{x}} + \mu \left[\frac{2\pi}{3a}(1 - n_1 - n_2) \right] \hat{\mathbf{y}},\end{aligned}\tag{S10}$$

with $\mu = \pm 1$. Exploiting these symmetries one can rewrite the summations over photonic indices as

$$\sum_{mn} \Delta_{\mathbf{K}mn}^{(v)} \cdots \rightarrow \frac{1}{6} \sum_{mn} \Delta_{\mathbf{K}mn}^{(v)} \sum_{j=0}^2 \sum_{\mu=\pm} \cdots\tag{S11}$$

where the factor of $1/6$ is introduced to avoid over counting. First, the constant (zeroth order) term in the expansion of $W_{\mathbf{q}}$ is

$$W_{\mathbf{K},\mathbf{k}}^{(0)} = \omega_0 - \Omega\mathcal{S} - \frac{\Omega}{6} \sum_{mn} \Delta_{\mathbf{K}mn}^{(0)} \sum_{j=0}^2 \sum_{\mu=\pm} 1 = \bar{\omega}_0.\tag{S12}$$

The linear (first order) term vanishes as

$$W_{\mathbf{K},\mathbf{k}}^{(1)} = \frac{\Omega a^2}{6} \sum_{mn} \Delta_{\mathbf{K}mn}^{(1)} \sum_{j=0}^2 \sum_{\mu=\pm} [(\mathbf{K} - \mathbf{G}_n^{j\mu})_x k_x + (\mathbf{K} - \mathbf{G}_n^{j\mu})_y k_y] = 0,\tag{S13}$$

which illustrates that the light-matter interaction does not introduce anisotropic warping of the Dirac cone to first order. Finally, the quadratic (second order) term is

$$\begin{aligned}W_{\mathbf{K},\mathbf{k}}^{(2)} &= -\frac{\Omega a^4}{6} \sum_{mn} \Delta_{\mathbf{K}mn}^{(2)} \sum_{j=0}^2 \sum_{\mu=\pm} (\mathbf{K} - \mathbf{G}_n^{j\mu})_x (\mathbf{K} - \mathbf{G}_n^{j\mu})_y k_x k_y \\ &\quad - \frac{\Omega a^4}{12} \sum_{mn} \Delta_{\mathbf{K}mn}^{(2)} \sum_{j=0}^2 \sum_{\mu=\pm} [(\mathbf{K} - \mathbf{G}_n^{j\mu})_x^2 k_x^2 + (\mathbf{K} - \mathbf{G}_n^{j\mu})_y^2 k_y^2] \\ &\quad + \frac{\Omega a^2}{12} \sum_{mn} \Delta_{\mathbf{K}mn}^{(1)} \sum_{j=0}^2 \sum_{\mu=\pm} (k_x^2 + k_y^2) \\ &= -D(k_x^2 + k_y^2),\end{aligned}\tag{S14}$$

The constant (zeroth order) term in the expansion of $F_{\mathbf{q}}$ vanishes at the high-symmetry points as

$$F_{\mathbf{K},\mathbf{k}}^{(0)} = -\frac{\Omega}{6} \sum_{mn} \Delta_{\mathbf{K}mn}^{(0)} \sum_{j=0}^2 \sum_{\mu=\pm} \phi_{nj\mu}^2 = 0,\tag{S15}$$

where $\phi_{nj\mu} = \exp(ia\mathbf{G}_n^{j\mu} \cdot \hat{\mathbf{y}})$. This shows that the CDPs remain pinned at the K and K' points and are completely robust against the coupling to the surrounding photonic environment. The linear (first order) term reads

$$\begin{aligned}F_{\mathbf{K},\mathbf{k}}^{(1)} &= -\frac{3\Omega a}{2}(1 - \mathcal{I})(k_x + ik_y) + \frac{\Omega a^2}{6} \sum_{mn} \Delta_{\mathbf{K}mn}^{(1)} \sum_{j=0}^2 \sum_{\mu=\pm} \phi_{nj\mu}^2 [(\mathbf{K} - \mathbf{G}_n^{j\mu})_x k_x + (\mathbf{K} - \mathbf{G}_n^{j\mu})_y k_y] \\ &= -\bar{v}(k_x + ik_y),\end{aligned}\tag{S16}$$

Finally, the quadratic (second order) term is given by

$$\begin{aligned}
F_{\mathbf{K},\mathbf{k}}^{(2)} &= \frac{3\Omega a^2}{8}(1 - \mathcal{I})(k_x - ik_y)^2 + \frac{\Omega a^2}{12} \sum_{mn} \Delta_{\mathbf{K}mn}^{(1)} \sum_{j=0}^2 \sum_{\mu=\pm} \phi_{nj\mu}^2 (k_x^2 + k_y^2) \\
&\quad - \frac{\Omega a^4}{12} \sum_{mn} \Delta_{\mathbf{K}mn}^{(2)} \sum_{j=0}^2 \sum_{\mu=\pm} \phi_{nj\mu}^2 [(\mathbf{K} - \mathbf{G}_n^{j\mu})_x^2 k_x^2 + (\mathbf{K} - \mathbf{G}_n^{j\mu})_y^2 k_y^2] \\
&\quad - \frac{\Omega a^4}{6} \sum_{mn} \Delta_{\mathbf{K}mn}^{(2)} \sum_{j=0}^2 \sum_{\mu=\pm} \phi_{nj\mu}^2 (\mathbf{K} - \mathbf{G}_n^{j\mu})_x (\mathbf{K} - \mathbf{G}_n^{j\mu})_y k_x k_y \\
&= \bar{t} (k_x - ik_y)^2.
\end{aligned} \tag{S17}$$

We can now identify the effective Hamiltonian $\bar{H}_{\mathbf{K}}^{\text{eff}} = \sum_{\mathbf{k}} \psi_{\mathbf{k}}^\dagger \bar{\mathcal{H}}_{\mathbf{K},\mathbf{k}}^{\text{eff}} \psi_{\mathbf{k}}$ in the matter sublattice space, where the effective Bloch Hamiltonian $\bar{\mathcal{H}}_{\mathbf{K},\mathbf{k}}^{\text{eff}}$ is given by equation (9) in the main text.