



UNIVERSITI PUTRA MALAYSIA

**MOLECULAR AND BIOLOGICAL CHARACTERIZATION OF TWO
VERY VIRULENT INFECTIOUS BURSAL DISEASE VIRUS
ISOLATES, UPM94/273 AND UPM97/61**

KONG LIH LING

FPV 2003 2

**MOLECULAR AND BIOLOGICAL CHARACTERIZATION OF TWO
VERY VIRULENT INFECTIOUS BURSAL DISEASE VIRUS ISOLATES,
UPM94/273 AND UPM97/61**

By

KONG LIH LING

**Thesis Submitted to the School of Graduate Studies,
Universiti Putra Malaysia, in Fulfilment of the Requirement for
the Degree of Master of Science**

February 2003



**Dedicated with love and gratitude
to:**

**My dearest parents, fiance, family
and five lovely nieces**



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Master of Science

MOLECULAR AND BIOLOGICAL CHARACTERIZATION OF TWO VERY VIRULENT INFECTIOUS BURSAL DISEASE VIRUS ISOLATES, UPM94/273 AND UPM97/61

By

KONG LIH LING

February 2003

Chairman: Associate Professor Dr. Abdul Rahman Omar, Ph.D.

Faculty: Veterinary Medicine

An atypical very virulent (vv) strain (UPM94/273) and typical vv strain (UPM97/61) of infectious bursal disease virus (IBDV) isolated in Malaysia, were characterized both *in vivo* and at the molecular level. Comparison of the deduced amino acid sequences with other serotype 1 and 2 sequences revealed 16 amino acid residues, which were conserved only in the vvIBDV. Among the 16 unique amino acid differences, 8 were in VP1 (146 Asp, 147 Asn, 242 Glu, 390 Met, 393 Asp, 562 Pro, 687 Pro and 695 Arg), 3 were in VP2 (222 Ala, 256 Ile and 294 Ile), 2 were in VP3 (990 Val and 1005 Ala) and 3 were in VP4 (685 Asn, 715 Ser and 751 Asp). The importance of these unique amino acid residues is not known but they could affect the virulence of vvIBDV. The UPM94/273 also demonstrated 6 unique amino acid residues at segment



A at positions Ser254, Glu270, Lys588, Ser745, Phe838 and Lys863 and 8 unique amino acid residues at segment B at positions Ala92, Ser100, Val208, Asp253, Asp560, Asn565, Gly750 and Gly876. In addition, these amino acid substitutions have not been reported before in vIBDV and were found only on variant, classical and/or serotype 2 strains. However, the VP5 region of both vIBDV strains was conserved. The UPM97/61 demonstrated 7 unique amino acid substitutions at segment A and 4 unique amino acid substitutions at segment B. However, none of the amino acids changes have been reported elsewhere in other IBDV strains. Although the actual functions of the amino acid substitutions are not known, the unusual amino acid substitutions at segment A and/or B of both isolates may be important in virus virulence. Alignments of the nucleic acid and amino acid sequences of segment A and B followed by distance analysis allowed the generation of phylogenetic trees. Phylogenetic analysis based on segment A and B revealed that all the vIBDV strains including UPM94/273 isolate can be clustered in a group that is distinct from classical, variant, attenuated and serotype 2 strains. However, the tree branching patterns were quite different between segment A and segment B. In addition, the vIBDV strains showed several conserved amino acid substitutions at segment B as found in the Australian 002-73 and serotype 2 strains. These findings indicate that probably a genetic reassortment may have played an important role in the emergence of vIBDV. Flow cytometry and real time PCR assays, indicated that chickens infected with UPM97/61 induced higher



percentages of apoptotic cells but lower level of viral load whereas UPM94/273 induced lower percentages of apoptotic cells but higher level of viral load, suggesting a negative correlation between viral load and apoptosis. These results indicated that UPM97/61 was more virulent than UPM94/273.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Master Sains

**PENCIRIAN MOLEKUL DAN BIOLOGIK ISOLAT-ISOLAT UPM94/273
DAN UPM97/61 VIRUS PENYAKIT BURSA BERJANGKIT**

Oleh

KONG LIH LING

Februari 2003

Pengerusi: Profesor Madya Dr. Abdul Rahman Omar, Ph.D.

Fakulti: Perubatan Veterinar

Virus penyakit bursa berjangkit amat virulen (vIBDV) yang tidak khusus (UPM94/273) dan yang khusus (UPM97/61) diperolehi dari Malaysia, telah dicirikan *dalam vivo* dan berdasarkan molekul. Perbandingan jujukan asid amino dengan jujukan serotip 1 dan 2 yang lain telah menunjukkan 16 asid amino residu yang cuma terdapat di vIBDV. Perbezaan antara 16 asid amino yang unik ini, 8 terdapat di VP1 (146 Asp, 147 Asn, 242 Glu, 390 Met, 393 Asp, 562 Pro, 687 Pro dan 695 Arg), 3 terdapat di VP2 (222 Ala, 256 Ile dan 294 Ile), 2 terdapat di VP3 (990 Val dan 1005 Ala) dan 3 terdapat di VP4 (685 Asn, 715 Ser dan 751 Asp). Kepentingan residu-residu asid amino ini adalah tidak diketahui tetapi ia boleh mempengaruhi virulen vIBDV. UPM94/273 juga menunjukkan 6 asid amino yang unik pada kedudukan Ser254, Glu270,

Lys588, Ser745, Phe838 dan Lys863 mana setiap 2 berada di VP2, VP3 dan VP4. Tambahan pula, penggantian asid amino pada kedudukan Gly254Ser, Ala270Glu, Glu588Lys dan Asp745Ser cuma terdapat di varian, klasik dan/atau serotip 2. Walaubagaimanapun, bahagian VP5 untuk kedua-dua wIBDV adalah kekal. UPM97/61 menunjukkan 7 penggantian asid amino yang unik di segmen A dan 4 penggantian asid amino yang unik di segmen B. Tetapi, perubahan asid amino ini tidak pernah dilaporkan pada strain-strain IBDV yang lain. Walaupun fungsi sebenar penggantian asid amino adalah tidak diketahui, tetapi penggantian asid amino yang luar biasa ini pada segmen A dan/atau B untuk kedua-dua isolat mungkin penting dalam virulen virus. Susunan untuk jujukan asid nukleik dan asid amino untuk segmen A dan B diikuti oleh analisis jarak membenarkan pembentukan pokok filogenetik. Analisis filogenetik berdasarkan segmen A dan B menunjukkan semua strain wIBDV termasuk isolat UPM94/273 IBDV boleh dikumpulkan dalam satu kumpulan di mana adalah berbeza daripada strain-strain klasik, varian, akenuat dan serotip 2. Walaupun begitu, corak-corak untuk cabang pokok adalah berlainan antara segmen A dan segmen B. Tambahan pula, strain-strain wIBDV menunjukkan beberapa penggantian asid amino yang kekal dalam segmen B seperti yang dijumpai dalam strain-strain Australian 002-73 dan serotip 2. Penemuan ini menyatakan bahawa kemungkinan penyusunan semula genetik yang mungkin memainkan peranan yang penting dalam kemunculan wIBDV. Cara-cara aliran sitometri dan PCR masa sebenar, menyatakan bahawa

ayam-ayam yang dijangkiti dengan UPM97/61 mengakibatkan peratusan sel-sel apoptosis yang lebih tinggi tetapi takat muatan virus yang lebih rendah manakala UPM94/273 mengakibatkan peratusan sel-sel apoptosis yang lebih rendah tetapi takat muatan virus yang lebih tinggi, mencadangkan satu perhubungan yang negatif wujud di antara muatan virus dan apoptosis. Keputusan ini menyatakan bahawa UPM97/61 adalah lebih virulen daripada UPM94/273.



ACKNOWLEDGEMENTS

I would like to express my heartiest gratitude and appreciation to Dr. Abdul Rahman Omar, chairman of the supervisory committee for providing invaluable advice and untiring assistance in this master research. His constant guidance, participation in laboratory works and mostly for sharing knowledge on molecular technique that enabled me to accomplish my work. Moreover, I would like to express my sincere thanks and appreciation to Professor Dr. Aini Ideris, Associate Professor Dr. Mohd Hair Bejo, Associate Professor Dr. Seow Heng Fong , members of the supervisory committee for their constructive suggestion, proper guidance and encouragement throughout my study period.

I am grateful to Maha Abdullah for helping and guiding me in handling flow cytometry technique; guidance from Dino and Gabrielle in real time PCR approach, and also Siaw Liung and Tzer Miin for their helpful guidance in SPSS statistical analysis. Special thanks also go to the staffs in Biologics Lab, Puan Rodiah Husin and Mr. Mohd Redha Izwan B. Adnani; and friends Sheau Wei, Siti, Shila, Aini, Balkis, Lee Kim, Su Fun, May Ling, Keng Fei, Zul, Thapa, Do Yew, Sok Fang and every body for always being so willing to render assistance throughout the course of my study.



I certify that an Examination Committee met on 25th February 2003 to conduct the final examination of Kong Lih Ling on her Master of Science thesis entitled "Molecular and Biological Characterization of Two Very Virulent Infectious Bursal Disease Virus Isolates, UPM94/273 and UPM97/61" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulation 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

DATIN KHATIJAH YUSOFF, Ph.D.

Professor,
Department of Biochemistry and Microbiology,
Faculty of Science and Environmental Studies,
Universiti Putra Malaysia,
(Chairman)

ABDUL RAHMAN OMAR, Ph.D.

Associate Professor,
Department of Veterinary Pathology and Microbiology,
Faculty of Veterinary Medicine,
Universiti Putra Malaysia.
(Member)

AINI IDERIS, Ph.D.

Professor,
School of Graduate Studies,
Universiti Putra Malaysia.
(Member)

MOHD HAIR BEJO, Ph.D.

Associate Professor,
Department of Veterinary Pathology and Microbiology,
Faculty of Veterinary Medicine,
Universiti Putra Malaysia.
(Member)

SEOW HENG FONG, Ph.D.

Associate Professor,
Department of Clinical Laboratory Science,
Faculty of Medicine and Health Sciences,
Universiti Putra Malaysia.
(Member)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor/Deputy Dean,
School of Graduate Studies,
Universiti Putra Malaysia.

Date: 2 MAY 2003

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Science. The members of the Supervisory Committee are as follows:

ABDUL RAHMAN OMAR, Ph.D.

Associate Professor,
Department of Veterinary Pathology and Microbiology,
Faculty of Veterinary Medicine,
Universiti Putra Malaysia.
(Chairman)

AINI IDERIS, Ph.D.

Professor,
School of Graduate Studies,
Universiti Putra Malaysia.
(Member)

MOHD HAIR BEJO, Ph.D.

Associate Professor,
Department of Veterinary Pathology and Microbiology,
Faculty of Veterinary Medicine,
Universiti Putra Malaysia.
(Member)

SEOW HENG FONG, Ph.D.

Associate Professor,
Department of Clinical Laboratory Science,
Faculty of Medicine and Health Sciences,
Universiti Putra Malaysia.
(Member)



AINI IDERIS, Ph.D.
Professor/Dean,
School of Graduate Studies,
Universiti Putra Malaysia.

Date: **12** JUN 2003

DECLARATION

I here declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



KONG LIH LING

Date: 2 / 5 / 2003

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL SHEETS	x
DECLARATION FORM	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF ABBREVIATIONS	xx

CHAPTER

I	INTRODUCTION	1
II	LITERATURE REVIEW	8
	Infectious Bursal Disease	8
	Infectious Bursal Disease Virus (IBDV)	9
	Viral Proteins	10
	Antigenic and Pathotypic Variation	12
	Virulence Markers	13
	Pathogenesis and Immunosuppression	15
	Diagnosis	17
	Prevention and Control	20
	Reverse Transcription Polymerase Chain Reaction	23
	Phylogenetic Analysis	24
	Flow Cytometric Detection of Apoptosis	26
	Real-Time Quantitative PCR	28
III	AMPLIFICATION AND CLONING OF INFECTIOUS BURSAL DISEASE VIRUS BIASEGMENTED GENOME	31
	Introduction	31
	Materials and Methods	33
	IBDV Isolation	33
	Propagation of Viruses On Specific Pathogen Free (SPF) Embryonated Chicken Eggs .	33
	Extraction of RNA	34
	Determination of RNA Concentration and Purity	35
	First-strand Complementary DNA (cDNA) and PCR	35
	Determination of PCR Product Concentration and Purity	38
	Agarose Gel Electrophoresis	38



	Purification of PCR Products	39
	Cloning of the Full Length of VP1, VP3, VP4 and VP5 Genes TOPO® Cloning Reaction and One Shot® Chemical Transformation	40
	Subculture and Analysis of Positive Colonies	41
	Preparation of Stock Culture	42
	Plasmid Extraction and Purification.....	42
	Restriction Enzyme Analysis of the Plasmid	44
	Results	45
	Amplification of IBDV Genome	45
	Analysis of Recombinant Plasmid	46
	Restriction Enzyme Digestion Analysis	46
	Discussion	61
IV	SEQUENCE AND PHYLOGENETIC ANALYSIS OF IBDV ...	63
	Introduction	63
	Materials and Methods	65
	Cloning and Sequencing	65
	Sequence Assembly and Analysis	65
	Construction of Phylogenetic Tree	66
	Results	69
	Sequence Analysis of Segment A	69
	VP5 Protein	69
	Precursor Polyprotein (NH ₂ -VP2-VP4-VP3-COOH)	71
	Sequence Analysis of Segment B	79
	Phylogenetic Analysis	90
	Percentage of Homology	93
	Discussion	95
V	FLOW CYTOMETRIC AND REAL TIME QUANTITATIVE PCR ANALYSIS OF IBDV	102
	Introduction	102
	Materials and Methods	104
	Viruses	104
	SPF Embryonated Chicken Eggs	104
	Chorioallantoic Membrane (CAM) Preparation	104
	IBDV Inoculation	105
	Titration of IBDV	105
	SPF Chickens	105
	Experimental Design	105
	Single Cell Suspension Preparation	106
	Viability (Trypan Blue) Test	107
	Propidium Iodide Staining	107
	RNA Extraction	108
	Primer Designing	108
	Conventional RT-PCR Amplification	109



SYBR Green 1 Optimization	109
Real-Time Quantitative PCR	109
Statistical Analysis	111
Results	111
Flow Cytometric Analysis	111
Propidium Iodide Analysis	111
PCR Amplification	118
Real-Time Quantitative PCR Analysis	118
SYBR Green I Optimization	118
Standard Curve	118
Comparison of Viral RNA Levels	120
Discussion	126
VI GENERAL DISCUSSION AND CONCLUSION	131
BIBLIOGRAPHY	138
APPENDICES	162
A. Percent Embryo Lethal Dose	162
B. Nucleotide and Amino Acid Sequence Alignment.....	163
C. Phylogenetic Tree.....	208
D. Genetic Distances.....	210
E. Statistical Analysis for Flow Cytometry and Real Time Quantitative PCR	212
VITA	230
PUBLICATION	231



LIST OF TABLES

Table		Page
3.1	Primers used to amplify VP3, VP4, VP5 genes and segment B	36
4.1	Characteristics of IBDV strains used in multiple alignment analysis of segment A sequence.....	67
4.2	Characteristics of IBDV strains used in multiple alignment analysis of segment B sequence.....	68
4.3	Comparison of Amino Acid Substitution at Different Position Between UPM94/273 and Other Published Strains in Segment A of IBDV.....	78
4.4	Comparison of Amino Acid Substitution at Different Position Between Malaysian Strains and Other Published Strains in Segment B of IBDV.....	89
5.1	Percentage of apoptotic cells determined by flow cytometry in IBDV infected and uninfected bursa lymphoid cells.....	117
5.2	Quantitation of viral RNA in bursa lymphoid cells from chickens infected with vaccine strain D78, vIBDV strains UPM94/273 and UPM97/61 based on cycle threshold values.....	121
5.3	Mortality of SPF chickens following infection with different IBDV strains.....	123

LIST OF FIGURES

Figure		Page
3.1	RT-PCR Product of the Expected UPM94/273 IBDV VP3 Gene Using Specific Primer.....	47
3.2	RT-PCR Product of the Expected UPM94/273 IBDV VP4 Gene Using Specific Primer.....	48
3.3	RT- PCR Product of the Expected UPM94/273 IBDV VP5 Gene Using Specific Primer.....	49
3.4	RT- PCR Product of the Expected 3 Fragments for VP1 Gene of UPM94/273 and UPM97/61 IBDV Using Specific Primers.....	50
3.5	PCR Screening for Recombinant VP3 Gene of UPM94/273 IBDV Isolate.....	51
3.6	PCR Screening for Recombinant VP4 Gene of UPM94/273 IBDV Isolate.....	52
3.7	PCR Screening for Recombinant VP5 Gene of UPM94/273 IBDV Isolate.....	53
3.8	PCR Screening for Recombinant Fragment 1 of VP1 Gene of UPM94/273 and UPM97/61 IBDV Isolates.....	54
3.9	PCR Screening for Recombinant Fragment 2 of VP1 Gene of UPM94/273 and UPM97/61 IBDV Isolates.....	55
3.10	PCR Screening for Recombinant Fragment 3 of VP1 Gene of UPM94/273 and UPM97/61 IBDV Isolates.....	56
3.11	Purified Plasmids Digested With <i>EcoR1</i> Enzyme for VP3 Gene of UPM94/273 IBDV Isolate.....	57
3.12	Purified Plasmids Digested With <i>EcoR1</i> Enzyme for VP4 Gene of UPM94/273 IBDV Isolate.....	58
3.13	Purified Plasmids Digested With <i>EcoR1</i> Enzyme for VP5 Gene of UPM94/273 IBDV Isolate.....	59
3.14	Purified Plasmids Digested With <i>EcoR1</i> Enzyme for 3 Fragments of VP1 Gene of UPM94/273 and UPM97/61 IBDV Isolates.....	60



4.1	Nucleotide Sequence and Translation of Amino Acid of VP5 Gene of UPM94/273 IBDV Isolate.....	70
4.2	Nucleotide Sequence and Translation of Amino Acid of Precursor Polyprotein, VP2-VP4-VP3 Genes of UPM94/273 IBDV Isolate.....	72
4.3	Nucleotide Sequence and Translation of Amino Acid of Segment B, VP1 Gene of UPM94/273 IBDV Isolate.....	80
4.4	Nucleotide Sequence and Translation of Amino Acid of Segment B, VP1 Gene of UPM97/61 IBDV Isolate.....	84
4.5	Phylogenetic Tree Construction Based on Amino Acids Sequence of the Segment A IBDV Isolates.....	91
4.6	Phylogenetic Tree Construction Based on Amino Acids Sequence of the Segment B IBDV Isolates.....	92
5.1	Detection of apoptosis in bursal lymphocytes by flow cytometry for propidium iodide staining. Bursal lymphocytes were isolated from uninfected control chickens at (A) day 1, (B) day 2, (C) day 3, (D) day 7 and (E) day 14.....	112
5.2	Detection of apoptosis in bursal lymphocytes by flow cytometry for propidium iodide staining. Bursal lymphocytes were isolated from vaccine strain D78 infected chickens at (A) day 1, (B) day 2, (C) day 3, (D) day 7 and (E) day 14.....	113
5.3	Detection of apoptosis in bursal lymphocytes by flow cytometry for propidium iodide staining. Bursal lymphocytes were isolated from atypical vvIBDV strain UPM94/273 infected chickens at (A) day 1, (B) day 2, (C) day 3, (D) day 7 and (E) day 14.....	114
5.4	Detection of apoptosis in bursal lymphocytes by flow cytometry for propidium iodide staining. Bursal lymphocytes were isolated from typical vvIBDV strain UPM97/61 infected chickens at (A) day 1, (B) day 2, (C) day 3, (D) day 7 and (E) day 14.....	115
5.5	RT-PCR products of the 120 bp VP2 gene.....	119
5.6	Standard curve of serial 10-fold dilutions of quantitative standard from 10^{-1} to 10^{-5} mg/ml UPM94/273 IBDV stock virus.....	122



5.7	A linear relationship was observed between the amount of input RNA and the CT values for the IBDV infected lymphoid cells, labeled as unknown samples.....	122
5.8	Quantitation of viral RNA in bursa lymphoid cells from chickens infected with vaccine strain D78, vvIBDV strains UPM94/273 and UPM97/61 based on log concentration...	124
5.9	(A) Fluorescence vs. temperature plot of data collected during the melt curve run protocol between 54°C-95°C following amplification of vvIBDV strains with SYBR Green I. (B) Melt curve data collected after vvIBDV strains amplification with designated primers produced a product peak at 88°C.....	125



LIST OF ABBREVIATIONS

AGPT	Agar gel precipitin test
BLAST	Basic local alignment search tool
bp	Basepair
BSA	Bovine serum albumin
CAM	Chorioallantoic membrane
cDNA	Complementary deoxyribonucleic acid
°C	Degree Celcius
CE	Chicken embryo
CMX-Ros	Chloromethyl-X-rosamine
CT	Threshold cycle
DEPC	Diethyl pyrocarbonate
DH ₂ O	Distilled water
DMSO	Dimethylsulphoxide
DNA	Deoxyribonucleic acid
dNTP	Deoxynucleoside triphosphate
ds	Double strand
DTT	Dithiothreitol
dUTP-FITC	dUTP flurorescein isothiocyanate
<i>E. coli</i>	<i>Escherichia coli</i>
EDTA	Ethylene diamine tetra acetic acid
EID ₅₀	Embryo infective dose fifty
ELISA	Enzyme linked immunosorbent assay
HVT	Herpes virus of turkey
IBD	Infectious bursal disease
IBDV	Infectious bursal disease virus
IFN	Interferon
IPNV	Infectious pancreatic necrosis virus
kb	Kilobase
kDa	Kilodalton
LB	Luria-Bertani
M	Molar
MC-540	Merocyanine-540
MDA	Maternally derived antibody
MgSO ₄	Magnesium sulfat
ml	Millilitre
mM	Millimolar
µg	Microgram
NaCl	Sodium Chloride
Ial	Sodium iodide
NaOH	Sodium hydroxide
NCBI	National centre for biotechnology information
ng	Nanogram
NJ	Neighbour-joining
OD	Optical density
ORF	Open reading frame



PBS	Phosphate buffered saline
PCR	Polymerase chain reaction
pi	Post infection
PI	Propidium iodide
pmol	Picamol
PTC	Peltier thermal cyler
QC-PCR	Quantitative competitive PCR
QGDPT	Quantitative gel diffusion precipitin test
RdRp	RNA dependent RNA polymerase
RFLP	Restriction fragment length polymorphism
RNA	Ribonucleic acid
RT-PCR	Reverse-transcriptase PCR
RT	Room temperature
SDS	Sodium dodecyl sulfate
SPF	Specific pathogen free
SPSS	Statistical program for social science
TAE	Tris-acetate-EDTA
TCVN	Tissue culture virus neutralization
Tris	2-amino-2-(hydroxymethyl)-1, 3 propanediol
TUNEL	Terminal deoxynucleotidyl transferase mediated nick and labeling
UPGMA	Unweighted pair group method with arithmetic mean
UPM	Universiti Putra Malaysia
UV	Ultraviolet
VNF	Virus neutralizing factor
VP	Viral protein
vv	Very virulent
(w/v)	Weight/volume
X-gal	5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside

Amino Acid	Single/Three Letter Amino Acid Code
Alanine	A Ala
Arginine	R Arg
Asparagine	N Asn
Aspartic Acid	D Asp
Glutamine	Q Gln
Glutamic Acid	E Glu
Glycine	G Gly
Isoleucine	I Ile
Leucine	L Leu
Lycine	K Lys
Methionine	M Met
Phenylalanine	F Phe
Proline	P Pro
Serine	S Ser
Threonine	T Thr
Tryptophan	W Trp
Valine	V Val

CHAPTER I

INTRODUCTION

Infectious bursal disease (IBD) is an acute contagious viral disease of young chickens (Kibenge *et al.*, 1988; Lasher *et al.*, 1994). The etiological agent, IBD virus (IBDV), has a predilection for the cells of the bursa of Fabricius where the virus infects actively dividing and differentiating lymphocytes of the B-cell lineage (Burkhardt *et al.*, 1987). Thus, IBD is a fatal immunosuppressive disease causing heavy losses to the poultry industry (Etteradossi *et al.*, 1998).

The first outbreak of IBD was reported in commercial chicken flocks in Delaware, USA (Cosgrove, 1962). The IBDV strains, which were isolated during this outbreak, are now referred to as classical serotype I isolates. Later on, a second serotype – serotype II of IBDV was identified (McNulty and Saif, 1988). These isolates are apathogenic and are recovered mainly from turkeys (Ismail *et al.*, 1988). Based on antigenic variation and virulence, serotype I isolates can be divided into several groups : classical virulent, attenuated, antigenic variant and very virulent (vv) strains (Cao *et al.*, 1998). Since 1985, antigenic variants of serotype I IBDV isolates had been recovered from flocks with selection pressure of field vaccination against classical IBDV serotype I (Snyder, 1990). Although being antigenic variant, these isolates have only minor amino



acid changes and do not form any separate serotype. Nevertheless, these changes occur at the VP2 conformation-dependent antigenic epitopes that are responsible for stimulating virus neutralizing antibodies (Bayliss *et al.*, 1990). In 1991, IBDV isolates, which were able to break through levels of maternal antibodies that were normally protective, were reported in Europe (Chettle *et al.*, 1989). These isolates, the so called very virulent IBDV (vvIBDV), cause more severe clinical signs during an outbreak with mortality approaching 100% in susceptible flocks, and are now found almost world-wide (VandenBerg, 2000).

The emergence of highly virulent strains of IBDV has complicated the immunization programs against the disease. Early vaccination may result in failure due to interference with the maternal antibody, whilst its delay may cause field virus infections. Therefore, it is important to characterize the antigenicity and the virulence of IBDV in both vaccine and field strains in the control of the disease. The effectiveness in the latter is also dependent on the diagnostic methods used. The disease can be diagnosed based on virus isolation, electron microscopy, immunofluorescence, virus neutralization, monoclonal antibody assays, and/or enzyme-linked immunosorbent assay (Lukert and Saif, 1991; Wu *et al.*, 1992; Liu *et al.*, 1994). However, these methods have one or more disadvantages such as being time consuming, labour intensive, expensive and of low sensitivity (Wu *et al.*, 1992).

Recently, more sensitive and specific molecular methods have been used to diagnose and characterize IBDV infections (Jackwood and Nielsen, 1997; Moody *et al.*, 2000; Boot *et al.*, 2001). The reverse transcriptase polymerase chain (RT-PCR) has been widely used to detect IBDV (Tham *et al.*, 1995; Jackwood and Nielsen, 1997). RT-PCR followed by restriction fragment length polymorphism (RFLP) has also been used to detect and differentiate IBDV strains (Jackwood and Sommer, 1997; Hoque *et al.*, 2001). RT-PCR RFLP profiles of the amplified hypervariable region of the VP2 gene have been used to diagnose and identify molecular differences in the IBDV strains isolated in different parts of the world (Jackwood and Sommer, 1999; Hoque *et al.*, 2001). In these studies, it was found that all vvIBDV isolates have a conserved *Ssp1* and *Taq1* sites at the hypervariable region of the VP2 gene. A study has also been carried out to develop a PCR method for the detection of IBDV based on colorimetric technique (Phong, 2002).

Generally the severity of IBDV infections has been assessed in terms of mortality or the degree of bursal damage, and it has been difficult to assess viral load because virulent strains of IBDV do not replicate in tissue culture (Moody *et al.*, 2000). A quantitative competitive PCR (QC-PCR) assay has been developed to monitor IBDV RNA extracted from infected bursae (Wu *et al.*, 1997). However, this protocol is labour intensive and the technique has limitations (Souaze *et al.*, 1996). Difficulties in using the RT-PCR technique for quantification

