



UNIVERSITI PUTRA MALAYSIA

**CLINICOPATHOLOGICAL CHANGES IN ADJUVANT INDUCED
ARTHRITIS IN CANINE STIFLE JOINT**

SITI NORHAYATI BT. ISMAIL

FPV 2001 4

**CLINICOPATHOLOGICAL CHANGES IN ADJUVANT INDUCED ARTHRITIS
IN CANINE STIFLE JOINT**

SITI NORHAYATI BT. ISMAIL

**MASTER OF VETERINARY SCIENCE
UNIVERSITI PUTRA MALAYSIA**

2001



**CLINICOPATHOLOGICAL CHANGES IN ADJUVANT INDUCED ARTHRITIS
IN CANINE STIFLE JOINT**

BY

SITI NORHAYATI BT. ISMAIL

**Thesis Submitted in Fulfilment of the Requirement for the Degree of
Master of Veterinary Science in the Faculty of Veterinary Medicine
Universiti Putra Malaysia**

August 2001



Dedicated to,

Mummy and Abah,

For your loves, support and trust,

Along, Cik Yong, Isma and Cici

Thanks for everything...

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in
fulfilment of the requirement for the degree of Master of Veterinary Science

CLINICOPATHOLOGICAL CHANGES IN ADJUVANT INDUCED ARTHRITIS IN CANINE STIFLE JOINT

By

SITI NORHAYATI BT. ISMAIL

August 2001

Chairman : Rashid Ibrahim, Ph.D.

Faculty : Veterinary Medicine

Osteoarthritis is the most common joint disease and cause of physical disability in man and animals. It is a complex disease with unknown etiology. Intra-articular injection of 1 ml. Freund's adjuvant was inoculated into twenty-five adult Mongrel dogs weighing between 10-15kg. Osteoarthritis was induced in the left stifle joint, while the right joint act as a control. The dogs were evaluated for clinical evidence of joint heat, effusion and pain, and gait abnormalities. Radiographs were obtained for soft tissue swelling, osteophytosis and degenerative changes. At the end of each trial period (week 1, 2, 3, 4, 5) the dogs were euthanised. The left stifle joint were opened, examined grossly and the articular cartilage and synovial membrane were harvested and fixed for histopathologic and electron microscopic studies.



Clinical signs of joint swelling and pain upon palpation, weight bearing lameness and reduced range of motion were observed within week 1 post-inoculation. These signs were positively correlated with the acute pathological lesions in the synovial membrane and articular cartilage; and in radiograph evaluation. Radiograph study revealed evidences of soft tissue swelling, increased intra-articular space, osteophytes formation; the clinical signs that were suggestive of degenerative changes in osteoarthritis. However, plain radiograph was found to be not informative enough in the early stage of osteoarthritis. Gross changes during post mortem revealed, swelling of the adjacent soft tissue, hypertrophy of the joint capsule and synovial membrane, and joint effusion. These were signs of inflammation of the joint tissues and it was believed that the inflammatory process was one of the major factors in the development of degenerative joint disease. Lameness evaluation was positively correlated with gross examination but negatively correlated with radiograph examination.

In histopathology study, there were signs of inflammation in the synovial membrane and formation of synovial pannus, which was thought to be related with the development of degenerative changes on the articular cartilage. Hyperplasia of intima and subintima layer, edema and congestion; flaking and erosion in articular cartilage, were observed as early as week 1 p.i. Under scanning electron microscopy, cartilage fibrillation and erosion, were observed as early as week 1 p.i. Significant positive correlation between the histological changes in articular cartilage with changes in the synovial membrane suggested that changes in the synovium preceded changes in the articular

cartilage. The synovial membrane was highly vascularised, causes respond to injury more promptly. In articular cartilage, it took time to heal since healing depending on the depth of lesion.

In this study, the pathogenesis of osteoarthritis was divided into three stages: the onset phase, which was observed within one week post-induction; the second phase or the intermediate stage and the end phase. Each structure, cells and tissues was found to have their own role in the production of osteoarthritis. Study on the pathogenesis must emphasize on these structures and cascade of events that occur during the production of osteoarthritis, which will aid in the treatment of osteoarthritis. The clinical, morphological and microstructure changes that occurred in osteoarthritis had been characterised, but the role of each in the aetio-pathogenesis of osteoarthritis is still not rigidly defined.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk Ijazah Master Sains Veterinar

**PERUBAHAN PATOLOGI KLINIKAL DI DALAM INDUKSI ARTRITIS
ADJUVAN PADA LUTUT ANJING**

Oleh

SITI NORHAYATI BT. ISMAIL

Ogos 2001

Pengerusi : Rashid Ibrahim, PhD.

Fakulti : Perubatan Veterinar

Osteoarthritis adalah penyakit yang selalu menyerang sendi dan menyebabkan ketidakstabilan fizikal dalam manusia dan haiwan. Ia adalah penyakit yang kompleks dan puncanya tidak diketahui. Suntikan intra-rawan sebanyak 1 ml. adjuvan Freund's telah diberikan kepada dua puluh lima ekor anjing Mongrel dewasa yang mempunyai berat 10-15 kg. Osteoarthritis telah dihasilkan di dalam lutut kiri anjing sementara lutut kanan bertindak sebagai kawalan. Anjing tersebut telah dimantau secara klinikal untuk mengesan kehangatan sendi, efusi dan kesakitan, dan ketidaknormalan pergerakan. Radiograf telah diambil untuk mengesan bengkakan tisu, osteopitosis dan perubahan degenerasi. Pada akhir setiap eksperimen (minggu 1, 2, 3, 4 dan 5), anjing-anjing tersebut dimatikan. Lutut kiri anjing-anjing tersebut telah diperiksa secara mata kasar, sampel sendi tulang rawan dan membran sinovial

telah diambil dan diawetkan untuk pemeriksaan histopatologi dan mikroskop elektron.

Tanda-tanda klinikal seperti pembengkakan lutut, kesakitan, ketempangan dan kesukaran pergerakan telah diperhatikan dalam minggu pertama selepas induksi. Tanda-tanda tersebut berhubung kait secara positif dengan lesi patologi akut di dalam membran sinovial dan tulang rawan; dan ujian radiograf. Pemeriksaan radiograf pula menunjukkan tanda-tanda kebengkakan tisu, tekanan tinggi antara rawan sendi, pembentukan osteopit; adalah tanda-tanda klinikal yang menyokong penghasilan degenerasi dalam osteoarthritis. Namun begitu, gambar radiograf didapati tidak begitu meyakinkan dalam pembentukan awal osteoarthritis. Pemerhatian mata kasar semasa bedah siasat menunjukkan pembengkakan tisu keliling sendi, ketebalan kapsul sendi dan membran sinovial. Tanda-tanda tersebut adalah tanda-tanda inflamasi pada tisu sendi lutut dan dipercayai bahawa proses inflamasi adalah salah satu proses penting dalam pembentukan penyakit degenerasi sendi. Ujian ketempangan berhubung kait secara positif dengan ujian mata kasar, tetapi berhubungkait secara negatif dengan ujian radiograf.

Pemeriksaan histopatologi menunjukkan tanda-tanda inflamasi di dalam membran sinovial dan pembentukan sinovial panus yang dipercayai mempunyai kaitan dalam pembentukan degenerasi di dalam sendi rawan. Hiperplasia di dalam lapisan sinovial intima dan subintima, edema dan kongesi; kelupasan dan hakisan rawan, kelihatan seawal minggu pertama selepas induksi. Di dalam imbasan mikroskop electron, pada permukaan rawan,

fibrilasi dan kehausan rawan dapat diperhatikan seawal minggu pertama selepas induksi. Hubungkait positif di antara perubahan lesi histologi dan rawan sendi yang didapati di dalam membran sinovial, meyakinkan bahawa pembentukan lesi membran synovial akan menyebabkan pembentukan lesi pada rawan sendi. Membran synovial dipenuhi pembuluh, dan tindakbalas terhadap kecederaan adalah lebih cepat. Bagi rawan sendi pula, ia mengambil masa untuk sembuh dan bergantung kepada kedalaman lesi.

Dalam kajian ini, patogenesis osteoarthritis dapat dibahagikan kepada tiga peringkat; peringkat permulaan, di mana perubahan diperhatikan dalam masa seminggu selepas induksi, peringkat kedua ataupun pertengahan dan peringkat akhir. Setiap struktur, sel dan tisu didapati mempunyai fungsi masing-masing dalam pembentukan osteoarthritis. Kajian mengenai patogenesis mestilah mengambil kira pada setiap struktur dan aliran proses yang berlaku semasa pembentukan osteoarthritis, yang dapat menolong dalam mengubati penyakit osteoarthritis. Perubahan klinikal, morfologi dan mikrostruktur yang berlaku semasa osteoarthritis telah dikaji, namun fungsi setiap satu dalam etio-patogenesis osteoarthritis masih belum dapat diterangkan dengan pasti.

ACKNOWLEDGEMENTS

All praise to Almighty Allah, the Merciful and the Benevolent. Had it not been due to His will and favour, the completion of this study would not have been possible.

I would like to express my deepest and sincere gratitude and appreciation to my supervisor, Dr. Rashid Ibrahim, for his invaluable guidance, advise, supervision, and support throughout the course of this study.

I wish to express my sincere gratitude to Associate Prof. Dr. Mohd Zamri Saad and Dr. Md. Zuki B. Abu Bakar, who are my co-supervisor for their valuable advice, support and continuous encouragement towards the completion of this work. Special thanks and sincere appreciation to Dr. Ungku Chulan Ungku Mohsin for his resourceful comments and suggestions during completing this thesis.

My sincere thanks to Dr. Noor Bakry Hj. Mat Darus, managing director of Vet-Fine (M) Sdn. Bhd.; the head department and supporting staff from the Department of health, Dewan Bandaraya Kuala Lumpur, for providing the experimental animals. I would like to thank the Institute Bioscience staff, Ms. Azilah Ab. Jalil, Mr. Ho Oi Kuan, Ms. Suleka and Ms. Faridah Akmal. Not forgetting, our University Veterinary Hospital supporting staff; Pn. Noraini Othman (surgery), Mr. Osman Asnawi (Radiology), Mr. Palaniandy Malyandy and Mr Arumugam Iyasamy; from histology laboratory, Mr. Jamil Samad; from



post mortem laboratory, Mr. Ghazali Md. Yusoff, Mr. Noraziman Sulaiman and Mr. Apparau Somanaidu; from clinical pathology laboratory, Mr. Mohd Halmi Othman and Mr. Abdullah Misron; and lastly Mr. Zaid Othman and Mr. Chamadre Vengadasamy for their help throughout the course of this study.

I have also been very fortunate in receiving assistance from a number of my colleagues and friends. Many of them went out of their way to assist me and it would not be possible to name all of them. However I wish to express my sincere gratitude to Dr. Loqman Mohamad Yusof (veterinary surgeon in university veterinary hospital and a lecturer), my colleagues; Dr. Zurina Ramli, Dr. Anun Man, Dr. Zamirah Zainal Abidin, Dr. Marina Hassan, Dr. Yuslan Sanuddin, Dr. Ronald Francis and Dr. Wan Mastura Shaik Mossadeq for their help and support in materializing my research.

To one and all whom I may have inadvertently left out, thank you.

Last but not least, to my mother, father and my family, thank you for their support and trust.



I certify that an Examination Committee met on 13th August 2001 to conduct the final examination of Siti Norhayati Bt. Ismail on her Master of Veterinary Science thesis entitled "Clinicopathological Changes in Adjuvant Induced Arthritis in Canine Stifle Joint" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

SHANTHI GANABADI, Ph.D.
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Chairman)

RASHID IBRAHIM, Ph.D.
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

MOHD ZAMRI SAAD, Ph.D.
Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

MD. ZUKI ABU BAKAR, Ph.D.
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)



MOHD GHAZALI MOHAYIDIN, Ph.D.
Professor/Deputy Dean of Graduate School,
Universiti Putra Malaysia

Date: 25 SEP 2001

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Master of Veterinary Science.



AINI IDERIS, Ph.D.
Professor/Dean of Graduate School,
Universiti Putra Malaysia

Date: 08 NOV 2001

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at UPM or other institutions.



SITI NORHAYATI BT. ISMAIL

Date: 19-9-2001

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL SHEETS	xii
DECLARATION FORM	xiii
LIST OF TABLES	xvi
LIST OF FIGURES	xvii
LIST OF PLATES	xviii
LIST OF ABBREVIATIONS	xxi

CHAPTER

1.0	INTRODUCTION	1
	1.1 General Introduction	1
	1.2 Osteoarthritis Study	2
	1.3 Study on Joint Diseases in Animals and Humans	3
	1.4 Objectives	4
2.0	LITERATURE REVIEW	5
	2.1 Structures and Functions of Stifle Joint	5
	2.1.1 Anatomy and Physiology of the Stifle Joint	5
	2.1.2 Articular Cartilage	7
	2.1.3 Joint Capsule and Synovial Membrane	11
	2.2 Animal Models in Arthritis Research	14
	2.2.1 Rat Air Pouch	16
	2.2.2 Arthritis Induction	18
	2.2.2.1 Physical Induction	18
	2.2.2.2 Surgical Induction	19
	2.2.2.3 Articular Irritants	20
	2.3 Morphological Study	26
	2.3.1 Gross Examination	26
	2.3.2 Scanning Electron Microscopy Study	27
	2.4 Clinical Study	32
	2.4.1 Lameness Evaluation	32
	2.4.2 Radiographic Evaluation	36
	2.5 Microstructure Study	38
	2.5.1 Histopathological Study	38



3.0	GENERAL MATERIALS AND METHODS	44
3.1	Experimental Animals	44
3.2	Experimental Design	44
3.3	Induction of Arthritis	45
4.0	CLINICAL ASSESSMENT AND GROSS CHANGES ON THE STIFLE JOINT FOLLOWING EXPERIMENTAL OSTEOARTHRITIS	48
4.1	Introduction	48
4.2	Materials and Methods	51
4.2.1	Clinical Assessment	51
4.2.2	Lameness Evaluation	51
4.2.3	Radiographic Assessment	52
4.2.4	Gross Necropsy	53
4.2.5	Statistical Analysis	54
4.3	Results	55
4.3.1	Lameness Evaluation	55
4.3.2	Radiographic Evaluation	56
4.3.3	Gross Examination	56
4.3.4	Correlation Between Clinical Evaluation and Gross Changes	57
4.4	Discussion	72
4.4.1	Summary	78
5.0	MICROSCOPIC STUDY ON THE ARTICULAR CARTILAGE AND SYNOVIAL MEMBRANE OF THE STIFLE JOINT FOLLOWING AN EXPERIMENTAL OSTEOARTHRITIS	80
5.1	Introduction	80
5.2	Materials and Methods	83
5.2.1	Histopathology	83
5.2.2	Scanning Electron Microscopy	85
5.2.3	Statistical Analysis	87
5.3	Results	87
5.3.1	Histopathology	87
5.3.2	Scanning Electron Microscopy Study	88
5.3.3	Correlation Between Scanning Electron Microscopy Study and Histopathological Study	89
5.4	Discussion	100
5.4.1	Summary	105
6.0	GENERAL DISCUSSION	106
6.1	Conclusion	115
	BIBLIOGRAPHY	117
	APPENDICES	140
	VITA	145



LIST OF TABLES

Table		Page
4.1	Lameness scoring system	52
4.2	Radiographic evaluation scoring	53
4.3	Gross lesion scoring for articular cartilage, synovium and fibrous capsule	54
4.4	Results for clinical evaluation and gross changes	58
5.1	Lesion score for microscopic changes in synovial membrane	84
5.2	Lesion score of microscopic changes on articular cartilage	85
5.3	Lesion score on the articular surface under SEM	86
5.4	Results of lesion score in the articular cartilage and synovial membrane	90



LIST OF FIGURES

Figure		Page
2.1	Schematic drawing of articular cartilage histology showing its layers and fibril arrangement. (A) Surface membrane. (B) Tangential zones. (C) Intermediate zones. (D) Radial zone. (E) Calcified zone. (F) Subchondral zone. (G) Tidemark. (H) Fibrils or 'wickets'. Source: Piermattei and Flo, 1997	8
2.2	A schematic drawing of a longitudinal section of a knee joint of an adult man. The plane of the section that is illustrated in the main drawing is indicated in the inset (upper right). Source: Ham, 1974	14
2.3	A schematic drawing of a cross section of normal articular cartilage. Source: Johnston, 1997	41
3.1	Summary of materials and methods	47
4.1	Graph of mean clinical score for lameness evaluation	59
4.2	Graph for mean clinical score for lameness and radiographic evaluation and mean lesion score for gross changes	59
5.1	The mean lesion scoring following microscopic study	91

LIST OF PLATES

Plate		Page
4.1	Photograph of the stifle joint; (a) Right stifle joint; normal and non-induced (b) Left stifle joint. Day 1 p.i. the stifle joint is swollen, and presence of haematoma (arrow) at the site of injection.	60
4.2	Photograph of the stifle joint; (a) Right stifle joint; normal and non-induced. (b) Left stifle joint. Day 3 p.i., the stifle joint is swollen (arrow), heat and pain were detected upon palpation.	61
4.3	Photograph of the left stifle joint. Weight bearing lameness was evidence on day 1 p.i.	62
4.4	Radiograph of the stifle joint. (a) Right stifle joint; normal and non-induced. (b) Left stifle joint. At week 1 p.i., there are evidence of osteophyte formation (big arrow head) appear as bony outgrowth and soft tissue swelling (small arrow head). Lateral view.	63
4.5	Radiograph of the stifle joint. (a) Right stifle joint; normal and non induced. (b) Left stifle joint. Increased intra-articular space (black arrow) and swelling of the fibrous capsule (white arrow) were evidenced in week 2 p.i. Anterio-Posterior view	64
4.6	Radiograph of the stifle joint. (a) Right stifle joint; normal and non-induced. (b) Left stifle joint. Increased intra-articular space (black arrow) and slight soft tissue swelling (white arrow) were evidence in week 3 p.i. Anterio-Posterior view	65
4.7	Radiograph of the stifle joint. (a) Right stifle joint; normal and non-induced. (b) Left stifle joint. At week 5 p.i., there is a small amount of periosteal new bone formation (osteophyte formation), which appears as slight radio-dense area at the medial and lateral tibia (small arrow), an increased intra-articular space and osteolysis (arrow head). Anterio-Posterior view.	66
4.8	Radiograph of the stifle joint. (a) Right stifle joint; normal and non-induced (b) Left stifle joint. At week 5 p.i., there is evidence of osteophyte formation (arrow) appear as lips of new bone around the edge of the joint. Lateral view	67

4.9	Photograph of the stifle joint. (a) Right stifle joint; normal and non-induced. (b) Left stifle joint. Thickening of the articular cartilage especially at the troclear groove (TG) and troclear ridge (TR) and hypertrophy of the synovial membrane (black arrow) and joint capsule (white arrow) were evidence at week 1 p.i.	68
4.10	Photograph of the stifle joint. (a) Right stifle joint; normal and non-induced. (b) Left stifle joint. Thickening of articular cartilage especially at the trochlear groove (TG) were evidence at week 3 p.i. Hypertrophy of the synovial membrane (black arrow) and joint capsule (white arrow) were also observed.	69
4.11	Photograph of the stifle joint. (a) Right stifle joint; normal and non-induced. (b) Left stifle joint at week 4 p.i. showing normal articular cartilage compared with (a), however the synovial membrane (black arrow) and fibrous capsule (white arrow) is still swollen .	70
4.12	Photograph of the stifle joint. (a) Right stifle joint; normal and non-induced. (b) Left stifle joint at week 5 p.i. showing normal looking articular cartilage (AC) but with slightly swollen soft tissue (arrow).	71
5.1	Photomicrograph of the synovial membrane at week 1 p.i. The arrow is showing the thickened intima layer (S), which is three to four layers thicker. The border is irregular and diffusely ulcerated. Notice the marked vascular congestion (bv) and oedema (O). H&E x100.	92
5.2	Photomicrograph of the synovial membrane at week 1 p.i. In response to arthritis, the synovium proliferates with villous like projections (arrow). H&E x100.	92
5.3	Photomicrograph of the synovial membrane at week 2 p.i. showing hyperplasia (H) of the subintima layer with mononuclear inflammatory cells (arrow). H&E x100	93
5.4	Photomicrograph of the synovial membrane at week 4 p.i. The synovial subintima shows generalised and severe hyperplasia (H) with the presence of lymphocytes (arrow), indicating chronic synovitis. H&E x100	93
5.5	Photomicrograph of the articular cartilage at week 1 p.i. showing flaking and erosion (arrow). The increased density from zone 1 (A) to tidemarks (T) is evidenced. H&E x100	94

- 5.6 Photomicrograph of the articular cartilage at week 2 p.i. showing flaking and mild fibrillation of the articular surface. H&E x40 94
- 5.7 Photomicrograph of the articular cartilage at week 3 p.i. showing reduced density between zone 1 (A) and tidemarks (T). H&E x100 95
- 5.8 Photomicrograph of the articular cartilage at week 5 p.i. The big arrow is pointing at the superficial fissures or laceration. There was also some clustering of the chondrocytes (small arrow) observed. H&E x100 95
- 5.9 Electron micrograph showing the normal articular cartilage of a dog. The perichondrium (P) has numerous depressions that can serve as reservoirs for synovial fluids. The perichondrium (P) is folded, to show the collagen fibres (F), in the chondrocyte layer. SEM x 330. 96
- 5.10 Electron micrograph of the articular cartilage with severe changes on the surface (lesion score 3). The cartilage shows severe damage with visible chondrocytes layer (C), observed on week 1 p.i. SEM x 1,100 96
- 5.11 Electron micrograph of the articular cartilage with severe lesions with score 4. The damage on the chondrocytes layer (C), observed on week 2 p.i. is severe. SEM x 2,200 97
- 5.12 Electron micrograph of the articular cartilage with moderate lesions with the score of 2. There are moderate destruction (D) leading to erosion of the articular surface, observed on week 4 to 5 p.i. SEM x 1,900 97
- 5.13 Electron micrograph showing the normal synovial membrane of the adipose type. The dome-shaped synoviocyte layer (S), standing out well above the subintimal layer. SEM x750 98
- 5.14 Electron micrograph showing the normal synovial membrane of the fibrous type, is folded and has a surface mat (M) of fibres. SEM x1,000 98
- 5.15 Electron micrograph showing the normal synovial membrane of the areolar type. SEM x1,000 99

LIST OF ABBREVIATIONS

%	percent
μm	micrometer
<	Less than
DBKL	Dewan Bandaraya Kuala Lumpur
<i>et al.</i>	And others (Latin: et alii)
H&E	haematoxylin-eosin
HA	Hyaluronic acid
IL-1	interleukin 1
ILT-6	interleukin-6
kg	kilogram
mg/kg	milligram perkilogram
MIA	Sodium monoiodoacetate
ml	mililitre
mm	milimeter
NSAIDs	non-steroidal anti-inflammatory drugs
OA	osteoarthritis
OCD	osteocondritis dissecans
OsO ₄	osmium tetroxide
P	probability
p.i.	Post induction / post inoculation
PG	proteoglycan
PGE ₂	prostaglandin
PSGAG	Polysulphated Glycoaminoglycans
SEM	Scanning electron microscopy
TNF-α	Tumor necrosis factor



CHAPTER 1

INTRODUCTION

1.1 General Introduction

Arthritis in a simple definition is inflammation of the joint. Osteoarthritis, also known as degenerative joint disease is not a single disease or process, but rather the clinical and pathological outcome of a range of disorders (Nuki, 1999) characterized by progressive deterioration of the articular cartilage, synovitis and joint effusion (McIlwraith, 1996).

The purpose of joint is to support the greatest stability to the body during weight bearing and motion. Painless and full range of joint motion are needed for normal ambulation and performance of daily living chores (Leach and Jacobs, 1990). Interruption of normal joint mechanics leads to painful osteoarthritis and physical incapacity thereby reducing an individual's quality of life and increasing the burden on others (Bennet, 1990). Thus, diseases of joint are said to constitute the greatest single cause of disability encountered by the medical profession (McIlwraith, 1996).

Osteoarthritis is one of the most frequently encountered joint diseases in dogs (Bennet, 1984). Developmental diseases such as patella luxation and osteochondrosis of the femoral condyles, and traumatic and degenerative diseases such as rupture of the cranial cruciate ligament or primary degenerative joint disease are seen frequently in small animal practices (Payne and



Constantinescu, 1993). Although the cause of osteoarthritis varies, initial changes in many animals include inflammation of the synovial membranes and joint capsule without cartilage damage (Sherman *et al.*, 1999).

In dogs and cats, osteoarthritis is not idiopathic or primary. It is usually secondary to trauma, unstable joints, mal-alignment or conformation defects, or congenital conditions such as osteochondritis dissecans (OCD) and hip dysplasia (Payne and Constantinescu, 1993). Thus, proper diagnostic and management of joint disease depend on the understanding of basic anatomy and physiology of the musculoskeletal system (McIlwraith, 1996).

1.2 Osteoarthritis Study

Osteoarthritis (OA) or degenerative joint disease is a chronic disease involving cartilage degeneration and pain (Simmons *et al.*, 1999). It is an important orthopedic condition that affects diarthrodial joints and results in high morbidity (Anderson *et al.*, 1993). The most common joint that is affected with osteoarthritis is the stifle joint. In dogs, it has been identified as a common joint disease (Sherman *et al.*, 1999), accounting for approximately 37% of all lameness in this species of animal (Bennet, 1980). In dogs of more than 1-year old, osteoarthritis may affect as many as 20% of the dog (Anderson *et al.*, 1993).

Arthritis in humans and animals is characterised by joint degeneration, which is usually accompanied by intra-articular inflammation. Until recently, most emphasis in arthritis research has been focussed on intra-articular events