



UNIVERSITI PUTRA MALAYSIA

**THE ISOLATION OF KOJIC ACID PRODUCING FUNGI AND THE
OPTIMIZATION OF KOJIC ACID PRODUCTION**

MADIHAH MD SALLEH

FSMB 1996 4

**THE ISOLATION OF KOJIC ACID PRODUCING FUNGI AND THE
OPTIMIZATION OF KOJIC ACID PRODUCTION**

BY

MADIHAH MD SALLEH

**Thesis Submitted in Fulfilment the Requirement for the Degree of Master
of Science in the Faculty of Food Science and Biotechnology,
Universiti Pertanian Malaysia**

April 1996



SPECIALLY DEDICATED TO;

HAJJAH KALTHOM ABDUL MAJID (EMAK)

HAJI SALLEH BIN SATAK (ABAH)

MOHD NAJIB SALLEH (ADIK)

HAJJAH KAMARIAH, ISMAIL, ISHAK AND ILIAS

HAJJAH MARIAM, MUNA AND ADAM

***"THANKS FOR YOUR LOVING SUPPORT DURING THE
DIFFICULT TIME"***



ACKNOWLEDGEMENTS

I give thank to Merciful ALLAH for giving me the strengths to complete this study.

I also wish to express my deepest appreciation and gratitude to my committee members; Professor Dr. Mohammed Ismail Abdul Karim, Dr. Arbakariya Ariff, Mr. Ali Hassan and Associate Professor Dr Gulam Rusul for their constructive criticisms, guidance, support and encouragement throughout my graduate study and during the preparation of thesis. Special thanks also due to my former supervisor, Dr Baharuddin Abdul Ghani who is currently working with Land and General Bhd. for his invaluable suggestion and encouragement during my graduate study.

I would also like to express my sincere appreciation to the JICA experts - Professor Sugio, Professor Mitsugi, Associate Professor Dr. Kanzaki and Professor Tano for their attention and suggestion to this project.

Special thanks go to Miss Hasni , Mrs. Fadzillah and Miss Rosmah for their invaluable friendship and help for the last few years.

My thanks are also due to laboratory assistants especially Mr. Abdul Hadi, Mr Rosli, Mr. Azman, Miss Jamaliah and Mr. Chan for their assistance in analysis work, photography and preparation of this thesis.

I would also like to forward my sincere thanks to all research assistants and post graduate students in bioprocess laboratory for their encouragement and moral support.

Finally, I would like to thank my parents for their constant support and for being so patient throughout my graduate study.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	III
LIST OF TABLES	VII
LIST OF FIGURES	IX
LIST OF PLATES	XIII
LIST OF ABBREVIATIONS	XIV
ABSTRACT	XV
ABSTRAK	XVIII
CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	4
Properties of Kojic Acid	4
Applications of Kojic Acid	5
Kojic Acid Production	8
Selection and Screening	9
Strain Improvement	13
Effect of Cultural Conditions on Kojic Acid Production	18
Effect of Medium Composition on Kojic Acid Production	26
Techniques of Fermentation	38
Biosynthetic Pathways of Kojic Acid During Fermentation	48
Concluding Remarks	55



III MATERIALS AND METHODS.....	58
Equipments	58
Chemicals	61
Media.....	63
Experimental Design	64
Isolation, Screening and Identification of Kojic Acid Producer.....	65
Optimization of Culture Condition and Medium Composition for Kojic Acid Production	71
Optimization of Aeration Condition for Kojic Acid Production	74
Effect of Yeast-Extract Feeding During Kojic Acid Production in a Fed-Batch Culture using Two Phases Dissolved Oxygen Control	78
Analytical Method	80
Kojic Acid Analysis.....	80
Reducing Sugar Analysis	81
Determination of Non Reducing Sugar	82
Determination of Starch Concentration	82
Determination of Pellets Diameter.....	82
Dry Cell Weight Determination	83
Total Nitrogen Determination	83
Specific Oxygen Uptake Rate (QO_2) Determination	83
IV RESULTS AND DISCUSSIONS	86
Isolation, Screening and Identification of Kojic Acid Producer.....	86
Isolation and Screening of Kojic Acid Producer	86
Selection of Medium and Fermentation Mode	93
Monospores Isolation, Screening and Maintenance of Microorganism.....	101
Identification of Strain 44 - 1.....	109
Optimization of Medium and Cultural Condition for Kojic Acid Production in Shake Flask Culture.....	109
Effect of Initial Culture pH	109

Effect of Calcium Carbonate and Methanol.....	116
Effect of Methanol Concentration.....	122
Effect of Carbon Sources.....	125
Effect of Glucose Concentration.....	128
Effect of Nitrogen Sources.....	131
Effect of Yeast-Extract Concentration.....	135
Production of Kojic Acid Using Optimized Cultural Conditions	138
Aeration Control Strategy for the Improvement of Kojic Acid.....	140
Kojic Acid Production in Fermentation Without Dissolved Oxygen Control	140
Kojic Acid Production in Fermentation with Single Phase Dissolved Oxygen Control	143
Kojic Acid Production in Fermentation with Two Phases Dissolved Oxygen Control	150
 Effect of Yeast Extract Feeding During Kojic Acid Production Phase in Fed-Batch Culture	 153
V CONCLUSIONS AND SUGGESTIONS FOR FURTHER WORK	156
REFERENCES.....	164
APPENDICES	176
BIOGRAPHICAL SKETCH	189



LIST OF TABLES

Table		Page
1	Applications of Kojic Acid	6
2	Toxins Produced by Yellow - Green <i>Aspergilli</i>	11
3	Screening of <i>Aspergillus</i> and <i>Penicillium</i> Species for Aflatoxin and Kojic Acid Production	12
4	Mutagens Effective Against Fungi and Their Action	15
5	Comparison of Sexual and Parasexual Mechanism for Recombination.....	17
6	Optimum Temperature for Kojic Acid and Aflatoxin Production	19
7	Influence of pH, Carbon and Nitrogen Sources During Kojic Acid Fermentation	22
8	Effect of Aeration on Certain Products	24
9	Order of Preference of Carbon Sources for Kojic Acid Production	27
10	Order of Preference of Nitrogen Sources for Kojic Acid Production	30
11	Order of Strength of Inhibitors on Kojic Acid Production	32
12	Stimulants Used for Kojic Acid and Other Organic Acids Fermentation.....	34
13	Composition of Isolation Medium (M1 Medium)	63



14	Composition of Screening Medium (M2 Medium)	63
15	Composition of Growth Medium for Plate and Slant Culture (M3 Medium)	64
16	Composition of Selected Growth and Production Media	68
17	Dissolved Oxygen Tension Control Strategy in Batch Culture	77
18	Sources of Isolated Strains	87
19	Screening Result of Locally Isolated Strains	88
20	Effect of Glucose Concentration on Growth of <i>A flavus</i> Link and Kojic Acid Production	131
21	Effect of Nitrogen Sources and Concentration on Growth of <i>A flavus</i> Link and Kojic Acid Production	133
22	Effect of Yeast-extract Concentration on Growth of <i>A flavus</i> Link and Kojic Acid Production	137
23	Comparison of Performance Parameters of Batch and Fed-Batch Fermentations with Different DOT Control Strategies	149



LIST OF FIGURES

Figures	Page
1	Structure of Kojic Acid..... 4
2	A Typical Time Course of Kojic Acid Fermentation in a Stirred Tank Fermenter 43
3	Biosynthesis Pathway of Kojic Acid Formation by <i>Aspergillus flavus</i> 53
4	Flowchart of Experimental Design of Kojic Acid Production by <i>Aspergillus flavus</i> Link St 44-1..... 63
5	Fermenter Set up for Batch Culture 75
6	Fermenter Set up for Fed-Batch Culture 79
7	Correlation Between HPLC and Ferric Chloride Methods for Kojic Acid Quantification in Fermentation Broth 81
8	Dynamic Gassing Out Technique for Determination of Specific Oxygen Uptake Rate During Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link 85
9	Scanning Result of Culture Filtrate of Strain 44-1 after React with FeCl ₃ 91
10	Scanning Result of Standard Kojic Acid After React with FeCl ₃ 91
11	Schematic Diagram of Qualitative Analysis of Kojic Acid Production by Several Strains Using Thin Layer Chromatography (TLC) Plate..... 92



12	Kojic Acid Production by Two Different Strains of Fungus in Shake Flask Culture.....	93
13	Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link Strain 44-1 in Two Different Type of Media Using (A) Agitated and (B) Static Shake Flask Culture	96
14	Preliminary Study on the Effect of Different Carbon Sources on Kojic Acid Production by <i>Aspergillus flavus</i> Link	98
15	Preliminary Study on the Effect of Different Nitrogen Sources on Kojic Acid Production by <i>Aspergillus flavus</i> Link	100
16	Screening of Monospores Isolated from F1 and F2 of <i>Aspergillus flavus</i> Link : Kojic Acid Production after 9 Days of Cultivation in 16 mm Diameter Test	104
17	Screening of Monospores Isolation : Effect of 4% Methanol and 0.5% CaCO ₃ on Kojic Acid Production by <i>Aspergillus flavus</i> Link (F1-1, F1-5 and F1-15).....	106
18	Screening of Monospores Isolation : Kojic Acid Production by <i>Aspergillus flavus</i> Link (F1 - 1) in Shake Flask Culture after 9 Days of Cultivation	108
19	Screening of Monospores Isolation : Kojic Acid Production by <i>Aspergillus flavus</i> Link (F1 - 1 - 28) in Shake Flask Culture After 9 Days of Cultivation	108
20	Effect of Different Initial Culture pHs on the Maximum Concentration of Cell (<i>X</i>), Kojic Acid (<i>P</i>) and <i>P/X</i> After 15 Days of Cultivation in Shake Flask Culture.....	110
21	Profile of Culture pH During Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link in Shake Flask Culture at Different Initial Culture pHs (A) Initial Culture pH of Between 2 - 5 (B) Initial Culture pH of Between 5.5 - 8	114



22	Profile of Glucose Consumption During Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link in Shake Flask Culture at Different Initial Culture pHs (A) Initial Culture pH between 2 - 5.5 (B) Initial Culture pH of between 5.5 - 8	115
23	Effect of Methanol and CaCO ₃ on (A) Kojic Acid Production and Glucose Consumption (B) pH Profile and Cell Growth	118
24	Effect of the Addition of Methanol into a Medium on Maximum Concentration of Cell (<i>X</i>), Kojic Acid (<i>P</i>) <i>P/X</i> and yield after 21 Days of Cultivation in Shake Flask Culture.....	123
25	Effect of Pellet Size on Kojic Acid Production by <i>Aspergillus flavus</i> Link in Shake Flask Culture. In all Cases, Kojic acid was Measured at 21 Days of Cultivation in Shake Flask Culture	124
26	Time Courses of Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link Using Different Type of Carbon Sources	126
27	Time Courses of Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link in Different Glucose Concentrations	129
28	Time Courses of Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link Using Different Type of Nitrogen Sources.....	132
29	Time Courses of Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link in Different Yeast Extract Concentrations.....	136
30	Time Course of Kojic Acid Fermentation by <i>Aspergillus flavus</i> Link in Optimized Glucose - Salt Medium in Shake Flask Culture	139
31	Time Course of Kojic Acid Fermentation Without Dissolved Oxygen Control. Air Flow Rate was Fixed at 1 L/ min	141



32	The Profile of DOT During Kojic Acid Fermentation with Different Oxygen Control Strategies	144
33	The Profiles of Specific Oxygen Uptake Rate During Kojic Acid Fermentation at 50% and 80% Oxygen Saturation	145
34	Time Course of Batch Kojic Acid Fermentation in which DOT were Controlled at 30%, 50% and 80% Saturations	147
35	Time Course of Batch Kojic Acid Fermentation with Two Phases DOT Control in which DOT was Controlled at 80% Saturation during Growth Phase and 30% During Production Phase	152
36	Time Course of Fed-Batch Kojic Acid Fermentation with Two Phases DOT Control in which DOT was Controlled at 80% Saturation during Growth Phase and 30% during the Production Phase	154

LIST OF PLATES

Plate		Page
1	Strain 44 - 1 (<i>Aspergillus flavus</i> Link).....	89
2	Strain 34 - 1	89
3	Monospores Isolation of Strain 44 - 1 (F1)	103
4	Monospores Isolation of Strain 44 - 1 (F2)	103
5	Pellets of <i>Aspergillus flavus</i> Link in Control Medium (A) and with the Addition of 4% Methanol (B)	121
6	Pellets of <i>Aspergillus flavus</i> Link in Medium with the Addition of 4% Methanol Under 40X Magnification.....	121



LIST OF ABBREVIATIONS

h	:	Hour
QO_2	:	Specific Oxygen Uptake Rate (mmol O_2/ g Cell)
P	:	Product Concentration (g/L)
X	:	Cell Concentration (g/L)
t	:	Time (hour)
DOT	:	Dissolved Oxygen Tension (%)



Abstract of thesis submitted to the Senate of Universiti Pertanian Malaysia
as fulfilment of the requirements for the degree of Master of Science.

**THE ISOLATION OF KOJIC ACID PRODUCING FUNGI AND THE
OPTIMIZATION OF KOJIC ACID PRODUCTION**

BY

MADIHAH MD SALLEH

APRIL 1996

Chairman : Professor Mohammed Ismail Abdul Karim, PhD.

Faculty : Food Science and Biotechnology

A local strain of fungus, which was found to be a high kojic acid producer has been isolated from Morning glory flower (*Bixa orellana*) using glucose-peptone medium. The pure strain was obtained through several steps of monospore isolation procedures using spread plate technique and identified by International Mycological Institute as *Aspergillus flavus* Link.

Optimization of medium composition and cultural conditions for kojic acid production by this fungus were carried out in shake flask. The development of dissolved oxygen tension (DOT) control strategies aimed at improving kojic acid production by this strain was carried out using 1.5 L stirred tank reactor with automatic pH and DOT control systems. Initial culture pH 3 was found to



be optimum for kojic acid production. This strain was able to grow and produce kojic acid in various carbon sources such as glucose, starch, sucrose and xylose. However, the highest production of kojic acid was obtained at 100 g/L (w/v) glucose. In addition, 5 g/L of yeast-extract was found to be the best nitrogen source for fungal growth and enhancement of kojic acid production. Addition of 4% (v/v) of methanol to glucose yeast-extract medium increased kojic acid production by two times. The optimized medium for kojic acid production for this strain was proposed and the medium consisted of 100 g/L glucose, 1.0 g/L KH_2PO_4 , 0.5 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5.0 g/L yeast-extract and 4% methanol. Using this optimal medium the maximum kojic acid production in batch fermentation using shake flask was 39 g/L. This fermentation gave yield and productivity of 0.04 g/g and 0.07 g/L/h, respectively and is comparable to that reported in the literature for industrial strain.

The DOT control strategy for the improvement of the kojic acid production by this strain was also developed. By controlling DOT at 80% saturation during growth phase and then switched to 30% saturation during production phase, increased the production of kojic acid by about two times as compared to uncontrol fermentation in the stirred tank fermenter which only produced maximum concentration of kojic acid of 15 g/L.



The effect of the addition of yeast-extract during the fermentation was also investigated using a constant fed-batch culture. The addition of 15 g/L yeast-extract at constant flow rate of 3.2 ml/h during the production phase caused excessive mycelial growth and decreased kojic acid production.

Abstrak tesis yang dikemukakan kepada Senat Universiti Pertanian Malaysia sebagai memenuhi keperluan Ijazah Master Sains

**PEMENCILAN FUNGI YANG MENGHASILKAN ASID KOJIK DAN
PENGOPTIMUMAN PENGHASILAN ASID KOJIK**

OLEH

MADIHAH MD. SALLEH

APRIL 1996

Pengerusi : Professor Mohammed Ismail Abdul Karim, PhD.

Fakulti : Sains Makanan Dan Bioteknologi

Kulat tempatan yang berupaya menghasilkan asid kojik yang tinggi telah dipencilkan daripada bunga Kembang Pagi (*Bixa orellana*) dengan menggunakan media glukosa-pepton. Strain tulen yang didapati melalui beberapa peringkat pemencilan spora tunggal dengan menggunakan teknik plat sebaran telah dikenalpasti oleh Institut Antarabangsa Mikologi sebagai *Aspergillus flavus* Link.

Pengoptimuman komposisi media dan keadaan kultur untuk penghasilan asid kojik oleh kulat ini telah dijalankan dalam kelalang penggoncang. Strategi pengawalan kepekatan oksigen terlarut (DOT) dengan tujuan untuk



meningkatkan asid kojik oleh strain ini telah dijalankan dengan menggunakan fermenter berpengaduk dengan sistem kawalan pH dan oksigen terlarut secara automatik. pH awalan optimum bagi penghasilan asid kojik adalah 3. Strain ini berupaya tumbuh dan menghasilkan asid kojik dalam pelbagai sumber karbon seperti glukosa, kanji, sukrosa dan xilosa. Walaubagaimanapun, penghasilan asid kojik tertinggi didapati pada 100 g/L glukosa. Selain daripada itu, 5 g/L yis-ekstrak adalah sumber nitrogen terbaik untuk pertumbuhan kulat dan peningkatan penghasilan asid kojik. Penambahan 4% metanol ke dalam media glukos yis-ekstrak meningkatkan penghasilan asid kojik sebanyak dua kali ganda. Media optimum untuk penghasilan asid kojik oleh strain ini adalah terdiri daripada 100 g/L glukosa, 1.0 g/L KH_2PO_4 , 0.5 g/L $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$, 5.0 g/L yis-ekstrak dan 4% metanol. Penggunaan media optimum dalam kultur sesekelompok dengan menggunakan kelalang penggoncang menghasilkan maksimum asid kojik 39.9 g/L. Fermentasi ini juga memberikan 0.04 g/g kadar penghasilan serta 0.07 g/L/h produktiviti dan hasil kajian ini adalah sama seperti strain industri yang telah dilaporkan oleh kajian bahan bertulis.

Strategi pengawalan oksigen terlarut untuk meningkatkan penghasilan asid kojik telah dijalankan. Pengawalan oksigen terlarut pada ketepuan 80% semasa fasa pertumbuhan dan diubah ke 30% semasa fasa penghasilan meningkatkan penghasilan asid kojik dua kali ganda berbanding dengan



fermentasi tanpa kawalan di dalam fermenter berpengaduk yang menghasilkan kepekatan asid kojik maksimum 15 g/L.

Kesan penambahan yis-ekstrak semasa fermentasi juga dikaji dengan menggunakan kultur suapan sesekelompok tetap. Penambahan 15 g/L yis - ekstrak pada kadar aliran tetap 3.2 ml/j semasa fasa penghasilan menyebabkan peningkatan pertumbuhan kulat dan penurunan penghasilan asid kojik.

CHAPTER I

INTRODUCTION

Kojic acid is secreted by several microorganisms such as *Aspergillus oryzae* and *Aspergillus flavus*. This crystalline substance was first isolated by Saito in 1907 from the mycelia of *Aspergillus oryzae* grown on steamed rice. The chemical structure was determined as 5-hydroxy-2-hydroxymethyl- δ pyrone by Yabuta (1924). Although kojic acid can be synthesized from chemical conversion of various substrates, commercially it is produced by aerobic fermentation of *Aspergillus* species.

Kojic acid has several economic uses in various fields. In medical field, kojic acid is used as an antibacterial and antifungal agents. In the food industries, kojic acid is used as an antispeck and antimelanosis (blackening) agents for agricultural products. Since Malaysia has many agricultural products, the use of kojic acid will be economically important in the post harvest process. In addition, kojic acid is also used as a chelating agent and activator in insecticide production. Recently, a new application of kojic acid is found in the cosmetic industry. It is used as a whitening agent and ultraviolet filter in skin care products.



Although kojic acid has been produced and applied industrially, attempts to improve kojic acid production are still being intensively studied. Two main areas which are normally considered for the improvement of kojic acid production are the improvement of strain and development of the fermentation process. Screening of high kojic acid producer from various sources and also improvement of strain through mutation process had been conducted in the last few years. Although works on optimization of medium composition and environmental condition for kojic acid production by microorganisms have been studied extensively, the effect of each nutrient component on kojic acid production had not been well described.

Recently, the use of immobilized viable cells as an approach to enhance kojic acid production has also been reported (Kwak et al., 1991). However, work on the improvement of the production of kojic acid using different free cell fermenter operations such as fed-batch culture in bioreactor is not reported. Information on the influence of aeration condition on kojic acid production is also scarce.

The objectives of the research is to (1) screen and isolate potential kojic acid producer from local sources including flowers, fruits, fermented food and hot spring (2) optimize the culture medium condition using various nitrogen and carbon sources and stimulant, (3) study on the effect of dissolved oxygen

tension (DOT) during kojic acid fermentation in batch culture using without DOT control, single phase DOT control and two phases DOT control and (4) investigate the effect of the addition of nitrogen source (yeast-extract) during kojic acid production phase in fed-batch culture.

CHAPTER II

LITERATURE REVIEW

Properties of Kojic Acid

Kojic acid is a monocyclic pyrone consisting of carbon ring with two double bonds (Figure 1). The hydroxyl group at position carbon 5 has weakly acidic characteristics that are capable of producing salts with few metals. The side chain of C5 also behaves as a primary alcohol whose reactivity was enhanced by the adjacent oxygen atom in the nucleus (Beelik, 1956). Kojic acid forms a complex with ferric ions to produce a reddish purple colour which has maximum absorption at 500 nm (Bently, 1957; Crueger et al., 1984). This colorization is widely used as a principle method for the quantitative determination of kojic acid.




Figure 1 : Structure of Kojic Acid

(Source : Beelik, 1956)