



UNIVERSITI PUTRA MALAYSIA

**EXTRACT OF *GELIDIELLA ACEROSA* (S-ACT-1) AS A
CAPACITATION AGENT FOR *IN VITRO*
FERTILISATION IN GOAT**

ALAWATTAGE DON NIMAL CHANDRASIRI

FPV 1997 6

**EXTRACT OF *Gelidiella acerosa* (S-ACT-1) AS A CAPACITATION AGENT
FOR *IN VITRO* FERTILISATION IN GOAT**

By

ALAWATTAGE DON NIMAL CHANDRASIRI

**Dissertation Submitted in Fulfilment of the Requirements for
the Degree of Doctor of Philosophy in the Faculty of
Veterinary Medicine and Animal Science,
Universiti Pertanian Malaysia.**

March 1997



DEDICATION

This thesis is exclusively dedicated to the memory of my late father

Mr. A.D. Gunatissa

and

my beloved mother

Mrs . R.A.D.C. Rupasinghe



ACKNOWLEDGEMENTS

From the beginning of my study to the conclusion of this final product I have come across success, failure, suffering and confidence, all combined to understand the real spirit of a Doctor of Philosophy. I owe my deep appreciation to many individuals and organisations that helped me in one way or another during this arduous journey. First of all I would like to extend my utmost appreciation and gratitude to Professor Dr. M.R. Jainudeen for his invaluable guidance, advice, constructive criticisms and comments throughout this study. I could never forget his excellent untiring support extended to me in the preparation of this thesis.

I wish to express my sincere gratitude and deep appreciation to Professor Dr. Tengku Azmi Ibrahim and Dr. Abdul Wahid Haron, the other two members of the supervisory committee for their valuable suggestions and for providing animals and other facilities to carry out the experiments at Universiti Pertanian Malaysia (UPM.)

I gratefully acknowledge the encouragement given by Dr. M.C.L.De Alwis, the Additional Director of the Department of Animal Production and Health (D.A.P.& H.), Sri Lanka. I would have neither commenced nor completed this study without his moral support throughout this study. My sincere thanks are also due to Professor Viranjane Gunawardene, Department of Veterinary Pre-Clinical Studies and Dr. H. Abeygunawardene, Senior Lecturer, Department of Veterinary Clinical Studies, Faculty of Veterinary Medicine and Animal Science, University of Peradeniya, Sri Lanka for their guidance.



My thanks are due to Dr. Sirimal Premakumara, Department of Zoology, Faculty of Science, University of Colombo, Sri Lanka for generously donating the major test material (S-ACT-1) to conduct the study. My thanks are also due to the Deputy Director (Animal Breeding), Officer in Charge and his staff at the Central Artificial Insemination Station, of the Department of Animal Production and Health, Sri Lanka for providing the goats and other facilities to collect and evaluate semen and the Medical Research Institute, Colombo for providing hamsters. I am also grateful to Mr. Yap Keng Chee, Mrs. Haliza Hasim and Mr. Abu Bakar Dahari at the Department of Veterinary Clinical Studies, Faculty of Veterinary Medicine and Animal Science, UPM and Miss R.M.S. Malkanthi, Mr. A.S.B. Ratnayake and Mr. R.G. Senaratne at the Veterinary Research Institute, Sri Lanka for their technical assistance.

Special thanks are due to Professor R.O. Thattil, Department of Crop Science, Faculty of Agriculture, University of Peradeniya and Dr. M.G. Jeyaruban, Veterinary Research Institute, Sri Lanka for their assistance during data analysis. I am grateful to the Agricultural Research Policy (ARP), Sri Lanka for providing financial support for the overseas study component and Council for Agricultural Research Policy (CARP) for providing financial support for the local research component. Last but not least my deepest gratitude is due to my beloved wife and two little daughters, Vidumini and Nethmini for their love, understanding and sacrifice during my stay outside the country.



TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iii
LIST OF TABLES	viii
LIST OF FIGURES	x
LIST OF PLATES	xi
LIST OF ABBREVIATIONS	xii
ABSTRACT.....	xiv
ABSTRAK	xvi
CHAPTER	
I INTRODUCTION.....	1
II LITERATURE REVIEW	4
Introduction	4
<i>In Vivo</i> Fertilisation	4
Spermatogenesis.....	4
Folliculogenesis	9
Oocyte maturation.....	10
Fertilisation	14
<i>In Vitro</i> Fertilisation (IVF)	21
Success rates	22
Uses of IVF	23
Embryo Production through <i>In Vitro</i> Techniques	26
<i>In Vitro</i> Maturation of Oocytes	26
Different IVM Systems	34
Assessment of <i>In Vitro</i> Maturation	37
Preparation of Sperm for IVF	38
Capacitation Media:	46
Acrosome Reaction	47
<i>In Vitro</i> Culture of embryos	55
Co-Culture of Embryos	56



III	ACROSOME REACTION OF GOAT SPERMATOZOA	64
	Introduction	64
	Objectives.....	65
	Materials and Methods	65
	Semen Donors.....	65
	Room and Glassware Sterility.....	66
	Preparation of Semen Samples	66
	Dual Stain Procedure	69
	Examination of the Spermatozoa	69
	Results	71
	Discussion.....	78
IV	ASSESSMENT OF SPERM CAPACITATION IN GOAT	84
	Introduction	84
	Materials and Methods	85
	Sterility of Glassware and Room	85
	Media Preparation	85
	Preparation of Zona-free Ova from Hamster	86
	Collection of Semen from Goat	88
	Preparation of Fertilisation Droplets.....	88
	<i>In Vitro</i> Fertilisation of Hamster Ova	88
	Statistical Analysis.....	89
	Results	90
	Discussion.....	95
V	<i>IN VITRO</i> OOCYTE MATURATION, FERTILISATION AND EMBRYO CULTURE IN GOAT	100
	Introduction	100
	Objectives.....	101
	Materials and Methods	101
	Animals.....	101
	Room and Glassware Sterility.....	101
	Recovery of Oocytes	101
	<i>In vitro</i> Maturation.....	103
	Results	108



	Recovery of Oocytes.....	108
	Maturation.....	108
	Cleavage Rate	108
	Embryo Culture and Development	112
	Discussion.....	122
	Collection and Transport of Ovaries	122
	Oocyte Recovery.....	122
	Criterion for Fertilisation	123
	Capacitation Agents	124
	Conclusions.....	125
VI	GENERAL DISCUSSION.....	127
	Capacitation Agents.....	128
	Problems Associated with IVF	129
	Factors Affecting IVF.....	129
VII	SUMMARY AND CONCLUSIONS.....	134
	BIBLIOGRAPHY.....	138
	APPENDIX-A.....	162
	APPENDIX-B	169
	APPENDIX-C	177
	BIOGRAPHICAL SKETCH	181



LIST OF TABLES

Table		Page
1	Staining Characteristics of Goat Spermatozoa after Dual Staining Procedure.....	72
2	Analysis of Variance and F Test for Capacitation Agents on LA(-) Spermatozoa	74
3	Analysis of Variance and F Test for Capacitation Agents on LA (-) Spermatozoa	76
4	Analysis of Variance and F Test for Effect of Capacitation Agents on DA(-) Spermatozoa.....	77
5	Analysis of Variance and F Test for Capacitation Agents on DA(-) Spermatozoa	78
6	Penetration of Zona Free Hamster Oocytes by Goat Spermatozoa Capacitated with Heparin and S-ACT-1 at Different Incubation Times	92
7	Analysis of Variance and F Test for Effect of Capacitation Agents, Concentrations and Incubation Time	93
8	Summary of Steps for <i>In vitro</i> Fertilisation and Embryo Culture.....	107
9	Recovery of Follicular Oocytes from Slaughterhouse Goat Ovaries Using Two Different Recovery Techniques	108
10	Sperm Cleavage Rates of <i>In Vitro</i> Matured Goat Oocytes after Incubation with Various Concentrations of Heparin and S-ACT-1.	109
11	Analysis of Variance and F test for Cleavage Rate.....	111
12	Development of <i>In Vitro</i> Produced 2-cell Goat Embryos in Two Different Culture Media.....	115
13	Analysis of Variance and F test for % Survival at the 4- cell Stage	116
14	Analysis of Variance and F test for % Survival at the 8-cell Stage	118
15	Analysis of Variance and F test for % Survival of <i>Morulae</i>	119
16	Analysis of Variance and F-test for % Survival at the <i>Blastocyst</i> Stage.....	120



17	Composition of Biggers, Whitten and Whittingham (BWW medium) or Modified Kreb's Ringer's Solution.....	163
18	Composition of Sperm Capacitation Medium (Modified Ca ⁺⁺ free Tyrodes Medium)	164
19	Composition of Fertilisation Medium (Modified Tyrode's Medium).....	165
20	Composition of Wash II Medium (Modified Hapes-buffered Tyrode's Medium).....	166
21	Reproductive Cycle of Female Golden Hamsters (<i>Mesocricetus auratus</i>).....	167
22	Superovulation Schedule of Golden Hamsters (<i>Mesocricetus auratus</i>).....	168



LIST OF FIGURES

Figure		Page
1	Swim-up Preparations of Goat Spermatozoa for <i>In Vitro</i> Fertilisation.....	68
2	Staining Patterns of the Spermatozoa after Dual Staining Method.	70
3	Interaction between the Capacitation Agents and the Concentration on True Acrosome-Reacted (TAR) Goat Spermatozoa.....	75
4	Effect of Capacitation Agent and Concentration on Goat Sperm Penetration of Zona-free Hamster Ova.	94
5	Interaction between the Capacitation Agents and Incubation Period on the Penetration Rate.	95
6	Effect of Concentrations of the Capacitation Agents on Cleavage Rate of Oocytes in Goat.	112
7	Interaction of Capacitation Agent and Concentration on the Survival Rate of 4-cell Embryos.	117
8	Interaction between Capacitation Agents and Concentration on the Survival Rate of 8-cell Embryos.....	119
9	Interaction between the Capacitation Agent and Concentration on the Survival Rate of <i>Morulae</i>	120
10	Interaction between the Capacitation Agent and Concentration on the Survival Rate of <i>Blastocysts</i>	121
11	Relationship between the Survival Rate and the Developmental Stages of Goat Embryos Developed <i>In Vitro</i>	122



LIST OF PLATES

Plate		Page
1	Goat Spermatozoa Stained with Trypan Blue-Giemsa (40 x 10)	73
2	Zona Free Hamster Ova Penetration Test for Goat Sperm	91
3	<i>In vitro</i> Maturation of Goat Oocytes	110
4	<i>In Vitro</i> Culture of Goat Embryos.....	113
5	Embryos Derived from <i>In Vitro</i> Fertilisation.....	114



LIST OF ABBREVIATIONS

AI	Artificial insemination
AR	Acrosome reaction
BOEC	Bovine oviductal epithelial cell
BSA	Bovine serum albumin
BWW	Biggers, Whitten and Whittingham
CAIS	Central Artificial Insemination Station
COC	Cumulus ophorus complexes
COECM	Caprine oviductal epithelial cell monolayer
DA-	Dead sperm with detached acrosome
DA+	Dead sperm with intact acrosome
DAP&H	Department of Animal Production and Health
ESS	Oestrus sheep serum
ET	Embryo transfer
FAR	False acrosome reaction
FCS	Foetal calf serum
FSG	Fucose sulphated glycoconjugate
FSH	Follicular stimulating hormone
GAG	Glycoseaminoglycans
GV	Germinal vesicle
GVBD	Germinal vesicle break down
hCG	Human chorionic gonadotrophin



IU	International units
IVC	<i>In vitro</i> culture
IVF	<i>In vitro</i> fertilisation
IVM	<i>In vitro</i> maturation
LA-	Live sperm with detached acrosome
LA+	Live sperm with intact acrosome
LH	Luteinising hormone
MOET	Multiple ovulation and embryo transfer
MRI	Medical Reserach Institute
PBS	Phosphate buffered saline
PHE	Penicillamine, hypotaurine and epinephrine
PMSG	Pregnant mare serum gonadotrophin
POD	Post ovulatory discharge
S-ACT-1	Sperm activating factor 1
SOF	Synthetic oviductal fluid
SS	Sheep serum
TAR	True acrosome reaction
TCM 199	Tissue culture medium 199
UPM	Universiti Pertanian Malaysia
VRI	Veterinary Research Institute
ZHOPT	Zona-free hamster oocyte penetration test
ZP	Zona pellucida



Abstract of dissertation submitted to the Senate of Universiti Pertanian Malaysia in fulfilment of the requirements for the degree of Doctor of Philosophy

EXTRACT OF *Gelidiella acerosa* (S-ACT-1) AS A CAPACITATION AGENT FOR *IN VITRO* FERTILISATION IN GOATS

By

ALAWATTAGE DON NIMAL CHANDRASIRI

March, 1997

Chairman: Professor Tengku Azmi bin Tengku Ibrahim, Ph. D

Faculty: Veterinary Medicine and Animal Science

A study was conducted to determine if goat ovaries obtained from a slaughterhouse could be used for *in vitro* fertilisation (IVF). Three experiments were conducted to ascertain if a substance named S-ACT-1, extracted from a marine algae, (*Gelidiella acerosa*), with a chemical structure similar to heparin could be used for sperm capacitation.

The acrosome reaction was tested by a dual staining technique (Experiment 1) and the zona free hamster penetration test (Experiment 2). *In vitro* maturation of goat oocytes and fertilisation were assessed by the number of zygotes cleaving beyond the 2-cell stage (Experiment 3).

Freshly ejaculated semen from two fertile bucks were used for sperm capacitation using two defined media (DM-H and DM-Ca). Heparin and S-ACT-1 were tested at concentrations of 10, 20, 50 and 100 µg /ml. After incubating for 15 min. at 39° C with 5% CO₂ in air under maximum humidity, samples were stained with trypan blue and Giemsa stain (dual stain). The results indicated that the sperm treated with heparin and S-ACT-1 showed similar staining characteristics.



Concentration of the capacitation agent had a significant effect on the number of spermatozoa undergoing true acrosome reaction. The number of true acrosome reacted spermatozoa was significantly ($F= 6.69$) different between the treatment and control groups.

The zona free hamster ova penetration test showed that the sperm penetration rate was higher with heparin than S-ACT-1. Based on the results of Experiments 1 and 2, S-ACT-1 can be used as a sperm capacitation agent for IVF in goat at a concentration of 20 $\mu\text{g/ml}$ and 3 h of incubation. A total of 865 oocytes obtained from slaughtered goats were incubated for 27 h at 39° C and 5% CO_2 under maximum humidity. The maturation rate was 89.75%. Matured oocytes were inseminated with the goat spermatozoa capacitated with 10, 20, 50, 100 and 200 $\mu\text{g/ml}$ of heparin or S-ACT-1 and incubated for 24 h. The maximum cleavage rate was 42.2% (38 out of 90) for 10 $\mu\text{g/ml}$ heparin and 33.8% (23 out of 68) for 20 $\mu\text{g/ml}$ S-ACT-1. Two celled embryos were cultured either in synthetic oviductal fluid (SOF) medium or caprine oviductal epithelial cell monolayer (COECM) cultures up to the blastocyst stage. Higher survival rates were observed in the embryos fertilised with the sperm capacitated with 10 $\mu\text{g/ml}$ of heparin (21.1% or 4 out of 19) or 20 $\mu\text{g/ml}$ of S-ACT-1 (25% or 3 out of 12).



Abstrak disertasi yang dikemukakan kepada Senat Universiti Pertanian Malaysia
bagi memenuhi syarat-syarat untuk Ijazah Doktor Falsafah

**Akstrak *Gelidiella acerosa* (S-ACT-1) SEBAGAI AGEN KAPASITASI UNTUK
PERSENYAWAAN *IN VITRO* PADA KAMBING**

oleh

ALAWATTAGE DON NIMAL CHANDRASIIRI

Mac, 1997

Pengerusi: Profesor Tengku Azmi bin Tengku Ibrahim, Ph. D

Fakulti: Kedoktoran Veterinar dan Sains Peternakan

Satu kajian telah dijalankan untuk mengenalpasti samada ovari kambing yang diambil dari rumah sembelih boleh digunakan dalam persenyawaan *in vitro* (IVF). Tiga eksperimen dikendalikan untuk menentukan jika bahan yang dikenali sebagai S-ACT-I, diakstrak dari alga laut, (*Gelidiella acerosa*), mempunyai struktur kimia menyerupai heparin boleh digunakan dalam kapasitasi sperma.

Reaksi akrosom diuji dengan teknik pewarnaan berkembar (Eksperimen 1) dan ujian penembusan ova hamster tanpa zona (Eksperimen 2). Pematangan *in vitro* oosit kambing dan persenyawaannya dinilai melalui bilangan zaigot yang membelah melebihi peringkat 2-sel (Eksperimen 3).

Semen yang baru diejakulat dari dua kambing jantan subur digunakan untuk kapasitasi dengan menggunakan dua media (DM-H and DM-Ca). Heparin dan S-ACT-1 diuji pada kepekatan 10, 20, 50 dan 100µg /ml. Selepas diincubat selama 15 min. pada 39°C dengan 5% CO₂ dalam udara di bawah humiditi maksimum, kesemua sampel diwarnakan dengan pewarna tryphan biru dan Giernsa (pewarnaan berkembar). Keputusan menunjukkan sperma yang dirawat dengan heparin dan S-

ACT-1 memberikan ciri-ciri pewarnaan yang serupa. Kepekatan agen kapasitasi mempunyai kesan yang ketara terhadap bilangan spermatozoa yang mengalami reaksi akrosom sebenar. Bilangan spermatozoa yang mengalami reaksi akrosom sebenar mempunyai perbezaan yang ketara ($F= 6.69$) di antara kumpulan rawatan dan kawalan.

Ujian penembusan ova hamster tanpa zona menunjukkan kadar penembusan sperma yang tinggi dengan heparin berbanding S-ACT-1. Berdasarkan Eksperimen 1 dan 2, S-ACT-1 boleh digunakan sebagai agen kapasitasi yang berkesan untuk IVF kambing pada kepekatan $20\mu\text{g/ml}$ dan 3 jam inkubasi.

Sejumlah 865 oosit didapati dari kambing yang disembelih dan diinkubut selama 27 jam pada 39°C dan 5% CO_2 di bawah humiditi maksimum. Kadar pematangan adalah 89.75% . Oosit matang diinseminasi dengan spermatozoa kambing yang dikapasitasi dengan 10, 20, 50, 100 dan $200\mu\text{g/ml}$ heparin atau S-ACT-1 dan diinkubut selama 24 jam. Kadar penembusan maksima adalah 42.2% (38 dari pada 90) untuk $10\mu\text{g/ml}$ heparin dan 33.8% (23 dari pada 68) untuk $20\mu\text{g/ml}$ S-ACT-1.

Ova yang ditembusi dikultur dalam media cecair oviduk sintetik (SOF) dan monolayer sel epitelium oviduk kambing (COECM) sehingga ke peringkat blastosista. Kadar hidup yang tinggi diperhatikan pada embrio yang disenyawa dengan $10\mu\text{g/ml}$ of heparin (21.1% atau 4 dari pada 19) atau $20\mu\text{g/ml}$ S-ACT- 1 (25.0% atau 3 dari pada 12).

CHAPTER I

INTRODUCTION

Manipulation of reproduction has been accepted as the major tool of optimizing the productivity of farm animals. Since the 1940s, artificial insemination (AI) has made a significant contribution to the genetic improvement of farm animals. This is possible because a few highly selected males produce enough spermatozoa to inseminate thousands of females.

On the contrary, despite the presence of several thousands of oocytes in the ovary, only a few offspring are born during the lifetime of a female. Since the 1970s, efforts have been made to increase the number of offspring from superior animals through superovulation, multiple conceptions recovery and transfer of embryos into the recipients who are genetically less distinguished. This technique known as multiple ovulation and embryo transfer (MOET) can be regarded as the female counterpart of artificial insemination.

Several commercial companies for embryo transfer (ET) in farm animals are found worldwide. However, a major problem confronting ET is the lack of a cheap and reliable technique of inducing multiple ovulation. In recent years, researchers have turned their attention to seek ways of harvesting immature oocytes from developing follicles, promoting their maturation, fertilisation and embryo culture *in vitro* to obtain a large number of embryos for embryo transfer.



These *in vitro* techniques are gaining importance for several other reasons: to understand the process of fertilisation, extend the value of ovaries which contain thousands of oocytes which would otherwise be wasted under natural circumstances, and allow gene manipulation through which desirable characteristics could be amplified or new characteristics introduced.

In vitro techniques for sperm capacitation, oocyte maturation (IVM), fertilisation (IVF) and culture (IVC) systems could be used to produce a large numbers of embryos at the same stage of development. Reliable techniques for producing developmentally competent goat embryos *in vitro* using slaughter house ovaries would solve the major problem of non-availability of genetically superior breeding materials.

Though *in vitro* techniques have been practised for over two decades, they need refinement of sperm capacitation agents, their optimum concentration, and the incubation periods required to achieve maximum oocyte penetration rate. Suitable *in vitro* culture media are also lacking for embryos to develop to the blastocyst stage.

Among the sperm capacitation agents, heparin is most widely used. It is a synthetic compound and belongs to a group called glycoaminoglycans (GAG). Recently, a natural substance isolated from marine red algae found in coastal areas of Sri Lanka has been proved to possess human sperm motility activation property. This substance, named as S-ACT-1, has a structure quite similar to that of heparin. Whether this motility activating property could also induce capacitation of sperm is not known.

Therefore a series of experiments was conducted in Kandy, Sri Lanka to understand the process of sperm capacitation, the acrosome reaction and *in vitro* maturation, fertilisation and culture of oocytes. S-ACT-1 was tested against heparin as the capacitation agent. The goat was selected as the model animal mainly because of the availability of goat ovaries from the local slaughterhouse.

The objectives of this study were

1. to compare heparin and S-ACT-1 as agents for capacitation of goat sperm
2. to develop the IVF technique for production of embryos from oocytes recovered from goat ovaries collected at slaughterhouse.

The following experiments were conducted:

Experiment 1 Capacitation and acrosome reaction of goat spermatozoa (Chapter III)

Experiment 2 Assessment of sperm capacitation in goat (Chapter IV)

Experiment 3 *In vitro* oocyte maturation, fertilisation and culture of goat embryos (Chapter V)

CHAPTER II

LITERATURE REVIEW

Introduction

This review is divided into two parts. The first part reviews some fundamental process on the male and female gametes, their development and events leading to fertilisation *in vivo* and embryonic development. The second part reviews studies on *in vitro* fertilisation (IVF) in farm animals.

***In Vivo* Fertilisation**

Understanding how spermatozoa achieve fertilisation *in vivo* is of utmost importance to solve many problems associated with interaction between oocytes and spermatozoa *in vitro*. Fertilisation in mammals requires three critical events: sperm migration through cumulus cells, sperm attachment and migration through the ZP and sperm fusion with the ooplasm (Hafez, 1983).

Spermatogenesis

The seminiferous epithelium of the testis contains a series of developing germ cells which will be finally converted to spermatozoa (Hafez, 1993).



Spermatogenesis involves two processes: *spermatocytogenesis* and *spermiogenesis*. In *spermatocytogenesis*, spermatogonia situated at the periphery of the seminiferous epithelium divide several times to form spermatocytes which after undergoing meiosis to become spermatids. In *spermiogenesis*, spermatids contain haploid number of chromosomes and after a series of morphological and developmental changes, become spermatozoa, which are elongated in shape. All the developing germinal cells are closely associated with the Sertoli cells and with further development, sperm move from the basement membrane toward the lumen. The process of release of spermatozoa into the lumen is known as spermiation. Semen is the semi-gelatinous liquid containing sperm and secretions of the accessory organs of the male reproductive tract.

Maturation

In mammals, maturation of spermatozoon primarily occurs in the caput and corpus epididymis (Bedford and Hoskin, 1990). It is a complex and sequential process (Parks and Hammerstedt, 1985). Spermatozoon's maturation changes depend on the epididymal secretions and transport time through the epididymis. In the ram, the transport time in the epididymis is 16 days (Amann, 1981). At ejaculation, although spermatozoa show their normal morphologic characteristics and motility, they are still unable to penetrate and fertilise the ova (Burks and Saling, 1992).

The final stage of sperm maturation is completed in the female reproductive tract. It involves biochemical, physiological and some morphological changes just before fertilisation. The changes, which take place in the female reproductive tract will be reviewed in detail later.

During transit along the male tract, sperm will undergo some remodelling of membrane lipids (phospholipid) (Parks and Hammerstedt, 1985). Sperm cell proteins undergo changes during maturation. Modification of DNA protein complex, plasma membrane, mitochondria, fibrous and microtubular components and development of surface characteristics have been observed (Amann et al., 1993). Migration of some macromolecules along the cell surface has also been observed.

Maturation of spermatozoon is not completed until they acquire motility, fertilisability and viability. The spermatozoon attains progressive motility during passage through the epididymis (Bedford, 1975). With increase in progressive motility, the water content of the spermatozoon decreases and as a result the specific gravity increases (Hafez, 1983).

Morphology of the Spermatozoon

The general morphology of mammalian spermatozoon is similar (Hafez, 1993). But detailed sperm ultra-structure is considerably different from species to species. Basically, the spermatozoon consists of two main parts: the head and the tail. Average length of a mammalian spermatozoon is about 60 μm and the head is about 40 μm long, 4 μm wide and 0.5 μm thick. The nucleus which contains DNA and the tail which contains the apparatus necessary for cell motility mainly occupy the head. The tail can further be subdivided into the neck, the middle piece, the principal piece and the end piece. Shape of the head varies greatly from species to species. In most of mammals including goat, the head has a flattened pyriform shape. The nuclear chromatin is highly condensed and exhibits a rather uniform thickness except at the anterior end where it tapers gently and thickening slightly at the base.

The anterior two third of the head is covered with another structure known as the acrosomal cap. The acrosomal cap consists of three layers; the outer membrane, the inner membrane and a thicker, moderately electron-dense middle layer. The inner and outer acrosomal membranes are continuous with each other at the posterior margin of the cap forming an enclosed space. This space is filled with acrosomal material mainly proteolytic enzymes such as hyaluronidase, proacrosin, acrosin, proteases, esterases, and acid hydrolases. These enzymes are involved in the fertilisation process (Mann and Lutwak-Mann, 1981). The outer acrosomal membrane is closely associated with the plasmalemma or cell membrane, which covers the entire length of the sperm.

Along the anterior margin of the sperm head, the acrosomal cap is somewhat thickened and bulges out reaching its maximum thickness at the apex. This structure is known as the apical ridge. This is the widest part of the acrosome. The acrosomal content is nearly uniform and has a moderate density. The posterior portion of the acrosome is termed as the equatorial segment. This structure is important because this is the part of the spermatozoon, which initially fuses with the oocyte during fertilisation.

The area posterior to the equatorial segment consists of the plasmalemma and the nuclear membrane and is known as the post nuclear cap or post acrosomal cap. Anterior end of the post-nuclear cap is attached to the outer membrane of the equatorial segment. At the base of the sperm head, the post nuclear cap is closely in contact with the plasmalemma. The two nuclear membranes fuse together at the base of the head and closely contact with the nuclear material.