



UNIVERSITI PUTRA MALAYSIA

**ELEMENTAL ASSESSMENTS OF SOUTH CHINA SEA MARINE
SEDIMENTS ALONG THE EAST COAST OF PENINSULAR
MALAYSIA**

**KHADIJEH REZAE EBRAHIM SARAEE
FS 2010 25**

**ELEMENTAL ASSESSMENTS OF SOUTH CHINA SEA MARINE
SEDIMENTS ALONG THE EAST COAST OF PENINSULAR MALAYSIA**

KHADIJEH REZAE EBRAHIM SARAEE

**DOCTOR OF PHILOSOPHY
UNIVERSITI PUTRA MALAYSIA**

2010



**ELEMENTAL ASSESSMENTS OF SOUTH CHINA SEA MARINE
SEDIMENTS ALONG THE EAST COAST OF PENINSULAR MALAYSIA**

By

KHADIJEH REZAE EBRAHIM SARAEE

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia,
in Fulfilment of the Partial Requirements for the Degree of Doctor of
Philosophy**

February 2010



DEDICATION

This thesis is dedicated to:

“My Dear Lavin”



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment
of the requirement for the degree of Doctor of Philosophy

**ELEMENTAL ASSESSMENTS OF SOUTH CHINA SEA MARINE
SEDIMENTS ALONG THE EAST COAST OF PENINSULAR MALAYSIA**

By

KHADIJEH REZAE EBRAHIM SARAEE

November 2009

Chair: Professor Elias Saion, PhD

Faculty: Science

Malaysia is located in the west of the South China Sea and is divided into two major land masses, i.e., Peninsular Malaysia and East Malaysia on the Island of Borneo, with a total coast line of 4675 km (Peninsular Malaysia 2068 km; East Malaysia 2607 km). As most seas, the majority of pollution occurs in the South China Sea arise from the land namely waste that comes from large cities (sewage, industrial waste and hydrocarbons) and agricultural runoff (nutrients, pesticides and fertilizers). The aim of this research is to access information on horizontal elemental distribution of coastal surface sediments and on vertical elemental distribution of offshore core sediments and to determine the status of pollution in the surface and core marine sediments. The results of this research together with experimental data from other



researches on marine sediments on the east coast of Peninsular Malaysia may be used to establish a baseline data for this region.

All together 43 elements were identified and then classified according to heavy metals (As, Cd, Cu, Cr, Ni, Pb and Zn), trace elements (B, Ba, Be, Bi, Br Co, Cs, Ga, Ge, Hf, Li, Mo, Nb, Sc, Rb, Sb, Sn, Sr, Ta, V, W, Y and Zr), rear earth elements (Ce, Dy, Eu, La, Lu, Nd, Sb, Sm and Yb), major elements (Al, Ca, K, Fe, Mg, Mn Na, P, S and Ti), and actinides elements (Th and U) depending on their physical and chemical characteristics. The elemental concentrations of the surface sediments from 10 stations and the core sediments from 5 stations have been determined using the Instrumental Nuutron Activation Analysis (INAA) at Nuclear Agency Malaysia and the Inductively Coupled Plasma-Atomic Emission Spectroscopy (ICP-AES) at Geological Survey of Iran (GSI) laboratories. To know the status of contamination in heavy metals and trace elements, the enrichment factor, geoaccumulation index (I_{geo}) and modified degree of contamination (mC_d) methods were used to interpret the results. Where appropriate the linear regression analysis, pearson's correlation factors, first and second of the principal component analysis and the cluster analysis were performed to determine the relationships between obtained results.

For the surface elemental distributions, heavy metal Pb was found to be originated from anthropogenic sources at the stations at the larger river mouths of Kelantan, Pahang and Rompin rivers, possibly due to land-based anthropogenic activities, automotive emission and gas industries. However, the average I_{geo} and mC_d values indicate that the heavy metals of the surface sediments are uncontaminated in all sampling stations. The same situation was true for trace elements, except for Hf

which has the enrichment factor greater than 2 in EC8 station. The high concentration of Hf at this station was a natural occurring because of it has negative correlation with Zn in this station similarity with those in nature. Shale averages and chondrite normalized were used to normalize rare earth elements (REEs). Ce was found anthropogenic element, possibly due to occurrence of +4 oxidation states, while other REEs are of non-anthropogenic sources. Most REEs have lower concentrations at the larger river systems and Ce and Eu anomalies occurred in samples taken from the mouths of Kelantan and Pahang rivers. For major elements, Al, Ca, Mg, Na, and Ti were of anthropogenic sources, possibility due to tropical weathering. Both actinides Th and U were non-anthropogenic for all sampling stations except for Kuala Dungun possibility due to a radioactive related factory like among in that area.

For the vertical element distribution, heavy metals As, Cd and Pb; trace elements Br, Hf, Sr and Zr; major elements Ca and Na and actinide element Th were of anthropogenic sources. However, by considering the average I_{geo} and mC_d values, most of the layers in the east coast of Peninsular Malaysia were uncontaminated. Only Hf was the most polluted in affected stations and were classified as moderately to stronger contamination. The chondrite-normalized ratios of REEs showed LREEs enrichment and flat HREE depletion. No Ce anomaly was observed in sediments of all layers in all the stations. Overall, the results showed that the normalized patterns of REEs of the core sediments followed a general pattern of REEs of the world.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**ANALISIS UNSUR ENDAPAN LAUT DARI LAUT CHINA SELATAN
SEPANJANG PESISIRAN PANTAI TIMUR SEMENJUNG MALAYSIA**

Oleh

KHADIJEH REZAE EBRAHIM SARAEE

February 2010

Pengerusi: Profesor Elias Saion, PhD

Fakulti: Sains

Malaysia terletak di bahagian barat Laut China Selatan dan ia boleh dibahagikan kepada dua kawasan utama iaitu Semenanjung Malaysia dan Malaysia Timur di Pulau Borneo dengan panjang pantai 4675 km (2068 km di Semenanjung; 2607 km di Malaysia Timur). Seperti berlaku di kebanyakan laut, pencemaran di Laut China Selatan adalah bersumber daripada tanah dataran oleh sisa buangan daripada kotaraya (air kumbahan, sisa industri dan hidrokarbon) dan sisa pertanian (nutrien, racun serangga dan baja). Tujuan kajian ini adalah untuk mendapatkan maklumat tentang taburan mendatar unsur daripada endapan laut permukaan dan taburan mencancang daripada endapan laut teras dan seterusnya menentukan status pencemaran permukaan dan teras endapan laut. Keputusan daripada kajian ini bersama-sama dengan data eksperimen daripada kajian-kajian lain mengenai

endapan laut di pantai timur Semenanjung Malaysia boleh digunakan sebagai membena pengkalan data untuk kawasan ini.

Keseluruhannya 43 unsur telah dikenalpastikan dan dikelaskan menurut sebagai logam berat (As, Cd, Cu, Cr, Ni, Pb and Zn), unsur surih (B, Ba, Be, Bi, Br Co, Cs, Ga, Ge, Hf, Li, Mo, Nb, Sc, Rb, Sb, Sn, Sr, Ta, V, W, Y dan Zr), unsure nadir bumi (Ce, Dy, Eu, La, Lu, Nd, Sb, Sm dan Yb), unsur utama (Al, Ca, K, Fe, Mg, Mn Na, P, S dan Ti) dan unsur aktinides (Th dan U) yang ditentukan bergantung kepada ciri-ciri fizik dan kimia. Kepekatan unsur endapan laut permukaan daripada 10 station dan endapan laut teras daripada 5 station telah ditentukan dengan menggunakan Analisis Instrumentasi Pengaktifan Nuklear (INAA) di Agensi Nuklear Malaysia dan menggunakan Aruhan Terganding Plasma-Spektroskopi Penyinaran Atom (ICP-AES) di makmal Geological Survey of Iran (GSI). Untuk mengetahui tahap pencemaran logam berat dan unsur surih, kaedah faktor pengkayaan, indek pengumpulan geologi (I_{geo}) dan darjah pencemaran terubahsuai (mC_d) telah digunakan untuk mentafsirkan keputusan. Dimana-mana yang sesuai analisis regresi liner, faktor korelasi pearson, analisis dua komponen pertama dan analisis guggusan telah dilakukan untuk menentukan hubungan di antara keputusan-keputusan yang diperolehi.

Untuk taburan di permukaan unsur logam berat Pb adalah bersumber antropogen di station bertentangan dengan muara sungai besar iaitu sungai Kelantan, Pahang dan Rompin, kerana banyak aktiviti antropogen di dataran. Walau bagaimanapun, purata nilai I_{geo} and mC_d menunjukkan bahawa endapan di permukaan tidak mengalami pencemaran daripada logam berat di semua station. Keadaan yang sama juga berlaku kepada unsur surih, kecuali Hf yang didapati mempunyai faktor pengkayaan

melebihi 2 di station EC8. Kepekatan tinggi pada Hf di station tersebut berlaku secara semulajadi kerana ia mempunyai kolorasi negative terhadap Zn di station itu sama terdapat pada semulajadi. Purata nilai ternormal terhadap shale dan chondorite telah digunakan untuk unsur nadir bumi. Didapati bahawa Ce adalah antropogen kerana berkemungkinan terdapat keadaan pengoksidan +4 unsur tersebut, sedangkan unsur nadir bumi lain tidak antropogen. Kebanyakan unsur nadir bumi mempunyai kepekatan rendah dalam sistem sungai besar dan anormali Ce dan Eu hanya berlaku pada sampel bertentangan muara sungai Kelantan dan Pahang. Unsur utama Al, Ca, Mg, Na, dan Ti adalah antropogen, berkemungkinan kerana keadaan luluhhawa tropika. Kedua aktinides Th dan U adalah tidak antropogen di semua station kecuali di Kuala Dungun berkemungkinan daripada kilang berkaitan bahan radioaktif seperti among di kawasan ini.

Untuk taburan unsur seranjang logam berat As, Cd dan Pb; unsur surih Br, Hf, Sr dan Zr; unsur major Ca dan Na dan unsur aktinide Th didapati adalah antropogen. Unsur-unsur lain adalah tidak antropogen. Bagaimanapun, dengan mengambil purata nilai I_{geo} and mC_d kebanyakan lapisan sampel teras bagi pantai timur Semenanjung Malaysia sebenarnya tidak mangalami pencemaran. Kecuali Hf adalah unsur paling tercemar di station yang terlibat dan dikelaskan di antara pencemaran serdahana dan pencemaran berat. Untuk unsur nadir bumi nisbah ternormal terhadap chondrite menunjukkan pengkayaan unsur nadir bumi ringan dan pengurangan unsur nadir bumi berat. Tiada anormali Ce terdapat pada semua lapisan sampel endapan teras di semua station. Keseluruhannya daripada keputusan kajian ini menunjukkan corak unsur nadir bumi bagi endapan teras adalah sama seperti ditempat-tempat lain di dunia.

ACKNOWLEDGEMENTS

First and foremost, I would like to express my gratitude to my supervisor Prof. Dr. Elias. Saion for all his advice and support during the time I spent at the Universiti Putra Malaysia.

I would like to thank my committee members Prof. Dr. Abdul Halim B Shaari, Prof. Dr. Sidek B Hj AB Aziz, Dr. Abdul Khalik Haji Wood and Dr. Khoo Kok Siong for their constructive comments and criticisms.

Thanks are expressed to the Nuclear Agency Malaysia for their continues supporting and the usage of their facilities. I am very much thankful and especial thanks to Mr. Md Suhaimi Elias for his help in collecting data in the laboratories. Especial thanks to the staff of counting room of research reactor for their help.

I would like to extend my great thanks to the staff of the Department of Physics, Universiti Putra Malaysia.

Special thanks and deep appreciation goes to Dr. Mohammad Reza Abdi for his valued advice and help.

I would like to express my deepest gratitude to my sister, parents and brothers and engineer Dokouhaki for their prayers and unending encouragement.



Last but not least, to my sweet heart, my dearest in my life, my daughter Lavin for creating real love in my heart and encouraging me to do my study.



I certify that a Thesis Examination Committee has met on 11 February 2010 to conduct the final examination of Khadijeh Rezaee Ebrahim Saraee on her Doctor of Philosophy thesis entitled "Elemental Assessments of South China Sea Marine Sediments Along the East Coast of Peninsular Malaysia" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student Khadijeh Rezaee Ebrahim Saraee.

Members of the Thesis Examination Committee were as follows:

Halimah Mohamed Kamari, PhD

Lecturer
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Zainal Abidin Sulaiman, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Yap Chee Kong, PhD

Associate Professor
Faculty of Science
Universiti Putra Malaysia
(Internal Examiner)

Datuk. Sukiman Suliman, PhD

Professor
Faculty of Science
Universiti Kebangsaan Malaysia
(External Examiner)



BUJANG BIN KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 April 2010



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Elias Saion, PhD

Professor
Faculty of Science
Universiti Putra Malaysia
(Chairman)

Abdul Halim B Shaari

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Sidek B HJ AB Aziz

Professor
Faculty of Science
Universiti Putra Malaysia
(Member)

Khoo Kok Siong

Nucleas Science Programme School of Applied Physics
Universiti Kebangsaan Malaysia
(Member)

Abdul Khalik Hj Wood, PhD

Agensi Nuklear Malaysia
(Nuklear Malaysia)
Malaysia
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 13 May 2010



DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

Rezaee 

KHADIJEH REZAAE EBRAHIM SARAEE

Date: 10 February 2010

TABLE OF CONTENTS

| | Page |
|------------------------------|--------------|
| DEDICATION | i |
| ABSTRACT | ii |
| ABSTRAK | v |
| ACKNOWLEDGEMENTS | viii |
| APPROVAL | x |
| DECLARATION | xii |
| LIST OF TABLES | xvii |
| LIST OF FIGURES | xxiii |
| LIST OF ABBREVIATIONS | xxvi |

CHAPTER

| | | |
|----------|--|----|
| 1 | INTRODUCTION | |
| | 1.1 General Introduction | 1 |
| | 1.2 Sources of Contamination and Pollution in Marine Sediments | 4 |
| | 1.2.1 Heavy Metals | 5 |
| | 1.2.2 Major elements | 6 |
| | 1.2.3 Trace Elements | 7 |
| | 1.2.4 Rare Earth Elements | 8 |
| | 1.2.5 Actinides | 9 |
| | 1.3 Dification of the Study Area | 10 |
| | 1.4 Significant of the Study | 12 |
| | 1.5 Problem Statement | 13 |
| | 1.6 Scope of the Study | 14 |
| | 1.7 Study Objectives | 15 |
| | 1.8 Outline | 16 |
| 2 | LITERATURE REVIEW | |
| | 2.1 Introduction | 17 |
| | 2.1.1 REEs Normalized in Chondritic Meteorites | 17 |
| | 2.1.2 REEs Normalized in Shale Materials | 22 |
| | 2.2 Marine Sediment Studies in the Coastal of Malaysia | 24 |
| | 2.2.1 Heavy Metals | 25 |
| | 2.2.2 Trace Elements | 33 |
| | 2.2.3 Rare Earth Elements | 36 |
| 3 | THEORY | |
| | 3.1 Introduction | 42 |



| | | |
|----------|---|-----|
| 3.2 | Neutron Energy Classification | 42 |
| 3.3 | Neutron Interaction | 44 |
| 3.3.1 | Neutron Cross Section | 48 |
| 3.4 | Neutron Sources | 50 |
| 3.4.1 | Triga Mark II Nuclear Research Reactor | 52 |
| 3.5 | Neutron Activation Analysis | 55 |
| 3.5.1 | Principle of NAA | 57 |
| 3.5.2 | INAA-Comparative Method | 59 |
| 3.6 | Semiconductor Detectors | 62 |
| 3.6.1 | HPGe Detector for Gamma Rays | 63 |
| 3.6.2 | Interaction of Gamma Rays with Matter | 64 |
| 3.6.3 | Gamma Ray Spectroscopy | 67 |
| 3.7 | Principle of Inductively Coupled Plasma | 70 |
| 3.7.1 | Interferences | 73 |
| 4 | MATERIALS AND METHODS | |
| 4.1 | Introduction | 74 |
| 4.2 | Standard Reference Material | 75 |
| 4.3 | Marine Sediments Sampling | 76 |
| 4.3.1 | Location of Sampling | 76 |
| 4.3.2 | Collection of Surface Sediment Samples | 79 |
| 4.3.3 | Collection of Core Sediment Samples | 80 |
| 4.3.4 | Preparation of Sediment Samples for INAA | 81 |
| 4.3.5 | Irradiation of Sediment Samples for INAA | 82 |
| 4.3.6 | Preparation of Standard Solutions | 83 |
| 4.3.7 | Sample Digestion for ICP-AES Equipment | 86 |
| 4.3.8 | ICP-AES Elemental Analysis | 87 |
| 4.4 | Measurement of Gamma Rays | 88 |
| 4.4.1 | Energy Resolution of Ge Detector | 89 |
| 4.4.2 | Detector Efficiency | 91 |
| 4.4.3 | Setup and Settings | 92 |
| 4.4.4 | Analysis of HPGe Detector Energy Spectra | 93 |
| 4.4.5 | Lower Limit of Detection | 95 |
| 4.5 | Photo Peak Selection | 96 |
| 4.6 | Statistical Analysis | 100 |
| 4.6.1 | Hierarchical Clustering Algorithms | 100 |
| 4.6.2 | Principle Component Analysis | 101 |
| 4.6.3 | Pearson Correlation Analysis | 102 |
| 4.6.4 | Post Hoc and One-Way ANOVA | 102 |
| 4.7 | Methods for Estimating Pollutant Impact | 103 |
| 4.7.1 | Enrichment Factor (EF) | 104 |
| 4.7.2 | Geoaccumulation Index (I_{geo}) | 105 |
| 4.7.3 | Modified Degree of Contamination (mC_d) | 106 |
| 5 | RESULTS AND DISCUSSION | 109 |
| 5.1 | Introduction | 110 |
| 5.2 | Quality Assurance and Quality Control | 111 |
| 5.2.1 | Validity of Results | |



| | | |
|-------|--|-----|
| 5.2.2 | Limitation of Detection | 115 |
| 5.3 | Heavy Metals in the Surface Sediments of the East Coast Peninsular Malaysia | 116 |
| 5.3.1 | Variation of Heavy Metals in the Surface Sediments | 117 |
| 5.3.2 | Enrichment Factor | 120 |
| 5.3.3 | Assessment Results Using Geoaccumulation Index and Modified Degree of Contamination | 123 |
| 5.3.4 | Comparison of Heavy Metal Concentrations | 125 |
| 5.3.5 | Classification of the Stations Using Hierarchical Cluster | 128 |
| 5.3.6 | Principal Component Analysis | 130 |
| 5.3.7 | Pearson's Correlation between Heavy Metal Concentrations | 133 |
| 5.3.8 | Comparison Mean and Range of Heavy Metal Concentrations in the Surface Sediments of the East Coast of Peninsular Malaysia with those in other Coasts | 134 |
| 5.4 | Heavy Metals in the Core sediments of the East Coast of Peninsular Malaysia | 135 |
| 5.4.1 | Enrichment Factor (<i>EF</i>) | 140 |
| 5.4.2 | Variation of Heavy Metals in the Core Sediment | 144 |
| 5.4.3 | Statistical Analyses | 145 |
| 5.4.4 | Comparison Heavy Metals in the Surface and Core Sediments with the International Guidelines | 152 |
| 5.5 | Trace Elements in the Surface Sediments of the East Coast of Peninsular Malaysia | 154 |
| 5.5.1 | Enrichment Factor | 158 |
| 5.5.2 | Assessment Results using Geoaccumulation Index and Modified Degree of Contamination | 160 |
| 5.5.3 | Comparison of Trace Elements Concentrations in the Surface Sediment | 163 |
| 5.5.4 | Cluster Analysis | 171 |
| 5.5.5 | Principal Component Analysis | 176 |
| 5.5.6 | Pearson Correlation between Trace Element Concentrations | 178 |
| 5.5.7 | Comparison the Trace Element Concentrations of in the Surface Sediments with those in the other Coasts of Malaysia | 180 |
| 5.6 | Trace Elements in the Core Sediments of the East Coast of Peninsular Malaysia | 182 |
| 5.6.1 | Total Trace Element Concentrations | 182 |
| 5.6.2 | Methods for Estimating Pollutant Impact | 188 |
| 5.7 | REEs in the Surface Sediments of the East Coast of Peninsular Malaysia | 193 |
| 5.7.1 | Enrichment Factor | 199 |
| 5.7.2 | Chondrite-Normalized REE Pattern | 201 |
| 5.7.3 | The Shale-Normalized REE Pattern | 202 |
| 5.7.4 | Comparison the Mean Concentration of REEs in the Surface Sediments with that Those in the Other | 205 |



| | | |
|----------|---|-----|
| | Coasts of Malaysia | |
| 5.8 | REEs in the Core Sediment of the East Coast of the Peninsular Malaysia | 207 |
| 5.8.1 | Enrichment Factor | 215 |
| 5.8.2 | The Shale Normalized REE Pattern | 217 |
| 5.8.3 | Chondrite - Normalized REE Pattern | 219 |
| 5.8.4 | Comparison the Mean Concentration of REEs in the Surface Sediment of the East Coast of Peninsular Malaysia with Those in the Other Coasts | 220 |
| 5.9 | Actinides in the Surface Sediment along the East Coast of Peninsular Malaysia | 222 |
| 5.10 | Actinides in the Core Marine Sediment along the East Coast of Peninsular Malaysia | 224 |
| 5.11 | Major Elements in the Surface Sediments of the East Coast of Peninsular Malaysia | 229 |
| 5.11.1 | Enrichment Factor | 235 |
| 5.11.2 | Cluster Analysis | 236 |
| 5.11.3 | Principal Components Analysis | 237 |
| 5.11.4 | Pearson Correlation between Major Element Concentrations in the Surface Sediments of the East Coast of Peninsular Malaysia | 239 |
| 5.11.5 | Comparison the Mean Concentration of Major Elements in the Surface Sediment of the East Coast of Peninsular Malaysia with Those in the Other Coasts | 241 |
| 5.12 | Major Elements in the Core Marine Sediments along the East Coast of Peninsular Malaysia | 242 |
| 5.12.1 | Enrichment Factor | 244 |
| 6 | CONCLUSION | |
| 6.1 | Conclusion and Future Works | 246 |
| 6.2 | Future Works and Recommendations | 249 |
| | REFERENCES | 250 |
| | APPENDICES | 268 |
| | BIODATA OF STUDENT | 289 |

LIST OF TABLES

| Table | | Page |
|-------|--|------|
| 2.1 | Concentration values (mg/kg dry weight) of rare earth elements in mg/kg | 19 |
| 2.2 | Comparison between the mean concentrations (mg/kg dry weight except for Al, Fe and Ti %) in the grab and core sediments of Penang Island (Data sited from Wood, et al. 2004) | 32 |
| 2.3 | Range and mean concentrations (mg/kg dry weight except for Al, Fe and Ti %) Trace elements of grab samples sediments of the Johor Straits (Data sited from Wood et al., 1997) | 34 |
| 2.4 | Elemental concentration (mg/kg dry weight except for Al, Ca, Fe K, Mg and Ti %) ranges and means of trace elements in the grab and core sediments of the Malacca straits (Data sited from Al-Zahrany et al., 2007) | 35 |
| 2.5 | Range and mean concentrations (mg/kg dry weight) of REEs in the grab samples sediments of the Johor Straits (Data sited from Wood et al., 1997) | 37 |
| 2.6 | Comparison concentration means (mg/kg dry weight) in the grab and core sediments of the Penang Island (Data sited from Wood, et al. 2004) | 38 |
| 2.7 | Range and mean concentrations (mg/kg dry weight) of REEs in the grab and core sediments of the Malacca straits (Data sited from Al-Zahrany et al., 2007) | 39 |
| 2.8 | Some literatures related to elemental distribution of different coasts of Malaysia | 41 |
| 4.1 | The longitude and latitude of sampling sites | 77 |
| 4.2 | The irradiation, cooling and counting times for first and second counting of radioactivity measurements, and the determined isotopes for each cycle by comparative method | 83 |
| 4.3 | Standard sources used for short-lived radionuclide | 85 |
| 4.4 | Standard sources used for medium-lived radionuclide | 85 |
| 4.5 | Standard sources used for long-lived radionuclide | 86 |



| | | |
|------|--|-----|
| 4.6 | Gamma-Ray energy lines used for analysis, half-life, and intensity of gamma-ray energy, nuclear reaction and thermal cross section of reaction for short-lived radio nuclides (Browne, E., et. al., 1978; IAEA-TCS-4, 1992; Vertes, A., et. al., 2003a) | 97 |
| 4.7 | Gamma-Ray energy lines used for analysis, half-life, and intensity of gamma-ray energy, nuclear reaction and thermal cross section of reaction for medium-lived radio nuclides (Browne, E., et. al., 1978; IAEA-TCS-4, 1992; Vertes, A., et. al., 2003a) | 98 |
| 4.8 | Gamma-Ray energy lines used for analysis, half-life, and intensity of gamma-ray energy, nuclear reaction and thermal cross section of reaction for long-lived radio nuclides (Browne, E., et. al., 1978; IAEA-TCS-4, 1992; Vertes, A., et. al., 2003a) | 99 |
| 4.9 | Muller's classification for geoaccumulation index | 106 |
| 4.10 | Hakanson (1980) classification of the modified degree of contamination (Abraham et al., 2007) | 108 |
| 5.1 | Comparison of determined concentrations of elements in the IAEA-Soil-7 by two INAA and ICP-AES methods with the certified values | 112 |
| 5.2 | Comparison of determined concentrations of elements in the IAEA-SL-1 by INAA method with the certified values | 114 |
| 5.3 | Limitation of detection (LOD) for each element in mg/kg | 115 |
| 5.4 | Heavy metal concentrations, in mg/kg, measured in the surface sediments of the east coast of Peninsular Malaysia | 116 |
| 5.5 | Enrichment factors (EF) of heavy metals in the surface sediments of the east coast of peninsular Malaysia, normalized with respected to the iron content in the continental shales | 121 |
| 5.6 | Enrichment factors (EF) of heavy metals in the surface sediments of the east coast of peninsular Malaysia, normalized with respected to the iron content in the EC8 station | 122 |
| 5.7 | Contamination factors, modified degree of contamination (mC_d) using average shale baseline values for heavy metals in the surface sediments of the east coast of Peninsular Malaysia | 123 |
| 5.8 | Index of geoaccumulation (I_{geo}) of heavy metals in the surface sediments compared to the average shale baseline values | 124 |
| 5.9 | Separations of significant mean As concentrations (mg/kg) in the | 126 |



| | | |
|------|--|-----|
| | sediment samples using LSD test (Least Significant Difference) | |
| 5.10 | Separations of significant mean Cu concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 127 |
| 5.11 | Separations of significant mean Pb concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 127 |
| 5.12 | Separations of significant mean Zn concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 127 |
| 5.13 | Correlation matrix (r) for heavy metals in the surface sediments (n=10) (significant values are expressed in bold) | 132 |
| 5.14 | Mean and range of heavy metal concentrations (mg/kg dry weight) in the surface sediments of the east coast of Peninsular Malaysia with those in Straits of Malacca, Straits of Johor, Pinang Island, average shales and crustal material | 134 |
| 5.15 | Heavy metal concentrations (mg/kg dry weight) measured in the core sediments of the east coast of Peninsular Malaysia | 136 |
| 5.16 | Enrichment factors (EF) of anthropogenic heavy metals in the core sediments of the east coast of peninsular Malaysia, normalized with respect to the iron content in the continental shales | 141 |
| 5.17 | Correlation matrix (r) for heavy metals in the core sediments (n=47) (significant values are expressed in bold) | 144 |
| 5.18 | Regression equations of Fe with heavy metals As, Cd, Cr, Cu, Hg, Ni, Pb and Zn | 148 |
| 5.19 | Regression equations of Al with heavy metals As, Cd, Cr, Cu, Hg, Ni, Pb and Zn | 149 |
| 5.20 | Mean and range of heavy metal concentrations in the core sediments of the east coast of Peninsular Malaysia with those in Straits of Malacca, Straits of Johor, Pinang Island, average shales and crustal material | 150 |
| 5.21 | Comparison between heavy metals in the studied area with Canadian Sediment Quality Guidelines for the Protection of Aquatic Life | 151 |
| 5.22 | Comparison between heavy metals in the studied area with Technical Guidance for Screening Contaminated Sediments (New York) | 152 |

| | | |
|------|---|-----|
| 5.23 | Comparison between heavy metals in the studied area with Australian and New Zealand Guidelines for Fresh and Marine Water Quality | 153 |
| 5.24 | Trace element concentrations (mg/kg dry weight) measured in the surface sediments of the east coast of Peninsular Malaysia | 155 |
| 5.25 | Enrichment factors (EF) of trace elements in the surface marine sediments of the east coast of peninsular Malaysia, normalized with respected to the iron content in the continental shales | 158 |
| 5.26 | Contamination factors, modified degree of contamination (mC_d) using average shale baseline values for trace elements in the surface sediments of the east coast of Peninsular Malaysia | 160 |
| 5.27 | Index of geoaccumulation (I_{geo}) of trace elements in the surface sediments compared to the average shale baseline values | 161 |
| 5.28 | Separations of significant mean Co concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 165 |
| 5.29 | Separations of significant mean Cs concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 165 |
| 5.30 | Separations of significant mean Ga concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 166 |
| 5.31 | Separations of significant mean Sc concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 166 |
| 5.32 | Separations of significant mean Li concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 167 |
| 5.33 | Separations of significant mean Nb concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 167 |
| 5.34 | Separations of significant mean Ta concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 168 |
| 5.35 | Separations of significant mean V concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 168 |
| 5.36 | Separations of significant mean W concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 169 |
| 5.37 | Separations of significant mean Zr concentrations (mg/kg) in the sediment samples using LSD test (Least Significant Difference) | 169 |
| 5.38 | Correlation matrix (r) for trace elements in the surface sediments | 178 |



(n=10)

| | | |
|------|---|-----|
| 5.39 | Mean and range of trace element concentrations (mg/ kg dry weight) in the surface sediments of the east coast of Peninsular Malaysia with those in Straits of Malacca, Straits of Johor, Pinang Island, average shales and crustal material | 180 |
| 5.40 | Trace element concentrations (mg/kg dry weight), measured in the core sediments of the east coast of Peninsular Malaysia | 182 |
| 5.41 | Enrichment factors (EF) of anthropogenic trace elements in the core sediments of the east coast of peninsular Malaysia, normalized with respected to the iron content in the continental shales | 190 |
| 5.42 | Modified degree of contamination using shale average baseline values for trace elements in core sediments from the east coast of Peninsular Malaysia | 191 |
| 5.43 | Total concentration (mg/kg) of \sum REE, \sum LREE, \sum HREE, and ratios of, La/Sm, La/Yb, Ce/La, Eu/Sm, Yb/Sm, and Ce/Ce* in the surface sediments of the east coast of Peninsular Malaysia and corresponding average values | 193 |
| 5.44 | Enrichment factors (EF) of REEs in the surface sediments of the east coast of peninsular Malaysia, normalized with respected to the scandium content in the continental shales | 199 |
| 5.45 | Mean and range of REE concentrations (mg/kg dry weight) in the surface sediments of the east coast of Peninsular Malaysia with those in Straits of Malacca, Straits of Johor, Pinang Island, average shales and crustal material | 205 |
| 5.46 | REE concentrations (mg/kg dry weight) measured in the core sediments of the east coast of Peninsular Malaysia | 207 |
| 5.47 | Total concentration (mg/kg dry weight) of \sum LREE, \sum HREE, and ratios of \sum LREE/ \sum HREE, Eu/Sm, Ce/La, La/Yb and Ce/Ce* in the core sediments and corresponding average values | 212 |
| 5.48 | Mean and range of REE concentrations (mg/kg dry weight) in the core sediments of the east coast of Peninsular Malaysia with those in Straits of Malacca, Straits of Johor, Pinang Island, average shales and crustal material | 220 |
| 5.49 | Total concentration (mg/kg) of U and Th, and ratios of (U/Al)/U/Al) _{Shale} , (Th/Al)/(Th/Al) _{Shale} and Th/U in the surface sediments corresponding average values | 222 |



| | | |
|------|---|-----|
| 5.50 | Mean and range of Th and U concentrations (mg/kg dry weight) in the surface sediments of the east coast of Peninsular Malaysia with those in Straits of Malacca, Straits of Johor, Pinang Island, average shales and crustal material | 223 |
| 5.51 | Total concentration of U and Th, and ratios of $(U/Al)/(U/Al)_{Shale}$, $(Th/Al)/(Th/Al)_{Shale}$ and Th/U in the core sediments corresponding average values | 225 |
| 5.52 | Major element concentrations, in mg/kg, measured in the surface sediments of the east coast of Peninsular Malaysia | 230 |
| 5.53 | Enrichment factors (EF) of major elements in the surface sediments of the east coast of peninsular Malaysia, normalized with respected to the iron content in the continental shales | 235 |
| 5.54 | Correlation matrix (r) for major elements in the surface sediments (n=10) | 239 |
| 5.55 | Mean and range of major element concentrations in the surface sediments of the east coast of Peninsular Malaysia with those in Straits of Malacca, Straits of Johor, Pinang Island, average shales and crustal material | 241 |
| 5.56 | Major element concentrations (mg/kg dry weight), measured in the core sediments of the east coast of Peninsular Malaysia | 242 |