

A Putative Proline-rich Protein of *B. napus*

Parameswari Namasivayam^{1,*} and David Hanke²

¹Department of Cell and Molecular Biology, Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia, 43400 UPM, Serdang, Selangor, Malaysia

²Department of Plant Sciences, University of Cambridge, Downing Street, Cambridge CB2 3EA, UK

*E-mail: parameswari@biotech.upm.edu.my

ABSTRACT

Proline-rich proteins are among the major protein components of plant cell walls. So far, two different proline-rich cell wall proteins have been described in *Brassica napus*. This paper reports a study on expression and sequence analysis of a novel class of a proline-rich putative protein, tentatively designated Ae4. The largest ORF of Ae4 encodes 166 amino acid residues without the start and stop codons. Ae4 is a partial length cDNA. The Ae4 gene expression was investigated and the results demonstrate that it accumulates in all vegetative tissues tested as well as in the embryogenic culture of *Brassica napus*. However, expression of Ae4 was undetectable in the non-embryogenic and cytokinin-treated embryogenic tissues. These results indicate that the Ae4 gene might play a role in somatic embryo formation.

Keywords: Proline-rich protein, PRP, *Brassica napus*, oilseed rape, somatic embryogenesis

INTRODUCTION

Proline-rich proteins (PRPs) are one class of structural cell wall protein members (Showalter, 1993). All the PRPs are characterised by the repeating occurrence of Pro-Pro repeats contained within a variety of larger repeat units. These proteins lack the SerPro4 repetitive element defined for extensins (Jose-Estanyol and Puigdomenech, 2000). The most extensively studied PRPs are from soybean that contains the PPVYK motif or variations (Hong *et al.*, 1987, 1989; Datta *et al.*, 1989; Wyatt *et al.*, 1992). Members of the PRP gene family were shown to be developmentally regulated and their expression tissue/organ specific (Hong *et al.*, 1989; Lindstrom and Vodkin 1991; Jose-Estanyol *et al.*, 1992). In general, PRPs are thought to have a structural role in the cell wall (Cassab and Varner, 1988). They have been also implicated in plant defence reactions (Chen and Varner, 1985; Ebener *et al.*, 1993), nodule morphogenesis (Franssen *et al.*, 1987; Wilson *et al.*, 1994), and are expressed during somatic embryogenesis (Aleith and Reichter, 1990; Gyorgyevy *et al.*, 1997; Yasuda *et al.*, 2001).

To date, two different proline-rich cell wall proteins have been described in *Brassica napus* (*B. napus*). In this paper, we report the sequence and expression analysis of an additional proline-rich putative protein, tentatively designated as Ae4. The amino acid sequence of Ae4 has been deduced from the nucleic acid sequence of a copy DNA (cDNA), isolated previously from a subtracted library of *B. napus* embryogenic culture (Namasivayam *et al.*, 2006b).

MATERIALS AND METHODS

Sources of Plant Materials

Plants of *Brassica napus* ssp. *oleifera* cv. Primor were grown from seeds, in pots with soil in the Botanic Garden, Cambridge. Sources and preparation of plant materials for the pre-embryogenic, mature embryogenic and non-embryogenic *Brassica napus* ssp. *oleifera* cv. Primor culture was as described in Namasivayam *et al.* (2006a,b). The cytokinin-treated embryogenic tissue was generated from hypocotyls of embryoids grown for 20 days on MS media containing 10^{-4} M kinetin, 2% (w/v) sucrose and

* Corresponding Author

0.8% (w/v) agar. Various organs/tissues such as young leaves, stem, buds, flowers, siliques, roots, stamens, carpels, petals and sepals from mature *B. napus* plants were harvested, immediately frozen in liquid nitrogen and stored at -80°C until isolation of total RNA.

Sequence Analysis

The *Ae4* cDNA sequence has been submitted to the GenBank under the accession number AY570239. Sequence analysis was carried out using BLAST 2.0 (Basic Local Alignment Search Tool; Altschul *et al.*, 1997), accessible from the internet (<http://www.ncbi.nlm.nih.gov/BLAST>). Alignments of the protein sequence with several closely related genes was carried out using the CLUSTAL W program from the Biology Workbench version 3.2, accessible from the internet (<http://biowb.sdsc.edu/CGI/BW.cgi>). Other sequence analyses were performed using Biology Workbench version 3.2 to compute molecular weight (MW), hydrophobicity and isoelectric point (pI) determination.

Total RNA Isolation

Total RNA from various frozen tissues/organs of the mature plant and tissue culture materials were extracted using the acid guanidium thiocyanate-phenol-chloroform extraction method described by Chomczynski & Sacchi (1987). The concentration of RNA in each sample was determined spectrophotometrically (Sambrook *et al.*, 1989).

Northern Blot Analysis

Equal amounts of total RNA (10 (g per lane) were resolved on 1.3% (w/v) agarose-formaldehyde denaturing gel and blotted onto HybondTM-XL nylon membrane (Amersham Biosciences). Hybridisation was carried out at 65°C using standard techniques (Sambrook *et al.*, 1989). The entire *Ae4* sequence was used as probe labelled with [³²P- α]-dCTP using the Prime-IT[®] II Random Primer Labeling Kit (Stratagene). Washes were carried out at room temperature in the first wash buffer (40 mM sodium phosphate pH 7.2, 1% (w/v) SDS and 1 mM EDTA) for 10 min and followed by second wash in 40 mM sodium phosphate pH 7.2, 5% (w/v) sodium dodecyl sulphate (SDS) and 1 mM ethylenediaminetetraacetic acid (EDTA) at 65°C for 15 min. The hybridisation signals were captured by a Phosphorimager Typhoon 8600

(Amersham Pharmacia Biotech). After removal of probe, the same blot was hybridised with radiolabelled Arabidopsis *Actin2/7* cDNA probe as a loading control.

RT-PCR

Equal amounts of DNase-treated total RNA (200 ng) from each tissue sample was added individually to a sterile 0.2 ml polymerase chain reaction (PCR) tube and the volume adjusted to 13.5 μl with DEPC-treated sterile deionised water (SDW). Oligo (dT₁₈) (1 μl of 20 pmoles/ μl) was added to the tube and the reaction mix incubated for 10 min at 70°C . Following brief centrifugation, the following reagents were added: 4 μl 5 x first strand buffer (Promega), 0.5 μl 'RNase Out' ribonuclease inhibitor (40 U/ μl) (Invitrogen), 0.5 μl 10 mM Bioline dNTPs mix and 0.5 μl MMLV-RT RNase H minus (200 U/ μl) (Promega) and incubated at 37°C for 1 h. Later, the reaction mix was heat deactivated before using for PCR reactions. PCR reactions were performed in 12.5 μl reactions with the following components: 2 μl of the RT product, 1 x Bioline PCR buffer (Mg²⁺- free), 1.5 mM MgCl₂, 0.4 mM dNTP mix, 2.5 pmoles of forward primer (5' GGACTATAAATTGGTGTGGAGGTTTCA 3') and reverse primer (5' TATTTATAGT CCTCCCGTAATGCCA - 3') respectively, and 1.5 U BioTaq DNA polymerase. An internal control was prepared using actin2 primers (forward primer: 5'-CCATTCTTGCTTCCCTCAG-3' and reverse primer: 5'(-GACGTAAGTAAAAACCCAG-3') and containing all the components as above to test for equal loading of the template. Also, a negative control without template was included. Amplification was performed as follows: 95°C for 3 min; followed by 35 cycles at 94°C for 30 s, 65°C for 1 min, 72°C for 1 min; and a final extension at 72°C for 3 min. The annealing temperature used for *Actin2* was 60°C . Reverse transcription polymerase chain reaction (RT-PCR) products were separated on a 2% (w/v) agarose gel and the gel was photographed. The agarose gel containing the PCR amplified products was blotted (Southern, 1975) and hybridised with a labelled specific probe (*Ae4* cDNA).

RESULTS

Sequence Analysis

The BNPE AE4 clone contains a cDNA insert of 678 bp, excluding a poly A tail. The longest open reading frame (ORF) encodes 166 amino

acids starting from nucleotide no. 3 and there is no start and stop codons, as shown in Fig. 1. Therefore, it is unlikely to be a full length clone. The predicted amino acid sequence is rich in proline (44.89 %), lysine (10.2 %), threonine (10.22 %), valine (8.89 %), serine (6.67 %), tyrosine (5.78 %) and glutamine (5.78 %). The predicted protein fragment represents a calculated molecular weight of 17.7 kDa and has a predicted pI value of 10.5. Hydropathy analysis indicated that 13 residues at the N terminal end of the predicted protein fragment are hydrophobic and the other regions are highly hydrophilic. The polypeptide is primarily composed of two repeat units: a 10-mer repeat unit (P P I/V K/M P P P V Q K/Q) and a 7-mer repeat unit (P P T P I/S/T Y S). Protein database search revealed that there is no significant similarity between the predicted amino acid sequence and protein sequences deposited in

the GenBank. However, comparison of the *Ae4* nucleotide sequence with nucleotide sequences in the GenBank showed that it is homologous (82% identical) to a genomic fragment of Arabidopsis in chromosome 2 (*At2g27380*) which encodes a putative proline-rich protein. Also, a few hits to *B. napus* seed EST sequences with a homology of 80% to 95% were found in the database. A comparison of the amino acid sequence of this clone with the Arabidopsis putative proline-rich protein and translated EST sequences is depicted in Fig. 2.

Expression Analysis of Ae4

The pattern of expression of *Ae4* in organs and tissues was investigated using Northern analysis and RT-PCR. The Northern analysis on the organ/tissue specific-blot failed to detect a distinct band for *Ae4* transcripts except for a very faint smear observed in each sample lane

```

3   caacaccaacttatagcctcctatcaaaccaccaccagtgcaaaagcctccaactccc   62
   Q H Q L I A L L S N H H P V Q K P P T P

63   acttatagtcctccgataaagccaccaccagtgacagaagcctccaacaccgacctacagt   122
   T Y S P P I K P P P V Q K P P T P T Y S

123  ccaccaggttaaaccaccaccagtgaaagcctccaacacctatttatagtcctccgtaaatg   182
   P P V K P P P V K P P T P I Y S P P V M

183  ccaccaccagtgcaacaacctccgacaccatcttatagtcctcctgtaaaaccaccacca   242
   P P P V Q Q P P T P S Y S P P V K P P P

243  gtgcaaaaacctccaacaccacttatagtcacctgttaaaccaccacctgtgcaaaag   302
   V Q K P P T P T Y S P P V K P P P V Q K

303  cctccaactccaacttacagtcctcctatcaaaccaccaccggtgcaaaaacctccaaca   362
   P P T P T Y S P P I K P P P V Q K P P T

363  ccaacttatagcctcctatcaaaccaccacctgtgcaaaagcctccgacgccacttat   422
   P T Y S P P I K P P P V Q K P P T P T Y

423  agtccacctgttaaaccaccaccggtccagaagcctccacaccaacttacagtcctcct   482
   S P P V K P P P V Q K P P T P T Y S P P

483  atcaaacacctccagtgaaacctccaacaccaatttatagtcggccagtgaaaccacca   542
   I K P P P V K P P T P I Y S P P V K P P

543  cccgtgcaaaagcctccaacccaacgtacagcccaccaattaaaccacctccagtaaaa   602
   P V Q K P P T P T Y S P P I K P P P V K

603  cctccgacaccaacttatagtcctcctgtaaaaccacctccagtgcaaaagcctccgacg   662
   P P T P T Y S P P V K P P P V Q K P P T

663  ccacttatagtccac   678
   P T Y S P

```

Fig. 1: Nucleotide and deduced amino acid sequence of clone *Ae4* (Genbank accession no. AY570239). The ORF is underlined. This is a truncated clone without the start and stop codons.

CD825663 Translated	-----	
CD830533 Translated	-----	
NP_180307	1 MRVPLIDFLRFLVLILSLSGASVAADATVKQNFNKYETDSGHAHPPPIYG	50
Ae4	-----	
CD825663 Translated	-----	
CD830533 Translated	-----	
NP_180307	51 APPSYTTPPPPIYSPPPIYPPPIQKPPPTYSPPPIYPPPIQKPPPTYSPPPIY	100
Ae4	-----	
CD825663 Translated	-----	
CD830533 Translated	-----	
NP_180307	101 PPPIQKPPPTYSPPPIYPPPIQKPPPTYSPPPIYPPPIQKPPPTPSYSPV	150
Ae4	-----	
CD825663 Translated	-----	
CD830533 Translated	-----	
NP_180307	151 KPPPVQMPPTYSPPPIKPPPVHKPPPTYSPPPIKPPPVHKPPPTIYSPPI	200
Ae4	-----	
CD825663 Translated	-----	
CD830533 Translated	-----	
NP_180307	201 KPPPVHKPPPTIYSPPIKPPPVHKPPPTYSPPVKPPPVHKPPPTIYSPPI	250
Ae4	-----	
CD825663 Translated	-----P	
CD830533 Translated	-----	
NP_180307	251 IKPPPVHKPPPTIYSPPVKPPVQTPPTIYSPPVKPPPVHKPPPTIYSP	300
Ae4	-----	
	1 -----QHQLIALLS-----	9
CD825663 Translated	1 PVKPPPVQKPPPTIYSPPVKPPPVQKPPPTIYSPPIKPPPVQKPPPTIYS	50
CD830533 Translated	1 --MPPPVQPPPTIYSPPVKPPPVQKPPPTIYSPPVKPPPVQKPPPTIYS	48
NP_180307	301 PVKSPPVQKPPPTIYSPPIKPPPVQKPPPTIYSPPIKPPPV--KPPPTIYS	350
Ae4	10 --NHHPVQKPPPTIYSPPIKPPPVQKPPPTIYSPPVKPPPV--KPPPTIYS	57
CD825663 Translated	51 PEIKPPPVQKPPPTIYSPPEIKPPPVQKPPPTIYSPPVKPPPV--NPPAPIY	99
CD830533 Translated	100 PEIKPPPVQKPPPTIYSPPEIKPPPVQKPPPTIYSPPVKPP-----	139
NP_180307	351 PEVKPPPVHKPPPTIYSPPEVKPPPVHKPPPTIYSPPVKPPPIQKPPPTTY	400
Ae4	58 PEVMPPPVQQPPTIYSPPEVKPPPVQKPPPTIYSPPVKPPPVQKPPPTTY	107
CD825663 Translated	100 SPPVKPPPVQPPPTPSYSP-----PVKPPPVQKPPPTTY	133
CD830533 Translated	140 -----PVQKPPPTTY	149

A Putative Proline-rich Protein of *B. napus*

NP_180307	401	SPPIKPPPPLQKPPTPTYSPPIKLPVVKPPTPIYSPVVKPPVHKPPTFIY	450
Ae4	108	SPPIKPPPVPQKPPTPTYSP-----PIKPPVQKPPTFIY	141
CD825663 Translated	134	SPPVKKPPPQKPPTPTYSPPIKPPPQ-----KPPTFIY	167
CD830533 Translated	150	SPPIKPPPV-KPPTPTYSPPIKPPPQ-----KPPTFIY	183
NP_180307	451	SPPVKKPPPVHKPPTPTYSPPIKPPPVKPPTPTYSPVQPPPQKPPTFIY	500
Ae4	142	SPPVKKPPPQKPPTPTYSPPIKPPPVKP-----PFIY	174
CD825663 Translated	168	SPIKPPPVKPPTPTYSPVKKPPPQKPPTPTYSPPIKPPVVKTSKTNL	218
CD830533 Translated	184	SPPV-----	187
BWB4372	501	SPPVKKPPIQKPPTPTYSPPIKPPPV-KPPTPTYSPPIKPPVHKPPT--	550
Ae4	175	SPPVKKPPPQKPPTPTYSPPIKPPPV-KPPTPTYSPVKKPPPQKPPT--	222
CD825663 Translated	219	PTYKATTQCNNL-----	230
CD830533 Translated		-----	
NP_180307	551	PTYSPPIKPPPPIHKPPTPTYSPPIKPPPVHKPPTPTYSPPIKPPPVHKPP	600
Ae4	223	PTYSP-----	227
CD825663 Translated		-----	
CD830533 Translated		-----	
NP_180307	601	TPTYSPPIKPPPVHKPPTPTYSPPIKPPPVHKPPTPTYSPPIKPPPVHKP	650
Ae4		-----	
CD825663 Translated		-----	
CD830533 Translated		-----	
NP_180307	651	PTPTYSPPIKPPPQKPPTPTYSPVKKPPPQLPPTPTYSPVKKPPPQV	700
Ae4		-----	
CD825663 Translated		-----	
CD830533 Translated		-----	
NP_180307	701	PPTPTYSPVKKPPPQVPTPTYSPPIKPPPQVPTPTTPSPQGGYGT	750
Ae4		-----	
CD825663 Translated		-----	
CD830533 Translated		-----	
NP_180307	751	PPPYAYLSHPIDIRN	761
Ae4		-----	

Fig. 2: Alignment of predicted amino acid sequence of *Ae4* with the *Arabidopsis* putative proline rich protein and translated EST sequences

Shaded sequences denote identical amino acids and gaps introduced in the alignment are marked with dashes. The amino acid sequences were obtained from GenBank: *Arabidopsis* At2g27380 (GenBank accession no. NP_180307), *Brassica seed* EST clone BN25061G23 (GenBank accession no. CD825663) and *Brassica napus* seed EST clone BN40045N17 (GenBank accession no. CD830533)

(Fig. 3A, top panel). However, hybridisation with the *Actin2/7* cDNA probe (Fig. 3A, bottom panel) showed clear signals, suggesting that the RNA was not degraded. To verify these results, a RT-PCR approach was employed using *Ae4* gene-specific primers and equal amount of cDNA from leaves, stems, buds, flowers, siliques, roots and carpels. After 30 amplification cycles, no product was visible on the ethidium bromide stained gel (Fig. 3C, top panel). Therefore, the gel was blotted and the RT-PCR gel blot was hybridised with a ³²P-labelled *Ae4* cDNA. An autoradiograph of this blot showed the presence of an approximately 370 bp band (as expected)

in all lanes with various intensities of hybridisation (Fig. 3C, bottom panel), suggesting differential expression of *Ae4* in different organs/tissues of the mature *B. napus* plant. Also, there was an additional faint band at approximately 600 bp in lane 2, suggesting the presence of another isoform of the *Ae4* gene or possibly an unspliced *Ae4* transcript.

Northern analysis of the tissue culture blot (Fig. 3B, top panel) detected expression of *Ae4* transcripts in the embryogenic culture, both in the pre-embryogenic (lane PEC) and mature embryogenic tissue (lane MEC). Relative to the actin control, expression in both these tissues




Fig. 3: Expression analysis of *Ae4* in *Brassica napus*

(A) Tissue/organ-specific and (B) tissue culture RNA gel blots containing 10 µg of total RNA per lane were first hybridised to ³²P-labelled *Ae4* and then to an *Arabidopsis Actin2/7* cDNA (control). Lanes: L, leaves; S, stem; B, buds; F, flowers; Si, siliques; R, roots; St, stamens; Ca, carpels; Pe, petals; Se, sepals; NEC, non-embryogenic tissue; PEC, pre-embryogenic tissue; MEC, mature embryogenic tissue; CK-EC, cytokinin-treated tissue.

(C) Top panel shows results of RT-PCR analysis of *Ae4* gene expression. (C) Bottom panel is the RT-PCR gel blot probed with ³²P-labelled *Ae4* cDNA. Lanes: M, 1 kb Bioline DNA ladder; L, leaves; S, stems; B, buds; F, flowers; Si, siliques; R, roots; Ca, carpels.

seems to be approximately at the same level. By contrast, *Ae4* transcripts were not expressed at detectable levels in the non-embryogenic tissue (lane NEC) and the cytokinin-treated embryogenic tissue (lane CK-EC).

DISCUSSION

Ae4 Encodes a Partial Length Protein with Proline-rich Domain

The *Ae4* cDNA encodes a partial length protein having a proline-rich domain. Both the amino acid composition and the presence of repeating motifs of proline are characteristics of proline-rich cell wall proteins (Jose-Estanyol and Puigdomenech, 2000). The 10-mer motif of the putative *AE4* protein is unique in its sequence and belongs to a group of long repeat elements (Gyorgyey *et al.*, 1997). The repeating motifs do not correspond to any of the common motifs previously identified in proline-rich proteins and extensins (Showalter, 1993, Gyorgyey *et al.*, 1997; Jose-Estanyol and Puigdomenech, 2000). Therefore, *Ae4* may encode a novel class of proline-rich proteins (PRPs). Two PRP genes have already been isolated from *Brassica napus*, one that accumulated during pod development (Coupe *et al.*, 1993) and the other one induced by cold treatment (Goodwin *et al.*, 1996). However, the *Ae4* sequence is not similar to either of them.

Expression Analysis of *Ae4*

The expression of *Ae4* transcripts in all vegetative and floral tissues examined corresponds to the observations by Fowler *et al.* (1999) in *Arabidopsis*. They reported that *AtPRP2* and *AtPRP4* transcripts were most abundant in the aerial parts of the plant, namely in leaves, stems, flowers and siliques. Also, *AtPRP4* expression was detected in the early stages of lateral root formation. Since most PRPs are members of a multigene family, it is likely that the same will be true for those of *B. napus*.

Based on Northern analysis, *Ae4* was detected in the pre-embryogenic but not non-embryogenic tissue and this suggests that the encoded protein may be associated with somatic embryogenesis. There have been a few PRP transcripts that have been shown to accumulate during somatic embryogenesis, such as in carrot (Aleith and Richter, 1990; Holk *et al.*, 1996; Yasuda *et al.*, 2001) and *Medicago sativa* (Gyorgyey *et al.*, 1997).

A proline-rich protein encoded by the *DC 2.15* gene was identified as one of the genes that is differentially expressed during induction of somatic embryogenesis in carrot cell suspension culture (Aleith and Richter, 1990). The expression of this gene during somatic embryogenesis was detectable from 3 days after induction, and transcript abundance increased until the heart-shape stage (Aleith and Richter, 1990). This observation was further supported by promoter studies of the *DC 2.15* gene (Holk *et al.*, 1996). Also, Gyorgyey *et al.* (1997) reported that *MsPRP5*, a cDNA clone encoding a small proline-rich protein is preferentially expressed in alfalfa dedifferentiated callus cells. They proposed that the proline-rich protein may cause structural changes of the cell wall required for certain switches in function by plant cells.

He *et al.* (2002) reported that expression of a soybean PRP gene was inhibited by treatment with kinetin. This is consistent with our observation from the Northern analysis of the tissue culture blot that *Ae4* transcripts could not be detected in cytokinin-treated tissue. This suggests that cytokinin treatment has suppressed *Ae4* transcription to undetectable levels, which correlates with the suppression of secondary embryogenesis (Loh *et al.*, 1983). Alternatively, *Ae4* transcripts could be down regulated due to the low rate of secondary embryogenesis in the cytokinin-treated tissue.

Potential Roles of *Ae4* in *Brassica napus*

Proline-rich proteins have been thought to provide strength and rigidity in the cell wall by forming covalently cross-linked networks with cell wall components (Showalter, 1993). PRPs have a relatively high content of tyrosine and lysine residues which have been implicated as the substrate for the peroxidase-mediated insolubilisation of PRPs in soybean via isodityrosine crosslinks (Kleis-San Francisco and Tierney, 1990; Bradley *et al.*, 1992; Brisson *et al.*, 1994). Insolubilisation of PRPs in the cell wall occurs as a rapid response to wounding and treatment with fungal elicitors. PRPs are rapidly insolubilized within the cell wall in response to physical damage, treatment with fungal elicitors, and pathogen infection (Kleis-San Francisco and Tierney, 1990; Bradley *et al.*, 1992; Brisson *et al.*, 1994), indicating an active role in plant defence reactions. Thus, it was proposed that the PRPs

contribute to the cell wall structure of specific cell types based on their patterns of gene expression during plant development and induction by biotic and abiotic stresses. The enhanced expression of *Ae4* transcripts in the pre-embryogenic and mature embryogenic tissues suggests that *Ae4* may have a role during embryo formation. In carrot cultures, it was suggested that two PRPs encoded by No.93 and *DC 2.15* might act as extracellular signal factors during the development of somatic embryos (Yasuda *et al.*, 2001). However, there is no clear evidence for PRPs as signalling molecules that induce somatic embryogenesis.

Assuming that *Ae4* encodes a proline-rich protein, we propose that the expression of *Ae4* transcripts preferentially in embryogenic tissue could be possibly to provide mechanical strength to the embryonic cells that will protect the cell during later phase of embryo development in vitro. More experiments such as isolation of the full-length sequence of *Ae4*, immunolocalization and transgenic studies, especially promoter analysis, are required to provide more information on possible biological functions of the *Ae4* gene. Also, the inducibility of the *Ae4* gene in response to abiotic and biotic stresses in *B. napus* should be tested to explore its regulation.

REFERENCES

- ALEITH, F. and RICHTER, G. (1990). Gene expression during induction of somatic embryogenesis in carrot cell suspensions. *Planta*, *183*, 17-24.
- ALTSCHUL, S.F., MADDEN, T.L., SCHAFFER, A.A., ZHANG, J., ZHANG, Z., MILLER, W. and LIPMAN, D.J. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Research*, *25*, 3389-402.
- BRADLEY, D.J., KJELLBOM, P. and LAMB, C.J. (1992). Elicitor and wound-induced oxidative cross-linking of a proline-rich plant cell wall protein: a novel, rapid defense response. *Cell*, *70*, 21-30.
- BRISSON, L.F., TENHAKEN, R. and LAMB, C. (1994). Function of oxidative cross-linking of cell wall structural proteins in plant disease resistance. *Plant Cell*, *6*, 1703-1712.
- CASSAB, G.I. and VARNER, J.E. (1988). Immunocytolocalisation of extensin in developing soybean seed coats by immunogold-silver staining and by tissue printing on nitrocellulose paper. *Journal of Cell Biology*, *105*, 2581-8.
- CHEN, J. and VARNER, J.E. (1985). An extracellular matrix protein in plants - characterisation of a genomic clone for carrot extensin. *EMBO Journal*, *4*, 2145-2151.
- CHOMCZYNSKI, P. and SACCHI, N. (1987). Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Analytical Biochemistry*, *162*, 156-9.
- COUPE, S.A., TAYLOR, J.E., ISAAC, P.G. and ROBERTS, J.A. (1993). Identification and characterization of a proline-rich mRNA that accumulates during pod development in oilseed rape (*Brassica napus* L.). *Plant Molecular Biology*, *23*, 1223-32.
- DATTA, K., SCHMIDT, A. and MARCUS, A. (1989). Characterization of two soybean repetitive proline-rich proteins and a cognate cDNA from germinated axes. *Plant Cell*, *1*, 945-52.
- EBENER, W., FOWLER, T.J., SUZUKI, H., SHAVER, J. and TIERNEY, M.L. (1993). Expression of DcPRP1 is linked to carrot storage root formation and is induced by wounding and auxin treatment. *Plant Physiology*, *101*, 259-65.
- FRANSSSEN, H.J., NAP, J.P., GLOUDEMANS, T., STIEKEMA, W., VANDAM, H., GOVERS, F., LOUWERSE, J., VANKAMMEN, A. and BISSELING, T. (1987). Characterisation of cDNA for nodulin-75 of soybean - A gene product involved in early stages of root nodule development. *Proceedings of the National Academy of Sciences, USA*, *84*, 4495-4499.
- FOWLER, T.J., BERNHARDT, C. and TIERNEY, M.L. (1999). Characterization and expression of four proline-rich cell wall protein genes in Arabidopsis encoding two distinct subsets of multiple domain proteins. *Plant Physiology*, *121*, 1081-1091.
- GOODWIN, W., PALLAS, J.A. and JENKINS, G.I. (1996). Transcripts of a gene encoding a putative cell wall plasma membrane linker protein are specifically cold-induced in *Brassica napus*. *Plant Molecular Biology*, *31*, 771-781.

- GYORGYEY, J., NEMETH, K., MAGYAR, Z., KELEMEN, Z., ALLIOTTE, T., INZE, D. and DUDITS, D. (1997). Expression of a novel type small proline-rich protein gene of alfalfa is induced by 2,4-dichlorophenoxyacetic acid in dedifferentiated callus cells. *Plant Molecular Biology*, *34*, 593-600.
- HE, C.Y., ZHANG, J.S. and CHEN, S.Y. (2002). A soybean gene encoding a proline-rich protein is regulated by salicylic acid, an endogenous circadian rhythm and by various stresses. *Theoretical and Applied Science*, *104*, 1125-1131.
- HONG, J.C., NAGAO, R.T. and KEY, J.L. (1987). Characterization and sequence analysis of a developmentally regulated putative cell wall protein gene isolated from soybean. *Journal of Biological Chemistry*, *262*, 8367-76.
- HONG, J.C., NAGAO, R.T. and KEY, J.L. (1989). Developmentally regulated expression of soybean proline-rich cell wall protein genes. *Plant Cell*, *1*, 937-43.
- HOLK, A., KALDENHOFF, R. and RICHTER, G. (1996). Regulation of an embryogenic carrot gene (DC 2.15) and identification of its active promoter sites. *Plant Molecular Biology*, *31*, 1153-1161.
- JOSE-ESTANYOL, M. and PUIGDOMENECH, P. (2000). Plant cell wall glycoproteins and their genes. *Plant Physiology and Biochemistry*, *38*, 97-108.
- JOSE-ESTANYOL, M., RUIZ-AVILA, L. and PUIGDOMENECH, P. (1992). A maize embryo-specific gene encodes a proline-rich and hydrophobic protein. *Plant Cell*, *4*, 413-23.
- KLEIS-SAN FRANCISCO, S.M. and TIERNEY, M.L. (1990). Isolation and characterization of a proline-rich cell wall protein from soybean seedlings. *Plant Physiology*, *94*, 1897-1902.
- LINDSTROM, J.T. and VODKIN, L.O. (1991). A soybean cell wall protein is affected by seed colour genotype. *Plant Cell*, *3*, 561-71.
- LOH, C.S., INGRAM, D.S. and HANKE, D.E. (1983). Cytokinins and the regeneration of plantlets from secondary embryoids of winter oilseed rape, *Brassica napus* ssp. *oleifera*. *New Phytologist*, *95*, 349-358.
- NAMASIVAYAM, P., SKEPPER, J. and HANKE, D. (2006a). Identification of a potential structural marker for embryogenic competency in the *Brassica napus* spp. *oleifera* embryogenic tissue. *Plant Cell Reports*, *25*, 887-895.
- NAMASIVAYAM, P. and HANKE, D. (2006b). Identification of differentially expressed sequences in pre-embryogenic tissue of oilseed rape by suppression subtractive hybridisation (SSH). *Plant, Cell and Organ Culture*, *86*(3), 417-421.
- SAMBROOK, J., FRITSCH, E.F. and MANIATIS, T. (1989). *Molecular Cloning: A Laboratory Manual* (2nd ed.). New York, USA: Cold Spring Harbor Press.
- SHOWALTER, A.M. (1993). Structure and function of plant cell wall proteins. *Plant Cell*, *5*, 9-23.
- SOUTHERN, E.M. (1975). Detection of specific sequences among DNA fragments separated by gel electrophoresis. *Journal of Molecular Biology*, *98*, 503.
- WILSON, R.C., LONG, F., MARUOKA, E.M. and COOPER, J.B. (1994). A new proline-rich early nodulin from *Medicago truncatula* is highly expressed in nodule meristematic cells. *Plant Cell*, *6*, 1265-75.
- WYATT, R.E., NAGAO, R.T. and KEY, J.L. (1992). Patterns of soybean proline-rich protein gene expression. *Plant Cell*, *4*, 99-110.
- YASUDA, H., NAKAJIMA, M., ITO, T., OHWADA, T. and MASUDA, H. (2001). Partial characterization of genes whose transcripts accumulate preferentially in cell clusters at the earliest stage of carrot somatic embryogenesis. *Plant Molecular Biology*, *45*, 705-712.

