

Cloning, extracellular expression and characterization of predominant β -CGTase from *Bacillus* sp. G1 in *E. coli*.

ABSTRACT

The cyclodextrin glucoamylase (CGTase, EC 2.4.1.19) gene from *Bacillus* sp. G1 was successfully isolated and cloned into *Escherichia coli*. Analysis of the nucleotide sequence revealed the presence of an open reading frame of 2,109 bp and encoded a 674 amino acid protein. Purified CGTase exhibited a molecular weight of 75 kDa and had optimum activity at pH 6 and 60 degrees C. Heterologous recombinant protein expression in *E. coli* is commonly problematic causing intracellular localization and formation of inactive inclusion bodies. This paper shows that the majority of CGTase was secreted into the medium due to the signal peptide of *Bacillus* sp. G1 that also works well in *E. coli*, leading to easier purification steps. When reacted with starch, CGTase G1 produced 90% β -cyclodextrin (CD) and 10% γ -CD. This enzyme also preferred the economical tapioca starch as a substrate, based on kinetics studies. Therefore, CGTase G1 could potentially serve as an industrial enzyme for the production of β -CD.

Keyword: *Bacillus* sp. G1; Cyclodextrin; Cyclodextrin glucoamylase; Extracellular expression; Predominant β -CGTase; Signal peptide.