

Novel approach of vaccination against *Brucella abortus* 544 based on a combination of fusion proteins, human serum albumin and brucella abortus Lipopolysaccharides.

ABSTRACT

Lipopolysaccharide (LPS) of *Brucella abortus* is an essential component for developing the subunit vaccine against brucellosis. *B. abortus* LPS was extracted by n-butanol, purified by ultracentrifugation and detoxified by alkaline treatment. Pyrogenicity and toxicity of *B. abortus* LPS and detoxified-LPS (D-LPS) were analyzed and compared with LPS of *E. coli*. Different groups of mice were immunized intraperitoneally with purified *B. abortus* LPS, D-LPS, a combination of LPS with human serum albumin (LPS-HSA) and *B. abortus* S19 bacteria; besides, control mice were inoculated with sterile saline. Two doses of vaccine were given 4 weeks apart. Mice were challenged intraperitoneally with virulent *B. abortus* 544 strain 4 weeks after the second dose of vaccine. Sera and spleens of mice were harvested 4 weeks after challenge. LPS-*B. abortus* was 10,000-fold less potent in LAL test and 100-fold less potent in eliciting fever in rabbits than in *E. coli* LPS. And D-LPS was very less potent in LAL test and eliciting fever in rabbits ordinary LPS. The antibody titer of anti-LPS immunoglobulin G (IgG) was higher than D-LPS. However, mice immunized with either LPS, D-LPS or LPS-HSA vaccines showed a significant protection against infection of the spleen ($p < 0.01$). There was no significant difference between mice immunized with LPS and D-LPS in terms of protection ($p < 0.99$). Therefore, it was concluded that D-LPS and LPS-HSA for *B. abortus* can be used as safer and more potent vaccines than ordinary LPS-*B. abortus* vaccine.

Keyword: Novel approach of vaccination; *Brucella abortus* 544; Fusion proteins; Human serum albumin; *Brucella abortus* lipopolysaccharides.