

Optimization of an induction strategy for improving interferon- α 2b production in the periplasm of *Escherichia coli* using response surface methodology.

ABSTRACT

Induction strategies for the periplasmic production of recombinant human IFN- α 2b (interferon- α 2b) by recombinant *Escherichia coli* Rosetta-gami 2(DE3) were optimized in shake-flask cultures using response surface methodology based on the central composite design. The factors included in the present study were induction point, which related to the attenuation of the cell culture, IPTG (isopropyl beta-D-thiogalactoside) concentration and induction temperature. Second-order polynomial models were used to correlate the abovementioned factors to soluble periplasmic IFN- α 2b formation and percentage of soluble IFN- α 2b translocated to the periplasmic space of *E. coli*. The models were found to be significant and subsequently validated. The proposed induction strategies consisted of induction at an attenuation of 4 (measured as D600), IPTG concentration of 0.05 mM and temperature of 25 degrees C. The optimized induction strategy reduced inclusion-body formation as evidenced by electron microscopy and yielded 323.8 ng/ml of IFN- α 2b in the periplasmic space with translocation of 74% of the total soluble product. In comparison with the non-optimized condition, soluble periplasmic production and the percentage of soluble IFN- α 2b translocated to the periplasmic space obtained in optimized induction strategies were increased by approx. 20-fold and 1.4-fold respectively.

Keyword: *Escherichia coli*; Induction strategy; Interferon- α 2b (IFN- α 2b); Periplasm; Response surface methodology (RSM); Shake-flask culture.