



UNIVERSITI PUTRA MALAYSIA

**DEVELOPMENT OF A WATER-DISPERSIBLE PHYTOSTEROL
NANODISPERSION SYSTEM AND ITS APPLICATION IN SOY MILK
STORAGE**

LEONG WAI FUN

FSTM 2010 17

**DEVELOPMENT OF A WATER-DISPERSIBLE PHYTOSTEROL
NANODISPERSION SYSTEM AND ITS APPLICATION
IN SOY MILK STORAGE**

By

LEONG WAI FUN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfillment of the Requirements for the Degree of Doctor of Philosophy**

November 2010



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in Fulfillment of the requirement of the degree of Doctoral of Philosophy

**DEVELOPMENT OF A WATER-DISPERSIBLE PHYTOSTEROL
NANODISPERSION SYSTEM AND ITS APPLICATION
IN SOY MILK STORAGE**

By

LEONG WAI FUN

November 2010

Chairman: Associate Professor Tan Chin Ping, PhD

Faculty: Food Science and Technology

This work was aimed to develop a stable water-dispersible phytosterol nanodispersion system. In the first part of this work, the formation and characterization of phytosterol nanodispersions prepared using Tween 20 was investigated. The experiment demonstrated the feasibility of phytosterol nanodispersion production using hexane as organic phase through an emulsification-evaporation technique. The mean particle diameter of phytosterol nanoparticles produced was 50 nm in diameter and had a spherical shape. The dispersed phase ratio, conventional homogenization parameters and the homogenization pressure showed significant ($p < 0.05$) effects on the final phytosterol particles size and their distribution profiles. High-pressure homogenization caused significant phytosterol loss ($p < 0.05$).

Two response surface methodology (RSM) processes were applied to optimize the processing and formulation parameters for preparing phytosterol nanodispersions. The



optimized processing parameters were 15.25 min of mixing time, 7000 rpm of mixing speed and a homogenization pressure of 42.4 MPa. The corresponding responses for the optimized preparation conditions were a mean particle size (PS) of 52 nm and a phytosterol concentration (Phyto) of 336 mg/l. The optimized formulation parameters determined were a phase ratio (PR) of 3.54 and a mixture ratio (MR) of 0.19, and the corresponding optimized responses were a PS of 55.4 nm and 87.6% phytosterol concentration. The PS showed no significant ($p > 0.05$) change over a period of 8 weeks of storage at 4 °C.

The Tween 20 was replaced by four different types of sucrose fatty acid esters (SEs), namely sucrose palmitate (P-1570), sucrose laureate (L-1695), sucrose steareate (S-1570) and sucrose oleate (OWA-1570). The physicochemical properties of SE-stabilized water-dispersible phytosterol nanodispersions were examined. The PS and the %Phyto of the prepared phytosterol nanodispersions ranged from 2.8 to 259.9 nm and from 230.4 to 504.6 mg/l. All of the prepared phytosterol nanodispersions exhibited pseudoplastic flow behavior, with a low yield stress ranging from 0.630 to 9.183 mPas and a low consistency coefficient of 0.608 to 88.710 mPas. Less than 1.5 ppm of hexane residues in the prepared nanodispersions was detected. Sucrose esters P-1570, L-1695 and S-1570 were found to be appropriate for use in preparing phytosterol nanoparticles with small PS at a monomodal distribution, with high clarity. The high phytosterol-loaded nanodispersions prepared with co-solvents ethanol and L-1695 had small spherical PS of approximately 5 nm, with low viscosity and high clarity. The solvent residue levels in the final prepared nanodispersions were acceptable.



L-1695 was selected for further optimization of the production of L-1695-stabilized water-dispersible phytosterol nanodispersions through RSM. The optimized parameters were 5.5% of Ph (phytosterol concentration), 1.0% of L (L-1695 concentration), 3 C (homogenization cycle), and P(homogenization pressure) of 37 MPa. The corresponding responses for the optimized condition were a PS of 3 nm and a %Ph of 90.4%. The optimized phytosterol nanodispersions had a polydispersity index of 0.550 at a monomodal distribution. The pH value and hexane and ethanol residues concentration were 6.45, 48.2 $\mu\text{l/l}$ and 930.3 $\mu\text{l/l}$, respectively. The optimized nanodispersions were stable to heat treatment up to 121 °C, chilling at 4 and 10 °C and freezing with a cryoprotectant at – 4 and – 20 °C.

The stability of the optimized phytosterol nanodispersions and phytosterol-fortified soy milk (SMP) over a 12-week period was investigated. The storage resulted in increases in PS and reduced the total phytosterol concentration of the autoclaved phytosterol nanodispersions. Adding phytosterol nanodispersions increased the mean particle size of the soy milk. The fortified phytosterol nanoparticles became entrapped in the fat droplets of the soy milk. The stability of the SMP depended on the stability of the soy milk. The fortification of phytosterol nanodispersions in soy milk was feasible.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Doktor Falsafah

**PEMBANGUNAN NANOSERAKAN FITOSTEROL TERSERAK DALAM AIR
DAN APLIKASINYA DALAM PENYIMPANAN SUSU SOYA**

Oleh

LEONG WAI FUN

November 2010

Pengerusi: Profesor Madya Tan Chin Ping , PhD

Fakulti: Sains dan Teknologi Makanan

Kajian ini bertujuan untuk membangunkan nanoserakan fitosterol terserak dalam air untuk penggunaan makanan. Bahagian pertama kajian ini melibatkan kajian permulaan untuk menyiasat pembentukan dan perwatakan nanoserakan fitosterol yang disediakan dengan menggunakan Tween 20 sebagai agen pengemulsi. Eksperimen menunjukkan kemungkinan penghasilan nanoserakan fitosterol dengan menggunakan pelarut organik heksana melalui kaedah pengemulsian-penyejatan. Diameter purata partikel nanoserakan yang dihasilkan berukuran 50 nm dan berbentuk sfera. Nisbah fasa serakan, parameter penghomogenan biasa dan tekanan penghomogenan menunjukkan kesan ketara ($p < 0.05$) ke atas saiz dan pengagihan partikel fitosterol. Selain itu, penghomogenan bertekanan-tinggi menyebabkan kehilangan fitosterol yang ketara ($p < 0.05$).

Due kaedah permukaan gerak balas (RSM) telah digunakan untuk pengoptimuman parameter pemprosesan dan formulasi bagi penyediaan nanoserakan fitosterol.

Parameter pemrosesan yang optimum ialah 15.25 min masa pengaulan, 7000 rpm halaju pengacauran and tekanan penghomogenan ialah 42.4 MPa. Parameter-parameter pemrosean pada keadaan optimum tersebut menghasilkan saiz purata partikel (PS) bernilai 52 nm and kepekatan fitosterol (Phyto) bernilai 336 mg/l. Parameter formulasi yang optimum ialah nisbah serakan (PR) bernilai 3.54 and nisbah campuran (MR) bernilai 0.19. Formulasi optimum ini telah menghasilkan PS bernilai 55.4 nm dan 87.6% kepekatan fitosterol. Tiada perbezaan ketara ($p < 0.05$) pada PS selama 8 minggu penyimpanan pada suhu 4°C.

Seterusnya, Tween 20 telah digantikan dengan ester sukrosa (SEs) iaitu palmitat sukrosa (P-1570), lauriat sukrosa (L-1695), stearat sukrosa (S-1570) dan oliat sukrosa (OWA-1570). Kajian ke atas sifat fisikokimia nanoserakan fitosterol yang disediakan dengan SEs telah dijalankan. PS dan %fitosterol dalam nanoserakan fitosterol yang dihasilkan adalah berukuran 2.8 hingga 259.9 nm dan 230.4 hingga 504.6 mg/l masing-masing. Semua nanoserakan fitosterol yang disediakan menunjukkan sifat pengaliran pseudoplastik dengan hasil stress yang rendah bernilai 0.630 hingga 9.183 mPa and nilai ketekalan yang rendah di antara 0.608 hingga 88.710 mPas. Kurang daripada 1.5 ppm sisa heksana telah dikesan dalam nanoserakan yang telah disediakan. Ester sukrosa P-1570, L-1695 dan S-1570 didapati bersesuaian untuk digunakan dalam penyediaan nanoserakan fitosterol dengan PS yang kecil, corak serakan monomodel dan kejernihan yang tinggi. Nanoserakan fitosterol dengan kepekatan fitosterol yang tinggi telah disediakan dengan sepelarut etanol and L-1695 dan menghasilkan PS yang kecil dan sfera pada ukuran kira-kira 5 nm, kekentalan yang rendah dan kejernihan yang tinggi.

Kepekatan sisa pelarut pada nanoserakan fitosterol akhir adalah pada tahap yang munasabah.

L-1695 telah dipilih untuk kajian pengoptimuman bagi menghasilkan nanoserakan fitosterol terserak dalam air melalui RSM. Parameter optimum ialah 5.5% Ph (kepekatan fitosterol), 1.0% L(kepekatan L-1695), 3C (pusingan penghomogenan) dan 37 MPa P (tekanan penghomogenan). Parameter optimum ini telah menghasilkan PS berukuran 3 nm and %Ph bernilai 90.4%, Nanoserakan fitosterol yang dioptimumkan mengandungi indek poly-serakan berukuran 0.550 pada corak serakan monomodel. Nilai pH, kepekatan sisa heksana dan etanol, ialah 6.45, 48.2 $\mu\text{l/l}$ dan 930.3 $\mu\text{l/l}$ masing-masing. Nanoserakan fitosterol yang dioptimumkan adalah stabil terhadap rawatan pemanasan sehingga suhu 121°C, penyejukan pada 4 dan 10°C, dan pembekuan dengan cryoprotectant pada -4 dan -20°C.

Kestabilan nanoserakan fitosterol yang telah dioptimumkan dan kestabilan susu soya yang telah ditambah dengan nanoserakan fitosterol (SMP) telah dikaji selama 12 minggu. Penyimpanan telah meningkatkan PS dan mengurangkan kepekatan fitosterols pada nanoserakan yang telah diautoklaf. Penambahan nanoserakan fitosterol dalam susu soya meningkatkan PS susu soya. Nanopartikel fitosterol yang ditambahkan dalam susu soya telah terperangkap dalam titisan lemak susu soya. Sehubungan dengan ini, kestabilan SMP bergantung pada kestabilan susu soya. Penambahan nanoserakan fitosterol dalam susu soya adalah boleh diterima.

ACKNOWLEDGEMENTS

I wish to express a huge amount of gratitude to Assoc. Prof. Dr. Tan Chin Ping, my primary supervisor, for his support, guidance and encouragement throughout this study. I am extremely grateful for the effort he continues to spend on my research, manuscripts and thesis. Special thanks should go to my cosupervisors, Prof. Yaakob Che Man, Assoc. Prof. Dr. Lai Oi Ming and Dr. Kamariah Long, for their constant support and encouragement.

I cannot thank the Graduate School of Universiti Putra Malaysia enough for the Graduate Research Fellowship, which permitted me to concentrate on my studies without financial worries. I would also like thank Assoc. Prof. Dr. Misni Misran and Ms. Tan Hsiao Wei of the Colloid Research Group of Department of Chemistry, University Malaya, for providing the analytical instruments and technical support for particle sizing and zeta-potential analysis. A sincere gratitude is also extended to Mrs. Aminah Jusoh of Unit Microscopy, Institute BioScience, Universiti Putra Malaysia, for helping with the TEM analysis.

I cannot forget to thank all my laboratory mates for their concern and assistance throughout my study. Many thanks are also due to the faculty members, staff and technicians for their invaluable assistance and support. Finally, I would like to express my appreciation to my family, my parents, my siblings and all my friends for their unconditional love and encouragement.



I certify that a Thesis Examination Committee has met on 23 November 2010 to conduct the final examination of Leong Wai Fun on her thesis entitle “Development of water-dispersible phytosterol nanodispersion system and its application in soy milk ” is accordance with Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Examination Committee were as follows:

Sharifah Kharidah Syed Muhammad

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Hasanah Mohd. Ghazali

Dean
School of Graduate Studies
University Putra Malaysia
(Internal Examiner)

Mohammad Reza Mozafari

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Internal Examiner)

Qingrong Huang

Lecturer
School of Environmental and Biological Sciences
The State University of New Jersey
(External Examiner)

BUJANG BIN KIM HUAT, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Tan Chin Ping, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Chairman)

Yaakob Bin Che Man, PhD

Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Lai Oi Ming

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Kamariah Long, PhD

Deputy Director
Bio-processing Units,
Biotechnology ReserachCentre
Malaysian Agricultural Research and Development Institute (MARDI)
(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:



DECLARATION

I declare that the thesis is my original work except for quotations and citations, which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.

LEONG WAI FUN

Date: 23 November 2010



TABLE OF CONTENT

	Page
ABSTRACT	ii
ABSTRAK	v
ACKNOWLEDGEMENTS	viii
APPROVAL	ix
DECLARATION	xi
LIST OF TABLES	xvii
LIST OF FIGURES	xx
LIST OF ABBREVIATIONS	xxiii
CHAPTER	
1 GENERAL INTRODUCTION	1
2 LITERATURE REVIEW	
Introduction of phytosterols	7
Phytosterols and health	8
Phytosterols in foods	11
Safety of phytosterols	15
Emulsion	15
Nanodispersion	16
Formation of nanodispersions	17
Preparation of nanodispersions	20
Application, bioavailability and safety of nanodispersions	25
Emulsifiers	26
Hydrophile-lipophile Balance (HLB)	28
Sucrose fatty acid esters	29
Stability of nanodispersions	32
Creaming or sedimentation	33
Coalescence	34
Flocculation	35
Ostwald ripening	35
Characterization of nanodispersions	39
Particle size and particle-size distribution	39
Morphology of the particles	41
Rheological properties	43
Appearance	45
3 PREPARATION, CHARACTERIZATION AND OPTIMIZATION OF PHYTOSTEROL NANODISPERIONS STABILIZED BY TWEEN 20	
Introduction	48
Materials and Methods	
Materials	53



Methods	53
Part 1: Preparation and characterization of Tween 20 stabilized phytosterol nanodispersions	
Solubility of phytosterols in organic solvents	53
Preparation of phytosterol nanodispersions	54
Analysis of particle size and its polydispersity	55
Gas Chromatography analysis of phytosterols content	
Sample preparation	56
Preparation of standards and internal standards	57
Gas chromatography analysis of phytosterol	57
Calculation of phytosterol content	58
Transmission electro microscopy (TEM)	58
Statistical analysis	59
Part 2: Optimization of processing parameters	
Experimental design	59
Preparation of phytosterol nanodispersions	61
Statistical analysis	62
Optimization and verification	62
Part 3: Optimization of formulation parameters	
Experimental design	63
Preparation of phytosterol nanodispersions	64
Optimization and verification	65
Results and Discussion	
Part 1: Preparation and characterization of Tween 20 stabilized phytosterol nanodispersions	
Solubility of phytosterol	66
Effect of organic phase types on the physicochemical properties of phytosterol nanodispersions	67
Effect of phase ratio on the particle size of phytosterol nanodispersions	70
Effect of conventional homogenization parameters on the physicochemical properties of the phytosterol nanodispersions	73
Effect of high-pressure homogenization parameters on the physicochemical properties of the phytosterol nanodispersions	76
TEM analysis	81
Summary	81
Part 2: Optimization of processing parameters	
General	83
Fitting the model	84
The main and the interaction effects of the independent variables	86
Optimization of processing parameters for the production of phytosterol nanodispersions	89
Model verification	90



	Summary	90
Part 3:	Optimization of formulation parameters	
	General	91
	Fitting the model	92
	Main and the interaction effects of the independent variables	93
	Optimization of processing parameters for the production of phytosterol nanodispersions	98
	Model verification	98
	Stability of optimized nanodispersions during storage	99
	Summary	100
4	EFFECT OF SUCROSE FATTY ACIDS ESTERS ON WATER-DISPERSIBLE PHYTOSTEROL NANODISPERSIONS	
	Introduction	103
	Materials and methods	
	Materials	105
	Methods	
	Solubility of phytosterols in organic solvent	106
	Preparation of phytosterol nanodispersions	107
	Analysis	108
	Determination of solvent residual	
	HS-SPME- analysis	108
	Gas chromatography analysis of organic solvent residue	109
	Viscosity measurement	
	Transmittance measurement	110
	Statistical analysis	110
		111
	Result and Discussion	
	Part 1: Effect of P-1570, L-1695, S-1570 and OWA-1570 on particle characteristic and flow properties of phytosterols nanodispersions	111
	Particle size and its distribution	112
	Retention of phytosterol	115
	Hexane residue	117
	Flow property	118
	Transmittance spectrum	121
	TEM analysis	124
	Part 2: Effect of phytosterol concentration on P-1570 and L-1695 stabilized phytosterols nanodispersions	
	Solubility of phytosterols in the mixture organic phase	125
	Particle size and distribution profile, and the phytosterols retention.	126
	Solvent residue	131
	Flow property	134
	Transmittance spectrum	136
	Part 3: preparation and characterization of L-1695 stabilized phytosterol	



nanodispersions	
Effect of phase ratio	138
Effect of high pressure homogenization parameters	139
Effect of conventional homogenization parameters	141
Summary	143
5 OPTIMIZATION OF SUCROSE LAUREATE-STABILIZED WATER-DISPERSIBLE PHYTOSTEROL NANODISPERSIONS	
Introduction	145
Materials and methods	
Materials	147
Methods	
Experimental design	148
Preparation of phytosterol nanodispersions	149
Analysis	150
Zeta potential measurement	151
pH measurement	151
Effect of temperature on particle size and distribution	151
Statistical analysis	152
Optimization and verification	152
Result and Discussion	
Model fitting	153
The main effects and the interaction effects of the independent variables	156
<i>Optimization of production of phytosterol nanodispersions</i>	160
Model verification	162
Characteristic of the prepared optimized phytosterol nanodispersions	163
Effect of thermal treatment on the optimized phytosterol nanodispersion	164
Summary	169
6 STORAGE STABILITY EVALUATION OF THE OPTIMIZED SUCROSE LAUREATE-STABILIZED PHYTOSTEROL NANODISPERSIONS AND ITS APPLICATION IN SOY MILK	
Introduction	172
Materials and methods	
Materials	174
Methods	
Preparation of phytosterol nanodispersions	175
Preparation of soymilk	175
Formulation and packaging	176
Analysis	177



Particle size growth ratio	177
Cloudiness measurement	177
Statistical analysis	178
Result and discussion	
Particle size and distribution	179
Solvent residue	189
Phytosterol concentration	191
Cloudiness	192
Flow properties	194
Summary	199
7 CONCLUSION AND RECOMMENDATIONS	200
BIBLIOGRAPHY	206
APPENDICES	225
BIODATA OF STUDENT	230
LIST OF PUBLICATION	231
LIST OF PAPERS PRESENTED AT TECHNICAL MEETINGS	232

