



UNIVERSITI PUTRA MALAYSIA

USING SEAWATER FOR WEED MANGEMENT IN TURFGRASS

MD. KAMAL UDDIN

ITA 2010 1

USING SEAWATER FOR WEED MANGEMENT IN TURFGRASS

BY

MD. KAMAL UDDIN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia in
Partial Fulfilment of the Requirements for the Degree of Doctor of Philosophy**

September 2010



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in partial fulfilment of the requirement for the degree of Doctor of Philosophy

USING SEAWATER FOR WEED MANGEMENT IN TURFGRASS

By

MD. KAMAL UDDIN

September 2010

Chairman: Associate Professor Abdul Shukor Juraimi, PhD

Institute: Tropical Agriculture

Long term use of herbicide can pose serious threat to environment. Alternative strategies to chemical herbicide in weed control have received increasing attention now a days. Salty water could be used to control weed in salt tolerant turfgrass areas. A series of experiments were conducted to examine the response of turfgrass species, common local weed species and soil microbial communities to salinity stress both singly and in combination with reduced rates of recommended herbicides. The plant species selected for evaluation in these studies were based on a detailed preliminary survey conducted in local turfgrass areas.

Four salinity levels (0, 24, 48 and 72 dSm⁻¹) were imposed on 16 turfgrass entries in glasshouse experiments to identify potential salt tolerant turf species. Tolerance ranking was based on 50% reduction of shoot and root dry weights, leaf firing and turf quality. Predicted salinity levels for 50% shoot and root growth reductions were, 17.0 – 39.8 and



$18.8 - 49.4 \text{ dS m}^{-1}$, respectively. *Paspalum vaginatum* proved to be superior in salt tolerance, and was closely followed by *Zoysia japonica* and *Zoysia matrella*.

In the field survey of four different turfgrass areas, 79 weed species belonging to 16 families were recorded. Maximum numbers of weed species were found on residential lawns, while least numbers were recorded on golf course putting greens; athletic fields and sod farms ranked intermediate. *Cyperus aromaticus* yielded the highest relative abundance value in golf course putting greens (71.5), followed by athletic fields (45.5) and residential lawns (21.7). *Fimbristylis dichotoma* and *Cyperus compressus* yielded the highest relative abundance values on residential lawns (35.0) and sod farms (30.4), respectively.

In studies evaluating the use of saline solutions for weed control, the same four salinity levels (0, 24, 48 and 72 dS m^{-1}) were applied to all 79 weed species, along with *P. vaginatum* (as a control). The results on injury ratings for salt tolerant weeds were categorized as highly susceptible, moderately susceptible and extremely tolerant. The results showed that 44 weed species were very susceptible and found to be effectively controlled (100%) at salinities of 24 and 48 dS m^{-1} , while 30 species were found to be effectively controlled with the 72 dS m^{-1} salinity treatment. The five most serious weeds viz. *Eragrostis atrovirens*, *Sporobolus diander*, *Cyperus aromaticus*, *Cyperus rotundus* and *Emilia sonchifolia* were found to be extremely tolerant, and were not controlled even at the highest salinity level of 72 dS m^{-1} .

In the physiological study, three most tolerant and three least tolerant turf species were subjected to seven salinity levels (0, 8, 16, 24, 32, 40 and 48 dS m⁻¹) treatments and were assessed for leaf proline, chlorophyll and relative water contents. Leaf firing, plant biomass, and concentrations of Na, K, Ca, and Mg in shoots and roots were also determined. Root structure injury was observed under a scanning electron microscope (SEM). In general, trends of proline concentrations were found to increase with increasing salinity levels. However, there was an extremely higher rate of increase in salt tolerant species than in sensitive ones. *Zoysia japonica* was the least Na accumulating at all salinity levels followed by *P. vaginatum* and *Z. matrella*, while *D. didactyla* was the highest Na accumulating species followed by *Cynodon dactylon* ‘satiri’ and *Cynodon dactylon* ‘tifdrawf’. *Paspalum vaginatum* always had greater K contents at all salinity levels followed by *Z. japonica* and *Z. matrella*. On the other hand, the lowest K contents were recorded in *C. dactylon* ‘tifdwarf’ followed by *C. dactylon* ‘satiri’ and *D. didactyla*. Salinity caused root cortex cells to collapse, in *C. dactylon* ‘tifdwarf’, *C. dactylon* ‘satiri’ and *D. didactyla* at highest salinity level (48 dS m⁻¹), but not in *P. vaginatum*, *Z. japonica* and *Z. matrella*. Soil microbes were more or less affected by application of either herbicides, sea water or in combination. However, salt water treatment in combination with reduced herbicide are more eco-friendly than other options.

Sensitivity to seawater plus reduced rates of herbicide was evaluated for *E. atrovirens*, *S. diander*, *C. aromaticus*, *C. rotundus* and *E. sonchifolia* as well as for turfgrasses viz. *P. vaginatum*, *C. dactylon* ‘satiri’, and *Z. japonica*. The weed species *S. diander*, *C.*

aromaticus, *C. rotundus* and *E. sonchifolia* except *E. atrovirens* were fully controlled when treated with $\frac{3}{4}$ recommended trifloxsulfuron-sodium with sea water, $\frac{3}{4}$ recommended trifloxsulfuron-sodium with $\frac{3}{4}$ sea water, $\frac{1}{2}$ recommended trifloxsulfuron-sodium with sea water, $\frac{3}{4}$ recommended quinclorac with sea water and $\frac{3}{4}$ recommended quinclorac with $\frac{3}{4}$ sea water. *Eragrostis atrovirens* showed maximum 48% injury when treated with $\frac{3}{4}$ recommended trifloxsulfuron-sodium and sea water. Among the tested turfgrasses, *P. vaginatum* showed only 8% injury to sea water in combination with $\frac{3}{4}$ recommended quinclorac, indicating greater salt tolerance. *Zoysia japonica* also showed no more than 14% injury when treated with sea water in combination with $\frac{3}{4}$ recommended trifloxsulfuron-sodium or quinclorac. *Cynodon dactylon* ‘satiri’ had up to 21% salt injury with $\frac{3}{4}$ sea water in combination with $\frac{3}{4}$ recommended trifloxsulfuron –sodium.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai
memenuhi keperluan untuk ijazah Doktor Falsafah

**MENGGUNAKAN AIR LAUT UNTUK PENGURUSAN RUMPAI
DI KAWASAN RUMPUT TURF**

Oleh

MD. KAMAL UDDIN

September 2010

Pengerusi : Professor Madya Abdul Shukor Juraimi, PhD

Institut : Pertanian Tropika

Penggunaan racun rumpai dalam jangka masa panjang memberikan kesan buruk kepada alam sekitar. Kini keadaan alternatif kepada racun rumpai dalam pengawalan rumpai semakin mendapat perhatian. Air masin mungkin boleh digunakan untuk mengawal rumpai bagi penanaman rumput turf yang toleran kepada kemasinan. Satu siri eksperimen telah dijalankan untuk menguji tindak balas spesies rumput turf, spesies rumpai tempatan dan mikrob tanah terhadap tekanan kemasinan secara tunggal dan dalam kombinasi dengan racun rumpai pada kadar rendah. Spesies tumbuhan yang dipilih untuk kajian ini adalah berdasarkan kepada keputusan kajian awal di kawasan rumput turf tempatan.

Di dalam eksperimen rumah kaca, 16 rumput turf telah dirawat dengan empat kadar kemasinan ($0, 24, 48$ dan 72 dS m^{-1}) untuk mengenalpasti spesies rumput turf yang tahan terhadap kemasinan. Tahap ketahanan adalah berdasarkan kepada 50%

pengurangan pertumbuhan pucuk dan akar, lecur daun dan kualiti rumput turf. Tahap kemasinan yang telah dijangka untuk 50% pengurangan pertumbuhan pucuk dan akar adalah masing-masing pada kadar 17.0-39.8 dan 18.8-49.4 dS m⁻¹. *Paspalum vaginatum* didapati lebih tahan kepada kemasinan diikuti oleh *Zoysia japonica* dan *Zoysia matrella*.

Bancian di lapangan mendapati 79 spesies rumput daripada 16 famili telah direkodkan di empat jenis kawasan rumput turf yang dikaji. Bilangan spesies rumput paling tinggi telah dicatatkan di laman kawasan perumahan manakala spesies rumput paling rendah didapati di ‘putting green’ Padang golf; bilangan spesies di padang olahraga dan nurseri turf adalah sederhana. *Cyperus aromaticus* mencatatkan populasi relatif yang paling tinggi di kawasan ‘putting green’ (71.5) diikuti dengan padang olahraga (45.5) dan laman kawasan perumahan (21.7). Populasi relatif paling tinggi bagi *Fimbristylis dichotoma* dan *Cyperus compressus* terdapat di kawasan perumahan (35.0) diikuti dengan nurseri turf (30.4).

Dalam kajian pengawalan rumput menggunakan air masin, kesemua empat kadar kemasinan yang sama (0, 24, 48 dan 72 dS m⁻¹) di uji ke atas kesemua 79 spesies rumput dan *Paspalum vaginatum* yang dijadikan sebagai kawalan. Rumput di bahagikan kepada tiga kumpulan berdasarkan kepada tahap kecederaan, iaitu tidak tahan masin, sederhana tahan dan paling tahan. Keputusan kajian menunjukkan 44 spesies rumput yang sangat tidak tahan terhadap kemasinan dapat dikawal secara berkesan (100%) pada kadar 24 dan 48 dS m⁻¹, manakala 30 spesies dapat dikawal secara berkesan hanya pada kadar 72 dS m⁻¹. Lima spesies rumput yang paling serius iaitu *Eragrostis atrovirens*, *Sporobolus*

diander, *Cyperus aromaticus*, *Cyperus rotundus* dan *Emilia sonchifolia* dikenalpasti sebagai yang paling tahan kerana tidak dapat dikawal walaupun pada kadar kemasinan yang paling tinggi iaitu 72 dS m^{-1} .

Dalam kajian fisiologi, tujuh kadar kemasinan ($0, 8, 16, 24, 32, 40$ dan 48 dS m^{-1}) telah digunakan ke atas tiga spesies rumput turf paling tahan dan tiga spesies yang paling tidak tahan kemasinan dengan menilai kandungan prolin, klorofil dan relatif air daun. Lecur daun, biomass, kepekatan Na, K dan Mg pucuk dan akar juga di tentukan. Kecederaan struktur akar diamati menggunakan mikroskop pengimbas elektron. Pada umumnya, kepekatan prolin meningkat selari dengan peningkatan kadar kemasinan. Semakin tahan spesies rumpai terhadap kemasinan, semakin tinggi kandungan prolin. Pengumpulan Na paling rendah pada semua kadar kemasinan di catatkan oleh *Z. japonica* diikuti dengan *P. vaginatum* dan *Z. matrella*, sementara *D. didactyla* mempunyai kandungan Na paling tinggi diikuti dengan *C. dactylon* ‘satiri’ dan *C. dactylon* ‘tifdwarf’. *P. vaginatum* adalah spesies paling tinggi kandungan K pada semua kadar kemasinan, diikuti oleh *Z. japonica* dan *Z. matrella*, manakala kandungan K paling rendah adalah pada spesies *C. dactylon* ‘tifdwarf’ diikuti dengan *C. dactylon* ‘satiri’, dan *D. didactyla*. Kemasinan tinggi (48 dS m^{-1}) telah memusnahkan sel kortek akar rumput turf *C. dactylon* ‘tifdwarf’, *C. dactylon* ‘satiri’ dan *D. didactyla*, kecuali species *P. vaginatum*, *Z. japonica* dan *Z. matrella*. Penggunaan racun rumpai air laut atau kombinasi kedua-duanya didapati mempengaruhi kehadiran mikrob tanah. Walau bagaimanapun, kombinasi air laut dengan racun rumpai kadar rendah didapati lebih mesra alam berbanding pilihan lain.

Kajian kepekatan terhadap kombinasi air laut dan racun rumpai kadar rendah telah dilakukan ke atas spesies rumpai *E. atrovirens*, *S. diander*, *C. aromaticus*, *C. rotundus* and *E. sonchifolia* dan rumput turf *P. vaginatum*, *C. dactylon* ‘satiri’, and *Z. japonica*. Keputusan kajian menunjukkan kesemua spesies rumpai kecuali *E. atrovirens* dapat dikawal sepenuhnya apabila dirawat dengan campuran $\frac{3}{4}$ daripada kadar disyorkan racun rumpai trifloxsulfuron–sodium bersama air laut, $\frac{3}{4}$ daripada kadar disyorkan racun rumpai trifloxsulfuron–sodium bersama $\frac{3}{4}$ kadar air laut, $\frac{1}{2}$ daripada kadar disyorkan racun trifloxsulfuron–sodium bersama air laut, $\frac{3}{4}$ daripada kadar disyorkan racun rumpai quinclorac bersama air laut dan $\frac{3}{4}$ daripada kadar disyorka racun rumpai quinclorac bersama $\frac{3}{4}$ air laut. *Eragrostis atrovirens* menunjukkan kecederaan paling maksima 48% apabila dirawat dengan $\frac{3}{4}$ daripada kadar disyorkan racun rumpai trifloxsulfuron–sodium bersama air laut. Di antara spesies rumput turf yang telah diuji, *P. vaginatum* menunjukkan ketahanan yang tinggi terhadap air laut dan kombinasi air laut dan $\frac{3}{4}$ daripada kadar racun rumpai quinclorac disyorkan, dengan nilai kecederaan paling maksimum hanya 8%. *Zoysia japonica* pula menunjukkan kecederaan yang tidak lebih dari 14% apabila di rawat dengan air laut atau kombinasi air laut bersama $\frac{3}{4}$ daripada kadar racun rumpai trifloxsulfuron–sodium atau quinclorac disyorkan. Kecederaan maksimum iaitu 21% telah dicatatkan oleh *C. dactylon* ‘satiri’ dengan kombinasi $\frac{3}{4}$ kadar air laut dan $\frac{3}{4}$ kadar racun rumpai trifloxsulfuron–sodium disyorkan.

ACKNOWLEDGEMENTS

All praise be to almighty ALLAH, The Exalted.

I would like to express my sincere gratitude to Associate Professor Dr. Abdul Shukor Juraimi, the Chairman of the Supervisory Committee for his invaluable supervision, productive discussion, constructive criticisms and continuous encouragement throughout the study. I am much grateful and indebted to Professor Dr. Mohd Razi Ismail, Associate Professor Dr. Radziah Othman and Associate Professor Dr. Anuar Abd. Rahim, members of the supervisory committee, for their constructive suggestions and guidance in all phases of this research project. I express my deep sense of respect to all of the lecturers and professors in the Department of Crop Science for their valuable suggestions and constructive comments.

I would like to thank En. Mohd Yunos Abdul Wahab for his assistance in glasshouse and field works. I am grateful to the administrative staff of the Department of Crop Science, Soil Microbiology laboratory and Department of Land Management Faculty of Agriculture.

I am indebted to my late parents, wife, daughters, brothers and sisters for their support, deep love and tolerance during the whole period of the study.

Thankfully acknowledge the assistance of Graduate Research Fellowship for financial support and facilitating my study and University Putra Malaysia for providing research facilities under the Science fund project under Ministry of Agriculture: 02-01-04-SF0169.

I certify that an Examination Committee has met on 2 September, 2010 date of viva voce to conduct the final examination of Md. Kamal Uddin on his Doctor of Philosophy thesis entitled “Using Seawater for Weed Management in Turfgrass” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U. (A) 106] 15th March 1998. The Committee recommends that the student be awarded the Doctor of Philosophy.

Members of the Examination Committee were as follows:

Adam b Puteh, PhD

Associate Professor

Department of Crop Science

Faculty of Agriculture

Universiti Putra Malaysia

(Chairman)

Rosli b Mohamad, PhD

Professor

Department of Crop Science

Faculty of Agriculture

Universiti Putra Malaysia

(Internal Examiner)

Yahya bin Awang, PhD

Associate Professor

Department of Crop Science

Faculty of Agriculture

Universiti Putra Malaysia

(Internal Examiner)

Jahid Ata Cheema, PhD

Professor

Department of Agronomy

University of Agriculture, Faisalabad

Pakistan

(External Examiner)

SHAMSUDDIN SULAIMAN, PhD

Professor and Deputy Dean

School of Graduate Studies

Universiti Putra Malaysia

Date: 26 November 2010



This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement of the degree of Doctor of Philosophy. The members of the Supervisory Committee were as follows:

Abdul Shukor Juraimi, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Chairman)

Mohd Razi Ismail, PhD

Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Member)

Anuar Abd. Rahim, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Member)

Radziah Othman, PhD

Associate Professor

Faculty of Agriculture

Universiti Putra Malaysia

(Member)

HASANAH MOHD GHAZALI, PhD

Professor and Dean

School of Graduate Studies

Universiti Putra Malaysia

Date:



DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any other degree at Universiti Putra Malaysia or other institutions.

MD. KAMAL UDDIN

Date:



TABLE OF CONTENTS

	Page
ABSTRACT	ii
ABSTRAK	vi
ACKNOWLEDGEMENTS	x
APPROVAL	xi
DECLARATION	xiii
LIST OF TABLES	xx
LIST OF FIGURES	xxiii
GLOSSARY OF ABBREVIATIONS	xxiv

CHAPTER

1 INTRODUCTION	1
2 LITERATURE REVIEW	5
2.0 Importance of turfgrass	5
2.1 Natural and taxonomic distribution of turfgrasses	6
2.2 Morphology of turfgrass species	7
2.3 Effect of weeds on growth and quality of turfgrass	8
2.4 Turf grasses in Malaysia and their common weeds	9
2.5 Weed control methods in turfgrass	9
2.5.1 Cultural practices	11
2.5.2 Mechanical (Physical) Control	11
2.5.3 Chemical control	13
2.5.4 Biological control	13
2.6 General aspects of salinity problems in plants	14
2.7 Salinity in water and soil	15
2.8 Tuning mechanisms in plants under saline environment	16
2.9 Salt tolerant turfgrass species	18
2.10 Effect of salinity on turfgrass morphology	19
2.11 Effect of salinity on Turfgrass physiology	20
2.12 Effect of salinity on proline content	21
2.13 Description of trifloxisuluron-sodium and quinclorac herbicide	22
2.14 Use of sea water for weed control in turfgrass species	23
2.15 Use of herbicides for weed control in turfgrass	24
2.16 Effect of salinity on microbial populations in soil	26
2.17 Effect of herbicides on soil microorganisms	28

3 SCREENING OF POTENTIAL SALT TOLERANT TURFGRASS SPECIES

3.0	Introduction	29
3.1	Materials and Methods	30
3.1.1	Site description	30
3.1.2	Duration of the experiment	30
3.1.3	Selection of turfgrass species	30
3.1.4	Soil properties and media preparation	31
3.1.5	Planting material collection and turfgrass establishment	32
3.1.6	Cultural practices	32
3.1.7	Salt media preparation and salinity treatment	33
3.1.8	Layout of the experiment	33
3.1.9	Determination of leaf firing and turf quality	33
3.1.10	Relative (%) shoot and root dry weight	34
3.1.11	Determination of chlorophyll content	34
3.1.12	Statistical analysis	35
3.2	Results	35
3.2.1	Leaf firing	35
3.2.2	Turf quality	37
3.2.3	Shoot dry weight	38
3.2.4	Root dry weight	41
3.2.5	Chlorophyll-a	44
3.2.6	Chlorophyll-b	44
3.2.7	Total chlorophyll	45
3.2.8	Chlorophyll a/b ratio	46
3.3	Discussion	47
3.4	Conclusion	49

4 SURVEY OF WEED SPECIES IN TURFGRASS AREAS IN MALAYSIA

4.0	Introduction	51
4.1	Materials and Methods	52
4.1.1	Sites surveyed	52
4.1.2	Sampling scheme	54
4.1.3	Data collected	54
4.1.4	Community coefficient	57
4.1.5	Statistical Methods	57
4.2	Results	58
4.2.1	Weed species taxonomy	58
4.2.2	Variation in weed populations by turfgrass area	58
4.2.3	Species frequency	64
4.2.4	Distribution uniformity	65
4.2.5	Site density	70
4.2.6	Relative abundance	70
4.2.7	Community coefficients	70
4.3	Discussion	74

4.4	Conclusion	78
5	SCREENING OF TURF WEED SPECIES FOR SALT INJURY	
5.0	Introduction	79
5.1	Materials and Methods	80
5.1.1	Site description	80
5.1.2	Experimental period	80
5.1.3	Weed species used in the experiment	80
5.1.4	Soil media preparation	82
5.1.5	Planting material collection and establishment	83
5.1.6	Cultural practices	83
5.1.7	Salt media preparation and salinity treatments	83
5.1.8	Evaluation of visual plant injury	84
5.1.9	Determination of shoot dry weight	84
5.1.10	Determination of root dry weight	84
5.1.11	Statistical analysis	84
5.2	Results	85
5.2.1	Effect of salt water on plant injury	85
5.2.2	Effect of salinity treatments on shoot dry weight	98
5.2.3	Effect of salinity treatments on root dry weight	101
5.3	Discussion	104
5.4	Conclusion	109
6	COMPARATIVE STUDY ON PHYSIOLOGICAL AND GROWTH RESPONSES OF TURFGRASS CULTIVARS TO SALINITY	
6.0	Introduction	110
6.1	Materials and Methods	111
6.1.1	Site description	111
6.1.2	Duration of the experiment	111
6.1.3	Selection of test species	111
6.1.4	Soil media preparation	112
6.1.5	Collection and establishment of turf grass species	112
6.1.6	Cultural practices	112
6.1.7	Salt media preparation and salinity treatment	113
6.1.8	Layout of the experiment	113
6.1.9	Determination of leaf firing	113
6.1.10	Determination of shoot dry weight	113
6.1.11	Determination of root dry weight	114
6.1.12	Determination of chlorophyll content	114
6.1.13	Determination of relative water content	115
6.1.14	Determination of proline content	115
6.1.15	Root histology using scanning electron microscopy	116
6.1.16	Chemical analysis	116
6.1.17	Statistical analysis	117
6.2	Results	117
6.2.1	Leaf firing (%)	118

6.2.2	Shoot dry weight	119
6.2.3	Root dry weight	119
6.2.4	Leaf proline content	120
6.2.5	Leaf Relative Water Content (RWC)	122
6.2.6	Leaf Chlorophyll content	123
6.2.7	Shoot sodium concentration (Na)	125
6.2.8	Shoot potassium concentration (K)	126
6.2.9	Shoot potassium/sodium (K/Na) ratio	128
6.2.10	Shoot calcium concentration (Ca)	129
6.2.11	Shoot magnesium concentration (Mg)	130
6.2.12	Pearson correlation matrix	132
6.2.13	Root sodium concentration (Na)	132
6.2.14	Root potassium concentration (K)	133
6.2.15	Root potassium/sodium (K/Na) ratio	135
6.2.16	Root calcium concentration (Ca)	135
6.2.17	Root magnesium concentration (Mg)	136
6.2.18	Root cell histology	137
6.3	Discussion	138
6.4	Conclusion	143
7	EFFECT OF SEA WATER WITH REDUCED HERBICIDE RATES ON SALT TOLERANT TURFGRASS WEED SPECIES AND SOIL MICROBIAL COMMUNITIES IN TURFGRASS	
7.0	Introduction	144
7.1	Materials and Methods	146
7.1.1	Glasshouse experiment	146
7.1.2	Field experiment	149
7.2	Results	156
7.2.1	Glasshouse study – Trifloxsulfuron-sodium	156
7.2.2	Glasshouse study - Quinclorac	173
7.2.3	Field study	187
7.3	Discussion	220
7.4	Conclusion	226
8	GENERAL DISCUSSION AND RECOMMENDATIONS	227
REFERENCES		236
APPENDICES		256
BIODATA OF STUDENT		270