

Improved mannan-degrading enzymes' production by *Aspergillus niger* through medium optimization.

ABSTRACT

The effect of different carbon and nitrogen sources on the production of mannan-degrading enzymes, focussing on β -mannanase, by *Aspergillus niger* was investigated using shake flask culture. The β -mannanase activity obtained during growth of *A. niger* on guar gum (GG, 1495 nkat mL⁻¹) was much higher than those observed on other carbon substrates, locust bean gum (1148 nkat mL⁻¹), α -cellulose (10.7 nkat mL⁻¹), glucose (8.8 nkat mL⁻¹) and carboxymethylcellulose (4.6 nkat mL⁻¹). For fermentation using GG as a carbon source, bacteriological peptone gave the highest β -mannanase activity (1744 nkat mL⁻¹) followed by peptone from meat (1168 nkat mL⁻¹), yeast extract (817 nkat mL⁻¹), ammonium sulphate (241 nkat mL⁻¹), ammonium nitrate (113 nkat mL⁻¹) and ammonium chloride (99 nkat mL⁻¹) when used as a nitrogen source. The composition of bacteriological peptone and initial pH of the medium were further optimized using response surface methodology (RSM). Medium consisted of 21.3 g L⁻¹ GG and 57 g L⁻¹ peptone with initial culture pH of 5.5 was optimum for β -mannanase production (2063 nkat mL⁻¹) by *A. niger*. The β -mannanase production obtained in this study using *A. niger* was significantly higher than those reported in the literature.

Keyword: *Aspergillus niger*; Hemicellulose; Mannanase; Carbon and nitrogen; Carbon substrates.