

Breadfruit starch-wheat flour noodles: preparation, proximate compositions and culinary properties

¹*Akanbi, T.O., ¹Nazamid, S., ²Adebowale, A.A., ¹Farooq, A. and ³Olaoye, A.O.

¹Department of Food Science, Faculty of Food Science and Technology, Universiti Putra Malaysia, 43400 Serdang, Selangor, Malaysia

²Department of Food Science, University of Agriculture, Abeokuta, Ogun State, Nigeria

³Division of Food Sciences, School of Biosciences, University of Nottingham, Sutton Bonnington Campus, Leicester, United Kingdom

Abstract: Proximate compositions, culinary and sensory properties of noodles prepared from proportionate combinations of breadfruit starch and wheat flour were investigated. Breadfruit starch (BS) isolated from matured breadfruit (*Artocarpus altilis*) was used to produce noodles in combination with hard red wheat flour (WF) at a ratio of 100% WF:0% BS, 80% WF:20% BS, 60% WF:40% BS, 40% WF:60% BS, 20% WF:80% BS. The protein, fat, ash, crude fibre and moisture contents of the Breadfruit starch-Wheat flour (BSWF) noodles prepared from the above blends ranged from 0.65 to 10.88%, 0.35 to 3.15%, 1.28 to 2.25%, 1.18 to 1.45% and 4.65 to 5.45%, respectively. The contents of protein, fat, ash and crude fibre increased as the percentage breadfruit starch decreased. However, values of moisture content did not follow the same trend, instead higher values were found for 100% BS:0% WF (5.35%) and 20% BS:80% WF (5.45%). The cooking yield of the BSWF noodles ranged from 21.02 (60% BS:40% WF) to 23.75 g (100% BS:0% WF), cooking loss ranged from 5.49 (20% BS:80% WF) to 9.19% (100% BS:0% WF), while swelling index ranged from 3.1 (20% BS:80% WF) to 3.4 (100% BS:0% WF). Throughout the study, noodles produced from blends of 20% breadfruit starch and 80% wheat flour showed superior proximate, culinary and sensory attributes.


Keywords: Breadfruit starch, wheat flour, noodle preparation, culinary properties, sensory evaluation

Introduction

Breadfruit is found in many parts of the world, especially in the tropical countries. Being a rich source of carbohydrates, lipids and proteins, some varieties have been studied and valued for their nutritional properties (Rincon *et al.*, 2004). Breadfruit (Figure 1) has been processed into many forms such as starch (Akanbi *et al.*, 2009) and flour (Olaoye *et al.*, 2008) for its utilization in the food industries. Starches are of great importance in the food industry. They have been used as thickeners or filler in products like custard, soup, baby foods, ice cream and pharmaceuticals (Ihekoronye and Ngoddy, 1985). They have also been used in salad dressings (Singhal and Kulkarni, 1990), cookies preparation (Kulp *et al.*, 1991), coating in dehydrated fruits, thickening in pie filling, soup mixer and instant puddings (Ihekoronye and Ngoddy, 1985).

Wheat flour has been extensively used for producing alimentary pastes such as macaroni, spaghetti and other noodle forms. Noodles are important foods throughout the world especially in

Asian countries such as China, Korea, Malaysia, the Philippines and Thailand. Noodles can be generally classified into Chinese types (Ramen), Japanese types (Soba), Korean types (Naengmyon), rice and starch noodles (Nagao, 1996). However, no information is available on the direct use of breadfruit starch for the production of noodles either as major raw material or in combination with other raw materials such as wheat flour.



Source: www.healthalternatives2000.com

Source: www.australian-tropical-foods.com

Figure 1. Typical breadfruit (*Artocarpus altilis*)

Starch (basically amylose and amylopectin) is a predominant component of wheat flour, as it helps to improve the appearance and structure of wheat flour foods. Breadfruit starch has also been reported to contain high amylose (22.52%) and amylopectin

*Corresponding author.

Email: akanbitaiwo@gmail.com

(77.48%) (Akanbi *et al.*, 2009). Studies have shown that moderately high amylose foods are helpful in reducing risk factors for diabetes and cardiovascular diseases (Behall and Howe, 1995). High amylopectin presence in food has also been reported to increase human insulin levels (Behall *et al.*, 1988). A study that was approved by the Human Studies Committees of the United States Department of Agriculture (USDA) and Georgetown University showed that the insulin levels of 12 women increased after being fed with cornstarch crackers containing 70% amylopectin (Behall *et al.*, 1988). Furthermore, diets high in complex carbohydrates including high levels of starches according to Behall *et al.* (1989) have been reported to normalize blood insulin and lipid levels in carbohydrate sensitive, diabetic, and hyperlipidemic individuals. Therefore, noodles produced from the blends of breadfruit starch and wheat flour with moderately high amylose and amylopectin might be explored as functional foods for normalizing the blood insulin levels and imparting other health benefits. Hence, the objective of this study is to produce noodles from breadfruit starch-wheat composite flour and investigate their proximate, culinary and sensory properties.

Materials and Methods

Materials

Breadfruit starch was previously isolated from breadfruits obtained from Ile-Ife Osun State, Nigeria (Akanbi *et al.*, 2009) and hard red wheat flour was obtained from Honey-well Flour Mill, Lagos, Nigeria.

Preparation of breadfruit starch – wheat flour Noodles

Breadfruit starch-wheat flour noodles were prepared using Chen *et al.* (2002) method. Following the combinations presented in Table 1, a total of 120 g of composite flour (breadfruit starch-wheat flour) was weighed. Carefully weighed 12 g of the composite flour was put into a clean beaker, mixed with 10 ml warm water (55°C) and then stirred completely after adding 24 ml of boiling water. The remaining 108 g of composite flour and 52 ml of warm water were added and stirred for 30 min at 60°C. The mixture was then kneaded with water to dough consistency before the noodle was injected using an “Extruder”– Lab-scale Cylindrical Extruder (100 – 500 g dough capacity) through the holes (1.5 cm diameter) of the stainless steel cylinder by gravity, using a well fitted 2.25 kg stainless steel piston into hot water (95 – 98°C), and heated for 50-70 seconds at this temperature before

transferring into cold water. When noodles were floated, they were taken out and dried. No additive was added during the whole procedure. The dried noodles were equilibrated at room temperature for 4 h, packed in clean pre-sterilized polyethylene bags and stored at room temperature before further analyses.

Table 1. Proximate composition of breadfruit starch - wheat flour noodles

Sample	Protein (%)	Fat (%)	Ash (%)	Fibre (%)	Moisture Content (%)
100% BS:0% WF	0.65 ^a ± 0.3	0.35 ^a ± 0.2	1.28 ^a ± 0.4	1.18 ^a ± 0.2	5.45 ^a ± 0.3
80% BS:20% WF	4.88 ^a ± 0.7	0.88 ^a ± 0.7	1.96 ^a ± 0.3	1.20 ^a ± 0.7	4.97 ^a ± 0.5
60% BS:40% WF	6.75 ^a ± 0.2	2.55 ^a ± 0.1	2.09 ^a ± 0.7	1.22 ^a ± 0.3	4.65 ^a ± 0.6
40% BS:60% WF	9.55 ^b ± 0.5	2.64 ^b ± 0.6	2.18 ^a ± 0.2	1.28 ^b ± 0.6	4.87 ^a ± 0.3
20% BS:80% WF	10.88 ^b ± 0.6	3.15 ^b ± 0.4	2.25 ^a ± 0.3	1.45 ^a ± 0.0	5.35 ^b ± 0.4

Results are means of three independent determinations. Mean values having different superscripts within the column are significantly different (p<0.05)
BS – Breadfruit Starch
WF – Wheat Flour

Proximate compositions of Breadfruit starch-wheat flour noodles

Moisture, ash, fibre and fat contents of the noodles were determined by the AOAC (1990) official methods and crude protein content by Kjeldahl method. Meanwhile, the proximate properties of the breadfruit starch have been previously determined (Akanbi *et al.*, 2009).

Culinary properties of breadfruit starch-wheat flour noodles

Cooking yield and cooking loss

Cooking yield and cooking loss of the noodles were determined as described in the AACC (American Association of Cereal Chemists, 2000) method. Ten grams of the noodles were added to a beaker containing about 150 ml boiling water. The beaker was covered with a watch glass and noodles cooked for 10 min with slight agitation. The cooked noodles were allowed to drain for 5 min and then weighed. The cooking yield was then calculated. The gruel was poured into a 200 ml volumetric flask and adjusted to volume with distilled water. Ten millilitre of the solution was pipetted into an aluminium dish and dried to a constant weight at 105°C. The cooking loss during cooking was calculated as given below.

$$\text{Cooking yield (g)} = \text{Weight of Noodles after cooking (g)} - \text{Weight before cooking (g)}$$

$$\text{Cooking loss (\%)} = \frac{\text{Weight of gruel and dish - Constant weight after drying} \times 100}{\text{Constant weight after drying}}$$

$$\text{Swelling index} = \frac{\text{Weight after cooking (g)}}{\text{Weight before cooking (g)}}$$

Sensory evaluation of breadfruit starch – wheat flour noodles

Uncooked noodles

Panellists were presented with five (5) samples of uncooked noodles and a reference sample “R” (the reference sample was a commercial brand of wheat noodles) to compare for appearance in terms of colour and also to compare for breakage, indicating the amount of difference that exists between the samples and the reference. A scale of 1 to 5 was used where 1 represents better than R and 5 represents extremely poorer than R.

Cooked noodles

Panellists were presented with five (5) samples of cooked noodles and a reference (R) to compare for colour, slipperiness, glossiness, taste, aroma and texture, predicting the difference that exists between the samples and the reference.

Statistical Analysis

All experiments were run in triplicate and data analyzed statistically using Microsoft Excel and SPSS v. 12.0.

Results and Discussion

Proximate properties of breadfruit starch - wheat flour noodles

Proximate properties of BSWF noodles are shown in Table 1. The proximate properties of all noodle samples varied significantly ($p < 0.05$) different. The mean protein and fat contents of BSWF noodles ranged from 0.65 (100% BS:0% WF) to 10.88% (20% BS:80% WF) and 0.35 (100% BS:0% WF) to 3.15% (20% BS:80% WF), respectively. This trend portends that the higher the percentage wheat flour in the blend, the higher the protein content. Results obtained for the protein and fat contents are not surprising because the protein and fat contents of wheat flour (Olaoye *et al.*, 2008) exceed those of breadfruit starch (Akanbi *et al.*, 2009). The result also supports the findings of Iwaoka *et al.* (1994) who reported that starch has very low fat content. The ash contents ranged from 1.28 (100% BS:0% WF) to 2.25% (20% BS:80% WF) which might be due to the fact that breadfruit starches are high in ash (Adebowale *et al.*, 2004; Akanbi *et al.*, 2009). The content of crude fibre was noted to be increased as the percentage wheat flour increased, ranging from 1.18 ± 0.00 (100% BS:0% WF) to 1.45 ± 0.00 (20% BS:80% WF). Olaoye *et al.* (2008) reported similar results for crude fibre content of blends of

wheat and breadfruit flour used for the production of baked products. Studies have shown that crude fibre is best obtained from foods than supplement and can reduce symptoms of chronic constipation (Odes *et al.*, 1993), heart diseases associated with high cholesterol (hyperlipidemia) (Odes *et al.*, 1993), diverticular disease (Gear *et al.*, 1979), and reduce the risk of colon cancer (Reddy, 1999). Hence, BSWF noodles could be acceptable in places where fibre diets and lower fatty foods are desired. Meanwhile, the moisture content of the noodles ranged from 4.65 (60% BS:40% WF) to 5.45% (100% BS:0% WF). These values are far lower than the Joint FAO/WHO standards for fried and non-fried instant noodles, which are 10 and 14%, respectively (FAO/WHO, 2003). An inference can therefore be drawn that BSWF noodles apart from being healthy, free of additives, also conform to the standards of world food and health regulatory agencies.

Culinary properties of breadfruit starch-wheat flour noodles

The results of culinary properties of BSWF noodles are presented in Table 2. There were significant ($P < 0.05$) differences observed in the cooking yield, cooking loss and swelling index of the noodles tested. The highest cooking yield of 23.75% was obtained from 100% BS:0% WF, while the lowest being 21.02% was from 60% BS:40% WF. The mean values of cooking loss ranged from 5.49 (20% BS:80% WF) to 9.19% (100% BS:0% WF). Meanwhile, the mean swelling indexes ranged from 3.04 (20% BS:80% WF) to 3.4 (100% BS:0% WF).

Table 2. Culinary properties of breadfruit starch – wheat flour noodles

Samples	Cooking yield (g)	Cooking Loss (%)	Swelling Index
100% BS:0% WF	23.75 ^a ± 0.4	9.19 ^a ± 0.4	3.40 ^a ± 0.5
80% BS:20% WF	22.00 ^b ± 0.5	8.17 ^b ± 0.5	3.28 ^b ± 0.5
60% BS:40% WF	21.02 ^c ± 0.4	6.23 ^c ± 0.5	3.17 ^b ± 0.3
40% BS:60% WF	22.03 ^b ± 0.3	6.01 ^d ± 0.3	3.11 ^c ± 0.5
20% BS:80% WF	21.27 ^c ± 0.2	5.49 ^c ± 0.3	3.04 ^d ± 0.2

Results are means of three independent determinations. Mean values having different superscripts within the column are significantly different ($p < 0.05$)
BS – Breadfruit Starch
WF – Wheat Flour

Sensory evaluation of breadfruit starch-wheat flour noodles

Tables 3 and 4 show the results obtained for the sensory evaluation of the uncooked and cooked BSWF noodles compared to the reference sample (R). The lower the mean values, the closer the noodles in attributes to the reference sample. There were significant ($p < 0.05$) differences observed for the colour and colour difference among the uncooked noodles (Table 3). The least mean scores for colour

Table 3. Mean sensory scores of uncooked breadfruit starch –wheat flour noodles

Sample	Colour	Colour difference	Ease of breakage	Ease of breakage difference
100% BS:0% WF	3.0 ^a ± 0.3	4.7 ^a ± 0.4	2.3 ^a ± 0.2	2.9 ^a ± 0.2
80% BS:20% WF	3.0 ^a ± 0.4	4.2 ^b ± 0.3	2.3 ^a ± 0.3	2.9 ^a ± 0.1
60% BS:40% WF	2.9 ^a ± 0.4	3.8 ^c ± 0.5	2.2 ^b ± 0.1	2.6 ^b ± 0.3
40% BS:60% WF	2.9 ^a ± 0.5	3.6 ^{bc} ± 0.3	2.0 ^c ± 0.5	2.4 ^c ± 0.6
20% BS:80% WF	2.5 ^b ± 0.2	3.2 ^d ± 0.5	1.8 ^d ± 0.2	2.3 ^d ± 0.5

Results are means of three independent determinations. Mean values having different superscripts within the column are significantly different ($p < 0.05$)

BS – Breadfruit Starch
WF – Wheat Flour

Table 4. Mean sensory scores of cooked breadfruit starch-wheat flour noodles

Sample	Colour	Colour difference	Slipperiness	Slipperiness difference	Gloss	Gloss difference	Taste	Taste difference	Aroma
100% BS:0% WF	3.0 ^a ± 0.5	4.9 ^a ± 0.8	1.4 ^d ± 0.5	1.5 ^c ± 0.6	1.4 ^d ± 0.5	1.4 ^d ± 0.5	3.0 ^a ± 0.5	4.9 ^a ± 0.8	3.0 ^a ± 0.7
80% BS:20% WF	3.0 ^a ± 0.3	4.9 ^a ± 0.8	2.0 ^e ± 0.7	2.3 ^d ± 0.7	2.0 ^c ± 0.7	2.3 ^c ± 0.7	3.0 ^a ± 0.4	4.6 ^a ± 0.6	3.0 ^a ± 0.5
60% BS:40% WF	3.0 ^a ± 0.3	4.5 ^b ± 0.6	2.3 ^b ± 0.6	2.7 ^c ± 0.8	2.2 ^{bc} ± 0.7	2.8 ^b ± 0.6	2.9 ^{ab} ± 0.3	4.3 ^b ± 0.4	2.9 ^a ± 0.4
40% BS:60% WF	2.9 ^a ± 0.4	4.2 ^c ± 0.4	2.5 ^{ab} ± 0.4	3.4 ^b ± 0.7	2.5 ^{ab} ± 0.5	3.4 ^a ± 0.5	2.8 ^b ± 0.5	3.9 ^c ± 0.3	2.4 ^b ± 0.7
20% BS:80% WF	2.5 ^b ± 0.2	3.4 ^d ± 0.5	2.7 ^a ± 0.6	3.7 ^a ± 0.8	2.7 ^a ± 0.5	3.7 ^a ± 0.8	2.4 ^c ± 0.7	3.0 ^d ± 0.7	2.0 ^c ± 0.3

Results are means of three independent determinations. Mean values having different superscripts within the column are significantly different ($p < 0.05$)

BS – Breadfruit Starch
WF – Wheat Flour

(2.5) and colour difference (3.2) were obtained for 20% BS:80% WF, while the highest scores (3.0 for colour and 4.7 for colour difference) were observed for 100% BS:0% WF. This indicates that as the percentage breadfruit starch decreases, the noodle's colour improves. Similar trends were observed for ease of breakage and ease of breakage difference although with different mean scores. There was significant ($p < 0.05$) difference in all sensory parameters for the cooked noodles (Table 4). For slipperiness, slipperiness difference, glossiness and glossiness difference, the lowest mean values were found to be 100% BS:0% WF. In terms of taste, taste difference and aroma, the mean scores decreased as the percentage breadfruit starch decreased, with 20% BS:80% WF having the lowest score. This implies that with reference to "R", Noodles from 20% BS:80% WF had better taste and aroma compared to the remaining samples.

Conclusion

The results of the present study revealed that beneficial noodles with health promoting factors can be produced from blends of wheat flour and breadfruit starch. Throughout the study, noodles produced from blends of 20% breadfruit starch and 80% wheat flour showed superior proximate, culinary and sensory attributes. Since both breadfruit starch and wheat flour are high in crude fibre which has been reported to reduce symptoms of chronic constipation, heart diseases associated with high cholesterol, diverticular disease and risk of colon cancer, therefore, an inference can be drawn that BSWF noodles are more than just noodles, but important functional foods.

Acknowledgments

Appreciation goes to all the staff of food processing laboratory of the University of Agriculture Abeokuta (UNAAB). Also, big thanks go to Mr. and Mrs. P.A. Akanbi (JP), Mrs. Abidemi Odelade (Nee Akanbi) and Mrs. Kehinde Akanbi (Nee Olunlade) for their financial and moral supports.

References

- AACC. Approved methods of the American association of cereal chemists (10th ed.), 2000. St. Paul, MN: American Association of Cereal Chemists.
- Adebawale, K.O., Olu-Owolabi, B.I., Olawumi, E.K. and Lawal O.S. 2004. Functional properties of native, physically and chemically modified breadfruit (*Artocarpus altilis*) starch. *India Crops and Products* 21: 343–351.
- Akanbi, T.O., Nazamid, S. and Adebawale, A.A. 2009. Functional and pasting properties of a tropical breadfruit (*Artocarpus altilis*) starch from Ile-Ife, Osun State, Nigeria. *International Food Research Journal* 16:151-157.
- AOAC Official Methods of Analysis, 15th edition. 1990. Association of Official Analytical Chemists (AOAC), Washington D.C.
- Behall, K.M. and Howe, J.C. 1995. Effect of long-term consumption of amylose vs amylopectin starch on metabolic variables in human subjects, *American Journal of Clinical Nutrition* 61:334–340.
- Behall, K.M., Scholfield, D.J. and Canary, J. 1988. Effect of starch structure on glucose and insulin responses in adults, *American Journal of Clinical Nutrition* 47:428–432.
- Behall, K.M., Scholfield, D.J., Yuhaniak, I. and Canary, J. 1989. Diets containing high amylase vs amylopectin starch: effects on metabolic variables in human

- subjects, *American Journal of Clinical Nutrition* 49: 337 – 344.
- Chen, B., Fu, X.G. and Lu, D.L. 2002. Improvement of predicting precision of oil content in instant noodles by using wavelet transforms to treat near-infrared spectroscopy, *Journal of Food Engineering* 53: 373–376.
- Chowdhury, F.A., Raman, Md. A. and Mian, A.J. 1997. Distribution of free sugars and fatty acids in jackfruit (*Artocarpus heterophyllus*). *Food Chemistry* 60:25-28.
- FAO/WHO Food Standards Programme CODEX Committee Alimentarius Commission, Rome Italy, 2003.
- Gear, J.S.S., Fursdon, P., Nolan, D.J., Ware, A., Mann, J.I., Brodribb, A.J.M. and Vessey, M.P. 1979. Symptomless diverticular disease and intake of dietary fibre. *The Lancet* 313(8115): 511-514.
- Ihekoronye, A.I. and Ngoddy, P.O. 1985. *Integrated Food Science and Technology for the Tropics*. Macmillan Publishers Ltd., London and Basingstoke. pp. 15-22
- Iwaoka, W., Yukio, H., Kasumi, U. and Takayuki, S. 1994. Volatile chemicals identified in fresh and cooked breadfruit. *Journal of Agriculture and Food Chemistry* 42(4): 975-976.
- Kasemsuwa, T., Bailey, T. and Jane, J. 1998. Preparation of clear noodles with mixtures of tapioca and high amylose starches. *Carbohydrate Polymer* 32: 301-312.
- Kim, Y.S. and Wiesenborn, D.P. 1996. Starch noodle quality as related to potato genotypes. *Journal of Food Science* 61:248-252.
- Kim, Y.S., Wiesenborn, D.P., Lorenzen, J.H. and Berglund, P. 1996. Suitability of edible bean and potato starches for starch noodles. *Cereal Chemistry* 73:302-308.
- Kulp, K., Olewink, M., Manhattan, K. and Lorenz, K. 1994. Starch functionality in cookie system. *Starch and Starke* 42 (4):53-57.
- Nagao, S. 1996. Processing technology of noodle products in Japan. In: *Pasta and Noodle Technology*. J. E. Kruger, R. B. Matsuo, and J. W. Dick, eds. American Association of Cereal Chemistry; St. Paul, MN, 169-194.
- Odes, H.S, Lazovski, H., Stern, I. and Madar, Z. 1993. Double-blind trial of a high dietary fiber, mixed grain cereal in patients with chronic constipation and hyperlipidemia. *Nutrition Research* 13(9): 979-985.
- Olaoye, O.A. and Onilude, A.A. 2008. Microbiological, proximate analysis and sensory evaluation of baked products from blends of wheat-breadfruit flours. *African Journal of Food, Agriculture, Nutrition and Development* 8: 192-203.
- Olatunji, O. and Akerele, A.J. 1978. Comparative rheological properties and bread qualities of wheat flour diluted with tropical tuber and Breadfruit flour. *Journal of Cereal Chemistry* 55(1): 1-6.
- Ready, B.S. 1999. Role of dietary fiber in colon cancer: an overview. *The American Journal of Medicine*, 106(1):16-19
- Rincon, F., Ros, G. and Collins, J.L. 1993. Mineral loss in cowpeas (*Vigna unguiculate* L.) by pressure heating in water. *Journal of Food Science* 58: 856–859.
- Singh, A., Kumar, S. and Singh, I.S. 1991. Functional properties of jackfruit seed flour. *Lebensmittel Wissenschaft und Technologie* 24: 373–374