



UNIVERSITI PUTRA MALAYSIA

***INCREASING RESISTANT STARCH CONTENT IN FISH CRACKERS
THROUGH VARIOUS COOKING-CHILLING CONDITIONS***

MOHD ZUHAIR BIN MOHD NOR

FK 2012 4

**INCREASING RESISTANT STARCH CONTENT IN FISH CRACKERS
THROUGH VARIOUS COOKING-CHILLING CONDITIONS**



By

MOHD ZUHAIR BIN MOHD NOR

**Thesis Submitted to the School of Graduates Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

August 2012

DEDICATION

To my beloved wife, Intan Syafinaz;

My family members and friends.

Thanks for your encouragement, patience and loving support.



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfilment of the partial requirement for the degree of Master of Science

**INCREASING RESISTANT STARCH CONTENT IN FISH CRACKERS
THROUGH VARIOUS COOKING-CHILLING CONDITIONS**

By

MOHD ZUHAIR BIN MOHD NOR

August 2012

Chairman : Rosnita A. Talib, PhD

Faculty : Engineering

Fish cracker is one of the favourite snacks in Malaysia. However, studies on resistant starch (RS) in it have been paid little attention. Resistant starch is a starch that goes through the small intestine without being digested, providing a similar effect as dietary fibre. This RS can potentially be formed during the cooking and chilling processes in the production of the fish cracker. Both processes involve starch gelatinization and retrogradation that can lead to the formation of RS, creating a healthy and valuable product. Thus, this work investigates the effect of varying cooking-chilling conditions on resistant starch (RS) content and other important quality characteristics in fish cracker products. Process conditions such as the number of repetitive cooking-chilling (RCC) cycles (1 to 4 cycles), cooking temperature (100 °C, 115 °C and 121 °C) and chilling duration (1 to 4 days) as well as the type of starch used in the formulation (tapioca, wheat and sago) were studied with the aim to enhance the fish crackers with an

appreciable amount of resistant starch without compromising the quality characteristics. The quality characteristics of fish crackers in terms of hardness and moisture content of the chilled fish cracker gels, the RS of the dried fish crackers as well as the linear expansion (LE), hardness and colour of the fried fish crackers were evaluated.

The results showed that up to four cycles of RCC increased the RS content in all products. Sago starch fish crackers cooked at 121 °C had the highest RS content in the dried samples. The repeated cooking-chilling cycles increased the extent of starch gelatinization with each successive cooking cycle and promoted retrogradation upon cooling, thus, promoting the formation of RS. However, initially cooking the fish crackers at 100 °C and exposing to a longer chilling duration of up to four days did not demonstrate any changes in the RS. Different combinations of cooking-chilling repetition, cooking temperatures and chilling durations produced varying RS content in the fish crackers.

As for other quality characteristics, increasing the RCC cycles produced chilled fish cracker gels that were less hard and had higher moisture content. Longer chilling durations gave the opposite results. The quality characteristics of the fried fish crackers, namely hardness and colour, were mostly dependent on their expansion ability during frying. Fish crackers containing a higher RS have a lower LE with a negative correlation of $R^2 = -0.777$, thus, fried products with a lower LE were higher in hardness values and darker in colour.

The findings of the present study demonstrated that the variations of cooking-chilling conditions were able to increase the RS content in fish crackers. The lab-produced fish crackers in this study were superior in RS content with an acceptable range of LE and hardness values compared to the commercial non-fried and instant fish crackers. Thus, the lab-produced fish crackers in this study offer better health benefits in term of RS with comparable LE and hardness values for the consumer.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

MENINGKATKAN KANDUNGAN KANJI RINTANG DI DALAM KEROPOK IKAN MELALUI PELBAGAI KEADAAN MEMASAK-MENYEJUK

Oleh

MOHD ZUHAIR BIN MOHD NOR

Ogos 2011

Pengerusi : Rosnita A. Talib, PhD

Fakulti : Kejuruteraan

Keropok ikan merupakan salah satu snek kegemaran di Malaysia. Sungguhpun begitu, kajian berkenaan kandungan kanji rintang (RS) di dalamnya jarang diberikan perhatian. Kanji rintang adalah kanji yang melalui usus kecil tanpa dicernakan, memberikan kesan yang serupa seperti gentian diet. Kanji rintang ini berpotensi untuk dihasilkan semasa proses memasak dan menyejuk di dalam pembuatan keropok ikan. Kedua-dua proses melibatkan gelatinisasi dan retrogradasi kanji yang membawa kepada pembentukan RS, seterusnya menghasilkan produk yang sihat dan bernilai. Oleh itu, penyelidikan ini mengkaji kesan kelainan keadaan memasak-menyujuk terhadap kandungan kanji rintang (RS) dan ciri kualiti penting yang lain di dalam produk keropok ikan. Keadaan proses seperti bilangan kitaran pusingan memasak-menyujuk (RCC) (1 hingga 4 kitaran), suhu memasak (100 °C, 115 °C dan 121 °C) dan tempoh penyejukan (1 hingga 4 hari) serta

jenis kanji yang digunakan di dalam formulasi (ubi kayu, gandum dan sagu) adalah diselidiki dengan tujuan bagi meningkatkan kandungan kanji rintang di dalam keropok ikan tanpa menjejaskan ciri kualitinya. Ciri kualiti bagi keropok ikan dinilai dalam bentuk nilai kekerasan dan kandungan kelembapan untuk gel keropok ikan, kandungan RS di dalam keropok ikan kering serta pengembangan lurus (LE), kekerasan dan warna untuk keropok ikan goreng.

Keputusan menunjukkan bahawa RCC sehingga empat pusingan meningkatkan kandungan RS di dalam semua produk. Keropok ikan diperbuat dari kanji sagu yang dimasak pada suhu 121 °C mempunyai kandungan RS paling tinggi di dalam sampel kering. Pengulangan pusingan memasak-menyejuk meningkatkan tahap penggelatinan kanji pada setiap pusingan memasak dan membantu retrogradasi apabila menyejuk, seterusnya menggalakan pembentukan RS. Namun begitu, keropok ikan yang dimasak pada suhu awalan 100 °C dan didedahkan kepada tempoh penyejukan yang lama sehingga empat hari tidak menunjukkan sebarang perubahan terhadap RS. Kombinasi pengulangan memasak-menyejuk, suhu memasak dan tempoh penyejukan yang berbeza menghasilkan keropok ikan dengan kandungan RS yang berlainan.

Untuk ciri kualiti yang lain, peningkatan kitaran RCC menghasilkan gel keropok ikan sejuk yang kurang keras dan mempunyai kandungan kelembapan yang lebih tinggi. Manakala pendedahan kepada tempoh penyejukan yang lebih lama memberi kesan yang bertentangan. Ciri kualiti bagi keropok ikan goreng iaitu nilai kekerasan dan warna adalah bergantung kepada kadar pengembangan keropok semasa penggorengan. Keropok ikan yang mengandungi kandungan RS yang tinggi mempunyai LE yang

rendah dengan kolerasi negatif $R^2 = -0.777$, seterusnya produk goreng dengan LE yang rendah adalah tinggi nilai kekerasannya dan lebih gelap dari segi warnanya.

Penemuan di dalam kajian ini menunjukkan bahawa kelainan keadaan memasak-menyujuk boleh meningkatkan kandungan RS di dalam keropok ikan. Keropok ikan yang dihasilkan di makmal melalui kajian ini adalah lebih baik dari segi kandungan RS serta mempunyai julat LE dan nilai kekerasan yang boleh diterima apabila dibandingkan dengan keropok ikan komersial kering dan segera. Seterusnya, keropok ikan yang dihasilkan di makmal melalui kajian ini memberikan faedah kesihatan yang lebih baik dari segi RS berserta nilai LE dan kekerasan yang berpatutan kepada pengguna.

ACKNOWLEDGEMENTS

Alhamdulillah, thanks to Almighty God for giving me strength to finish my research. I would like to express my sincere gratitude to my supervisor, Dr. Rosnita A. Talib for her expertise, understanding, patience and encouragement thorough my research project. This thesis would not have been possible without her guidance and unfailing help.

My deep appreciations are extended to my supervisory committee members, Dr. Noranizan Mohd Adzahan, Assc. Prof. Ir. Dr. Chin Nyuk Ling and Dr. Kamaruddin Hashim for their kind assistance, wise advices and constructive comments.

I am indebted to many of my friends for helping and supporting me throughout the difficulties time and caring they provided. I am grateful to all the staffs and laboratory technicians from department of Process and Food Engineering and Faculty of Food Science and Technology that have assisted me during my research work.

Lastly, and most importantly, I would like to thank my beloved wife, Intan Syafinaz and my family for their continuous support and endless love throughout my life.

I certify that a Thesis Examination Committee has met on 1 August 2012 to conduct the final examination of Mohd Zuhair bin Mohd Nor on his thesis entitle “Increasing resistant starch content in fish crackers through various of cooking-chilling conditions” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Johari Endan, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Siti Mazlina Mustapa Kamal, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Rosnah Shamsudin, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Cheow Chong Seng, PhD

Associate Professor
Faculty of Applied Sciences
Universiti Teknologi Mara
Malaysia
(External Examiner)

SEOW HENG FONG, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of **Master of Science**. The members of the Supervisory Committee were as follows:

Rosnita A. Talib, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Noranizan Mohd Adzahan, PhD

Senior Lecturer
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Chin Nyuk Ling, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

Kamaruddin Hashim, PhD

Director
Radiation Processing Technology Division
Malaysian Nuclear Agency (Nuclear Malaysia)
(Member)


BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

I declare that the thesis is my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously, and is not concurrently, submitted for any other degree at Universiti Putra Malaysia or at any other institution.



MOHD ZUHAIR BIN MOHD NOR

Date:



TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL SHEETS	x
DECLARATION FORM	xii
LIST OF TABLES	xvi
LIST OF FIGURES	xviii
LIST OF ABBREVIATIONS	xx
 CHAPTER	
1 INTRODUCTION	
1.1 Research Background	1
1.2 Problem Statements	2
1.3 Research Objectives	3
1.4 Outline of the Thesis	3
2 LITERATURE REVIEW	
2.1 Overview of Fish Crackers	6
2.2 Processing of Fish Crackers	7
2.2.1 Roles of Starch	10
2.2.2 Contribution of Other Raw Materials	11
2.2.3 Mixing	13
2.2.4 Cooking	14
2.2.5 Chilling	15
2.2.6 Drying	15
2.2.7 Frying	16
2.3 Quality Characteristics of Fish Crackers	17
2.3.1 Resistant Starch in Dried Fish Crackers	17
2.3.1.1 Formation and Structure of Resistant Starch Type 3	18
2.3.1.2 Health Benefits of Resistant Starch	21
2.3.1.3 Resistant Starch Intakes	23
2.3.1.4 Food Processing Application of Resistant Starch	24
- Heat Treatments	24
- Heating and Cooling Repetition	26
- Storage Conditions	28
- Effect of Starch Materials	30
- Effect of Presence of Non-Starch Materials	31
- Other Processing Applications	32
2.3.2 Texture Profiles and Moisture Content of Chilled Fish Cracker Gels	33
2.3.3 Linear Expansion and Texture Profiles of Fried Fish Crackers	33
2.3.4 Sensory Qualities and Colour Analysis of Fried Fish Crackers	34
2.4 Commercial Fish Crackers	35

3	METHODOLOGY	
3.1	Raw Materials	37
3.2	General Preparation of Fish Cracker	38
3.3	Treatment Approaches	40
3.3.1	Repetitive Cooking-Chilling (RCC) Cycles	40
3.3.2	Chilling Duration	42
3.3.3	Combination Treatments of Cooking and Chilling Process	42
3.4	Deep-frying of Selected Fish Crackers for Determination of Resistant Starch	43
3.5	Comparison of RS, LE and Hardness between Lab-Produced, Control and Commercial Fish Crackers	44
3.6	Quality Measurements of Dried Fish Crackers	
3.6.1	Resistant Starch	45
3.7	Quality Measurements of Chilled Fish Cracker Gels	
3.7.1	Moisture Content	47
3.7.2	Hardness	47
3.7.3	Observation of Physical Condition	48
3.8	Quality Measurements of Fried Fish Crackers	
3.8.1	Linear Expansion	50
3.8.2	Hardness	51
3.8.3	Colour	51
3.8.4	Resistant Starch	52
3.8.5	Morphology Observation	52
3.9	Data Analysis	52
4	EFFECT OF VARIOUS COOKING-CHILLING CONDITIONS ON RESISTANT STARCH AND IMPORTANT QUALITY CHARACTERISTICS OF FISH CRACKERS	
4.1	Effect of Cooking-Chilling Repetitions (RCC) on Fish Crackers by Different Starch Formulations at a Fixed Cooking Temperature of 100 °C	53
4.1.1	Resistant Starch Content of Dried Fish Crackers	53
4.1.2	Hardness and Moisture Content of Chilled Fish Cracker Gels	58
4.1.3	Linear Expansion, Hardness and Colour of Fried Fish Crackers	60
4.2	Effect of Different Cooking Temperatures in Repetitive Cooking-Chilling (RCC) Cycles on Sago Starch Based Fish Crackers	65
4.2.1	Resistant Starch Content of Dried Fish Crackers	65
4.2.2	Hardness and Moisture Content of Chilled Fish Cracker Gels	68
4.2.3	Linear Expansion, Hardness and Colour of Fried Fish Crackers	70
4.2.4	Physical Conditions of the Chilled Fish Cracker Gels	74
4.3	Effect of Chilling Duration on Fish Crackers Made From Sago Starch	76
4.3.1	Resistant Starch Content of Dried Fish Crackers	76
4.3.2	Hardness and Moisture Content of Chilled Fish Cracker Gels	79
4.3.3	Linear Expansion, Hardness and Colour of Fried Fish Crackers	81
4.4	Effect of Combination Treatments on Sago Starch Fish Crackers	84
4.4.1	Resistant Starch Content of Dried Fish Crackers	84
4.4.2	Hardness and Moisture Content of Chilled Fish Cracker Gels	85

4.4.3	Linear Expansion, Hardness and Colour of Fried Fish Crackers	87
4.5	Summary	90
5	A COMPARISON OF RESISTANT STARCH AND QUALITY CHARACTERISTICS IN SELECTED LAB-PRODUCED, CONTROL AND COMMERCIAL FISH CRACKERS	
5.1	Effect of Deep-Frying Process on Resistant Starch Content in Fish Crackers	92
5.2	Quality Characteristics Comparison between Lab-Produced, Control and Commercial Fish Crackers	95
5.2.1	Resistant Starch, Linear Expansion and Hardness Values Comparison	96
5.2.2	Morphology Observations on the Fish Crackers	99
5.3	Summary	106
6	CONCLUSIONS AND RECOMMENDATIONS	
6.1	Conclusions	107
6.2	Recommendations for Future Research	110
	REFERENCES	112
	APPENDICES	121
	BIODATA OF STUDENT	132
	LIST OF PUBLICATIONS	133