

Chemical characterization and antioxidant activity of three medicinal Apiaceae species

ABSTRACT

Centella asiatica ('Pegaga' = Malaysia) is well known plant that has been used as one of the ingredients in pharmaceutical and cosmetic industries. Using liquid chromatography with diode array and electrospray ionization/mass spectrometry (LC–DAD–ESI/MS) analysis, chemical profiling of three different Pegaga extracts (*C. asiatica*, *H. bonariensis* and *H. sibthorpioides*) revealed variations in their metabolite profile. Our findings showed that triterpenes of *C. asiatica* were characterized by the ursane-type triterpenes (madecassoside, asiaticoside, madecassic acid, and asiatic acid), while the two *Hydrocotyle* species consisted of oleanane-type triterpenes (barrigenol derivatives). Other variations are due to the difference in phenolic and flavonoid constituents. The three Pegaga extracts were also evaluated for their total phenolic content (TPC), 1,1-diphenyl-2-picrylhydrazyl (DPPH) free radicals, and xanthine oxidase inhibition (XOI) activities. The results showed that *C. asiatica* has the most potent antioxidant activity (TPC = 72.09 mg/100 g DW; DPPH = 72.99 µg/ml; XOI = 87.68 µg/ml) as compared to *H. bonariensis* (TPC = 28.55 mg/100 g DW; DPPH = 22.43 µg/ml; XOI = 32.23 µg/ml) and *H. sibthorpioides* (TPC = 56.23 mg/100 g DW; DPPH = 37.86 µg/ml; XOI = 14.73 µg/ml).

Keyword: *C. asiatica*; *H. bonariensis*; *H. sibthorpioides*; HPLC–DAD–ESIMS