

A persistent antimicrobial resistance pattern and limited methicillin-resistance-associated genotype in a short-term *Staphylococcus aureus* carriage isolated from a student population

ABSTRACT

The aim of the present study was to assess and compare the antimicrobial susceptibility pattern against a panel of antibiotics and molecular and methicillin resistance-associated genotypes of 120 carriage *S. aureus* isolates previously isolated from a student population at two isolation events within a one-month interval. The antibiotic susceptibility of isolates was determined using the Kirby-Bauer disc-diffusion method (cefoxitin by Etest). The MRSA was screened using polymerase chain reaction for the presence of the *mecA* gene. The *mecA*-positive isolates were subjected to staphylococcal cassette chromosome (SCC) *mec* typing, multilocus sequence typing (MLST) and eBURST analysis. All isolates were characterized for the presence of the PantónóValentine leukocidin (PVL) gene, an enterobacterial repetitive intergenic consensus-polymerase chain reaction (ERIC-PCR) pattern and the *spa* type. For the two occasions where *S. aureus* was isolated, the highest frequency of resistance was observed for penicillin (70% and 65%, respectively), with a lower rate against erythromycin and tetracycline (<12%). All isolates were susceptible to ciprofloxacin and gentamycin. As for methicillin resistance, eight isolates had minimum inhibitory concentrations (MIC) of resistant categories, but 10 isolates (8.33%) were positive for the *mecA* gene. The *mecA*-positive isolates belonged to SCC*mec* types I (n = 9) and V (n = 1). MLST was resolved for only three MRSA, ST508 (n = 1), ST88 (n = 1) and ST96 (n = 1). The results of the eBURST analysis showed that the MRSA isolates analyzed in the present study were potentially related to MRSA identified in other countries. Approximately half of the persistent *S. aureus* carriers harbored *S. aureus* of a similar *spa* type in the respective individuals during both isolation events. A persistent antimicrobial pattern and limited distinct MRSA were observed over the short study period. The latter frequently exhibited SCC*mec* type I, commonly associated with hospital-acquired (HA) characteristics, but further delineation is needed to justify the origins of these bacteria.

Keyword: Antimicrobial; MRSA; *Staphylococcus aureus*; *mecA* gene