

Assessment of differences on inflammatory and metabolic indicators between pre- and post-menopause women among hypertensive and/or diabetic patients

ABSTRACT

To assess the differences on inflammatory and metabolic indicators between pre-menopause and post-menopause women among hypertensive and/or diabetic type-2 women. A total of 236 obese women included in the study have chosen from Primary Health Care Centers in Gaza City, Palestine, through a cross-sectional study. Selection depended on health status hypertensive and/or diabetic type-2 (HT, T2DM, HT+T2DM). In HT group, post-menopause women had significant higher values than pre-menopause women on TC (200 ± 47 vs. 172.5 ± 55 mg dL⁻¹, $p<0.01$) and TG (166 ± 89 vs. 120.5 ± 82 mg dL⁻¹, $p<0.01$). In T2DM group, post-menopause women had significant higher values than pre-menopause women on SBP (132 ± 24 vs. 120 ± 20 mm Hg, $p<0.01$), TC (213 ± 40 vs. 185 ± 46 mg dL⁻¹, $p<0.05$) and TG (196 ± 118 vs. 136 ± 64 mg dL⁻¹, $p<0.05$). Finally, in HT+T2DM group, post-menopause women had significant higher value than pre-menopause women on SBP (144 ± 21 vs. 133 ± 14 mmHg, $p<0.05$), TC (214 ± 54 vs. 181 ± 55 mg dL⁻¹, $p<0.05$), TG (231 ± 83 vs. 158 ± 85 mg dL⁻¹, $p<0.05$), IL-6 (2.32 ± 1.34 vs. 1.71 ± 1.45 pg mL⁻¹, $p<0.05$) and BMI (36.48 ± 7.1 vs. 32.18 ± 5.6 kg m⁻², $p<0.05$). Post-menopause women diseased of HT and T2DM accompanied with higher rates of BMI are at risk for developing inflammatory and metabolic morbidities.