

Characterization of *Vibrio vulnificus* isolated from cockles (*Anadara granosa*): antimicrobial resistance, plasmid profiles and random amplification of polymorphic DNA analysis

ABSTRACT

Antibiotic susceptibility, plasmid profiles and random amplification of polymorphic DNA (RAPD) were used to study strains of *Vibrio vulnificus* isolated from cockles (*Anadara granosa*). Thirty-six isolates were analyzed. The prevalent biotypes were 1 (72.2% of the isolates) and 2 (27.8%). Among these, 21 strains of biotype 1 and two strains of biotype 2 contained plasmid DNA bands ranging in size from 1.4 to 9.7 MDa. Thirty-one (83.3%) were found to be resistant to one or more of the antimicrobial agents tested, however no specific correlation between antimicrobial resistance patterns and a single biotype was found. In addition, no particular plasmid profile was predictive of a particular pattern of antibiotic susceptibility. Two primers produced polymorphisms in all strains tested, producing bands ranging from 0.25 to 2.7 kb, indicating a high variability among both biotype 1 and biotype 2 of the *V. vulnificus* strains investigated. RAPD identity across biotypes was also observed among *Vibrio vulnificus* strains.

Keyword: Antimicrobial resistance; Plasmid; Random-amplified polymorphic DNA; *Vibrio vulnificus*