



UNIVERSITI PUTRA MALAYSIA

***BIOHYDROGEN PRODUCTION FROM SAGO HAMPAS
BY *Clostridium butyricum* A1***

MOHD AZWAN BIN JENOL

FBSB 2014 22



UPM
UNIVERSITI PUTRA MALAYSIA
BERILMU BERBAKTI

BIOHYDROGEN PRODUCTION FROM SAGO HAMPAS
BY *Clostridium butyricum* A1

By

MOHD AZWAN BIN JENOL

**Thesis Submitted to the School of Graduates Studies, Universiti Putra
Malaysia, in Fulfilment of the Requirement for the Degree of Master of
Science**

June 2014

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**BIOHYDROGEN PRODUCTION FROM SAGO HAMPAS BY
Clostridium butyricum A1**

By

MOHD AZWAN BIN JENOL

June 2014

Supervisor : Professor Suraini Abd-Aziz, PhD
Faculty : Biotechnology and Biomolecular Sciences

The hydrogen has been applied in fuel cell vehicle and expected to shift toward the technologies that produce no net greenhouse gas effects. Biohydrogen production from biomass is now attracting many researchers in developing a renewable, clean and environmental friendly biofuel. The availability of abundant biomass from various sources could possibly be an advantage for the production of biohydrogen as a competitive energy carrier in the future. There are vast choices of possible types of biomass that can be subjected as the carbon source for the production of biohydrogen including starch based and lignocellulosic biomass. Sago industry is one of the possible source of biomass since the industry is producing large quantities of starch and lignocellulosic biomass. Statistically, a single sago starch processing mill has produced 7 ton/day of sago *hampas*. Thus, this study aimed to produce biohydrogen from sago biomass by locally isolated biohydrogen producer and to optimize the production of biohydrogen using statistical approach.

The locally isolated biohydrogen producer *Clostridium butyricum* A1 was successfully isolated from landfill soil. This strain produced a biohydrogen yield of 1.90 mol H₂/mol glucose with productivity of 170 mL/L/h using pure glucose as substrate. The highest cumulative biohydrogen collected after 24 h of fermentation time was 2468 mL/L-medium. Biohydrogen fermentation using sago *hampas* hydrolysate generate higher biohydrogen yield (2.65 mol H₂/mol glucose) compared to sago pith residue (SPR) hydrolysate at 2.23 mol H₂/mol glucose. A higher biohydrogen productivity of 1757 mL/L/h was obtained when using sago *hampas* hydrolysate much higher when compared to pure glucose at 170 mL/L/h. In this study, the new isolate *C. butyricum* A1 together with the use of sago biomass as the substrate is a promising technology for future biohydrogen production. Optimization of biohydrogen production from sago *hampas* hydrolysate by *C. butyricum* A1 was conducted using four variables including temperature, sugar concentration, initial pH and inoculum size. This study has applied central composite design (CCD) and artificial neural network (ANN) as the optimization

step. As a result, three out of four variables have given significant effects on the production of biohydrogen from sago *hampas* hydrolysate; which are temperature, sugar concentration and pH. Using ANN, pH was found to be the most significant variable with the relative importance of 73.6%. The optimum conditions given by ANN with respect to optimized biohydrogen yield of 2.92 mol of H₂/mol of glucose are 39°C, pH 8, initial glucose concentration at 13 g/L and 13% (v/v) inoculum size. As conclusions, biohydrogen production from sago *hampas* by *C. butyricum* A1 has successfully conducted and optimized.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk ijazah Master Sains

**PENGHASILAN BIOHIDROGEN DARIPADA HAMPAS SAGU OLEH
Clostridium butyricum A1**

By

MOHD AZWAN BIN JENOL

Jun 2014

Penyelia : Professor Suraini Abd-Aziz, PhD

Fakulti : Bioteknologi dan Sains Biomolekul

Hidrogen telah diaplikasikan dalam kenderaan sel fuel dan dijangka akan menggerakkan teknologi yang tidak menghasilkan kesan gas rumah hijau. Penghasilan biohidrogen daripada biomas telah menarik minat ramai penyelidik dalam melestarikan penghasilan bahan api biologi yang boleh diperbaharui, bersih dan mesra alam. Ketersediaan biomas yang banyak daripada pelbagai sumber menjadi kelebihan pada penghasilan biohidrogen sebagai pembawa tenaga yang lebih berdaya saing pada masa hadapan. Pelbagai jenis biomas boleh dijadikan sumber karbon untuk penghasilan biohidrogen, termasuklah biomas berasaskan kanji dan lignoselulosa. Industri pemprosesan sagu adalah salah satu sumber biomas berpotensi kerana industri ini menghasilkan biomas berasaskan kanji dan lignoselulosa. Secara statistiknya, satu kilang pemprosesan sagu kanji telah menghasilkan sisa empulur sagu sebanyak 7 tan/hari. Tujuan kajian ini adalah untuk menghasilkan biohidrogen daripada biomas sagu oleh mikroorganisma penghasil biohidrogen pencilan tempatan dan meningkatkan penghasilan biohidrogen menggunakan pendekatan statistik.

Bakteria pencilan tempatan penghasil biohidrogen, *Clostridium butyricum* A1 telah berjaya dipencilkan daripada tanah tapak pelupusan. Bakteria ini telah berjaya menghasilkan biohidrogen sebanyak 1.90 mol H₂/mol glukosa dan 170 mL/L/j produktiviti dengan menggunakan glukosa tulen sebagai substrat. Biohidrogen kumulatif tertinggi yang telah dicatatkan selepas 24 j fermentasi adalah 2468 mL/L-media. Fermentasi biohidrogen menggunakan hidrolisat hampas sagu telah menghasilkan biohidrogen yang lebih tinggi (2.65 mol H₂/mol glukosa) berbanding hidrolisat sisa empulur sagu (SES) (2.23 mol H₂/mol glukosa). Kadar penghasilan biohidrogen yang tinggi juga telah dicatatkan dengan menggunakan hidrolisat hampas sagu iaitu 1757 mL/L/j berbanding dengan menggunakan glukosa tulen dengan produktiviti 170 mL/L/j. Dalam kajian ini, *C. butyricum* A1 bersama dengan penggunaan biomas sagu sebagai substrat menjanjikan satu teknologi yang berpotensi untuk penghasilan biohidrogen di masa hadapan. Pengoptimuman penghasilan biohidrogen daripada hidrolisat hampas sagu oleh *C.*

butyricum A1 telah dijalankan dengan berasaskan 4 pembolehubah, termasuklah suhu, kepekatan gula, pH awal dan saiz inokula. Kajian ini telah menggunakan *central composite design* (CCD) dan *artificial neural network* (ANN) di dalam proses pengoptimuman. Daripada hasil kajian, tiga dari empat pembolehubah telah memberikan kesan yang signifikan pada penghasilan biohidrogen daripada hidrolisat hampas sagu; iaitu suhu, kepekatan gula dan pH. Berdasarkan analisis menggunakan ANN, pH merupakan pembolehubah yang memberikan kesan yang paling ketara dengan kepentingan relatif sebanyak 73.6%. Keadaan yang optimum telah diberikan oleh ANN untuk penghasilan 2.92 mol H₂/mol glukosa ialah 39°C, pH 8, 13 g/L kepekatan glukosa dan saiz inokula 13% (v/v). Kesimpulannya, penghasilan biohidrogen daripada sisa empulur sagu oleh *C. butyricum* A1 telah berjaya dijalankan dan dioptimumkan.



ACKNOWLEDGEMENTS

Alhamdulillah, by the blessing of Allah S.W.T., I manage to complete the thesis entitled "Biohydrogen production from sago *hampas* by *Clostridium butyricum* A1" for my Degree of Master of Science. In completing this project, I would like to express my deep and appreciation to everyone that have helped me. First of all, I would like to thank my main supervisor, Professor Dr. Suraini Abd-Aziz, who is willing to guide, critic and encourage me throughout this project. Without her, this project would not be complete, as presented here.

I would also like to present my appreciation to my supervisor's committee, Assoc. Prof. Dr. Madihah Md Salleh and Dr. Phang Lai Yee, who willing to advise and encourage me in order to finish up my Master of Degree project. Also, to Dr. Mohamad Faizal Ibrahim, who is willing to guide and give a lot of assistances, in helping me throughout my experimental works. They are willing to spend the time for me throughout the project and thesis writing progress.

I am also grateful to all Environmental Biotechnology (EB) group members, Universiti Putra Malaysia, for helping me in terms of advice and suggestion. Also, I would like to express my gratitude to Universiti Putra Malaysia and Malaysia Education Ministry (MOE) for supporting my funding in my three years study.

Last but not least, I would like to express my deepest appreciation to all my family members for their moral supports, especially my beloved mum, Fauziah bte Mustapa, and my sisters, Suliana bte Jenol and Noraisyah bte Jenol. Their endless love, support, and encouragement have made me a better person.

I certify that a Thesis Examination Committee has met on Feb 2014 to conduct the final examination of Mohd Azwan Jenol on his thesis entitled “Biohydrogen Production from Sago Sampas by *Clostridium butyricum* A1” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(a) 106] 15 March 1998. The Committee recommends that the student be awarded the Degree of Master of Science.

Members of the Thesis Examination Committee were as follows:

Nor' Aini Abdul Rahman, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Wan Azlina Wan Ab Karim Ghani, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Umi Kalsom Md Salleh, PhD

Associate Professor
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Internal Examiner)

Kopli Bujang, PhD

Professor
Universiti Malaysia Sarawak
Malaysia
(External Examiner)



NORITAH OMAR, PhD

Associate Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 July 2014

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Suraini Abd-Aziz, PhD

Professor,
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Chairman)

Phang Lai Yee, PhD

Senior Lecturer
Faculty of Biotechnology and Biomolecular Sciences
Universiti Putra Malaysia
(Member)

Madiah Md Salleh, PhD

Associate Professor,
Faculty of Biosciences and Biomedic Engineering
Universiti Teknologi Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

DECLARATION

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work; quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____


Date: 02 August 2014


Name and Matric No.: Mohd Azwan Jenol (GS31880)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in Rule 41 Rules 2003 (Revision 2012-2013) were adhered to.

Signature : 
Name of Chairman
of Supervisory Committee : Prof. Dr Suraini Abd-Aziz (UPM)

Signature : 
Name of Member
of Supervisory Committee : Assoc. Prof. Dr Madihah Md Salleh

Signature : 
Name of Member
of Supervisory Committee : Dr Phang Lai Yee

TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiii
LIST OF ABBREVIATIONS	xiv
CHAPTER	
1 INTRODUCTION	1
2 LITERATURE REVIEW	5
2.1 Introduction	5
2.2 Biomass	6
2.2.1 Starch based biomass	6
2.2.2 Lignocellulosic biomass	9
2.3 Sago biomass in Malaysia	13
2.3.1 Sago waste	13
2.4 Overview of hydrogen	17
2.5 Hydrogen production	18
2.5.1 Steam reforming methane	19
2.5.2 Water hydrolysis	20
2.5.3 Fermentation process	20
2.6 Biohydrogen fermentation	20
2.6.1 Microorganisms	20
2.6.2 Substrates	21
2.6.3 Fermentation types	24
2.7 Factors affecting biohydrogen production	27
2.7.1 pH	27
2.7.2 Temperature	27
2.7.3 Substrate concentration	28
2.7.4 Inoculum size	28
2.7.5 Others	29
2.8 Statistical optimization	29
2.8.1 Response surface methodology	29
2.8.2 Artificial neural network	31
2.8.3 Others	32
3 MATERIALS AND METHODS	33
3.1 General experimental design	33
3.2 Substrate preparation	35
3.3 Identification of biohydrogen producer	35
3.3.1 DNA extraction	36
3.3.2 PCR amplification, sequencing and phylogenetic analysis	36

3.4	Cellulase production	36
3.5	Saccharification process	37
3.5.1	Sago <i>hampas</i>	37
3.5.2	Sago pith residue	38
3.6	Biohydrogen production	38
3.6.1	Inoculum preparation	38
3.6.2	Medium preparation	38
3.6.3	Fermentation process	39
3.7	Optimization of biohydrogen production	39
3.7.1	Central composite design	39
3.7.2	Artificial neural network	40
3.8	Analytical procedures	41
3.8.1	Determination of cellulose, hemicellulose and lignin content	41
3.8.2	Determination of starch content	43
3.8.3	Cell concentration	43
3.8.4	Amylase assay	43
3.8.5	Cellulase activity assay	44
3.8.6	Sugars determination	47
3.8.7	Biohydrogen analysis	47
3.9	The kinetic study	48
4	RESULTS AND DISCUSSION	49
4.1	Chemical composition of sago biomass	49
4.2	Saccharification of sago biomass	51
4.3	Identification and phylogenetic analysis of biohydrogen producer	53
4.4	Biohydrogen production using different types of carbon source	55
4.4.1	Glucose	55
4.4.2	Sago <i>hampas</i> hydrolysate	58
4.4.3	Sago pith residue hydrolysate	59
4.4.4	Comparison study	62
4.5	Optimization of biohydrogen production using statistical analyses	65
4.5.1	Central composite design approach	65
4.5.2	Artificial neural network approach	71
5	CONCLUSION AND RECOMMENDATIONS	73
	REFERENCES	76
	APPENDICES	99
	BIODATA OF STUDENT	105
	LIST OF PUBLICATIONS	106

LIST OF TABLES

Table		Page
2.1	The physicochemical characteristics of starchy-based wastewater	15
2.2	Four basic sections in the steam reforming methane plant	19
2.3	Overview of biohydrogen production from starch based and lignocellulosic biomass	23
2.4	An overview of biohydrogen production via dark and photo fermentation	25
3.1	The coded values for each variable for CCD	39
4.1	An overview of chemical composition of biomass	50
4.2	The hydrolysis of several types of biomass	51
4.3	Biohydrogen production by <i>Clostridium butyricum</i> A1 and <i>Clostridium butyricum</i> EB6 using synthetic glucose, sago <i>hampas</i> hydrolysate and sago pith residue hydrolysate	60
4.4	Summary of biohydrogen production using different types of carbon source	63
4.5	The ANOVA analysis for response surface quadratic model	68

LIST OF FIGURES

Figure		Page
2.1	The chemical structure of amylose	7
2.2	The molecular structure of amylopectin	8
2.3	Structure of lignocellulosic material	9
2.4	The chemical structure of native cellulose	10
2.5	The chemical structure of hemicellulose	11
2.6	The chemical structure of lignin	12
2.7	The schematic flowchart of sago starch extraction process	14
2.8	The hydrogen price from different production process	18
3.1	General experimental layout of this study	34
3.2	The dried sago <i>hampas</i>	35
3.3	The artificial neural network model	40
4.1	The phylogenetic tree of the anaerobic biohydrogen-producing strain and its close relatives based on the almost complete 16S DNA sequence	54
4.2	Biohydrogen production using 10 g/L of glucose by <i>Clostridium butyricum</i> A1 and <i>Clostridium butyricum</i> EB6	56
4.3	The total accumulated biohydrogen gas produced from sago <i>hampas</i> hydrolysate. <i>Clostridium butyricum</i> A1 and <i>Clostridium butyricum</i> EB6	58
4.4	The total accumulated biohydrogen production from SPR hydrolysate by <i>Clostridium butyricum</i> A1 and <i>Clostridium butyricum</i> EB6	61
4.5	Biohydrogen yield predicted by CCD.	69
4.6	Three dimensional surface graphs of model for biohydrogen production at the optimum point	70
4.7	The ANN model architecture for biohydrogen production.	71
4.8	Biohydrogen production predicted by ANN.	72

LIST OF ABBREVIATIONS

A.	<i>Aspergillus</i>
ADF	Acid Detergent Fiber
ADL	Acid Detergent Lignin
ANN	Artificial Neural Network
ANOVA	Analysis of Variance
BLAST	Basic Local Alignment Search Tool
BOD	Biological Oxygen Demand
BPNN	Back Propagation Neural Network
Btu	British Thermal Unit
C.	<i>Clostridium</i>
CCD	Central Composite Design
CCS	CO ₂ Capture and Storage
CDW	Cell Dry Weight
CH ₄	Methane
CO	Carbon Monoxide
CO ₂	Carbon Dioxide
COD	Chemical Oxygen Demand
DAS	Department of Agricultural Sarawak
DNA	Deoxyribonucleic Acid
DNS	3,5-Dinitrosalicylic Acid
dNTPs	Deoxyribonucleotide triphosphates
E.	<i>Enterobacter</i>
E.	<i>Escherichia</i>
EC	Enzyme Commission
EIA	Environmental Impact Assessment
GC	Gas Chromatography
g	G force
h	Hour
H ₂	Hydrogen
H ₂ S	Hydrogen Sulphite

HCl	Hydrochloric Acid
HHV	Higher Heating Value
HPLC	High Performance Liquid Chromatography
HRT	Hydraulic Retention Time
j	<i>Jam</i>
MgCl ₂	Magnesium Chloride
min	Minute
NaOH	Sodium hydroxide
NCBI	National Center for Biotechnology Information
NDF	Neutral Detergent Fiber
OPDC	Oil Palm Decantor Cake
OPEFB	Oil Palm Empty Fruit Bunches
PCR	Polymerase Chain Reaction
PDA	Potato Dextrose Agar
POME	Palm Oil Mill Effluent
PSA	Pressure Swing Adsorption
R ²	Correlation coefficient
RCM	Reinforced Clostridia Medium
rDNA	Ribosomal DNA
rpm	Rotation Per Minute
RSM	Response Surface Methodology
RSS	Residual Sum of Square
SAS	Sarawak Agriculture Statistic
sp.	Species
SPR	Sago Pith Residue
SRM	Steam Reforming of Methane
TKN	Total Kjeldahl Nitrogen
UPM	Universiti Putra Malaysia
VFA	Volatile Fatty Acid
VSS	Volatile Suspended Solid

CHAPTER 1

INTRODUCTION

Biohydrogen gas produced through fermentation process is an ideal alternative energy carrier that can be used as biofuel, due to its characteristics which are clean and sustainable bioenergy for the future. Biohydrogen is a simple gas with a molecular formula of H_2 and has high energy yield of 122 kJ/g, which is 2.75 times higher than fuels derived from petroleum (Claassen *et al.*, 2010; Chong *et al.*, 2009a; Pattra *et al.*, 2008). The combustion of hydrogen in engine as energy carrier is 50–70% more efficient than gasoline, hence the only byproduct generated is water (Pattra *et al.*, 2008; Fields, 2003), thus subsequently reduce the release of carbon dioxide into the atmosphere.

At present, the industrial production of hydrogen is through several processes including steam reforming of methane, electrolytic (water electrolysis) and thermochemical reactions (Levin and Chahine, 2010; Steinfeld, 2005). The major problems among these available technologies are ineffective cost and high energy consumption. Therefore, this situation leads to edge novel hydrogen uncompetitive to replace our current reliance on fossil fuel. Hence, innovative and novel biohydrogen production via fermentation using cheap feedstock has now being attempted because, it needs less land for set up and not being affected by weather conditions (Ni *et al.*, 2007).

In European countries, the “hyvolution” concept has been introduced in order to attain greener biohydrogen production. The concept is by exploiting bacteria that are capable in producing biohydrogen as byproduct during growth on biomass (Claassen *et al.*, 2010). Besides, utilization of biomass as a fermentation feedstock can reduce the cost of raw material which contributed around 50–70% of total biohydrogen production cost. By this mean, it will also help biomass related industries in managing their biomass waste and to generate extra profit (Hassan *et al.*, 2004).

Naturally, biomass present in various forms, including starch and lignocellulosic materials. The starch based biomass is composed of mainly starch, while the lignocellulosic biomass which is composed of 38–50% cellulose, 23–32% hemicellulose and 15–25% lignin (Perego and Bianchi, 2010). The polysaccharides of starch, cellulose and hemicellulose can be converted into simple sugars which are useful to be utilized as carbon sources for the fermentation process by microorganism. The bioconversion of biomass into simple sugars can be conducted through the hydrolysis process using specific enzyme. Utilization of carbon source recovered from the hydrolysis of biomass for biohydrogen production has been reported by several researchers including from palm oil mill effluent (Chong *et al.*, 2009a), oil palm empty fruit bunch (Inayat *et al.*, 2012), cassava wastewater (Sreethawong *et al.*, 2010) and sweet potato starch residue (Yokoi *et al.*, 2001) by exploiting various species of microorganisms.

Malaysia is blessed with favorable climate and abundant natural resources for commercial cultivation of crops such as sago palm. The Malaysian sago palm industry is one of the most important sago exporters in the world and exporting in the range of 55,000 to 65,000 tons/year of sago starch (Department of Statistic, Sarawak, 2012). The Department of Agriculture Sarawak (2010) has reported that the export value is increasing by 15–20% per year. In preliminary 2012, the export value of sago products was 48,314 tons compared to 2003 (40,780 tons) (Department of Statistic, Sarawak, 2012). Increment in production will significantly increase the number of waste generated from this industry, which may cause waste management problems and contribute to the environmental pollution. This industry has generated vast amount of wastes in the form of byproducts including sago bark, sago *hampas* and sago wastewater.

Sago *hampas* contains (on a dry weight basis) 58% starch, 23% cellulose, 9.2% hemicellulose and 3.9% lignin (Awg-Adeni *et al.*, 2012; Linggang *et al.*, 2012; Ozawa *et al.*, 1998). The enzymatic conversion of starch present in sago *hampas* into glucose produces another biomass which is known as sago pith residue (SPR). Linggang *et al.* (2012) has reported that SPR composed of (on a dry weight basis) 37% cellulose, 20% hemicellulose and 6% lignin and cellulose and hemicellulose can be hydrolyzed into a mixture of fermentable sugars using enzyme called cellulase. These types of biomass have the potential to be used as a carbon source in the production of valuable products. Thus, a study on the utilization of sago *hampas* and SPR for biohydrogen production was conducted. A new local isolated strain *Clostridium butyricum* A1 was employed to convert fermentable sugars into biohydrogen to be compared with a biohydrogen producer, *Clostridium butyricum* EB6 isolated by Chong *et al.* (2009a).

A variety of biohydrogen-producing microorganisms has been documented by several studies. To date, *Clostridium* sp. is widely employed for biohydrogen production since this species is capable in converting hexose sugar to biohydrogen with a theoretical yield of 4 mol H₂/mol hexose. This is higher than the biohydrogen produced by *Enterobacter* sp. that has a yield of 1 mol H₂/mol hexose (Kotay and Das, 2008). However, several studies have been done for *Enterobacter* sp. since this species can achieve higher productivity than other reported species (Prasertsan *et al.*, 2009; Kotay and Das, 2008; Chen *et al.*, 2005). Jamil *et al.* (2009) has been reported on production of biohydrogen by *Rhodopseudomonas palustris* PBUM001 with biohydrogen yield 0.66 mL H₂/mL POME. *Clostridium* sp. has the capability to produce a biohydrogen yield in the range of 1.4–2.8 mol H₂/mol glucose (Levin and Chahine, 2010; Lin and Tanaka, 2006; Chen *et al.*, 2005). The highest biohydrogen yield reported was 3.26 mol H₂/mol glucose by employing *C. butyricum* (Keskin and Hallenbeck, 2012). There are also several reports on the production of biohydrogen by *Thermotoga* sp. using biomass as substrate with a yield in the range of 1.1–2.0 mol H₂/mol glucose or other hexoses (Mars *et al.*, 2010; Evvyernie *et al.*, 2001).

However, biohydrogen production depends on various fermentation factors that correlate each other to improve biohydrogen yield. Due to the fact that, different substrate and Clostridia species employed in the fermentation produces different yield of biohydrogen, thus a tool that evaluate various fermentation factors at one time with statistical analysis are usually practiced. One of the most widely implemented tools is the response surface methodology (RSM) over the last two decades. RSM is an approach that implies the statistical technique based on the crucial technique of duplication and randomization, thus makes the optimization study easier (Baskar *et al.*, 2008). Based on the review, RSM is an effective tool to manipulate the large number of variables, therefore many reports have been used RSM to improve fermentation condition (Balusu *et al.*, 2005; Ezhumalai and Thangavelu, 2010). This tool has proven its ability to improve the biohydrogen yield using POME as the substrate, as conducted by Chong *et al.* (2009a; 2009b).

The artificial neural network (ANN) which is more advance and powerful statistical analysis tool is also able to optimize the fermentation parameters including for biohydrogen fermentation. One of the advantages of ANN is less time is required for development of extensive experimentation (Morteza *et al.*, 2013). This situation is due to the fact that, it could be conducted with limited numbers of experiments to predict the degree of non-linearity. It is also capable of learning complex relationships without requiring the knowledge of the model structure (Mingzhi *et al.*, 2009). According to Dutta *et al.* (2004), the ANN is a superior and more accurate tool in terms of modeling technique as compare to RSM because ANN depicts the nonlinearity of the model. Although, ANN has no ability to estimate the model equation similar to RSM, it still is able to estimate the response based on the trained data (Baş and Boyacı, 2007).

The study aims to produce the biohydrogen from sago biomass by *Clostridium butyricum* A1 with the specific objectives as follows:

1. To evaluate the suitability of sago *hampas* and sago pith residue for biohydrogen production by local isolates.
2. To compare biohydrogen production by *Clostridium butyricum* A1 and *Clostridium butyricum* EB6 using selected sago biomass.
3. To optimize biohydrogen production from selected sago biomass by selected *Clostridium* sp. using artificial neural network.



REFERENCES

- Aderibigbe, A.F., Anozie, A.N., Adejumo, L.A. and Owolabi, R.U. (2012). New Clues in Sciences.
- Afsar, N., Özgür, E., Gürkan, M., Akköse, S., Yücel, M., Gündüz, U. and Eroglu, I. (2011). Hydrogen productivity of photosynthetic bacteria on dark fermenter effluent of potato steam peels hydrolysate. *International Journal of Hydrogen Energy*, 36(1), 432-438.
- Agbor, V.B., Cicek, N., Sparling, R., Berlin, A. and Levin, D.B. (2011). Biomass pretreatment: fundamentals toward application. *Biotechnology Advances*, 29(6), 675-685.
- Ahn, Y., Park, E.J., Oh, Y.K., Park, S., Webster, G. and Weightman, A.J. (2005). Biofilm microbial community of a thermophilic trickling biofilter used for continuous biohydrogen production. *FEMS Microbiology Letters*, 249(1), 31-38.
- Alalayah, W.M., Kalil, M.S., Kadhum, A.A., Jahim, J.M. and Alauj, N.M. (2009). Effect of environmental parameters on hydrogen production using *Clostridium saccharoperbutylacetonicum* N1-4 (ATCC 13564). *American Journal of Environmental Science*, 5(1), 80-86.
- Annison, G. and Topping, D.L. (1994). Nutritional role of resistant starch: chemical structure vs physiological function. *Annual Review of Nutrition*, 14(1), 297-320.
- Apun, K., Jong, B. C., and Salleh, M. A. (2000). Screening and isolation of a cellulolytic and amylolytic *Bacillus* from sago pith waste. *The Journal of General and Applied Microbiology*, 46(5), 263-267.
- Ariffin, H., Hassan, M.A., Kalsom, M.U., Abdullah, N. and Shirai, Y. (2008b). Effect of physical, chemical and thermal pretreatments on the enzymatic hydrolysis of oil palm empty fruit bunch (OPEFB). *Journal of Tropical Agriculture and Food Science*, 36(2), 1-10
- Ariffin, H., Hassan, M.A., Shah, U.K.M., Abdullah, N., Ghazali, F.M. and Shirai, Y. (2008a). Production of bacterial endoglucanase from pretreated oil palm empty fruit bunch by *Bacillus pumilus* EB3. *Journal of Bioscience and Bioengineering*, 106(3), 231-236.
- Awg-Adeni, D. S., Bujang, K. B., Hassan, M. A. and Abd-Aziz, S. (2012). Recovery of glucose from residual starch of sago *hampas* for bioethanol production. *BioMed Research International*, DOI:10.1155/2013/935852.
- Awg-Adeni, D.S., Abd-Aziz, S., Bujang, K. and Hassan, M.A. (2010). Bioconversion of sago residue into value added products. *African Journal of Biotechnology* 9(14), 2016-2021.

- Aziz, A. A., Husin, M. and Mokhtar, A. (2002). Preparation of cellulose from oil palm empty fruit bunches via ethanol digestion: effect of acid and alkali catalysts. *Journal of Oil Palm Research*, 14, 9-14.
- Azwar, M. A. Hussain, K. and Ramachandran, B. (2006). The study of neural network-based controller for controlling dissolved oxygen concentration in a sequencing batch reactor. *Bioprocess and Biosystem Engineering*, 28(4), 251-265.
- Bahrin, E.K., Baharuddin, A.S., Ibrahim, M.F., Razak, N.A., Sulaiman, A., Abd-Aziz, S., Hassan, M.A., Shirai, Y. and Nishida, H. (2012). Physicochemical property changes and enzymatic hydrolysis enhancement of oil palm empty fruit bunches treated with superheated steam. *BioResources*, 7(2), 1784-1801.
- Balat, M. (2008). Potential importance of hydrogen as a future solution to environmental and transportation problems. *International Journal of Hydrogen Energy*, 33, 4013–4029.
- Balusu, R., Paduru, R.R., Kuravi, S.K., Seenayya, G. and Reddy, G. (2005). Optimization of critical medium components using response surface methodology for ethanol production from cellulosic biomass by *Clostridium thermocellum* SS19. *Process Biochemistry*, 40(9), 3025-3030.
- Banu, J. R., Kaliappan, S., and Beck, D. (2006). Treatment of sago wastewater using hybrid anaerobic reactor. *Water Quality Research Journal of Canada*, 41(1), 56-62.
- Barbosa, M.J., Rocha, J., Tramper, J. and Wijffels, R.H. (2001). Acetate as a carbon source for hydrogen production by photosynthetic bacteria. *Journal of Biotechnology*, 85(1), 25-33.
- Barros, A.R., Cavalcante de Amorim, E.L., Reis, C.M., Shida, G.M. and Silva, E.L. (2010). Biohydrogen production in anaerobic fluidized bed reactors: Effect of support material and hydraulic retention time. *International Journal of Hydrogen Energy*, 35(8), 3379-3388.
- Baş, D. and Boyacı, İ. H. (2007). Modeling and optimization II: Comparison of estimation capabilities of response surface methodology with artificial neural networks in a biochemical reaction. *Journal of Food Engineering*, 78(3), 846-854.
- Baskar, G., Muthukumaran, C., & Renganathan, S. (2008). Optimization of Enzymatic Hydrolysis of *Manihot esculenta* root starch by Immobilized α -amylase using Response Surface Methodology. *International Journal of Chemical and Biomolecular Engineering*, 1(3), 364-368.
- Berkers, L., Hilgsmann, S., Hamilton, C., Masset, J. and Thonart, P. (2010). Fermentative hydrogen production by *Clostridium butyricum* CWBI1009 and *Citrobacter freundii* CWBI952 in pure and mixed cultures. *Biotechnology Agronomy Society and Environmental*, 14(S2), 541-548.

- Bernfeld, P. (1955). Amylases, alpha and beta. *Methods in Enzymology I*, 149-158.
- Bertoft, E., Piyachomkwan, K., Chatakanonda, P. and Sriroth, K. (2008). Internal unit chain composition in amylopectins. *Carbohydrate polymers*, 74(3), 527-543.
- Bidlas, E., Du, T. and Lambert, R.J.W. (2008). An explanation for the effect of inoculum size on MIC and the growth/no growth interface. *International Journal of Food Microbiology*, 126, 140-152.
- Binford, L. and Ho, C-k. (1985). Taphonomy at a distance: Zhoukoudian, 'the cave home of Beijing man'?. *Current Anthropology*, 26, 413-442.
- Bishop, C.M. (1995) Neural networks for pattern recognition, 1st edn. *Oxford University Press, USA*. p. 482. (ISBN-13: 978-01 98538646).
- Boran, E., Özgür, E., van der Burg, J., Yücel, M., Gündüz, U. and Eroglu, I. (2010). Biological hydrogen production by *Rhodobacter capsulatus* in solar tubular photo bioreactor. *Journal of Cleaner Production*, 18, S29-S35.
- Boudet, A.M., Kajita, S., Grima-Pettenati, J. and Goffner, D. (2003). Lignins and lignocellulosics: a better control of synthesis for new and improved uses. *Trends in Plant Science*, 8(12), 576-581.
- Brayer, G.D., Luo, Y., and Withers, S.G. (1995). The structure of human pancreatic α -amylase at 1.8 Å resolution and comparisons with related enzymes. *Protein Science*, 4(9), 1730-1742.
- Bressan L., Collodi G. and Ruggeri F. (2009). Hydrogen generation for modern refineries. *Hydrocarbon Engineering, Foster Wheeler Italiana SpA, Corsico, Milan ,Italy*.
- Brodeur, G., Yau, E., Badal, K., Collier, J., Ramachandran, K.B. and Ramakrishnan, S. (2011). Chemical and physicochemical pretreatment of lignocellulosic biomass: a review. *Enzyme Research*, 2011.
- Bujang, K., Apun, K. and Salleh, M.A. (1996). A study in the production and bioconversion of sago waste. *Sago-The future Source of Food and Feed*. *Riau University Press, Indonesia*, 195-201.
- Chang, V.S. and Holtzapple, M.T. (2000). Fundamental factors affecting biomass enzymatic reactivity. In *Twenty-First Symposium on Biotechnology for Fuels and Chemicals* (pp. 5-37). Humana Press.
- Chen, C.Y., Yang, M.H., Yeh, K.L., Liu, C.H. and Chang, J.S. (2008). Biohydrogen production using sequential two-stage dark and photo fermentation processes. *International Journal of Hydrogen Energy*, 33(18), 4755-4762.

- Chen, W.H., Chen, S.Y., Khanal, S.K. and Sung, S. (2006). Kinetic study of biological hydrogen production by anaerobic fermentation. *International Journal of Hydrogen Energy*, 31, 2170-2178.
- Chen, W.M., Tseng, Z.J., Lee, K.S. and Chang, J.S. (2005). Fermentative hydrogen production with *Clostridium butyricum* CGS5 isolated from anaerobic sewage sludge. *International Journal of Hydrogen Energy*, 30, 1063-1070.
- Chittibabu, G., Nath, K. and Das, D. (2006). Feasibility studies on the fermentative hydrogen production by recombinant *Escherichia coli* BL-21. *Process Biochemistry*, 41(3), 682-688
- Chong, M.L., Abd-Rahim, R., Shirai, Y. and Hassan, M.A. (2009a). Biohydrogen production by *Clostridium butyricum* EB6 from palm oil mill effluent. *International Journal of Hydrogen Energy*, 34, 764-777.
- Chong, M.L., Rahman, N.A.A., Abd-Rahim, R., Abd-Aziz, S., Shirai, Y. and Hassan, M.A. (2009b). Optimization of biohydrogen production by *Clostridium butyricum* EB6 from palm oil mill effluent using response surface methodology. *International Journal of Hydrogen Energy*, 34, 7475-7482.
- Chong, M.L., Rahman, N.A.A., Yee, P.L., Abd-Aziz, S., Abd-Rahim, R., Shirai, Y. and Hassan, M.A. (2009c). Effects of pH, glucose and iron sulfate concentration on the yield of biohydrogen by *Clostridium butyricum* EB6. *International Journal of Hydrogen Energy*, 34, 8859-8865.
- Claassen, P.A.M., De Vrije, T., Budde, M.A.W., Koukios, E.G., Glynos, A. and Réczey, K. (2004, May). Biological hydrogen production from sweet sorghum by thermophilic bacteria. In *Proceedings 2nd World Conference on Biomass for Energy, Rome* (pp. 1522-1525).
- Claassen, P.A.M., Vrije, T.D., Koukios, E.G., Niel, E.W.J., Eroglu, I., Modigell, M., Friedl, A., Wukovits, W. and Ahrer, W. (2010). Non-thermal production of pure hydrogen from biomass: hyvolution. *Journal of Cleaner Production*, 18, S4-S8.
- Collet, C., Adler, N., Schwitzguébel, J.P. and Péringier, P. (2004). Hydrogen production by *Clostridium thermolacticum* during continuous fermentation of lactose. *International Journal of Hydrogen Energy*, 29(14), 1479-1485.
- Collodi, G. and Wheeler, F. (2010). Hydrogen production via steam reforming with CO₂ capture. *Chemical Engineering Transactions*, 19, 37-41.
- Cubillos, G., Arrué, R., Jeison, D., Chamy, R., Tapia, E., Rodríguez, J. and Ruiz-Filippi, G. (2010). Simultaneous effects of pH and substrate concentration on hydrogen production by acidogenic fermentation. *Electronic Journal of Biotechnology*, 13(1), 11-12.

- Dabrock, B., Bahl, H. and Gottschalk, G. (1992). Parameters affecting solvent production by *Clostridium pasterium*. *Applied Environmental Microbiology*, 58, 1123-1131.
- De Vrije, T. and Claassen, P.A.M. (2003). Dark hydrogen fermentations. *Bio-methane and Bio-hydrogen*, 103-123.
- de Vrije, T., Bakker, R.R., Budde, M.A.W., Lai, M.L., Mars, A.E. and Claassen, P.A.M. (2009). Efficient hydrogen production from the lignocellulosic energy crop *Miscanthus* by the extreme thermophilic bacteria *Caldicellulosiruptor saccharolyticus* and *Thermotoga neapolitana*. *Biotechnology for Biofuel*, 2, 12
- Delmer, D.P. and Amor, Y. (1995). Cellulose Biosynthesis. *Plant Cell*, 7(7), 987-1000.
- Department of Agricultural Sarawak (2010). Sarawak agricultural statistic: export of sago product 2001-10, DAS, Malaysia: (<http://www.doa.sarawak.gov.my/modules/web/page.php?id=504>).
- Department of Statistic, Sarawak (2012). Export of sago products 2003 – 2012. (<http://www.doa.sarawak.gov.my>)
- Divne, C., Ståhlberg, J., Teeri, T.T. and Jones, T.A. (1998). High-resolution crystal structures reveal how a cellulose chain is bound in the 50 Å long tunnel of cellobiohydrolase I from *Trichoderma reesei*. *Journal of Molecular Biology*, 275(2), 309-325.
- DOE, U. (2007). *International Energy outlook 2007*. Technical Report No. DOE/EIA-0484, <http://www.eia.doe.gov/oiaf/ieo/pdf/0484> (2007). pdf.
- Doremus, M.G., Linden, J.C. and Moreira, A.R. (1985). Agitation and pressure effects on acetone-butanol fermentation. *Biotechnology and Bioengineering*, 27(6), 852-860.
- Dutta, J.R., Dutta, P.K. and Banerjee, R. (2004). Optimization of culture parameters for extracellular protease production from a newly isolated *Pseudomonas* sp. using response surface and artificial neural network models. *Process Biochemistry*, 39(12), 2193-2198.
- Ebringerova, A., Hromadkova, Z. and Heinze, T. (2005). Hemicellulose. *Advance in Polymer Sciences*, 186, 1-67.
- EIA, U. (2011). Annual energy outlook 2011 with projections to 2035. *Washington DC: Energy Information Administration, United States Department of Energy*.
- El-Shafie, A. (2012). Neural network nonlinear modeling for hydrogen production using anaerobic fermentation. *Neural Computing and Applications*, 8, 1-9.

- El-Shafie, A., Abdelazim, T. and Noureldin, A. (2010b). Neural network modelling of time-dependent creep deformations in masonry structures. *Neural Computing and Applications* 19(4), 583–594.
- El-Shafie, A., Noureldin, A.E., Taha, M.R. and Basri, H. (2008). Neural network model for Nile river inflow forecasting based on correlation analysis of historical inflow data. *Journal of Applied Sciences*, 8(24), 4487-4499.
- El-Shafie, A., Osman, A., Noureldin, A. and Hussain, A. (2010a). Performance evaluation of a nonlinear error model for underwater range computation utilizing GPS sonobuoys. *Neural Computing and Applications*, 19(5), 272–283.
- El-Shafie, A.H., El-Shafie, A., El Mazoghi, H.G., Shehata, A. and Taha, M.R. (2011). Artificial neural network technique for rainfall forecasting applied to Alexandria, Egypt. *International Journal of Physical Sciences*, 6(6), 1306-1316.
- Eriksen, N.T., Nielsen, T.M. and Iversen, N. (2008). Hydrogen production in anaerobic and microaerobic *Thermotoga neapolitana*. *Biotechnology Letters*, 30, 103-109.
- Esteghlalian, A.R., Mansfield, S.D. and Saddler, J.N. (2002). Cellulases: Agents for fiber modification or bioconversion? The effect of substrate accessibility on cellulose enzymatic hydrolyzability. *Biotechnology Pulp Paper Industry: Program Biotechnology* 21, 21-36.
- Evyernie, D., Morimoto, K., Karita, S., Kimura, T., Sakka, K, and Ohmiya, K. (2001). Conversion of chitinous wastes to hydrogen gas by *Clostridium paraputrificum* M-21. *Journal of Biosciences and Bioengineering*, 91, 339-343.
- Ezeji, T.C., Qureshi, N. and Blaschek, H.P. (2007). Bioproduction of butanol from biomass: from genes to bioreactors. *Current Opinion in Biotechnology*, 18 (3): 220-227.
- Ezhumalai, S. and Thangavelu, V. (2010). Kinetic and optimization studies on the bioconversion of lignocellulosic material into ethanol. *BioResources*, 5(3), 1879-1894.
- Fabiano, B. and Perego, P. (2002). Thermodynamic study and optimization of hydrogen production by *Enterobacter aerogenes*. *International Journal of Hydrogen Energy*, 27(2), 149-156.
- Fan, Y., Li, C., Lay, J.J., Hou, H. and Zhang, G. (2004). Optimization of initial substrate and pH levels for germination of sporing hydrogen-producing anaerobes in cow dung compost. *Bioresource Technology*, 91(2), 189-193.
- Fang, H.H., Li, C. and Zhang, T. (2006). Acidophilic biohydrogen production from rice slurry. *International Journal of Hydrogen Energy*, 31(6), 683-692.

- Fang, H.H.P., Zhang, H. and Liu, H. (2002). Microbial diversity of mesophilic hydrogen producing sludge. *Applied Microbiology and Biotechnology*, 58, 112-118.
- Fang, J.M., Sun, R.C., Tomkinson, J. and Fowler, P. (2000). Acetylation of wheat straw hemicellulose B in a new non-aqueous swelling system. *Carbohydrate Polymer*, 41, 379-387.
- Fedorov, A.S., Tsygankov, A.A., Rao, K.K. and Hall, D.O. (1998). Hydrogen photoproduction by *Rhodobacter sphaeroides* immobilised on polyurethane foam. *Biotechnology Letters*, 20(11), 1007-1009.
- Fengel, D., and Wegener, G. (1989). Polyoses (hemicelluloses). *Wood: Chemistry, Ultrastructure, Reactions*, 106-131.
- Festucci-Buselli, R.A., Otoni, W.C. and Joshi, C.P. (2007). Structure, organization, and functions of cellulose synthase complexes in higher plants. *Brazil Journal of Plant Physiology* 19(1), 1-13.
- Fields, S. (2003). Making the best of biomass hydrogen for fuel cells. *Environmental Health Perspectives*, 1(11), A38-A41.
- Fuel Cell and Hydrogen Energy Association, (2013). Hydrogen energy review. Retrieved from <http://www.fchea.org/index.php?id=46>, on 15 November 2012.
- Gallucci, F., Comite, A., Capannelli, G. and Basile A. (2006). Steam reforming of methane in a membrane reactor: an industrial case study. *Industrial and Engineering Chemistry Research*. 45, 2994-3000.
- Ginkel, S.V. and Logan, B.E. (2005). Inhibition of biohydrogen production by undissociated acetic and butyric acids. *Environmental Science and Technology*, 39, 9351-9356.
- Goering, H.K. and Van Soest, P.J. (1970). Forage Fiber Analysis (apparatus, reagents, procedures and some applications) *USDA Agricultural Handbook* No. 379.
- Gowlett, J.A.J. (2006). The early settlement of northern Europe: fire history in the context of climate change and the social brain. *Comptes Rendus Palevol*, 5(1), 299-331.
- Grupe, H. and Gottschalk, G. (1986). Physiological events in *Clostridium acetobutylicum* during the shift from acidogenesis to solventogenesis in continuous culture and presentation of a model for shift induction. *Applied Environmental Microbiology*, 58(12), 3896-3902.
- Guo, W.Q., Ren, N.Q., Wang, X.J., Xiang, W.S., Ding, J., You, Y. and Liu, B.F. (2009). Optimization of culture conditions for hydrogen production by *Ethanoligenes harbinense* B49 using response surface methodology. *Bioresource Technology*, 100, 1192-1196.

- Ha, M.A., Apperley, D.C., Evans, B.W., Huxman, I.M., Jardine, W.G., Vietor, R.J., Reis, D., Vian, B. and Michael C.J. (1998). Fine structure in cellulose microfibrils: NMR evidence from onion and quince. *Plant*, 16(2), 183-90.
- Hamilton, C., Hilgsmann, S., Beckers, L., Masset, J., Wilmotte, A. and Thonart, P. (2010). Optimization of culture conditions for biological hydrogen production by *Citrobacter freundii* CWBI952 in batch, sequenced-batch and semicontinuous operating mode. *International Journal of Hydrogen Energy*, 35(3), 1089-1098.
- Hassan, M.A., Yacob, S. and Shirai, S.Y. (2004). Treatment of palm oil wastewaters, Wang, L.K., Hung Lo, Y.H.H. and Yapijakis, C. (ed.), Handbook of Industrial and Hazardous Waste Treatment, *Marcel Dekker Inc, New York*, p. 719-736.
- Hasyim, R., Imai, T., O-Thong, S. and Sulistyowati, L. (2011). Biohydrogen production from sago starch in wastewater using an enriched thermophilic mixed culture from hot spring. *International Journal of Hydrogen Energy* 36(21), 14162–14171.
- Heinze, T. and Liebert, T. (2001). Unconventional methods in cellulose functionalization. *Progress in Polymer Science*, 26(9), 1689-1762.
- Hendriks, A.T.W.M. and Zeeman, G. (2009). Pretreatments to enhance the digestibility of lignocellulosic biomass. *Bioresource Technology*, 100(1), 10-18.
- Hillmer, P. and Gest, H. (1977). H₂ metabolism in the photosynthetic bacterium *Rhodospseudomonas capsulata*: H₂ production by growing cultures. *Journal of Bacteriology*, 129(2), 724-731.
- Ho, K.L., Chen, Y.Y. Lee, D.J. (2010). Biohydrogen production from cellobiose in phenol and cresol-containing medium using *Clostridium* sp. R1. *International Journal of Hydrogen Energy*, 35(19), 10239-10244.
- Holladay, J.D., Hu, J., King, D.L. and Wang, Y. (2009). An overview of hydrogen production technologies. *Catalysis Today*, 139(4), 244-260.
- Horigome, T., Sakaguchi, E., Takamura, Y., Bintaro, M. H., Haryanto, B., Tandi, E. J. and Marangkey, M. P. (1991). The feeding value of pith and pith residue from sago palms. In 4. *International Sago Symposium, Kuching, Sarawak (Malaysia), 6-9 Aug 1990*. Ministry of Agriculture and Community Development.
- Ibrahim, M.F., Abd-Aziz, S., Razak, M.N.A., Phang, L.Y. and Hassan, M.A. (2012). Oil palm empty fruit bunch as alternative substrate for acetone butanol ethanol production by *Clostridium butyricum* EB6. *Applied Biochemistry and Biotechnology*, 166, 1615-1625.
- Ibrahim, M.F., Razak, M.N.A., Phang, L.Y., Hassan, M.A. and Abd-Aziz, S. (2013). Crude cellulase from oil palm empty fruit bunch by *Trichoderma asperellum* UPM1 and *Aspergillus fumigatus* UPM2 for fermentable sugars production. *Applied Biochemistry and Biotechnology*, doi 10.1007/s12010-013-0275-2.

- Ihme, M., Marsden, A. L. and Pitsch, H. (2008). Generation of optimal artificial neural networks using a pattern search algorithm: Application to approximation of chemical systems. *Neural Computation*, 20(2), 573-601.
- Ikram-ul-Haq, Javed, M.M., Khan, T.S. and Siddiq, Z. (2005). Cotton saccharifying activity of cellulases produced by co-culture of *Aspergillus niger* and *Trichoderma viride*. *Research Journal of Agriculture and Biological Sciences*, 1(3), 241-245.
- Inayat, A., Ahmad, M.M., Ahmad Mutalib, M.I. and Yusup, S. (2012). Process modeling for parametric study on oil palm empty fruit bunch steam gasification for hydrogen production. *Fuel Processing Technology*, 93(1), 26-34.
- Jayasinghearachchi, H.S., Singh, S., Sarma, P.M., Aginihotri, A. and Lal, B. (2010). Fermentative hydrogen production by new marine *Clostridium amygdalinum* strain C9 isolated from offshore crude oil pipeline. *International Journal of Hydrogen Energy*, 35(13), 6665-6673.
- Jones, D.T. and Woods, D.R. (1986). Acetone-butanol fermentation revisited. *Microbiological Reviews*, 50(4), 484-524.
- Kadam, S.K. and Demain, A.L. (1989). Addition of cloned β -glucosidase enhances the degradation of crystalline cellulose by the *Clostridium thermocellum* cellulose complex. *Biochemical and Biophysical Research Communications*, 161(2), 706-711.
- Kalia, V.C., Jain, S.R., Kumar, A. and Joshi, A.P. (1994). Fermentation of biowaste to H₂ by *Bacillus licheniformis*. *World Journal of Microbiology and Biotechnology*, 10(2), 224-227.
- Kalinci, Y., Hepbasli, A. and Dincer, I. (2009). Biomass-based hydrogen production: a review and analysis. *International Journal of Hydrogen Energy*, 34, 8799-8817.
- Kandra, L., Gyémánt, G., Remenyik, J., Ragunath, C. and Ramasubbu, N. (2003). Subsite mapping of human salivary α -amylase and the mutant Y151M. *FEBS Letters*, 544(1), 194-198.
- Karaca, F. and Özkaya, B. (2006). NN-LEAP: A neural network-based model for controlling leachate flow-rate in a municipal solid waste landfill site. *Environmental Modeling and Software*, 21(8), 1190-1197.
- Karim, A.A., Tie, A.P.L., Manan, D.M.A. and Zaidul, I.S.M. (2008). Starch from the sago (Metroxylon sagu) palm tree—properties, prospects, and challenges as a new industrial source for food and other uses. *Comprehensive Reviews in Food Science and Food Safety*, 7(3), 215-228.
- Kazim, A. and Veziroglu, T.N. (2001). Utilization of solar-hydrogen energy in the UAE to maintain its share in the world energy market for the 21st century. *Renewable Energy*, 24(2), 259-274.

- Keskin, T. and Hallenbeck, P.L. (2012). Fundamental of the fermentative production of hydrogen: Baskar C., Baskar S., Dhillon R.S. (ed.). Biomass conversion: the interface of biotechnology, chemistry and materials science. *Springer, US*, p. 313.
- Khaleb, N.A., Jahim, J.Md. and Ahmad Kamal, S. (2012). Biohydrogen production using hydrolysates of palm oil mill effluent (POME). *Journal of Asian Scientific Research*, 2(11), 705-710.
- Khanal, S.K., Chen, W.H., Li, L. and Sung, S. (2004). Biological hydrogen production: Effects of pH and intermediate products. *International Journal of Hydrogen Energy*, 29, 1123-1131.
- Kim, J. R., Dec, J., Bruns, M. A., & Logan, B. E. (2008). Removal of odors from swine wastewater by using microbial fuel cells. *Applied and Environmental Microbiology*, 74(8), 2540-2543.
- Kim, S.H., Han, S.K. and Shin, H.S. (2006). Effect of substrate concentration on hydrogen production and 16S rDNA-based analysis of the microbial community in a continuous fermenter. *Process Biochemistry*, 41(1), 199-207.
- Kleerebezem, R. and van Loosdrecht, M.C.M. (2007). Mixed culture biotechnology for bioenergy production. *Current Opinion In Biotechnology*, 18(3), 207-212.
- Kongjan, P., O-Thong, S., Kotay, M., Min, B. and Angelidaki, I. (2010). Biohydrogen production from wheat straw hydrolysate by dark fermentation using extreme thermophilic mixed culture. *Biotechnology and Bioengineering*, 105(5), 899-908.
- Kotay, S.M. and Das, D. (2007). Microbial hydrogen production with *Bacillus coagulans* IIT-BT S1 isolated from anaerobic sewage sludge. *Bioresource Technology*, 98(6), 1183-1190.
- Kotay, S.M. and Das, D. (2008). Biohydrogen as a renewable energy resource - prospects and potentials. *International Journal of Hydrogen Energy* 33(1), 258-263.
- Krishna, S.H., Prasanth, K., Chowdry, G.V. and Ayyanna, C. (1998). Simultaneous saccharification and fermentation of pretreated sugar-cane leaves to ethanol. *Process Biochemistry*, 38, 825-830.
- Kumar, A., Jain, S.R., Sharma, C.B., Joshi, A.P. and Kalia, V.C. (1995). Increased H₂ production by immobilized microorganisms. *World Journal of Microbiology and Biotechnology*, 11(2), 156-159.
- Kumar, N. and Das, D. (2000). Enhancement of hydrogen production by *Enterobacter cloacae* IIT-BT 08. *Process Biochemistry*, 35(6), 589-593.

- Kumaran, S., Sastry, C. and Vikineswary, S. (1997). Laccase, cellulase and xylanase activity during growth of *Pleurotus sajor-caju* on sago hampas. *World Journal of Microbiology and Biotechnology*, 13, 43-49.
- Lay, J.J., Li, Y.Y. and Noike, T. (1997). Influences of pH and moisture content on the methane production in high-solids sludge digestion. *Water Research*, 31(6), 1518-1524.
- Lay, C.H., Kuo, S.Y., Sen, B., Chen, C.C., Chang, J.S. and Lin, C.Y. (2012). Fermentative biohydrogen production from starch-containing textile wastewater. *International Journal of Hydrogen Energy*, 37(2), 2050-2057.
- Lay, J.J., Lee, Y.J. and Noike, T. (1999). Feasibility of biological hydrogen production from organic fraction of municipal solid waste. *Water Research*, 33, 2576-2586.
- Lee, Y.J., Miyahara, T. and Noike, T. (2002). Effect of pH on microbial hydrogen fermentation. *Journal of Chemical Technology and Biotechnology*, 77(6), 694-698.
- Levin, D.B. and Chahine, R. (2010). Challenges for renewable hydrogen production from biomass. *International Journal of Hydrogen Energy*, 35(10), 1-8.
- Levin, D.B., Islam, R., Cicek, N. and Sparling, R. (2006). Hydrogen production by *Clostridium thermocellum* 27405 from cellulosic biomass substrates. *International Journal of Hydrogen Energy*, 31(11), 1496-1503.
- Levin, D.B., Pitt, L. and Love, M. (2004). Biohydrogen production: prospects and limitations to practical application. *International Journal of Hydrogen Energy*, 29(2), 173-185.
- Li, C. and Fang, H.H.P. (2007). Fermentative hydrogen production from wastewater and solid wastes by mixed cultures. *Critical Reviews in Environmental Science and Technology*, 37(1), 1-39.
- Li, D. and Chen, H. (2007). Biological hydrogen production from steam exploded straw by simultaneous saccharification and fermentation. *International Journal of Hydrogen Energy*, 32(12), 1742-1748.
- Lin, C.Y. and Chang, R.C. (2004). Fermentative hydrogen production at ambient temperature. *International Journal of Hydrogen Energy*, 29(7), 715-720.
- Lin, C.Y. and Jo, C.H. (2003). Hydrogen production from sucrose using an anaerobic sequencing batch reactor process. *Journal of Chemical Technology and Biotechnology*, 78(6), 678-684.
- Lin, C.Y. and Lay, C.H. (2004). Carbon/nitrogen-ratio effect on fermentative hydrogen production by mixed microflora. *International Journal of Hydrogen Energy*, 29(1), 41-45.

- Lin, C.Y., Wu, C.C. and Hung, C.H. (2008). Temperature effects on fermentative hydrogen production from xylose using mixed anaerobic cultures. *International Journal of Hydrogen Energy*, 33(1), 43-50.
- Lin, P.Y., Whang, L.M., Wu, Y.R., Ren, W.J., Hsiao, C.J., Li, S.L. and Chang, J.S. (2007). Biological hydrogen production of the genus *Clostridium*: Metabolic study and mathematical model simulation. *International Journal of Hydrogen Energy*, 32(12), 1728-1735.
- Lin, Y. and Tanaka, S. (2006). Ethanol fermentation from biomass resources: current state and prospects. *Applied Microbiology and Biotechnology*, 69(6), 627-642.
- Linggang, S., Phang, L.Y., Wasoh, M.H. and Abd-Aziz, S. (2012). Sago pith residue as an alternative cheap substrate for fermentable sugars production. *Applied Biochemistry and Biotechnology*, 167(1), 122-131.
- Liu, G. and Shen, J. (2004). Effects of culture and medium conditions on hydrogen production from starch using anaerobic bacteria. *Journal of Biosciences and Bioengineering*, 98, 251-256
- Liu, H., Grot, S. and Logan, B.E. (2005). Electrochemically assisted microbial production of hydrogen from acetate. *Environmental Science and Technology*, 39(11), 4317-4320.
- Liu, I.C., Whang, L.M., Ren, W.J. and Lin, P.Y. (2011). The effect of pH on the production of biohydrogen by clostridia: thermodynamic and metabolic considerations. *International Journal of Hydrogen Energy*, 36. 439-449.
- Lo, Y.C., Chen, W.M., Hung, C.H., Chen, S.D. and Chang, J.S. (2008). Dark H₂ fermentation from sucrose and xylose using H₂-producing indigenous bacteria: feasibility and kinetic studies. *Water Research*, 42(4-5), 827-842.
- Lo, Y.C., Lu, W.C., Chen, C.Y. and Chang, J.S. (2010). Dark fermentative hydrogen production from enzymatic hydrolysate of xylan and pretreated rice straw by *Clostridium butyricum* CGS5. *Bioresource Technology*, 101(15), 5885-5891.
- Lora, J. H. and Glasser, W. G. (2002). Recent industrial applications of lignin: a sustainable alternative to nonrenewable materials. *Journal of Polymers and the Environment*, 10(1-2), 39-48.
- Luo, G., Karakashev, D., Xie, L., Zhou, Q. and Angelidaki, I. (2011). Long-term effect of inoculum pretreatment on fermentative hydrogen production by repeated batch cultivations: Homoacetogenesis and methanogenesis as competitors to hydrogen production. *Biotechnology and Bioengineering*, 108(8), 1816-1827.

- Maddox, I.S., Steiner, E., Hirsch, S., Wessner, S., Gutierrez, N.A., Gapes, J.R. and Schuster, K.C. (2000). The cause of “acid crush” and “acidogenic fermentations” during the batch acetone-butanol-ethanol (ABE) fermentation process. *Journal of Molecular Microbiology and Biotechnology*, 2(1), 95-100.
- Manners, D.J. (1989). Recent developments in our understanding of amylopectin structure. *Carbohydrate Polymers*, 11(2), 87-112.
- Mars, A.E., Veuskens, T., Budde, M.A.W., Doeveren, P.F.N.M., Lips, S.J.J., Bakker, R.R., de Vrije, G.J. and Claassen, P.A.M. (2010). Biohydrogen production from untreated and hydrolyzed potato steam peels by the extreme thermophiles *Caldicellulosiruptor saccharolyticus* and *Thermotoga neapolitana*. *International Journal of Hydrogen Energy*, 35, 7730-7737.
- Martínez, I. N., Morán, J. M. and Peña, F. J. (2006). Two-Step Cluster Procedure After Principal Component Analysis Identifies Sperm Subpopulations in Canine Ejaculates and Its Relation to Cryoresistance. *Journal of Andrology*, 27(4), 596-603.
- Masset, J., Calusinska, M., Hamilton, C., Hiligsmann, S., Joris, B., Wilmotte, A. and Thonart, P. (2012). Fermentative hydrogen production from glucose and starch using pure strains and artificial co-cultures of *Clostridium* spp. *Biotechnology for Biofuels*, 5(1), 35.
- McMillan, J.D. (1994). Pretreatment of lignocellulosic biomass. In: Himmel, M.E., Baker, J.O. and Overend, R.P. (Eds.), *Enzymatic conversion of biomass for fuels production*, ACS Symposium Series, vol. 566. ACS, Washington, DC; 1994. p. 292–324.
- McMillan, J.D. (2004). Biotechnological routes to biomass conversion. DOE/NASULGC Biomass & Solar Energy Workshops, August 3-4, 2004, *National Renewable Energy Laboratory*.
- Miller, G.L. (1959). “Use of dinitrosalicylic acid reagent for determination of reducing sugars,” *Analysis Chemistry*, 31, 426-428.
- Mingzhi, H., Ma, Y., Jinquan, W., and Yan, W. (2009). Simulation of a paper mill wastewater treatment using a fuzzy neural network. *Expert Systems with Applications*, 36(3), 5064-5070.
- Montgomery, D.C. and Myers, R.H. (1995). Response surface methodology: process and product optimization using designed experiments. *Raymond H.M. and Douglas C.M., A Wiley-Interscience Publications*.
- Morohoshi, N. (1991). Chemical characterization of wood and its components. In: Hon, D.N-S. and Shiraishi, N. (eds) *Wood and cellulosic chemistry*. Marcel Dekker, New York, p. 331-392.

- Morteza, A., Nakhjavani, M., Asgarani, F., Carvalho, F.L., Karimi, R. and Esteghamati, A. (2013). Inconsistency in albuminuria predictors in type 2 diabetes: a comparison between neural network and conditional logistic regression. *Translational Research*, 161(5), 397-405.
- Mosier, N., Wyman, C., Dale, B., Elander, R., Lee, Y.Y., Holtzapple, M. and Ladicsh, M. (2005). Features of promising technologies for pretreatment of lignocellulosic biomass. *Bioresource Technology*, 96(6), 673–86.
- Mu, Y., Wang, G. and Yu, H.Q. (2006). Response surface methodological analysis on biohydrogen production by enriched anaerobic cultures. *Enzyme and Microbial Technology*, 38(7), 905-913.
- Mullai, P., Yogeswari, M.K. and Sridevi, K. (2013). Optimisation and enhancement of biohydrogen production using nickel nanoparticles – a novel approach. *Bioresource Technology*, doi: 10.1016/j.biortech.2013.03.082.
- Mussatto, S.I., Fernandes, M., Milagres, A.M. and Roberto, I.C. (2008). Effect of hemicellulose and lignin on enzymatic hydrolysis of cellulose from brewer's spent grain. *Enzyme and Microbial Technology*, 43(2), 124-129.
- Myers, R. H. and Montgomery, D. C., (1995). Response surface methodology: process and product optimization using designed experiments. *John Wiley & Sons, New York*.
- Nagata, Y. and Chu, K.H. (2003). Optimization of a fermentation medium using neural networks and genetic algorithms. *Biotechnology Letters*, 25(21), 1837-1842.
- Nakamura, L.K. (1981). *Lactobacillus amylovorus*, a new starch-hydrolyzing species from cattle waste corn fermentation. *International Journal of Systematic Bacteriology*, 31(1), 56–63.
- Nasr, N., Elbeshbishy, E., Hafez, H., Nakhla, G. and El Nagggar, M.H. (2011). Biohydrogen production from thin stillage using conventional and acclimatized anaerobic digester sludge. *International Journal of Hydrogen Energy*, 36(20), 12761-12769.
- Nasr, N., Hafez, H., El Nagggar, M.H. and Nakhla, G. (2013). Application of artificial neural networks for modeling of biohydrogen production. *International Journal of Hydrogen Energy*, 38(8), 3189-3195.
- Nath, K. and Das, D. (2004). Improvement of fermentative hydrogen production: various approaches. *Applied Microbiology and Biotechnology*, 65(5), 520-529.
- Nazlina, H.M.Y., Nor Aini, A.R., Ismail, F., Yusof, M.Z.M. and Hassan, M.A. (2009). Effect of different temperature, initial pH and substrate composition on biohydrogen production from food waste in batch fermentation. *Asian Journal of Biotechnology*, 1(2), 42-50.

- Ni, M., Leung, D.Y.C., Leung, M.K.H. and Sumathy, K. (2006). An overview of hydrogen production from biomass. *Fuel Processing Technology*, 87, 461-472.
- Ni, M., Leung, M.K., Leung, D.Y. and Sumathy, K. (2007). A review and recent developments in photocatalytic water-splitting using TiO₂ for hydrogen production. *Renewable and Sustainable Energy Reviews*, 11(3), 401-425.
- Nikhil, G.N., Mohan, S.V. and Swamy, Y.V. (2014). Behavior of acidogenesis during biohydrogen production with formate and glucose as carbon source: substrate associated dehydrogenase expression. *International Journal of Hydrogen Energy*, 39, 7486-7495.
- Niu, K., Zhang, X., Tan, W.S. and Zhu, M.L. (2011). Effect of culture conditions on producing and uptake hydrogen flux of biohydrogen fermentation by metabolic flux analysis method. *Bioresource Technology*, 102, 7294-7300.
- Noureldin, A., El-Shafie, A. and Bayoumi, M. (2011). GPS/INS integration utilizing dynamic neural networks for vehicular navigation. *Information Fusion*, 12(1), 48-57.
- Oehlert, G.W. (2000). *A first course in design and analysis of experiments* (Vol. 1). New York: WH Freeman.
- Oh, Y.K., Park, M.S., Seol, E.H., Lee, S.J. and Park, S. (2003). Isolation of hydrogen-producing bacteria from granular sludge of an upflow anaerobic sludge blanket reactor. *Biotechnology and Bioprocess Engineering*, 8(1), 54-57.
- Oh, Y.K., Seol, E.H., Kim, M.S. and Park, S. (2004). Photoproduction of hydrogen from acetate by a chemoheterotrophic bacterium *Rhodopseudomonas palustris* P4. *International Journal of Hydrogen Energy*, 29(11), 1115-1121.
- Oh, Y.K., Seol, E.H., Lee, E.Y. and Park, S. (2002). Fermentative hydrogen production by a new chemoheterotrophic bacterium *Rhodopseudomonas Palustris* P4. *International Journal of Hydrogen Energy*, 27(11), 1373-1379.
- O-Thong, S., Hniman, A., Prasertsan, P. and Imai, T. (2011). Biohydrogen production from cassava starch processing wastewater by thermophilic mixed cultures. *International Journal of Hydrogen Energy*, 36(5), 3409-3416.
- Ozawa, T., Takahiro, O., and Osama, N. (1996). Hemicelluloses in the fibrous residue of sago palm. *Proceeding of the Sixth International Sago Symposium, Pekan Baru*.
- Ozawa, T., Ueno, T., Negishi, O., & Masaki, S. (1998). Chemical characteristics of hemicelluloses in the fibrous residue of sago palm [Metroxylon]. *Japanese Journal of Tropical Agriculture*, 42.
- Özgür, E., Afsar, N., de Vrije, T., Yücel, M., Gündüz, U., Claassen, P.A.M. and Eroglu, I. (2010c). Potential use of thermophilic dark fermentation effluents in

photofermentative hydrogen production by *Rhodobacter capsulatus*. *Journal of Cleaner Production*, 18, S23-S28.

- Özgür, E., Mars, A.E., Peksel, B., Louwerse, A., Yücel, M., Gündüz, U., Classeen, P.A.M. and Eroğlu, İ. (2010b). Biohydrogen production from beet molasses by sequential dark and photofermentation. *International Journal of Hydrogen Energy*, 35(2), 511-517.
- Özgür, E., Uyar, B., Öztürk, Y., Yücel, M., Gündüz, U. and Eroğlu, İ. (2010a). Biohydrogen production by *Rhodobacter capsulatus* on acetate at fluctuating temperatures. *Resources, Conservation and Recycling*, 54(5), 310–314.
- Özkaya, B., Visa, A., Lin, C.Y., Puhakka, J.A. and Yli-Harja, O. (2008). an artificial neural network based model for predicting H₂ production rates in a sucrose-based bioreactor system. *International Journal of Mathematical, Physical and Engineering Sciences*, 2(2).
- Palmqvist, E. and Hahn-Hägerdal, B. (2000a). Fermentation of lignocellulosic hydrolysates. I: inhibition and detoxification. *Bioresource Technology*, 74(1), 17-24.
- Palmqvist, E. and Hahn-Hägerdal, B. (2000b). Fermentation of lignocellulosic hydrolysates. II: inhibitors and mechanisms of inhibition. *Bioresource Technology*, 74(1), 25-33.
- Pan, C.M., Fan, Y.T., Xing, Y., Hou, H.W. and Zhang, M.L. (2008). Statistical optimization of process parameters on biohydrogen production from glucose by *Clostridium* sp. Fanp2. *Bioresource Technology*, 99(8), 3146-3154.
- Pandu, K. and Joseph, S. (2012). Comparisons and limitations of biohydrogen production processes: a review. *International Journal of Advance Engineering and Technology*, 2, 342-356.
- Passenier, A. and Lak, M. (2009). Scarcity and transition: Research questions for future policy. *The Hague, Ministry of Housing, Spatial Planning and the Environment and Ministry of Foreign Affairs*, 74.
- Patra, S., Sangyoka, S., Boonmee, M. and Reungsang, A. (2008). Bio-hydrogen production from the fermentation of sugarcane bagasse hydrolysate by *Clostridium butyricum*. *International Journal of Hydrogen Energy*, 33, 5256-5265.
- Patwardhan, A.D. (2008). Industrial wastewater treatment. technology and engineering, *PHI Learning PVT. Ltd., New Delhi*. ISBN-978-81-203-3350-5. pp: 235–236.
- Perego, C. and Bianchi, D. (2010). Biomass upgrading through acid–base catalysis. *Chemical Engineering Journal*, 161(3), 314-322.

- Perego, P., Fabiano, B., Ponzano, G.P. and Palazzi, E. (1998). Experimental study of hydrogen kinetics from agroindustrial by-product: optimal conditions for production and fuel cell feeding. *Bioprocess Engineering*, 19(3), 205-211.
- Perera, K.R.J., Ketheesan, B., Gadhamshetty, V. and Nirmalakhandan, N. (2010). Fermentative biohydrogen production: evaluation of net energy gain. *International Journal of Hydrogen Energy*, 35(22), 12224-12233.
- Phang, S. M., Miah, M. S., Yeoh, B. G., & Hashim, M. A. (2000). Spirulina cultivation in digested sago starch factory wastewater. *Journal of Applied Phycology*, 12(3-5), 395-400.
- Pin, C. and Baranyi, J. (2006). Kinetic of single cells: observation and modeling of a stochastic process. *Applied and Environmental Microbiology*, 72, 2163-2169.
- Potter, A.L. and Hassid, W.Z. (1948). Starch. II. Molecular Weights of Amyloses and Amylopectins from Starches of Various Plant Origins. *Journal of the American Chemical Society*, 70(11), 3774-3777.
- Prakasham, R.S., Brahmaiah, P., Sathish, T. and Sambasiva Rao, K.R.S. (2009). Fermentative biohydrogen production by mixed anaerobic consortia: impact of glucose to xylose ratio. *International Journal of Hydrogen Energy*, 34(23), 9354-9361.
- Prakasham, R.S., Sathish, T. and Brahmaiah, P. (2010). Biohydrogen production process optimization using anaerobic mixed consortia: A prelude study for use of agro-industrial material hydrolysate as substrate. *Bioresource Technology*, 101(14), 5708-5711.
- Prasertsan, P., O-Thong, S. and Birkeland, N. (2009). Optimization and microbial community analysis for production of biohydrogen from palm oil mill effluent by thermophilic fermentation process. *International Journal of Hydrogen Energy*, 34(17), 7448-7459.
- Qian, C.X., Chen, L.Y., Rong, H. and Yuan, X.M. (2011). Hydrogen production by mixed culture of several facultative bacteria and anaerobic bacteria. *Progress in Natural Science: Materials International*, 21(6), 506-511.
- Rajasimman, M. and Karthikeyan, C. (2007). Aerobic digestion of starch wastewater in a fluidized bed bioreactor with low density biomass support. *Journal of Hazardous Materials*, 143(1), 82-86.
- Rajeshwar, K., McConnell, R. and Licht, S. (2008). Solar hydrogen generation. toward a renewable energy future. *Springer-Verlag: New York*.
- Rajoka, M.I., Durani, I.S. and Khalid, A.M. (2004). Kinetics of improved production and thermostability of an intracellular β -glucosidase from mutant-derivative of *Cellulomonas biozotea*. *Biotechnology Letters*, 26, 281-285.

- Ramey, D. and Shang, T.Y. (2004). Production of butyric acid and butanol from biomass: final report, Environmental Energy Inc. Department of Chemical and Biomolecular Engineering, The Ohio State University.
- Razak, M.N.A., Ibrahim, M.F., Phang, L.Y., Hassan, M.A. and Abd-Aziz, S. (2012). Utilization of oil palm decanter cake for cellulase and polyoses production. *Biotechnology and Bioprocess Engineering*, 17(3), 547-555.
- Reeve, A. (1992). Starch hydrolysis: Process and equipment, Schenk, F.W. and Hebeda, R.E.(ed.). Starch hydrolysis products. *VCH Publishers Inc. New York*, pp. 79-120.
- Ren, N., Xing, D., Rittmann, B.E., Zhao, L., Xie, T. and Zhao, X. (2007). Microbial community structure of ethanol type fermentation in bio-hydrogen production. *Environment Microbiology*, 9(5), 1112-1125.
- Rifat, A., Paramaswari, S., Abdullah, N. and Sekaran, M. (2003). Optimization of laccase productivity during solid substrate fermentation of sago *hampas* by *Pycnoporus sanguineus*, *Malaysian Journal of Science*, 22, 35-42.
- Robledo-Narváez, P.N., Muñoz-Páez, K.M., Poggi-Varaldo, H.M., Ríos-Leal, E., Calva-Calva, G., Ortega-Clemente, L.A., Noemí Rinderknecht-Seijas, N., Estrada-Vázquez, C., Ponce-Noyola, M.T. and Salazar-Montoya, J.A. (2013). The influence of total solids content and initial pH on batch biohydrogen production by solid substrate fermentation of agroindustrial wastes. *Journal of Environmental Management*, 128, 126-137.
- Rodrigues, L., Teixeira, J., Oliveira, R. and Van Der Mei, H. C. (2006). Response surface optimization of the medium components for the production of biosurfactants by probiotic bacteria. *Process Biochemistry*, 41(1), 1-10.
- Roychowdhury, S., Cox, D. and Levandowsky, M. (1988). Production of hydrogen by microbial fermentation. *International Journal of Hydrogen Energy*, 13(7), 407-410.
- Rubin, E.M. (2008). Genomic of cellulosic biofuels. *Nature* 454, 841-845.
- Saha, B. C. (2003). Hemicellulose bioconversion. *Journal of Industrial Microbiology and Biotechnology*, 30(5), 279-291.
- Saint-Amans, S., Girbal, L., Andrade, J., Ahrens, K. and Soucaille, P. (2001). Regulation of carbon and electron flow in *Clostridium butyricum* VPI 3266 grown on glucose-glycerol mixtures. *Journal of Bacteriology*, 183(5), 1748-1754.
- Saratale, G.D., Chen, S.D., Lo, Y.C., Saratale, R.G. and Chang, J.S. (2008). Outlook of biohydrogen production from lignocellulosic feedstock using dark fermentation – a review. *Journal of Scientific and Industrial Research*, 67, 962-979.

- Saravanane, R., D.V.S. Murthy and K. Krishnaiah (2001). Anaerobic treatment and biogas recovery for sago wastewater management using a fluidized bed reactor. *Water Science and Technology*, 44, 141-147.
- Sarawak agriculture statistics 2010, exports of sago starch by country of destination 2006–2010, <http://www.doa.sarawak.gov.my/modules/web/page.php?id=504&se=sagostarch>.
- Saunders, J., Izydorczyk, M. and Levin D.B. (2011). Limitations and challenges for wheat-based bioethanol production, economic effects of biofuel production, Dr. Marco Aurelio Dos Santos Bernardes (Ed.), ISBN: 978-953-307-178-7, InTech, DOI: 10.5772/20258. Available from: <http://www.intechopen.com/books/economic-effects-of-biofuel-production/limitations-and-challenges-for-wheat-based-bioethanol-production>.
- Senic, G. (2002). Small-scale food processing enterprises in Malaysia, MARDI Malaysia: http://www.agnet.org/htmlarea_file/library/20110716000322/eb409.pdf.
- Shahrmi, Z., Sabaratnam, V., Rahman, N.A.A., Abd-Aziz, S., Hassan, M.A., Karim, M.I.A. (2008). Production of reducing sugars by *Trichoderma* sp. KUPM0001 during solid substrate fermentation of sago starch processing waste hampas. *Research Journal of Microbiology*, 3, 569-579.
- Shi, H., Gao, Y. and Wang, X. (2010). Optimization of injection molding process parameters using integrated artificial neural network model and expected improvement function method. *The International Journal of Advanced Manufacturing Technology*, 48(9-12), 955-962.
- Shin, H.S., Youn, J.H. and Kim, S.H. (2004). Hydrogen production from food waste in anaerobic mesophilic and thermophilic acidogenesis. *International Journal of Hydrogen Energy*, 29(13), 1355-1363.
- Singhal, R.S., Kennedy, J.F., GopalaKrishnan, S.M., Kaczmarek, A., Knill, C.J. and Akmar, P.F. (2008). Industrial production, processing and utilization of sago palm-derived products. *Carbohydrate Polymer*, 17(1), 1-20.
- Smith, A.M. (1999). Making starch. *Current Opinion Plant Biology*, 2, 223-229.
- Smith, J.L. Fratamico, P.M. and Novak, J.S. (2004). Quorum sensing: a primer for food microbiologists. *Journal of Food Protection*, 67, 1053.
- Souza, P.M. and Magalhaes, P.D.O.E. (2010). Application of microbial α -amylase production by *Bacillus* sp. using response surface methodology. *Process Biochemistry*, 40, 2291-2296.

- Sreethawong, T., Chatsiriwatana, S., Rangsunvigit, P. and Chavadej, S. (2010). Hydrogen production from cassava wastewater using an anaerobic sequencing batch reactor: Effects of operational parameters, COD:N ratio, and organic acid composition. *International Journal of Hydrogen Energy* 35(9), 4092-4102.
- Steinfeld, A. (2005). Solar thermochemical production of hydrogen – a review. *Solar Energy* 78, 603-615.
- Sun, Y. and Cheng, J.Y. (2002). Hydrolysis of lignocellulosic materials for ethanol from lignocellulosic materials: a review. *Bioresource Technology*, 83, 1-11.
- Taguchi, F., Yamada, K., Hasegawa, K., Saito-Taki, T. and Hara, K. (1996). Continuous hydrogen production by *Clostridium* sp. strain no. 2 from cellulose hydrolysate in an aqueous two-phase system. *Journal of Fermentation and Bioengineering*, 82, 180–183.
- Taherzadeh, M.J. and Kamiri, K. (2007). Enzyme-based hydrolysis processes for ethanol from lignocellulosic materials: a review. *Bioresource Technology*, 2(4), 707-38.
- Tangphatsornruang, S., Naconsie, M., Thammarongtham, C. and Narangajavana, J. (2005). Isolation and characterization of an α -amylase gene in cassava (*Manihot esculenta*). *Plant Physiology and Biochemistry*, 43(9), 821-827.
- Tao, Y., Chen, Y., Wu, Y., He, Y. and Zhou, Z. (2007). High hydrogen yield from a two-step process of dark- and photo-fermentation of sucrose. *International Journal of Hydrogen Energy*, 32, 200 – 206.
- Terracciano, J.S. and Kashket, E.R. (1986). Intracellular conditions required for initiation of solvent production by *Clostridium acetobutylicum*. *Applied Environmental Microbiology*, 52(1), 86-91.
- Tester, R.F. and Karkalas, J. (1996). Swelling and gelatinization of oat starches. *Cereal Chemistry*, 73(2), 271-277.
- Tsuru, T., Yamaguchi, K., Yoshioka, T. and Asaeda, M. (2004). Methane steam reforming by microporous catalytic membrane reactors. *American Institute of Chemical Engineers Journal*, 50(11), 2794-2805.
- Turner, J., Sverdrup, G., Mann, M.K., Maness, P.C., Kroposki, B. and Ghirardi, M. (2008). Renewable hydrogen production. *International Journal of Hydrogen Energy*, 32, 379–407.
- Ueno, Y., Haruta, S., Ishii, M. and Igarashi, Y. (2001). Microbial community in anaerobic hydrogen-producing microflora enriched from sludge compost. *Applied Microbiology and Biotechnology*, 57(4), 555-562.

- Ueno, Y., Sato, S. and Morimoto, M. (1996). Hydrogen production from industrial wastewater by anaerobic microflora in chemostate culture. *Journal of Fermentation and Bioengineering*, 82(2), 194-197.
- Umi Kalsom, M.S., Ariff, A.B. and Zulkifli, H.S. (1997). The treatment of oil palm empty fruit bunch fibre for subsequent use as substrate for cellulase production by *Chaetomium globosum* Kunze. *Journal of Bioresources Technology*, 62, 1-9.
- Ursu'a, A., Gandi'a, L.M. and Sanchis P. (2012). Hydrogen production from water electrolysis: current status and future trends. *Institute of Electrical and Electronic Engineers*, 100(2), 410-426.
- van der Maarel, M.J.E.C., van der Veen, B., Uitdehaag, J.C.M., Leemhuis, H. and Dijkhuizen, L. (2002). Properties and applications of starch converting enzymes of the α -amylase family. *Journal of Biotechnology*, 94, 137-155.
- van Ginkel, S. and Logan, B.E. (2005). Inhibition of biohydrogen production by undissociated acetic and butyric acids. *Environmental Science and Technology*, 39(23), 9351-9356.
- van Ginkel, S., Sung, S., and Lay, J.J. (2001). Biohydrogen production as a function of pH and substrate concentration. *Environmental Science and Technology*, 35(24), 4726-4730.
- Vikineswary, S., Shim, Y.L., Thambirajah, J.J. and Blakebrough, N. (1994). Possible microbial utilization of sago processing wastes. *Resources, Conservation and Recycling*, 11(1), 289-296.
- Walter, R.P., Morris, J.G. and Kell, D.B. (1987). The roles of osmotic stress and water activity in the inhibition of the growth, glycolysis and glucose phosphotransferase system of *Clostridium pasteurianum*. *Journal of General Microbiology*, 133(2), 259-66.
- Wang, C.C., Chang, C.W., Chu, C.P., Lee, D.J., Chang, B.V. and Liao, C.S. (2003). Producing hydrogen from wastewater sludge by *Clostridium bifermentans*. *Journal of Biotechnology*, 102(1), 83-92.
- Wang, J.L and Wan, W. (2008). Factors influencing fermentative hydrogen production: a review. *International Journal of Hydrogen Energy*, 34(2), 799-811.
- Wang, J.L. and Wan, W. (2009). Effect of temperature on fermentative hydrogen production by mixed culture. *International Journal of Hydrogen Energy*, 33, 587-593.
- Wang, L., Zhou, Q. and Li, F.T. (2006). Avoiding propionic acid accumulation in the anaerobic process for biohydrogen production. *Biomass Bioenergy*, 30(2), 177-182.

- Wang, W.J., Powell, A.D. and Oates, C.G. (1996). Sago starch as a biomass source: raw sago starch hydrolysis by commercial enzymes. *Bioresource Technology*, 55(1), 55-61.
- Wang, Z.L., Naterer, G.F., Gabriel, K.S., Gravelins, R. and Daggupati, V.N. (2010) comparison of sulphur-iodine and cooper-chlorine thermochemical hydrogen production. *International Journal of Hydrogen Energy*, 35, 4820-4830.
- Wei, D., Liu, X. and Yang, S.T. (2013). Butyric acid production from sugarcane bagasse hydrolysate by *Clostridium tyrobutyricum* immobilized in a fibrous-bed bioreactor. *Bioresource Technology*, doi:10.1016/j.biortech.2012.11.065.
- Wei, H., Bing, W., Xiaoye, L., Chunyu, L., Liran, Y. and Yongfeng, L. (2012). Fermentative hydrogen production from molasses in an activated sludge immobilized bioreactor. *International Journal of Energy and Engineering*, 2(1), 28–31.
- Wei, J., Liu, Z.T., and Zhang, X. (2010). Biohydrogen production from starch wastewater and application in fuel cell. *International Journal of Hydrogen Energy*, 35 , 2949–2952.
- Winter, C.J. (2009). Hydrogen energy-Abundant, efficient, clean: A debate over the energy-system-of-change. *International Journal of Hydrogen Energy*, 34, S1–S52.
- Wood, T.M. and Bhat, K.M. (1988). Methods for measuring cellulases activity, Wood, W.A. and Kellogg, S.T. (ed.). *Methods in Enzymology*, Academic Press Inc, London, 160, 87-112.
- Yang, H., Shao, P., Lu, T., Shen, J., Wang, D., Xu, Z. and Yuan, X. (2006). Continuous bio-hydrogen production from citric acid wastewater via facultative anaerobic bacteria. *International Journal of Hydrogen Energy*, 31(10) 1306–1313
- Yean, C.T. and Lan, S.Y. (1993). Sago processing wastes. In Yeoh et al. (eds) *Waste management in malaysia: current status and prospects for bioremediation. Ministry of Science, Technology and Environment of Malaysia*, pp. 159-167.
- Yinghuai, Z., Yuanting, K.T. and Hosmane, N.S. (2013). Applications of Ionic Liquids in Lignin Chemistry, *Ionic Liquids - New Aspects for the Future*, Dr. Jun-ichi Kadokawa (Ed.), ISBN: 978-953-51-0937-2, InTech, DOI: 10.5772/51161. Available from: <http://www.intechopen.com/books/ionic-liquids-new-aspects-for-the-future/applications-of-ionic-liquids-in-lignin-chemistry>.
- Yokoi, H., Saitsu, A., Uchida, H., Hirose, J., Hayashi, S. and Takasaki, Y. (2001). Microbial hydrogen production from sweet potato starch residue. *Journal of Biosciences and Bioengineering*, 91(1), 58-63.

- Yokoi, H., Tokushige, T., Hirose, J., Hayashi, S. and Takasaki, Y. (1997). Hydrogen production by immobilized cells of aciduric *Enterobacter aerogenes* strain HO-39. *Journal of Fermentation and Bioengineering*, 83(5), 481-484.
- Yokoi, H., Tokushige, T., Hirose, J., Hayashi, S. and Takasaki, Y. (1998). H₂ production from starch by mixed culture of *Clostridium buytricum* and *Enterobacter aerogenes*. *Biotechnology Letters*, 20, 143-147.
- Zamzuri, N.A. (2010). Screening and isolation of biobutanol producing bacteria, Undergraduate Thesis Faculty of Biotechnology and Biomolecular Sciences, Universiti Putra Malaysia. UPM/FBSB/TBP/0910/04.
- Jamil, Z., Annuar, M.S.M. and Ibrahim, S. (2009). Optimization of phototrophic hydrogen production by *Rhodospseudomonas palustris* PBUM001 via statistical experimental design. *International Journal of Hydrogen Energy*, 34(15), 7502-7512
- Zhang, H., Bruns, M.A. and Logan, B.E. (2006). Biological hydrogen production by *Clostridium acetobutylicum* in an unsaturated flow reactor. *Water Research*, 40(4), 728-734.
- Zhang, T., Liu, H., & Fang, H.H. (2003). Biohydrogen production from starch in wastewater under thermophilic condition. *Journal of Environmental Management*, 69(2), 149-156.
- Zhao, L., Montville, T.J., Schaffner, D.W. (2006). Evidence for quorum sensing in *Clostridium botulinum* 56A. *Letters in Applied Microbiology*, 42, 54-58.
- Zhao, X., Xing, D., Liu, B., Lu, L., Zhao, J. and Ren, N. (2012). The effects of metal ions and l-cysteine on *hyd A* gene expression and hydrogen production by *Clostridium beijerinckii* RZF-1108. *International Journal of Hydrogen Energy*, 37(18), 13711-13717.