

Malaysian Journal of Mathematical Sciences 10(3): 345–360 (2016)



MALAYSIAN JOURNAL OF MATHEMATICAL SCIENCES

Journal homepage: <http://einspem.upm.edu.my/journal>

## Weighted Block Runge-Kutta Method for Solving Stiff Ordinary Differential Equations

Saufianim Jana Aksah<sup>1</sup>, Zarina Bibi Ibrahim\*<sup>1</sup>, Yong Faezah Rahim<sup>2</sup>, and Siti Nur Iqmal Ibrahim<sup>1</sup>

<sup>1</sup>*Institute for Mathematical Research and Department of Mathematics, Faculty of Science, Universiti Putra Malaysia, Selangor, Malaysia*

<sup>2</sup>*Centre of Foundation Studies for Agricultural Science, Universiti Putra Malaysia, Selangor, Malaysia*

*E-mail: [zarinabb@upm.edu.my](mailto:zarinabb@upm.edu.my)*

*\*Corresponding author*

### ABSTRACT

In this paper, weighted block Runge-Kutta (WBRK) method is derived for solving stiff ordinary differential equations (ODEs). Implementation of weights on the method and its stability region are shown. Numerical results of the WBRK method are presented and compared with the existing methods to prove the ability of the proposed method to solve stiff ODEs. The results show that the WBRK method has better accuracy than the comparing methods.

**Keywords:** Runge-Kutta method, weights, block method, stiff ODEs, Centroidal mean.

# 1. Introduction

A differential equation (DE) is an equation that involves one or more derivatives of some unknown functions. Many problems in engineering, physical and social sciences are reduce to quantifiable form through the process of mathematical modelling which involved DEs. An example of modelling a real world problem using DE is in population change which is affected by births, death, immigration, and emigration. In most applicable situations, the exact solution of the DEs that model the problem is too complicated to be solved analytically. Hence, it is convenient to make approximations of the solution by numerical methods.

There are two main approaches for numerical techniques, namely, linear multistep method (LMM), and the one-step method. Adams method is widely known for the LMM users, while RK method has been used extensively in a one-step algorithm. The interest of using the one-step method, specifically RK method, over LMM is due to the advantage of RK method which requires no additional starting values and ability to readily change the steplength during computation. We use the explicit RK method which exhibits smaller computational cost than the implicit one which obtained through Taylor series as shown in Lambert (1973). Previously, it is known that implicit RK is more suitable to solve stiff problems however, Wu (1998) developed a sixth order A-stable explicit one-step method to show that an explicit method is efficient to solve stiff problems too.

A weighted fifth-order RK formulas for second-order differential equations proposed by Evans and Yaakub (1998) is an extension from Evans and Yaakub (1996) who introduced a new fifth order five stage Arithmetic Mean (AM) weighted RK (WRK) method. The implementation of weights on the method improves the efficiency of the method to solve problem of DEs. RK formula based on variety of means for solving the system of initial value problems (IVPs) as suggested by Murugesan et al. (2001) is then extended by Pushpam and Dhayabaran (2011) to solve stiff non-linear system. Ababneh and Rozita (2009) derived a weighted third order RK method based on Contraharmonic Mean (WRK3CoM) that is suitable to solve stiff problem. Sharmila and Amirtharaj (2011) used a modified weighted RK method based on Centroidal Mean (MWRK3CeM) for solving stiff IVPs, and shows that it is more effective than WRKCoM.

Many researchers especially mathematicians have extended the existing method or developed a new method to solve the problem efficiently by reducing the computational cost and obtained smaller error. The method that is

capable to evaluate functions in block of several of steps at once instead of one step at a time is a block method. Milne (1953) introduced a block method for numerical solutions of the first and higher order ODEs to be used as a starting value for a predictor-corrector algorithm. A block diagonally implicit RK (BDIRK) method by Cash (1983) is used to solve stiff and non-stiff problems of IVPs in ODEs. Then, Cash (1985) extended the method to form a block embedded explicit RK method which has the characteristics of standard explicit RK formula except for the function evaluation that is done in several steps at once. The derivation of BDIRK is also available in Rahim (2004). Majid and Suleiman (2007) developed a four-point fully implicit block method for solving first order ODEs by using variable stepsize. A block backward differentiation formula (BBDF) of variable step for solving stiff ODEs proposed by Ibrahim et al. (2007) have better accuracy with reduction of total steps and lesser computational time when compared with classical BDF by Suleiman (1979). The BBDF method is then improved in terms of its accuracy by several researchers, such as Yatim et al. (2011), Zawawi et al. (2012) and Ismail et al. (2014). Technique of partitioning introduced by Othman et al. (2007) which is based on BBDF proposed by Ibrahim et al. (2007) is proved to reduce the cost of the iteration scheme.

In this paper, we consider the following first order ODE of the form

$$\mathbf{y}' = A\mathbf{y} + \phi(x) \quad (1)$$

where  $\mathbf{y}^T = (y_1, y_2, \dots, y_m)$  and  $A$  is an  $m \times m$  matrix with eigenvalues  $\lambda_t$ ,  $t = 1, 2, \dots, m$ . According to Lambert (1993),

**Definition 1:** The system of first order ODE as shown in 1 is said to be stiff if

- 1)  $Re(\lambda_t) < 0, t = 1, 2, \dots, m$ .
- 2)  $max_t |Re(\lambda_t)| \gg min_t |Re(\lambda_t)|$  where  $\lambda_t$  are the eigenvalues of  $A$ .

As shown in Lambert (1973), stiffness is dependent only on the large ratio of the magnitudes of the negative real parts of the largest eigenvalue to that of the smallest one. Stiffness ratio is defined as  $S = \frac{max_t |Re(\lambda_t)|}{min_t |Re(\lambda_t)|}$ , hence stiff problem has  $S \gg 1$ .

Derivation of the WBRK method is shown in Section 2. The stability of WBRK is provided in Section 3. Section 4 presents the numerical results of the proposed method when tested with problems of stiff ODEs. Finally, we conclude findings of this research in Section 5.

## 2. Derivation of WBRK

Murugesan et al. (2001) extended the RK method based on variety of means which include AM, Geometric Mean (GM), Harmonic Mean (HaM), Heronian Mean (HeM), Root Mean Square (RMS), CeM, and CoM for solving system of IVPs. As shown in the paper, general form of CeM in terms of AM and GM used to derive the WBRK method is presented as follows.

$$\frac{4(AM)^2 - (GM)^2}{3(AM)}$$

or in general form of

$$\frac{2}{3} \left( \frac{x_1^2 + x_1x_2 + x_2^2}{x_1 + x_2} \right) \tag{2}$$

Since WBRK method is based on third order RK, we will use the MWRK3CeM method by Sharmila and Amirtharaj (2011) as a foundation of our proposed method. Standard form of MWRK3CeM with implementation of weights is shown as follows

$$y_{n+1} = y_n + \frac{2}{3}h \left( w_1 \frac{k_1^2 + k_1k_2 + k_2^2}{k_1 + k_2} + w_2 \frac{k_2^2 + k_2k_3 + k_3^2}{k_2 + k_3} \right) \tag{3}$$

where

$$\begin{aligned} k_1 &= f(x_n, y_n), \\ k_2 &= f(x_n + a_1h, y_n + ha_1k_1), \\ k_3 &= f(x_n + (a_2 + a_3)h, y_n + h(a_2k_1 + a_3k_2)). \end{aligned}$$

Equation (3) can be transformed into a block method by integrates forward over a step  $2h$  on the Taylor series expansion of  $k_i$  with  $i = 1, 2, 3$  which yields

$$\begin{aligned} k_1 &= f, \\ k_2 &= f + 2ha_1ff_y + 2h^2f^2a_1^2f_{yy} + \frac{4}{3}h^3f^3a_1^3f_{yyy}, \\ k_3 &= f + 2h(a_2 + a_3)ff_{yy} + 4h^2 \left( a_1a_3ff_y^2 + \frac{1}{2}(a_2 + a_3)^2f^2f_{yy} \right) + \\ & 8h^2 \left( \frac{1}{2}a_1^2a_3f^2f_yf_{yy} + a_1a_3(a_2 + a_3)f^2f_yf_{yy} + \frac{1}{6}(a_2 + a_3)^3f^3f_{yyy} \right) \end{aligned} \tag{4}$$




Figure 1: Block explicit RK method of constant stepsize

Figure 1 shows the computation works of WBRK method.

By substituting Equation (4) into Equation (3), we obtain an expression  $y_{n+2}$  in terms of the function with parameter  $a_i$  for  $i = 1, 2, 3$ , and its derivatives. The formula obtained involves a division of two series as follows

$$\sum_{i=1}^2 \frac{k_i^2 + k_i k_{i+1} + k_{i+1}}{k_i + k_{i+1}} \tag{5}$$

These problems can be solved by cross multiplying the series with the common denominator  $(k_1 + k_2)(k_2 + k_3)$ . Hence, it can be written as

$$y_{n+1} = y_n + \frac{U}{L} \tag{6}$$

with

$$U = 2(2h)(w_1(k_1^2 + k_1 k_2 + k_2^2)(k_2 + k_3) + w_2(k_2^2 + k_2 k_3 + k_3^2)(k_1 + k_2))$$

and

$$L = 3(k_2 + k_3)(k_1 + k_2)$$

The Taylor series expansion of  $y(x_{n+2})$ ,  $T$ , can be written as such

$$T = y_n + (2h)f + \frac{1}{2}(2h)^2 f f_y + \frac{1}{6}(2h)^3 (f f_y^2 + f^2 f_{yy})$$

The error of the method,  $E$ , is measured by the following expression

$$E = y(x_{n+2}) - y_{n+1}.$$

This yields

$$E = T - \frac{U}{L}. \tag{7}$$

Then, by comparing the coefficients of like terms for Equation (7) up to term  $h^3$  yields the following

$$hf^3 : 24w_1 + 24w_2 - 24 = 0 \tag{8}$$

$$h^2 f^3 f_y : 72w_1 a_1 + 24w_1 a_2 + 24w_1 a_3 + 72w_2 a_1 + 48w_2 a_2 + 48w_2 a_3 - 48a_1 - 24a_2 - 24a_3 - 24 = 0 \tag{9}$$

$$h^3 f^3 f_y^2 : 80w_1 a_1^2 + 80w_2 a_1^2 + 32w_2 a_2^2 + 32w_2 a_3^2 + 48w_1 a_1 a_2 + 96w_1 a_1 a_3 + 80w_2 a_1 a_2 + 176w_2 a_1 a_3 + 64w_2 a_2 a_3 - 24a_1^2 - 24a_1 a_2 - 72a_1 a_3 - 48a_1 - 24a_2 - 24a_3 - 16 = 0 \tag{10}$$

$$h^3 f^4 f_{yy} : 72w_1 a_1^2 + 24w_1 a_2^2 + 24w_1 a_3^2 + 72w_2 a_1^2 + 48w_2 a_2^2 + 48w_2 a_3^2 + 48w_1 a_2 a_3 + 96w_2 a_2 a_3 - 48a_1^2 - 24a_2^2 - 24a_3^2 - 48a_2 a_3 - 16 = 0 \tag{11}$$

Equations (8), (9), (10) and (11) are solved for the chosen weights to obtain a set of parameters. Following Evans and Yaakub (1998), the set of weights chosen must satisfy  $\sum_{i=1}^2 = 1$ . Hence, we randomly choose weights,  $w_1 = w_2 = \frac{1}{2}$ . On that account, we obtain the following set of parameters

$$a_1 = \frac{2}{3}, a_2 = -\frac{2}{9}, a_3 = \frac{8}{9}.$$

By substituting the weights and the set of parameters into Equation (3) results the following equation

$$y_{n+1} = y_n + \frac{1}{3}h \left( w_1 \frac{k_1^2 + k_1 k_2 + k_2^2}{k_1 + k_2} + w_2 \frac{k_2^2 + k_2 k_3 + k_3^2}{k_2 + k_3} \right), \tag{12}$$

$$y_{n+2} = y_n + \frac{2}{3}h \left( w_1 \frac{k_1^2 + k_1 k_2 + k_2^2}{k_1 + k_2} + w_2 \frac{k_2^2 + k_2 k_3 + k_3^2}{k_2 + k_3} \right),$$

where

$$k_1 = f(x_n, y_n),$$

$$k_2 = f\left(x_n + \frac{2}{3}h, y_n + \frac{2}{3}hk_1\right), \tag{13}$$

$$k_3 = f\left(x_n + \frac{2}{3}h, y_n + h\left(-\frac{2}{9}k_1 + \frac{8}{9}k_2\right)\right).$$

This completes the derivation.

### 3. Stability of WBRK

In this section, we analyze the stability of WBRK method. As referred in Rahim (2004), absolute stability can be defined as follows.

**Definition 2:** By letting  $z = h\lambda$ , then  $R(z)$  is known as the stability function of the method. Hence  $y_n \rightarrow 0$  as  $n \rightarrow \infty$  if and only if

$$|R(z)| < 1 \tag{14}$$

and the method is absolutely stable for those value of  $z$  for which Equation (14) holds. The region  $R_A$  of the complex  $z$ -plane which the equation holds is the region of absolute stability.

In order to construct the stability region of the method, we have to find the stability polynomial. By considering the following test equation,

$$\begin{aligned} y' &= f(x, y) \\ &= \lambda y \end{aligned} \tag{15}$$

We substitute Equation (15) into Equation (13) which gives us the following

$$\begin{aligned} k_1 &= f(x_n, y_n) = \lambda y_n, \\ k_2 &= f\left(x_n + \frac{2}{3}h, y_n + \frac{2}{3}hk_1\right) = \lambda y_n \left(1 + \frac{2}{3}h\lambda\right), \\ k_3 &= f\left(x_n + \frac{2}{3}h, y_n + h\left(-\frac{2}{9}k_1 + \frac{8}{9}k_2\right)\right) = \lambda y_n \left(1 + \frac{2}{3}h\lambda + \frac{16}{27}h^2\lambda^2\right). \end{aligned} \tag{16}$$

By substituting Equation (16) into Equation (12), and letting  $\frac{y_{n+1}}{y_n} = R(h\lambda)$ , we obtain the following

$$\begin{aligned} R(h\lambda) &= (13122 + 26244h\lambda + 26487h^2\lambda^2 + 16848h^3\lambda^3 + 6804h^4\lambda^4 + \\ &\quad 1920h^5\lambda^5 + 256h^6\lambda^6) \frac{1}{162(3 + h\lambda)(27 + 18h\lambda + 8h^2\lambda^2)} \end{aligned} \tag{17}$$

Then, we let  $z = h\lambda$  into Equation (17) to obtain

$$\begin{aligned} R(z) &= (13122 + 26244z + 26487z^2 + 16848z^3 + 6804z^4 + \\ &\quad 1920z^5 + 256z^6) \frac{1}{162(3 + z)(27 + 18z + 8z^2)} \end{aligned} \tag{18}$$

Equation (18) is known as stability function or stability polynomial of the method. Clearly,  $y_n \rightarrow 0$  as  $n \rightarrow \infty$  if and only if when  $|R(z)| < 1$  which satisfies the condition mentioned in **Definition 2**. Hence, the WBRK method is absolute stable. By using MAPLE, the stability regions of the method are presented as follows.




Figure 2: Stability region of WBRK method

Figure 2 shows that the stability regions of the method lies inside the closed region. We find the interval of absolute stability of the method by using MAPLE which is approximately  $[-1.94,0]$ . According to Burden et al. (2015), a method can be applied effectively to a stiff equation only if  $h\lambda$  is in the region of absolute stability of the method.

## 4. Numerical Results

In this section, we present the numerical results of WBRK method to check on the accuracy of the method. The method is tested with three ODEs of stiff type and we compare the maximum error (MAXE) and computational time (TIME) of the method with MWRK3CeM as shown in Sharmila and Amirtharaj (2011) and the third order RK method in Lambert (1973).

Table 1-3 tabulate the results while Figure 3-8 illustrate the graphical form between the  $\text{Log}(\text{MAXE})$  versus  $\text{Log}(h)$  and  $\text{Log}(\text{TIME})$ .

The methods are tested with stepsize,  $h=0.01, 0.0001, \text{ and } 0.000001$ .  $\lambda$  shows the eigenvalue(s) of the following test problems.



**Problem 1:**

$$y' = -15y, \quad y(0) = 1, \quad 0 \leq x \leq 1$$

with exact solution

$$y = e^{-15x}, \quad \lambda = -15$$

Source: Berland (2007).

**Problem 2:**

$$y' = -20(y - x) + 1, \quad y(0) = 1, \quad 0 \leq x \leq 10$$

with exact solution

$$y = e^{-20x} + x, \quad \lambda = -20$$

Source: Gear (1971).

**Problem 3:**

$$x' = 42y - 43x, \quad x(0) = 8, \quad 0 \leq x \leq 1$$

$$y' = -8y + 7x, \quad y(0) = 1,$$

with exact solution

$$x = 2e^{-t} + 6e^{-50t}, \quad \lambda = -1$$

$$y = 2e^{-t} - e^{-50t}, \quad \lambda = -50$$

Source: Huang (2005).

Table 1: Comparison of WBRK of weights,  $w_1 = w_2 = \frac{1}{2}$  with MWRK3CeM and RK3 for solving Problem 1

Stepsize, $h$	Method	MAXE( $y_{n+1}$ )
$10^{-2}$	WBRK	7.111898824E-05
	MWRK3CeM	1.536057468E-03
	RK3	2.184967940E-02
$10^{-4}$	WBRK	5.949740700E-11
	MWRK3CeM	1.381024156E-07
	RK3	1.842618470E-04
$10^{-6}$	WBRK	7.438494265E-15
	MWRK3CeM	1.380451309E-11
	RK3	1.839429394E-06

Table 2: Comparison of WBRK of weights,  $w_1 = w_2 = \frac{1}{2}$  with MWRK3CeM and RK3 for solving Problem 2

Stepsize, $h$	Method	MAXE( $y_{n+1}$ )
$10^{-2}$	WBRK	2.271357959E-04
	MWRK3CeM	2.803610572E-03
	RK3	2.840716293E-02
$10^{-4}$	WBRK	2.966545146E-08
	MWRK3CeM	2.430198235E-07
	RK3	2.250360654E-04
$10^{-6}$	WBRK	7.684324288E-10
	MWRK3CeM	7.684324288E-10
	RK3	2.244994309E-06

Table 3: Comparison of WBRK of weights,  $w_1 = w_2 = \frac{1}{2}$  with MWRK3CeM and RK3 for solving Problem 3

Stepsize, $h$	Method	MAXE( $y_{n+1}$ )
$10^{-2}$	WBRK	2.467754126E-02
	MWRK3CeM	1.331770888E-01
	RK3	2.966907540E-01
$10^{-4}$	WBRK	6.934759966E-08
	MWRK3CeM	9.247600947E-06
	RK3	1.859830420E-03
$10^{-6}$	WBRK	1.375306535E-10
	MWRK3CeM	9.214335961E-10
	RK3	1.849920277E-05

From Table 1-3, we can observed that the WBRK method obtained smaller MAXE when compared with the MWRK3CeM method and RK3 method. In comparison of the stepsize used, it shows that stepsize,  $h=0.000001$  obtained smaller MAXE.




Figure 3: Graph of Log(MAXE) vs Log(h) of WBRK method is compared with MWRK3CeM and RK3 for solving Problem 1




Figure 4: Graph of Log(MAXE) vs Log(TIME) of WBRK method is compared with MWRK3CeM and RK3 for solving Problem 1




Figure 5: Graph of Log(MAXE) vs Log( $h$ ) of WBRK method is compared with MWRK3CeM and RK3 for solving Problem 2




Figure 6: Graph of Log(MAXE) vs Log(TIME) of WBRK method is compared with MWRK3CeM and RK3 for solving Problem 2




Figure 7: Graph of Log(MAXE) vs Log( $h$ ) of WBRK method is compared with MWRK3CeM and RK3 for solving Problem 3




Figure 8: Graph of Log(MAXE) vs Log(TIME) of WBRK method is compared with MWRK3CeM and RK3 for solving Problem 3

Figure 3-8 illustrate the efficiency of WBRK method based on its execution time and stepsize used when compared with the MWRK3CeM and the RK3 method.

## 5. Conclusion

The derivation of WBRK based on CeM for solving stiff ODEs is shown in this paper. From the stability regions of the WBRK method, we can conclude that the method is absolute stable. The numerical results prove that WBRK method has better accuracy than the comparing methods. The results also show better accuracy when smaller stepsize is used. From the efficiency graph shown in Figure 3-8, we can observe that WBRK method performs better in terms of its execution time when compared with the MWRKCeM and the RK3 method. Hence, we can conclude that WBRK method can serve as an alternative solver for solving stiff ODEs problems.

## References

- Ababneh, O. Y. and Rozita, R. (2009). New third order runge-kutta based on contraharmonic mean for stiff problems. *Applied Mathematical Sciences*, 3(8):365–376.
- Berland, H. (2007). Explicit numerical methods experiencing instability on a stiff equation. An unpublished pre-print.
- Burden, R. L., Faires, J. D., and Burden, A. M. (2015). *Numerical Analysis (10th ed.)*. Cole-Thomson Learning, Brooks.
- Cash, J. R. (1983). Block runge-kutta method for the numerical integration of ivp. *Mathematics of Computation*, 40(161):175–191.
- Cash, J. R. (1985). Block embedded explicit runge-kutta methods. *Computers and Mathematics with Applications*, 11(4):395–409.
- Evans, D. and Yaakub, A. R. (1996). New fifth order weighted runge-kutta formula. *International Journal of Computational Mathematics*, 59(4):227–243.
- Evans, D. J. and Yaakub, A. R. (1998). Weighted fifth order runge-kutta formulas for second order differential equations. *International Journal of Computational Mathematics*, 70(2):233–239.

- Gear, C. W. (1971). *Numerical Initial Value Problems in Ordinary Differential Equations*. Prentice Hall Inc., New Jersey.
- Huang, S. J. Y. (2005). *Implementation of General Linear Methods for Stiff ODEs*. PhD thesis, University of Auckland, Auckland, New Zealand.
- Ibrahim, Z. B., Othman, K. I., and Suleiman, M. B. (2007). Variable step size block backward differentiation formula for solving stiff odes. *Proceedings of World Academy of Science, Engineering and Technology*, 2:785–789.
- Ismail, N., Ibrahim, Z. B., Othman, K. I., and Suleiman, M. (2014). 3-point block backward differentiation formulas for solving fuzzy differential equations. *Malaysian Journal of Mathematical Sciences*, 8:139–151.
- Lambert, J. D. (1973). *Computational Methods in Ordinary Differential Equations*. John Wiley and Sons, England.
- Lambert, J. D. (1993). *Numerical Methods for Ordinary Differential Equations: The Initial Value Problems*. John Wiley and Sons, New York.
- Majid, Z. A. and Suleiman, M. B. (2007). Implementation of four-point fully implicit block method for solving ordinary differential equations. *Applied Mathematics and Computation*, 184(2):514–522.
- Milne, W. E. (1953). *Numerical Solution of Differential Equations*. John Wiley and Sons, New York.
- Murugesan, K., Dhayabaran, D. P., Amirtharaj, E. C. H., and Evans, D. J. (2001). A fourth order embedded runge-kutta rkcm(4,4) method based on arithmetic and centroidal means with error control. *International Journal of Computational Mathematics*, 79(2):247–269.
- Othman, K. I., Ibrahim, Z. B. and Suleiman, M., and Majid, Z. (2007). Automatic intervalwise block partitioning using adams type method and backward differentiation formula for solving odes. *Applied Mathematics and Computation*, 188:1642–1646.
- Pushpam, A. E. K. and Dhayabaran, D. P. (2011). Comparison of single term walsh series technique and extended rk methods based on variety of means to solve stiff non-linear systems. *Recent Research in Science and Technology*, 3(9):22–30.
- Rahim, Y. F. (2004). *Block Diagonally Implicit Runge-Kutta Method for Solving Ordinary Differential Equation*. PhD thesis, Universiti Putra Malaysia, Serdang, Malaysia.

- Sharmila, R. G. and Amirtharaj, E. C. H. (2011). Implementation of a new third order weighted runge-kutta based on centroidal mean for solving stiff ivps. *Recent Research in Science and Technology*, 3(10):91–97.
- Suleiman, M. B. (1979). *Generalised Multistep Adams and Backward Differentiation Methods for the Solution of Stiff and Non-stiff Ordinary Differential Equations*. PhD thesis, University of Manchester, Manchester, England.
- Wu, X. Y. (1998). A sixth-order a-stable explicit one-step method for stiff systems. *Computers and Mathematics with Applications*, 35(9):59–64.
- Yatim, S. A. M., Ibrahim, Z., Othman, K. I., and Suleiman, M. B. (2011). Quantitative comparison of numerical method for solving stiff ordinary differential equations. *Mathematical Problems in Engineering*, 2011:1–12.
- Zawawi, I. S. M., Ibrahim, Z. B., Ismail, F., and Majid, Z. A. (2012). Diagonally implicit block backward differentiation formulas for solving ordinary differential equations. *International Journal of Mathematics and Mathematical Sciences*, 2012:1–8.