



UNIVERSITI PUTRA MALAYSIA

***EFFECT OF VARIOUS ACTIVE PACKAGING SYSTEM ON QUALITY AND
MICROBIAL STABILITY OF FRESH-CUT CANTALOUPE (*Cucumis melo L.*
var. *Reticulates cv. Glamour*)***

SYAHIDAH BINTI KAMARUDDIN

FK 2015 65



**EFFECT OF VARIOUS ACTIVE PACKAGING SYSTEM ON QUALITY AND
MICROBIAL STABILITY OF FRESH-CUT CANTALOUPE (*Cucumis melo* L.
var. *Reticulatus* cv. *Glamour*)**

By

SYAHIDAH BINTI KAMARUDDIN

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fulfilment of the Requirements for the Degree of Master of Science**

July 2015

COPYRIGHT

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the Master of Science

EFFECT OF VARIOUS ACTIVE PACKAGING SYSTEM ON QUALITY AND MICROBIAL STABILITY OF FRESH-CUT CANTALOUPE (*CUCUMIS MELO* L. VAR. *RETICULATES* CV. GLAMOUR)

By

SYAHIDAH BINTI KAMARUDDIN

July 2015

Chairman : Rosnah Bt Shamsudin, PhD
Faculty : Engineering

A study was conducted to determine the effect of initial packaging atmosphere by using different packaging system on the quality and microbial stability of fresh-cut Cantaloupe and to investigate the effect of different absorbers. Fresh-cut Cantaloupe was stored in two types of containers; specifically a rigid container and semirigid container. Each of containers was sealed under different conditions. These were Polypropylene (PP) film and Low-Density Polyethylene (LDPE) films. Between the two types of film, Low-Density Polyethylene (LDPE) film material was shown to have better quality of both containers, being the rigid and semirigid containers. It was found that the physico-chemical properties (firmness, colour, total soluble solid, titratable acidity) of fresh-cut Cantaloupe were maintained over 18 days of storage. Meanwhile, for pH value, there are no significant effects in the storage time for all type of packages. Microbial analysis (total plate count and yeast and mould) were found to increase over storage, but LDPE materials were found to have lower TPC compared to the other package. The packages did not exhibit any effects on the YM counts during the 18 days of storage. For the oxygen, carbon dioxide, respiration rate and ethylene production, the LDPE film materials could maintain the quality of about 11 days with acceptable level of oxygen (above 2%) and a moderate level of carbon dioxide (2%-12%). LDPE shows the lowest value of respiration rate and ethylene production in the rigid and semirigid container. The effects of absorbers were investigated after the selection of the seal package materials for rigid and semirigid containers. Two types of absorbers were used namely, oxygen absorber and ethylene absorber. There were four different conditions created in this experiment. The first container does not contain absorber. The second container contained an oxygen absorber. The third container contained ethylene absorber, and finally, the samples included both types of absorbers, being oxygen and ethylene absorbers. It was found that the physico-chemical of fresh-cut Cantaloupe (firmness, colour, total soluble solid, titratable acidity) was maintained over 18 days of storage for all types of conditions in an active packaging system. The microbial analysis (total plate count (TPC) and yeast and mould (YM)) were increased throughout the storage time and the storage can last until day 11 for microbial spoilage. It was found that all types of absorbers able to reduce the oxygen concentration of fresh-cut Cantaloupe but too little of oxygen may cause

anaerobic metabolism and production of off flavors and odors. The results showed that the absorbers had reduced the oxygen level of oxygen tremendously (below than 2%). The percentage of carbon dioxide had shown the same amount for both containers (2%-12% of CO₂). For respiration rate and ethylene production, the ethylene-oxygen absorbers had shown the lowest value for fresh-cut Cantaloupe for both containers. Therefore, the fresh-cut Cantaloupe packaged with LDPE film was found to maintain the quality and microbial stability up to 11 days of postharvest storage for rigid and semirigid container at 2±1°C. The fresh-cut Cantaloupe packaged with LDPE film can be consumed until day 11. However, the package associated with oxygen and ethylene absorbers can be only being consumed up to day 7.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia
Sebagai Memenuhi Keperluan untuk Ijazah Sarjana Sains

**KESAN KEPELBAGAIAN SISTEM PEMBUNGKUSAN AKTIF KE ATAS KUALITI DAN
KESTABILAN MIKROB BAGI TEMBIKAI SEGAR POTONG (*CUCUMIS MELO L. VAR.
RETICULATES CV. GLAMOUR*)**

Oleh

SYAHIDAH BINTI KAMARUDDIN

Julai 2015

Pengerusi :Rosnah Bt Shamsudin, PhD
Fakulti :Kejuruteraan

Satu kajian telah dijalankan untuk menentukan kesan atmosfera awal pembungkusan dengan menggunakan sistem pembungkusan yang berbeza terhadap kualiti dan kestabilan mikrob daripada buah tembikai segar potong dan untuk mengkaji kesan penyerap yang berbeza. Tembikai segar potong disimpan di dalam dua jenis bekas khususnya bekas tegar dan separa tegar. Dalam eksperimen yang pertama, setiap bekas telah dimeterai pada keadaan yang berbeza. Ini adalah filem polietilena(PP) dan filem berkepadatan rendah polietilena (LDPE). Antara kedua-dua jenis filem ini, bahan filem daripada berkepadatan rendah polietilena (LDPE) telah menunjukkan kualiti yang lebih baik bagi kedua-dua bekas yang tegar dan separa tegar. Ia telah mendapati bahawa sifat-sifat kimia fisiko (ketegasan, warna, jumlah kandungan pepejal larut,keasidan) tembikai segar potong dapat kekal selama 18 hari tempoh penyimpanan. Sementara itu, nilai pH, didapati tiada kesan yang ketara dalam masa penyimpanan bagi semua jenis pakej. Kestabilan mikrob iaitu kiraan jumlah plat (TPC) dan yis dan kulat(YM) telah meningkat sewaktu tempoh penyimpanan. Walau bagaimanapun, filem daripada berkepadatan rendah polietilena (LDPE) mempunyai TPC yang lebih rendah berbanding dengan pembungkusan yang lain. Semua jenis pembungkusan tidak memberikan apa-apa kesan untuk kiraan YM sepanjang tempoh 18 hari penyimpanan. Bagi oksigen, karbon dioksida, kadar respirasi dan pengeluaran etilena, filem daripada bahan berkepadatan rendah polietilena (LDPE) dapat mengekalkan kualiti kira-kira 11 hari dengan tahap oksigen (lebih 2%) dan tahap sederhana karbon dioksida (2%-12%). Filem daripada bahan berkepadatan rendah polietilena (LDPE) menunjukkan nilai terendah bagi kadar respirasi dan pengeluaran etilena bagi bekas yang tegar dan separa tegar. Kesan penyerap yang berbeza dikaji selepas pemilihan bahan pembungkusan filem bagi bekas yang tegar dan separa tegar. Dua jenis penyerap yang digunakan iaitu serapan oksigen dan serapan etilena. Terdapat empat keadaan yang berbeza di dalam eksperimen ini. Bekas pertama tidak mengandungi penyerap. Bekas kedua mengandungi penyerap oksigen. Bekas ketiga mengandungi penyerap etilena dan yang terakhir, sampel kajian

mengandung kedua-dua jenis penyerap iaitu penyerap oksigen dan penyerap etilena. Ia menunjukkan bahawa sifat-sifat kimia-fisiko (ketegasan, warna, jumlah kandungan pepejal larut, keasidan) dapat dikekalkan selama 18 hari tempoh penyimpanan bagi semua jenis keadaan dalam sistem pembungkusan yang aktif. Kestabilan mikrob (kiraan jumlah plat (TPC) dan yis dan kulat (YM)) telah meningkat sepanjang tempoh penyimpanan dan simpanan boleh dilanjutkan sehingga hari yang ke 11 bagi kerosakan mikrob. Semua jenis penyerap boleh mengurangkan kepekatan oksigen tembikai segar potong tetapi jika terlalu sedikit tahap oksigen, ia boleh menyebabkan metabolisme anaerobik dan kerosakan pada rasa dan bau. Hasil kajian menunjukkan bahawa serapan amat mengurangkan tahap oksigen (di bawah paras 1%). Peratusan karbon dioksida menunjukkan jumlah yang sama untuk kedua-dua bekas iaitu (2% - 12%). Untuk kadar respirasi dan pengeluaran etilena, serapan oksigen-etilena menunjukkan nilai yang terendah bagi tembikai segar potong untuk kedua-dua bekas. Oleh itu, tembikai segar potong yang dibungkus dengan LDPE didapati dapat mengekalkan kualiti dan kestabilan mikrob sehingga hari ke 11 penyimpanan lepas tuai untuk bekas tegar dan separa tegar pada $2\pm 1^{\circ}\text{C}$. Tembikai segar potong yang dibungkus dengan filem LDPE hanya boleh dimakan sehingga 11 hari. Manakala, bungkusan yang mengandungi penyerap oksigen dan etilena hanya boleh dimakan sehingga 7 hari sahaja.




ACKNOWLEDGEMENTS

In the name of Allah, the Benevolent, the Merciful.

I would like to express my most sincere gratitude and deepest appreciation to my main supervisor; Prof Madya Dr. Rosnah Bt Shamsudin (Faculty of Engineering) as well as my co-supervisor; Prof Madya Dr. Noranizan Bt Mohd Adzahan (Faculty of Food Technology), Dr. Zaulia Bt Othman (Horticulture Research Centre, MARDI) and Prof Madya Dr Anvarjohn Ahmedov (Faculty of Engineering) for their opinion, guidance, ideas, time, motivation, encouragement, and constructive comments for the accomplishment of this research.

My special thanks to the staff and technician at Department of Food and Process Engineering and Department of Food Technology for their help and guidance on how to use the instruments that is related to my research works. My sincere appreciation is also extended to my lab mates from UPM for their help, support and opinion.

Lastly, thanks to my beloved parents; Kamaruddin Bin Hashim and Jamaliah Binti Jamaludin, and also to my siblings; Nurul Shariah Kamaruddin, Shahrir Kamaruddin and Shafiz Kamaruddin for their support and motivation.



I certify that a Thesis Examination Committee has met on (August 2014) to conduct the final examination of (Syahidah Bt Kamaruddin) on her thesis entitled "Effect of Various Active Packaging System on Quality and Microbial Stability of Fresh-Cut Cantaloupe (*Cucumis Melo* L. Var. *Reticulatus* Cv. *Glamour*)" in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1998. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Yus Aniza Yusof, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Farah Saleena Taip, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Rosnita A. Talib, PhD

Senior Lecturer
Faculty of Engineering
Universiti Putra Malaysia
(Internal Examiner)

Aminah Abdullah, PhD

Professor
Universiti Kebangsaan Malaysia
Malaysia
(External Examiner)



ZULKARNAIN ZAINAL, PhD

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 22 September 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfillment of the requirement for the degree of Master of Science. The members of the Supervisory Committee were as follows:

Rosnah Shamsudin, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Chairman)

Noranizan Mohd Adzahan, PhD

Associate Professor
Faculty of Food Science and Technology
Universiti Putra Malaysia
(Member)

Zaulia Othman, PhD

Horticulture Research Centre
Malaysian Agricultural Research and Development Institute (MARDI)
(Member)

Anvarjon Ahmedov, PhD

Associate Professor
Faculty of Engineering
Universiti Putra Malaysia
(Member)

BUJANG KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____ Date: _____

Name and Matric No.: Syahidah Bt Kamaruddin (GS30067)



Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature: _____	Signature: _____
Name of Chairman of Supervisory Committee: <u>Rosnah Shamsudin, PhD</u>	Name of Member of Supervisory Committee: <u>Noranzan Mohd Adzahan, PhD</u>

Signature: _____	Signature: _____
Name of Member of Supervisory Committee: <u>Zaulia Bt Othman, PhD</u>	Name of Member of Supervisory Committee: <u>Anvarjon Ahmedov, PhD</u>



TABLE OF CONTENTS

	Page
ABSTRACT	i
ABSTRAK	iii
ACKNOWLEDGEMENTS	v
APPROVAL	vi
DECLARATION	viii
LIST OF TABLES	xii
LIST OF FIGURES	xiv
LIST OF ABBREVIATIONS	xvii
CHAPTER	
1	
INTRODUCTION	1
1.1 Problem Statements	3
1.2 Scope and Limitations of the study	4
1.3 Outline of the thesis	4
1.4 Objectives	4
2	
LITERATURE REVIEW	5
2.1 Overview of Cantaloupe fruits	5
2.2 Postharvest storage of Cantaloupe	6
2.3 The processing of fresh-cut Cantaloupe	6
2.4 Calcium treatments for fresh-cut fruits	9
2.5 Rigid and semirigid containers	11
2.6 Effects of packaging materials on quality	11
2.7 Effects of Modified Atmosphere Packaging (MAP) on quality	13
2.8 Effects of Active Packaging on quality	15
2.8.1 Oxygen absorbers	16
2.8.2 Ethylene absorbers	17
2.9. Physico-Chemical properties	18
2.9.1 Texture	18
2.9.2 Colour	19
2.9.3 pH, Total Soluble Solid (TSS) and Total Acidity (TA)	20
2.9.4 Microbial stability	22
2.9.5 O ₂ and CO ₂ levels	23
2.9.6 Respiration rate and Ethylene Production	23
3	
METHODOLOGY	26
3.1 Overview	26
3.2 Plant Material	28
3.3 Preparation of fresh-cut samples	28
3.4 Preparation of Calcium Lactate treatment	30
3.5 Preparation of packaging	30
3.6 Preparation of absorbers	33

3.7 Parameters	33
3.7.1 Texture	33
3.7.2 Color	33
3.7.3 pH, Total Soluble Solid (TSS), Total Acidity (TA)	34
3.7.4 Microbial stability	34
3.7.5 Gas measurement	35
3.7.6 Respiration rate and Ethylene production	35
3.8 Statistical Analysis	36
4	
RESULTS AND DISCUSSIONS	37
4.1 Effects of different types of packaging system on quality and microbial stability of fresh-cut Cantaloupe.	37
4.1.1 Texture	37
4.1.2 Colour (Lightness, Hue Angle, and Chromaticity)	38
4.1.3 pH, Total Soluble Solid (TSS) and Titratable Acidity (TA)	43
4.1.4 Microbial stability	47
4.1.5 Gases Measurement (Percentage of Oxygen and Carbon Dioxide)	50
4.1.5.1 Oxygen	50
4.1.5.2 Carbon Dioxide	52
4.1.6 Respiration Rate and Ethylene Production	54
4.1.6.1 Respiration Rate	54
4.1.6.2 Ethylene Production	55
4.2 Effect of different absorbers on quality and microbial stability of fresh-cut Cantaloupe	57
4.2.1 Texture	57
4.2.2 Colour (Lightness, Hue Angle and Chromaticity)	60
4.2.3 pH, Total Soluble Solid (TSS) and Titratable Acidity (TA)	64
4.2.4 Microbial stability	68
4.2.5 Gases Measurement (Percentage of Oxygen and Carbon Dioxide)	71
4.3.5.1 Oxygen	71
4.3.5.2 Carbon Dioxide	73
4.2.6 Respiration Rate and Ethylene Production	74
4.2.6.1 Respiration Rate	74
4.2.6.2 Ethylene Production	76
5	
CONCLUSIONS AND RECOMMENDATIONS	79
5.1 Conclusions	79
5.2 Recommendations for future work	80
REFERENCES	81
APPENDICES	101
BIODATA OF STUDENT	103
LIST OF PUBLICATIONS	104
LIST OF CONFERENCES	105

LIST OF TABLES

Table		Page
2.1	Application of calcium treatments for improving the quality and extending the shelf-life of fresh-cut fruits	10
2.2	Selective types of fresh-cut fruits using passive and active modified atmosphere packaging	14
2.3	Examples of active packaging technique which actively and constantly change the concentration of gases in the package headspace (sachet to be added in the package)	15
2.4	Respiration rate ranges of cantaloupe at various temperatures	24
3.1	The thickness of Polypropylene (PP) film and Low-Density Polyethylene (LDPE) film	31
4.1	Means for Firmness of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the rigid and semirigid containers during the 18 days of storage	38
4.2	Means for Colour (Lightness, Hue Angle, and Chromaticity) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in rigid containers during the 18 days of storage	40
4.3	Means for Colour (Lightness, Hue Angle, and Chromaticity) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in semirigid containers during the 18 days of storage	42
4.4	Means for pH, Total Soluble Solid (TSS), Titratable Acidity (TA) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in rigid containers during the 18 days of storage	45
4.5	Means for pH, Total Soluble Solid (TSS), Titratable Acidity (TA) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in semirigid containers during the 18 days of storage	46
4.6	Means for Firmness of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the rigid and semirigid containers during the 18 days of storage	59
4.7	Means for Colour (Lightness, Hue Angle, and Chromaticity) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in rigid containers during the 18 days of storage	61
4.8	Means for Colour (Lightness, Hue Angle, and Chromaticity) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in semirigid containers	63

during the 18 days of storage

- | | | |
|------|---|----|
| 4.9 | Means for pH, Total Soluble Solid (TSS), Titratable Acidity (TA) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in rigid containers during the 18 days of storage | 65 |
| 4.10 | Means for pH, Total Soluble Solid (TSS), Titratable Acidity (TA) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the samples in semirigid containers during the 18 days of storage | 67 |
| 4.11 | Means for percentage of Oxygen (O ₂) of fresh-cut Cantaloupe (2 ± 1 °C and 87 % RH) for the rigid and semirigid containers during the 18 days of storage | 72 |



LIST OF FIGURES

Figure		Page
2.1	Cantaloupe (<i>Cucumis Melo L. Var.Reticulates CV.Glamour</i>)	5
2.2	Typical fresh-cut process flow chart for fruits, vegetables and root crops	7
2.3	Ageless® formats available: sachet, pressure-sensitive label and card	16
2.4	Sachet of ethylene scavenger from Ethylene Control, Inc.	17
3.1	Flow chart of the overall methodology throughout the experiment	27
3.2	Procedure to prepare fresh-cut Cantaloupe	29
3.3	Procedure of calcium lactate treatment	30
3.4	Rigid and semirigid container	31
3.5	Diagram of packaging materials for fresh-cut Cantaloupe	32
3.6	Different types of absorbers (water, oxygen and ethylene)	33
3.7	Serial dilution of fresh-cut Cantaloupe samples	35
4.1	Total Plate Count (TPC) of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2±1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life	48
4.2	Total Plate Count (TPC) of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2±1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life	48
4.3	Yeast and Mould (YM) of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2±1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life	49
4.4	Yeast and Mould (YM) of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2±1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life	50

4.5	Percentage of oxygen of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days. The dashed line indicates the anaerobic condition developed below 2%	51
4.6	Percentage of oxygen of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days. The dashed line indicates the anaerobic condition developed below 2%	52
4.7	Percentage of carbon dioxide of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days	53
4.8	Percentage of carbon dioxide of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days	53
4.9	Respiration Rate ($\text{mlCO}_2/\text{kg h}$) of fresh-cut Cantaloupe in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days	54
4.10	Respiration Rate ($\text{mlCO}_2/\text{kg h}$) of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days	55
4.11	Ethylene production of fresh-cut Cantaloupe in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days	56
4.12	Ethylene production of fresh-cut Cantaloupe in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days	57
4.13	Total Plate Count (TPC) of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life	68
4.14	Total Plate Count (TPC) of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life	69
4.15	Yeast and Mould (YM) of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life	70
4.16	Yeast and Mould (YM) of fresh-cut Cantaloupe packaged in	70

Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days. The dashed line indicates the limit of microbial shelf life

4.17	Percentage of carbon dioxide of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days	73
4.18	Percentage of carbon dioxide of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days	74
4.19	Respiration Rate of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days	75
4.20	Respiration Rate of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days	76
4.21	Ethylene production of fresh-cut Cantaloupe packaged in Rigid (R) containers stored at 2 ± 1 °C and 87 % RH for 18 days	77
4.22	Ethylene production of fresh-cut Cantaloupe packaged in Semirigid (SR) containers stored at 2 ± 1 °C and 87 % RH for 18 days	78



LIST OF ABBREVIATIONS

a*	Greenness
ANOVA	One way analysis of variance
AP	Active Packaging
b*	Yellowness
C*	Chromaticity
C ₂ H ₄	Ethylene
CaCl ₂	Calcium chloride
Ca-Lactate	Calcium Lactate
CFU	Colony-Forming Unit
CO ₂	Carbon Dioxide
COO	Sample volume in ml
CRD	Completely Randomized Design.
cv	Cultivar
DMRT	Duncan's Multiple Range Test
DRBC	Dichloran Rose Bengal Chloramphenicol agar
EP1	Volume in ml NaOH up to endpoint (pH=8.5)
FDA	Food and Drug Administration
FID	Flame Ionization Detector
GC	Gas Chromatography
h°	Hue angle
H ₂ O ₂	Hydrogen Peroxide
HDPE	High-Density polyethylene
KMnO ₄	Potassium permanganate

L*	Lightness
LDPE	Low-Density Polyethylene
LLDPE	Linear Low-Density Polyethylene
MA	Modified Atmosphere
MAP	Modified Atmosphere Packaging
MGC	Mitsubishi Gas Chemical
NaOCl	Sodium Hypochlorite
NaOH	Sodium Hydroxide
O ₂	Oxygen
OTR	Oxygen Transmission Rate
PCA	Plate Count Agar
PE	Polyethylene
PET	Polyethylene Terephthalate
PP	Polypropylene
PPM	Part Per Million
PPO	Polyphenol Oxidases
PS	Polystyrene
PVC	Polyvinyl Chloride
R	Rigid
RH	Relative humidity
SR	Semirigid
SAS	Statistical Analysis Software
SSC	Soluble Solid Content
TA	Titrateable Acidity

TA	Texture Analyzer
TPC	Total Plate Count
TSS	Total Soluble Solid
USDA	United State Department of Agriculture
YM	Yeast and Mould



© COPYRIGHT UPM



CHAPTER 1

INTRODUCTION

Consumers of late have developed particular demands on the freshness, health, convenience and safety of their food (James and Ngarmark, 2010). Fresh-cut products, also recognised as minimally processed products, are highly delicate or perishable foods (Watada et al., 1996). These foods have been peeled, trimmed and cut into a ready-to-eat form (IFPA, 2000) and are in high demand for retail and food services because of their convenience, quality, safety and reduced waste (Watada et al., 1996; Luna-Guzmán, 1997; Robert et al., 2003; Olivas and Barbosa-Canovas, 2005). Changes in lifestyles have increased the demand for ready-to-eat food such as fresh processed fruits and vegetables (Benedetti et al., 2008). Within this context, the fruits are presented in a condition that is suitable for direct and instant consumption without the need to process it for consumption (Olivas and Barbosa-Canovas, 2005). When selecting the fruits, consumers gauge its appearance and avoid fruits that are bruised or damaged. Food packaging is one of the chief factors that affect the quality of fresh-cut fruit products. Packaging considerations are vital to food products as it will protect the food from the external environments, provide information about the food contained within the package and afford convenience to consumer food handling and food storage.

There have been many innovations in food packaging due to current consumer demands in advanced technology in terms of food packaging. Usually, for fresh-cut fruits, the treatments that have been used to extend the shelf life of food include using shock, ozone, chemical treatments sterilisers, ultraviolet radiation and heat. However these methods are not suitable for food packaging even though these technologies are popularly used in the United States and Europe. These methods can cause low consumer acceptability, reduce the freshness, and lessen the nutritional quality (Erturk and Picha, 2008; González -Aguilar et al., 2008). Another food packaging method that is commonly used to preserve the food is known as Modified Atmosphere Packaging (MAP). MAP of fresh-cut fruits and vegetables refer to the technique of sealing actively respiring produce in plastic film packages to modify the O₂ and CO₂ levels within the package atmosphere (Mir and Beaudry, 2004). MAP is effective in maintaining quality because of its effects on the modification of gas composition in the package headspace (Schlimme and Rooney, 1994; Jacxsens et al., 2002; Kim et al., 2003). The levels of O₂ and CO₂ within a package depend on the interaction between commodity respiration and the permeability properties of the packaging film or microperforations (Beaudry et al., 1992; Kader et al., 1997). There are two approaches for creating film barriers. The first approach employs continuous films that control the movement of O₂ and CO₂ into or out of the package. The second approach uses perforated films with small holes or microperforations as the primary of gas exchange. Both of the films show different permeability to O₂ and CO₂.

Cantaloupe is categorized as a climacteric fruit where it exhibits higher respiration and ethylene production rates, a quicker rate of softening and shorter shelf-life than the non-climacteric fruits. When O₂ levels are lowered, the respiration of the produce begins to decrease and generally continues to do so with lowering O₂ levels down to a level where anaerobic respiration takes place (Barrett et al., 2004). Microperforated

films are not suitable for fresh-cut Cantaloupe because perforated packages are for produce that has a high O₂ demand (Mir and Beaudry, 2004). Perforated packaging is more suitable for vegetables that have a high demand for O₂ (Gonzalez et al., 2008). Packaging films provide a wide range of different materials such as polypropylene (PP), polyethylene terephthalate (PET), polyethylene (PE) and polyvinyl chloride (PVC). Polypropylene is used largely in MAP, in two forms: continuous and perforated film (Mattoss et al., 2012). The selection of the plastic film packaging material strives to achieve equilibrium between the oxygen demand of the product (oxygen consumption by respiration) (Cantwell and Suslow, 1999). An important aspect in the application of modified atmosphere packaging is the selection of packaging materials (Barrett et al., 2004).

In general, there are three types of formats that are most commonly used in fresh-cut produce modified atmosphere packaging, namely rigid packaging, semirigid packaging and active packaging. Rigid packaging offers high impact strength, high stiffness, and high barrier properties, often exceeding industry standards for quality and durability (Anon, 2005). In rigid packaging, there are a number of designs commonly used in fresh-cut produce applications. The most common rigid container and tray includes clamshell, snap-on lid, sealable and easy-peel lidding film. These methods are utilized to close or seal the tray or container. Frequently these styles of rigid packages are referred to as natural aspiration packages, meaning that the final atmosphere in the package can vary according to how tight the lid is attached or "snapped" onto the container (Toivonen et al., 2009). For semi-rigid packaging, the materials are those which are neither rigid nor flexible in nature. There are four main groups of semi-rigid packages, namely paperboard packages, flexible pouches, plastic cups and trays (George, 2001).

Active packaging refers to the incorporation of certain additives into packaging film or within packaging containers (rigid and semirigid) with the aim of maintaining product shelf life (Day, 1989). Common examples of an active packaging include absorbent packaging (Toivonen et al., 2009). Absorbent packaging describes ability of packages to absorb liquids or gases produced by fresh-cut produce (Ozdemir and Floros, 2004). Active packaging with oxygen absorbers dynamically decreases the oxygen concentration in the package headspace, even down to 0.01%, thus slow down the oxidation processes, colour changes, yeast and mould growth, that are occurring in foods (Smith et al., 1990; Andersen and Rasmussen 1992). Accumulation of gases such as ethylene, the ripening hormone, can significantly reduce shelf life. These trends have initiated a huge amount of research in the field of active and intelligent packaging to provide the market with packaging technologies designed to keep produce fresh, since it is impossible to attain the optimum quality characteristics with passive plastic package for such highly deteriorative and metabolically active produce (Ozdemir and Floros, 2004). The active packaging technologies that have been developed for fruits and vegetables are oxygen absorbers, carbon dioxide emitters, moisture absorbers, ethylene absorbers, antimicrobial agent releasers, and others (Han and Floros, 2007). Oxygen absorber is an active element used in the packaging system to absorb residual oxygen that remains after the package is sealed that originated from the product respiration and the package permeability (Mehyar and Han, 2010). Controlling the oxygen concentration provides benefits in protecting the produce against the quality deterioration associated with oxygen, such as off-flavor formation, color change, nutritional value reduction, and safety losses (Sanjeev and Ramesh, 2006). Meanwhile, ethylene causes the increase in fruit respiration rate and textural and color changes in

climacteric fruits (Toivonen and Brummell, 2008). Ethylene absorber acts to remove the exogenous ethylene from the atmosphere surrounding the produce by oxidizing it to ethylene glycol, which later decomposes to carbon dioxide and water (Martinez-Romero and Bail'en, 2007). Carbon dioxide reduces the fruit respiration rate and blocks the synthesis of endogenous ethylene (Mehyar and Han, 2010). Several studies have been published on the use of oxygen absorbers, ethylene absorbers and their beneficial effects with fruit and vegetables. Bolin and Huxsoll (1989) showed that oxygen scavengers improve firmness of peaches. Charles et al. (2005) observed that oxygen absorbers did not modify the gas equilibrium composition and improved the quality of fresh endives, and Tarr and Clingeffer (2005) have reported that oxygen absorbers minimize color changes of dried vine fruit.

1.1 Problem Statements

Consumers generally purchase fresh-cut produce for convenience, freshness, nutrition, safety and the eating experience. Many soft fresh-cut fruit items need to be packaged in rigid and semirigid containers, possibly lidded with a plastic film that permits development of a modified atmosphere (Cantwell and Suslow, 1999). Initial atmospheric modification within a modified atmosphere package is a consequence of the respiratory O₂ uptake and CO₂ development of the packaged produce and the rate of gas transfer across the package (Al-Ati and Hotchkiss, 2002). Excessive levels of O₂ in a package may allow for cut surface discoloration to occur, while too little O₂ may cause anaerobic metabolism and production of off flavors and odors (Beaulieu and Gorny, 2001). Anaerobic conditions enable lactic acid bacteria to cause spoilage, such as souring of the product, gas production and slime formation (James and Ngarmsak, 2010). The purpose of an enclosed system is to provide a barrier to microorganisms and to prevent oxygen from degrading the food. Packaging film with the right permeability can create desirable modified atmosphere of fresh fruit and vegetables (Zhang and Chen, 1996). Senesi et al., (2000) found that after 7 days at refrigeration temperature, the environment within the package had become anaerobic and high in CO₂, stressing the importance of careful selection of a MAP film and initial gas atmosphere. The type of packaging, for example, flexible, rigid and semirigid plays a very important role for the fresh-cut Cantaloupe to retain the quality of the products. Flexible packages were commonly used in previous MAP studies. However, only a few studies used rigid and semirigid containers, and among those were trays overwrapped with films. Little information on quality changes of fresh-cut fruits in rigid and semirigid structures with different sealed packaging films are available in literature.

Incorporating oxygen and ethylene-absorbing materials in the container of the packaging material could prove to be more efficient (Huff, 2009). Packaging materials for fresh-cut fruits should have higher permeability to gases and ethylene, or contain gas absorbers to cope with high respiration and ethylene production (Mehyar and Han, 2010). However, there is no report on the effects of active packaging using ethylene and oxygen absorber combined with modified atmosphere on the quality and shelf life of fresh-cut Cantaloupe.

1.2 Scope and Limitations of the study

This study was conducted to determine the effects of different packaging systems on the quality and microbial stability of fresh-cut Cantaloupe. Fresh-cut cantaloupe was stored in two types of containers; specifically a rigid container and semirigid container. The container holds 350ml and polypropylene (PP) was used for both containers. Each container utilized different sealants, being polypropylene (PP) film and low-density polyethylene (LDPE) film. Between the two films, only the PP film or LDPE film was used to study the effects of different absorbers for the fresh-cut Cantaloupe. Two types of absorbers were used, namely oxygen absorber and ethylene absorber. There were four different conditions created in this experiment for comparison. In this study, Statistical Analysis System (SAS) 9.3 system was used for statistical analysis.

1.3 Outline of the thesis

This study is focused on the quality and microbial stability of fresh-cut Cantaloupe through the utilization of various active packaging systems used in this study. Chapter 2 introduces the attributes of Cantaloupe, its benefits in medication, the postharvest storage of Cantaloupe fruits, the processing of fresh-cut Cantaloupe, calcium treatments employed for fresh-cut fruits, the usage of rigid and semirigid containers, effects of the packaging materials, the usage and effects of modified atmosphere packaging (MAP) and active packaging on its quality. Chapter 2 also provides a literature review of published works from related research. The purpose of this chapter is to convey the information and ideas that have been established on the topic, solutions and methods as well as discussions. The research methodology was described in Chapter 3. The descriptions include materials, methods, procedures, equipments, and statistical analysis used for the entire experiment conducted. Chapter 4 discusses the results of analysis and discussions on the experiment conducted. The physico-chemical, microbial stability, gas measurement, respiration rate and ethylene production were discussed and evaluated. Lastly, chapter 5 summarises the main findings obtained in this research. The recommendations included are also suggested for future research.

1.4 Objectives

This research has two main objectives as listed below;

- 1) To determine the effects of initial packaging atmosphere by using different packaging systems on the quality and microbial stability of fresh-cut Cantaloupe.
- 2) To investigate the effects of different absorbers.

REFERENCES

- Abe, K. and Watada, A. E. (1991). Ethylene absorbent to maintain quality of lightly processed fruits and vegetables. *Journal of Food Science*, 58(6): 1589-1592.
- Abe, K., Tanase, M. and Chachin, K. (1998). Studies on physiological and chemical changes of fresh-cut bananas. I. Deterioration in fresh-cut green tip bananas. *Journal of the Japanese Society for Horticultural Science*, 67: 123-129.
- Abeles, F.B., Morgan, P.W. and Saltveit, M.E. (1992). *Ethylene in Plant Biology*, 2nd edition. New York: Academic Press.
- Abdul Raqeeb Ali Ahmed AL Eryani. Postharvest Quality Of Papaya Fruit (Carica Papaya) Associated With Applications Of Calcium And Chitosan. PhD Thesis, Universiti Putra Malaysia.
- Aday, M. S. and Caner, C. (2011). The applications of 'active packaging and chlorine dioxide' for extended shelf life of fresh strawberries. *Packaging Technology and Science*, 24: 123-136.
- Aday, M. S., Caner, C. and Rahvalı, F. (2011). Effect of oxygen and carbon dioxide absorbers on strawberry quality. *Postharvest Biology and Technology*, 62: 179-187.
- Aday, M. S. and Caner, C. (2012). The shelf life extension of fresh strawberries using an oxygen absorber in the biobased package. *Food Science and Technology*, 52(2): 102-109.
- Agar, I. T., Massantini, R., Hess-Pierce, B. and Kader, A. A. (1999). Postharvest CO₂ and ethylene production and quality maintenance of fresh-cut kiwifruit slices. *Journal of Food Science*, 64(3): 433-440.
- Aguayo, E., Escalona, V. and Artés, F. (Eds.). (2001). Proceedings: *The 8th International CA Conference on Minimally Processed 'Amarillo' Melon*. Ed. Oosterhaven, J and Peppelembos, H. Acta Horticulturae.
- Aguayo, A., Allende, A. and Artés, F. (2003). Keeping quality and safety of minimally fresh processed melon. *European Food Research and Technology*, 216(6): 494-499.
- Aguayo, E., Escalona, V. and Artés, F. (2004). Metabolic behaviour and quality changes of whole and fresh processed melon. *Journal of Food Science*, 69(4): 48-155.
- Ahvenainen, R. (1996). New approaches in improving the shelf life of minimally processed fruit and vegetables. *Trends in Food Science and Technology*, 7(6): 179-187.
- Ahvenainen, R. and Hurme, E. (1997). Active and smart packaging for meeting consumer demands for quality and safety. *Food Additives and Contaminants*, 14(6-7): 753-763.

- Ahvenainen, R. (2003). Active and intelligent packaging: an introduction. In: A. Raija (Ed.), *Novel Food Packaging Techniques* (pp. 5-21). Cambridge, UK: Woodhead Publishing Ltd
- Akelah, A. (2013). *Functionalized Polymeric Materials in Agriculture and the Food Industry*. Egypt: Springer Science
- Alam, M. S., Kaur, B., Gupta, K. and Kumar, S. (2013). Studies on refrigerated storage of minimally processed papaya (*Carica papaya L.*). *Agricultural Engineering International*, 15(4).
- Alandes, L., Pérez-Munuera, I. E., Llorca, A. and Quiles, I. H. (2009). Use of calcium lactate to improve structure of “Flor de Invierno” fresh-cut pears. *Postharvest Biology and Technology*, 53: 145-151.
- Al-Ati, T. and Hotchkiss, J.H. (2002). Application of packaging and modified atmosphere to fresh-cut fruits and vegetables, In: O. Lamikansa (Ed.), *Fresh-Cut Fruits and Vegetables: Science, Technology, and Market* (chap. 10) (pp. 305-338). Boca Raton, FL: CRC Press
- Al-Ati, T. and Hotchkiss, J. H. (2003). The role of packaging film permselectivity in modified atmosphere packaging. *Journal of Agricultural and Food Chemistry*, 51: 4133-4138.
- Ali, A., Muhammad, M., Sijam, K. and Siddiqui, Y. (2011). Effect of Chitosan coatings on the physicochemical characteristics of Eksotika II (*Carica papaya L.*) fruit during cold storage. *Food Chemistry*, 124: 620-626.
- Allahvaisi, S. (2012). Polypropylene in the Industry of Food Packaging, Polypropylene. In F.Dogan (Ed.), 2012, <http://www.intechopen.com/books/polypropylene/polypropylene-in-the-industry-of-food-packaging> (accessed 15 February 2013).
- Almenar, E., Del-Valle, V., Hernández-Munoz, P., Lagarón, J. M., Catalá, R. and Gavara, R. (2007). Equilibrium modified atmosphere packaging of wild strawberries. *Journal of the Science of Food and Agriculture*, 87: 1931-1939.
- Amaro, A. L., Beaulieu, J. C., Almeida, D. P. F. and Malcata, F.X. (Eds.). (2009). Proceedings: *International Conference on Environmentally Friendly and Safe Technologies for Quality of Fruit and Vegetables*, Universidade do Algarve, Faro: Portugal
- American Plastics Council, Understanding Plastic Film: Its Uses, Benefits and Waste Management Options by Headley Pratt Consulting, 1996, <http://plastics.americanchemistry.com/Understanding-Plastic-Film> (accessed 4 May 2013).
- Andersen, H. J. and Rasmussen, M. A. (1992). Interactive packaging as protection against photodegradation of the colour of pasteurized, sliced ham. *International Journal of Food Science and Technology*, 27:1-8.

- Andress, E. L., Cantaloupe Facts, 2003, <http://www.ext.colostate.edu/> (accessed 23 March 2013).
- Anonymous, Recycling Plastics, 1997,
<http://www.polymers.com/dotcom/home.html/>(accessed 14 May 2014).
- Anonymous, Introduction to Post-Harvest Technologies of Fruit and Vegetables, 1999,
http://www.unido.org/fileadmin/import/32102_14IntroPostharvestTech.6.pdf(a
ccessed 29 March 2014).
- Anonymous, Rigid Packaging, 2005,
<http://www.dow.com/polyethylene/na/en/application/rigid/> (accessed 23 April
2013).
- Arabsalmani, K. Evaluation of flowering fruiting and effect of seed extraction time on
seed quality characters of Cantaloupe (*Cucumis melo*) [dissertation]. Iran:
University of Tabriz; 1996. 65-80p.
- Arndt, G. W, BAM: Examination of Flexible and Semirigid Food Containers for
Integrity by U.S. Food and Drug Administration (FDA), 2001,
[http://www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm0727
03.htm](http://www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm072703.htm) (accessed 14 January 2015)
- Artés, F., Gómez, P. A. and Artés-Hernández, F. (2007). Physical, physiological and
microbial deterioration of minimally fresh processed fruits and vegetables.
Food Science and Technology International, 13: 177.
- Austin, J. W., Dodds, K. L., Blanchfield, B. and Farber, J. M. (1998). Growth and toxin
production by *Clostridium botulinum* on inoculated fresh-cut packaged
vegetables. *Journal of Food Protection*, 61(3):324-8.
- Bai, J. H., Saftner, R. A., Watada, A. E. and Lee, Y. S. (2001). Modified atmosphere
maintains quality of fresh- cut Cantaloupe (*Cucumis melo L.*). *Journal of Food
Science*, 66: 1207-1211.
- Bai, J. H., Saftner, R. A. and Watada, A. E. (2003). Characteristics of fresh-cut
honeydew (*Cucumis melo L.*) available to processors in winter and summer
and its quality maintenance by modified atmosphere packaging. *Postharvest
Biology and Technology*, 28: 349-359.
- Bail'en, G., Guillén, F., Castillo, S., Serrano, M., Valero, D. and Martínez-Romero, D.
(2006). Use of activated carbon inside modified atmosphere packaging to
maintain tomato fruit quality during cold storage. *Journal of Agricultural and
Food Chemistry*, 54: 2229-2235.
- Baner, A. and Piringer, O. (2008). *Preservation of Quality through Packaging. Plastic
Packaging: Interactions with Food and Pharmaceuticals, Second Edition.*
Germany: Wiley-VCH
- Barrett, D.M., Somogyi, L. and Ramaswamy, H.S. (2004), *Processing Fruits: Science
and Technology, Second Edition, Technology & Engineering.* Boca Raton, FL:
CRC Press

- Barrett, D. M., Beaulieu, J. C. and Shewfelt, R. (2010). Color, flavor, texture, and nutritional quality of fresh-cut fruits and vegetables: desirable levels, instrumental and sensory measurement, and the effects of processing. *Critical Reviews in Food Science and Nutrition*, 50: 369-389.
- Barry-Ryan, C., Martin-Diana, A., Rico, D. and Barat, J. (2007). Extending and measuring the quality of fresh-cut fruit and vegetables: a review. *Trends in Food Science and Technology*, 18:373-386.
- Barth, M., Hankinson, T. R., Zhuang, H. and Breidt, F. (2009). Microbiological Spoilage of Fruits and Vegetables. In W.H. Sperber, M.P. Doyle (Ed.), *Compendium of the Microbiological Spoilage of Foods and Beverages*. Food Microbiology and Food Safety (pp. 135-183). Lake Forest, IL: Springer Science.
- Beaudry, R.M., Cameron, A.C., Shirazi, A. and Dostal-Lange, D.L. (1992). Modified-atmosphere packaging of blueberry fruit: effect of temperature on package O₂ and CO₂. *Journal of the American Society for Horticultural Science*, 117:436-441.
- Beaudry, R. M. and Gran, C. D. (1993). Using a modified-atmosphere packaging approach to answer some postharvest questions: Factors affecting the lower oxygen limit. *Acta Hort*, 362:203-212.
- Beaudry, R. M. (2007). MAP as a Basis for Active Packaging. In C. L. Wilson (Ed.), *Intelligent and Active Packaging of Fruits and Vegetables* (pp. 31–55). Boca Raton, FL: CRC Press.
- Beaulieu, J. C. and Gorny, J. R. (2001). Fresh-cut fruits. In K. C. Gross., M. E. Saltveit, and C. Y.Wang (Ed.), *The Commercial Storage of Fruits, Vegetables, and Florist and Nursery Stocks* (pp. 1-49).USDA: Handbook 66.
- Beaulieu, J. C. and Baldwin, E. A. (2002). Flavor and Aroma of Fresh-cut Fruits and Vegetables. In O. Lamikanra (Ed.), *Fresh-Cut Fruits and Vegetables* (pp. 391-425). Boca Raton, FL: CRC Press.
- Beaulieu, J. C. and Lea, J. M. (2003). Aroma volatile differences in commercial orange-fleshed cantaloupes, the inbred parental lines, and stored fresh-cuts. *Acta Horticulturae*, 628: 809-815.
- Beaulieu, J. C., Ingram, D. A., Lea, J. M. and Bett-Garber, K. L. (2004). Effect of harvest maturity on the sensory characteristics of fresh-cut cantaloupe. *Journal of Food Science*, 69: 250-258.
- Beaulieu, J. C and Lea, J. M. (2007). Quality changes in cantaloupe during growth, maturation, and in stored minimally processed cubes prepared from fruit harvested at various maturities. *Journal of the American Society for Horticultural Science*, 132:720-8.

- Benedetti, B. C., Gómez, P., Martins, M., Conesa, A. and Artés, F. (2008). Effect of Pre-processing controlled atmosphere storage on the quality of fresh-cut Galia melons. *Food Science and Technology International*, 14: 13.
- Ben-Yehosiuu, S. (1961). The induction of the ripening process in avocado fruit. [Dissertation]. University of California, Los Angeles, California.
- Bett, K. L., Ingram, D. A., Grimm, C.C., Lloyd, S. W., Spanier, A. M., Miller, J. M., Gross, K. C., Baldwin, E. A. and Vinyard, B. T. (2001). Flavor of fresh-cut 'Gala' apples in modified atmosphere packaging as affected by storage time. *Journal of Food Quality*, 24(2): 141-156.
- Bett-Garber, K. L., Greene, J. L., Lamikanra, O., Ingram, D. A. and Watson, M. A. (2011). Effect of storage temperature variations on sensory quality of fresh-cut Cantaloupe melon. *Journal of Food Quality*, 34: 19-29.
- Bhattacharya, G, Served Fresh. Spotlight by Times Food Processing Journal, 2004, http://www.timesb2b.com/foodprocessing/dec03_jan04/spotlightml (accessed 18 July 2013)
- Bianco, V. V. and Pratt, H. K. (1977). Compositional changes in muskmelons during development and in response to ethylene treatment. *Journal of the American Society for Horticultural Science*, 102: 127-133.
- Bodelon, O. G., Blanch, M., Sanchez-Ballesta, M. T., Escribano, M. I. and Merodio, C. (2010). The effects of high CO₂ levels on anthocyanin composition, antioxidant activity and soluble sugar content of strawberries stored at low non-freezing temperature. *Food Chemistry*, 122: 673-678.
- Bolin, H. R. and Huxsoll, C. C. (1989). Storage stability of minimally processed fruit. *Journal of Food Biochemistry*, 13: 281-292.
- Boynton, B. B., Welt, B. A., Sims, C. A., Brecht, J. K., Balaban, M.O. and Marshall, M. R. (2005). Effects of low-dose electron beam irradiation on respiration, microbiology, color, and texture of fresh-cut cantaloupe. *Hort Science*, 15:802-807.
- Bozoglu, T. F., Deák, T. and Ray, B. (2001). *Novel Processes and Control Technologies in the Food Industry*. Antalya: IOS Press.
- Brewer, M. S. (1992). Reusing Food Packaging. . Is It Safe? In F. Weinbaum (Ed.), *Furtherance of Cooperative Extension Work University of Illinois at Urbana-Champaign* (pp. 1-10). University of Illinois: Cooperative Extension Service.
- Brody, A. I., Bugusu, B., Han, J. H., Sand, C. K. and Mchugh, T. H. (2008). Innovative Food Packaging Solutions. *Journal of Food Science*, 73(8).
- Brunazzi, G., Parisi, S. and Pereno, A. (2014). *The Importance of Packaging Design for the Chemistry of Food Products*. Italy: Springer Science

- Bryan Bruce Boynton. Determination of the effects of modified atmosphere packaging and irradiation on sensory characteristics, microbiology, texture and color of fresh-cut cantaloupe using modeling for package design. PhD Thesis, University Of Florida.2004.
- Cameron, A. C., Talasila, C. P. and Joles, D.W. (1995). Predicting film permeability needs for modified atmosphere packaging of lightly processed fruits and vegetables. *HortScience*, 30(1): 25-34.
- Caner, C., Aday, M. S. and Demir, M. (2008). Extending the quality of fresh strawberries by equilibrium modified atmosphere packaging. *European Food Research and Technology*, 227: 1575-1583.
- Cantwell, M. and Portela, S. (1997). Comparing varieties and storage method. *Fresh Cut*,14-18.
- Cantwell, P. and Hartz, T, Cantaloupe production in California by University of California, 2008, <http://anrcatalog.ucdavis.edu/pdf/7218.pdf> (accessed 3 July 2013).
- Cantwell, M and Suslow, T *Fresh-Cut Fruits and Vegetables: Aspects of Physiology, Preparation and Handling that Affect Quality*, Publication 3311, Postharvest Technology Horticultural Products, 1999, <http://ucanr.edu/datastoreFiles/608-357.pdf> (accessed 8 June 2015).
- Carrillo, L. A., Valdez, J. B., Rojas, R., Yahia, E. M. and Gomes, J. A. (1995). Ripening and quality of mangoes affected by coating with “Semperfresh”. *Acta Horticulturae*, 370: 203-216.
- Cha, D. S. and Chinnan, M. S. (2004). Biopolymer-based antimicrobial packaging: a review *Critical Reviews in Food Science and Nutrition*, 44: 223-237.
- Charles, F., Sanchez, J. and Gontard, N. (2003). Active modified atmosphere packaging of fresh fruits and vegetables: modeling with tomatoes and oxygen absorber. *Journal of Food Science*, 68: 1736-1742.
- Charles, F., Anchez, J. S., Gontard, N. (2005). Modeling of active modified atmosphere packaging of endives exposed to several postharvest temperatures. *Journal of Food Science*, 70: 443-449.
- Chia Su Ling. Effect of Ultraviolet Irradiation on Physicochemical, Microbial and Rheological Properties of Pineapple (*Ananas Comosus* L. Var Yankee) Juice. Master Thesis University Putra Malaysia.
- Chonhenchob, V., Chantarasomboon, Y. and Paul Singh, S. (2007). Quality changes of treated fresh-cut tropical fruits in rigid modified atmosphere packaging containers. *Packaging Technology Science*, 20: 27-37.
- Cia, P., Benato, E.A., Sigrist, J. M. M., Sarantopóulos, C., Oliveira, L. M. and Padula, M. (2006). Modified atmosphere packaging for extending the storage life of ‘Fuyu’ persimmon. *Postharvest Biology and Technology*, 42: 228-234.

- Cliff, M. A., Toivonen, P. M. A., Forney, C. F., Liu, P. and Lu, C. (2010). Quality of fresh-cut apple slices stored in solid and micro-perforated film packages having contrasting O₂ headspace atmospheres. *Postharvest Biology and Technology*, 58: 254-261.
- Clydesdale, F. M. (1993). Color as a factor in food choice. *Critical Reviews in Food Science and Nutrition*, 33(1): 83-101.
- Conte, A., Scrocco, C., Lecce, L., Mastromatteo, M. and Del Nobile, M.A. (2009). Ready-to-eat sweet cherries: Study on different packaging systems. *Innovative Food Science and Emerging Technologies*.
- Conway, W. S. and Sams, C. E. (1984). Possible mechanisms by which postharvest calcium treatment reduces decay in apples. *Phytopathology*, 74: 208-210.
- Corbo, M. R., Speranza, B., Campaniello, D., D'Amato, D. and Sinigaglia, M. (2010). Fresh-cut fruits preservation: current status and emerging technologies. In A. Mendex-Vilas, *Current Research, Technology and Education Topics in Applied Microbiology and Microbial Biotechnology* (pp.1143-1154).Spain: Formatex.
- Day, B.P.F. (1989). Extension of shelf-life of chilled foods. *European Food Drink Review*, 4: 47-56.
- Day, B. P. F. (1993). Fruit and vegetables. In: R. T Parry (Ed.), *Principles and applications of MAP of foods* (pp 114-33). New York, USA: Blackie Academic and Professional.
- Del-Valle, V., Hernandez-Munoz, P., Guarda, A. and Galotto, M. J. (2005). Development of a cactus-mucilage edible coating (*Opuntia ficus indica*) and its application to extend strawberry (*Fragaria ananassa*) shelf-life. *Food Chemistry*, 91(4): 751-756.
- Dong, X., Wrolstad, R. E. and Sugar, D. (2000). Extending shelf life of fresh-cut pears vegetables. *Sensory and Nutritive Qualities of Food*, 65:181-186.
- DOP, "Panduan Menanam Tembikai" by Department of Agriculture, Perak, 2010, http://www.pertanianperak.gov.my/jpp/index.php?option=com_content&view=article&id=358:panduan-menanam-tembikai&catid=68>manual-tanaman (accessed 1 August 2013).
- Doyle, M. P. (1990). Fruit and vegetable safety-microbiological considerations. *Hortscience*, 25: 1478-1481.
- Ergun, M., Jeong, J., Huber, D. J. and Cantliffe, D. J. (2007). Physiology of fresh-cut 'Galia' (*Cucumis melo* var. *reticulatus*) from ripe fruit treated with 1-methylcyclopropene. *Postharvest Biology and Technology*, 44: 286-292.
- Erol, A. and Sibel, T. (2003). A method for the measurement of the oxygen permeability and the development of edible films to reduce the rate of oxidative reactions in fresh foods. *Food Chemistry*, 80(3): 423-431.

- Erturk, E. and Picha, D. H. (2008). The effects of packaging film and storage temperature on the internal package atmosphere and fermentation enzyme activity of sweet potato slices. *Journal of Food Process Preservation*, 32: 817-838.
- Esturk, O., Ayhan, Z. and Gokkurt, T. (2014). Production and application of active packaging film with ethylene adsorber to increase the shelf life of broccoli (*Brassica Oleracea L. Var. Italica*). *Packaging Technology and Science*, 27: 179-191.
- Farber, J. M. (1991). Microbiological aspects of modified atmosphere packaging technology: a review. *Journal of Food Protection*, 54(1): 58-70.
- Floros, J. D. and Matsos, K. I. (2005). Introduction to modified atmosphere packaging. In: J. H Han (Ed.), *Innovations in food packaging* (pp. 159–172). Oxford, UK: Elsevier.
- Fonseca, S. C., Oliveira, F. A. R. and Brecht, J. K. (2002). Modelling respiration rate of fresh fruits and vegetables for modified atmosphere packages: a review. *Journal of Food Engineering*, 52: 99-119.
- Food and Drug Administration (FDA). Analysis and Evaluation of Preventive Control Measures for the Control and Reduction/Elimination of Microbial Hazards on Fresh and Fresh-Cut Produce: Chapter VI. Microbiological Safety of Controlled and Modified Atmosphere Packaging of Fresh and Fresh-Cut Produce, 2000, <http://www.fda.gov/Food/FoodScienceResearch/SafePracticesforFoodProcesses/ucm091016.htm> (accessed 8 September 2015)
- Food and Drug Administration (FDA). FDA advises consumers about produce safety, 2000, <http://www.cfsan.fda.gov/lrd/tpproduce.html> viewed (accessed 19 October 2014)
- Food and Drug Administration (FDA), FDA Survey of Domestic Fresh Produce. U.S. Food and Drug Administration by Center for Food Safety and Applied Nutrition, 2001, <http://www.cfsan.fda.gov/~dms/>(accessed 12 October 2013).
- Garcia, E. and Barrett, D. M. (2002). Preservative treatments for fresh-cut fruits and vegetables. In O. Laminkara, *Fresh-Cut Fruits and Vegetables: Science, Technology, and Market* (pp.91-117). Boca Raton: CRC Press.
- George Mateljan Foundation (GMF), Cantaloupe by World's Healthiest Foods, 2012, <http://www.whfoods.com/genpage.php?tname=foodspice&dbid=17>(accessed 21 April 2013).
- George, W. A. Jr, (NFPA) BAM: Examination of Flexible and Semirigid Food Containers for Integrity, 2001, <http://www.fda.gov/Food/FoodScienceResearch/LaboratoryMethods/ucm072703.htm> (accessed 16 May 2014).

- Gibe, A. J. G. and Kim, J. G. (2013). Influence of cutting size and packaging materials on the quality of fresh-cut winter squash (*var. Ajijimang*). *Agricultural Sciences*, 4(9): 477-482
- Gil, M. I., Aguayo, E. and Kader, A. A. (2006). Quality changes and nutrient retention in fresh-cut versus whole fruits during storage. *Journal of Agricultural and Food Chemistry*, 54(12): 4284-4296.
- González-Aguilar, G. A., Ruiz-Cruz, S., Cruz-Valenzuela, R., Rodríguez-Félix, A. and Wang, C.Y. (2004). Physiological and quality changes of fresh-cut pineapple treated with antibrowning agents. *Food Science and Technology*, 37: 369-376.
- González, J., Ferrer, A., Oria, R. and Salvador, M. L. (2008). Determination of O₂ and CO₂ transmission rates through microperforated films for modified atmosphere packaging of fresh fruits and vegetables. *Journal of Food Engineering*, 86(2):194-201.
- González-Aguilar, G. A., Ruiz-Cruz, S., Cruz-Valenzuela, R., Ayala-Zavala, J. F., De La Rosa, L. A. and Alvarez-Parrilla, E. (2008). New technologies to preserve quality of fresh-cut produce. In G. F. Gutiérrez-López, G. V. Barbosa-Cánovas, J. Welti-Chanes and E. Parada-Arias (Eds.), *Food Engineering: Integrated Approaches* (pp. 105–115). New York: Springer.
- Gorny, J. R. Modified Atmospheres Packaging and the Fresh-cut Revolution by Perishables Handling Newsletter, 1997, <http://ucanr.edu/datastoreFiles/234-159.pdf>(accessed 18 January 2013).
- Gorny, J. R., Gil, M. I. and Kader, A. A. (1998). Postharvest physiology and quality maintenance of fresh-cut pears. *Acta Horticulturae*, 464: 231-236.
- Gorny, J.R., Hess-Pierce, B., Kader, A.A. (1999). Quality changes in fresh-cut peach and nectarine slices as affected by cultivar, storage atmosphere and chemical treatments. *Journal of Food Science*, 64: 429-432.
- Gorny, J. 2001. A summary of CA and MA requirements and recommendations for fresh-cut (minimally processed) fruits and vegetables. Postharvest Horticultural Series No. 22. Davis, CA. Univ. Calif. Postharv. Outreach Program
- Gorny, J. R., Hess-Pierceb, B., Cifuentesb, R. A. and Kader, A. A. (2002). Quality changes in fresh-cut pear slices as affected by controlled atmospheres and chemical preservatives. *Postharvest Biology and Technology*, 24: 271-278.
- Gross, J. (1991). *Pigment in vegetables: Chlorophylls and Carotenoids*. New York: Van Nostrand Reinhold.
- Guynot, M. E., Sanchis, V., Ramos, A. J. and Marin, S. (2003). Mold-free shelf-life extension of bakery products by active packaging. *Journal of Food Science*, 68:2547-2552.
- Han, J. H. and Floros, J. D. (2007). Active packaging. In G. Tewari and V.K. Juneja (Ed.), *Advances in thermal and non-thermal food preservation* (pp. 167–183). Ames, LA: Blackwell Professional.

- Harker, F. R., Gunson, R. A. and Jaeger, S. R. (2003). The case for fruit quality: An interpretive review of consumer attitudes, and preferences for apples. *Postharvest Biology and Technology*, 28: 333-347.
- Harris, M. Food Packaging. (1999). *A Learning Cycle of Activities*. Washington: National Middle Level Science Teachers' Association (NMLSTA).a
- Hertzberg, R., Greene, J. and Vaughan, B. (2010). *Putting Food By: Fifth Edition*. London: Penguin Group.
- Hoffman, N. E. and Yang, S. F. (1982). Enhancement of wound-induced ethylene synthesis by ethylene in preclimacteric cantaloupe. *Plant Physiology*, 69: 317.
- Holcroft, D. M. and Kader, A. A. (1999). Controlled atmosphere-induced changes in pH and organic acid metabolism may affect color of stored strawberry fruit. *Postharvest Biology and Technology*, 17: 19-32.
- Hong, J. H. and Gross, K. C. (1998). Surface sterilization of whole tomato fruit with sodium hypochlorite influences subsequent postharvest behaviour of fresh-cut slices. *Postharvest Biology and Technology*, 13:51-58
- Howard, L. R., Yoo, K. S., Pike, L. M. and Miller Jr, G. H. (2006). Quality changes in diced onions stored in film packages. *Journal of Food Science*, 59(1): 110-112.
- Huang, D., Ou, B. and Prior, R. (2005). The chemistry behind antioxidant capacity assays. *Journal of Agricultural and Food Chemistry*, 53: 1841-1856.
- Huff, K. Active and intelligent packaging: innovations for the future [dissertation]. Virginia Polytechnic Institute and State University (Virginia Tech) Blacksburg; 2009. 13 p. Available from:<http://www.iopp.org/files/public/virginiatechkarleighhuff.pdf>(accessed 30 June 2013).
- International Fresh-cut Produce Association (IFPA), "Fresh-cut Produce: Get the Facts!", 2000, www.fresh-cuts.org(accessed 1 September 2013).
- International Fresh-cut Produce Association (IFPA), "Fresh-cut Facts", 2002, <http://www.creativeview.com/sites/ifpa/fcf.html> (accessed 10 July 2015).
- Ishaq, S., Rathore, H., Majeed, S., Awan. and Ali Shah, S. (2009). The studies on the physico-chemical and organoleptic characteristics of apricot (*Prunus armeniaca* L.) produced in Rawalakot, Azad Jammu and Kashmir during storage. *Pakistan Journal of Nutrition*, 8(6): 856-860.
- Jacobsson, A., Nielsen, T. and Sjöholm, I. (2004). Effects of type of packaging material on shelf-life of fresh broccoli by means of changes in weight, colour and texture. *European Food Research and Technology*, 218: 157-163.
- Jacxsens, L., Develieghere, F. and Debevere, J. (2002). Temperature dependence of shelf-life as affected by microbial proliferation and sensory quality of equilibrium modified atmosphere packaged fresh produce. *Postharvest Biology and Technology*, 26: 59-73.

- James, J. B. and Ngarmsak, T, *Processing of Fresh-Cut Tropical Fruits and Vegetables: A Technical Guide*. Food and Agriculture Organization of the United Nations Regional Office for Asia and the Pacific Bangkok, Rap Publication: Thailand. 2010.
- Kader, A. A., Zagory, D. and Kerbel, E. L. (1989). Modified atmosphere packaging of fruits and vegetables. *Critical Reviews in Food Science and Nutrition*, 28: 1-30.
- Kader, A. A. (1992). *Postharvest technology of horticultural crops*. Oakland CA: University of California Publication.
- Kader, A.A. (1997). A summary of CA requirements and recommendations for fruits other than apples and pears. In A.A. Kader (Ed.), *Fruits Other Than Apples and Pears* (pp. 1-36). California, USA: University of California Davis
- Kader, A. A. and Ben-Yehoshua, S. (2000). Effects of superatmospheric oxygen levels on postharvest physiology and quality of fresh fruits and vegetables. *Postharvest Biology and Technology*, 20: 1-13.
- Kartal, S. Aday, M. S. and Caner, C. (2012). Use of microperforated films and oxygen scavengers to maintain storage stability of fresh strawberries. *Postharvest Biology and Technology*, 71: 32-40.
- Kays, S.J. (1997). *Postharvest Physiology of Perishable Plant Products*, Van Nostrand Reinhold, New York, NY
- Kays, S. J. (1999). Preharvest factors affecting appearance. *Postharvest Biology and Technology*, 15: 233-247.
- Ke, D., Mateos, M., Siriphanich, J., Li, C. and Kader, A. A. (1993). Carbon dioxide action on metabolism of organic and amino acids in crisphead lettuce. *Postharvest Biology and Technology*, 3: 235-247.
- Kemble, J. M, Guide to the commercial production of muskmelon (cantaloupe) and related melons by Alabama cooperative extension system, 1996, <http://www.aces.edu/pubs/docs/A/ANR-0974/ANR-0974.pdf>(accessed 18 May 2014).
- Kim, J., Luo, Y. and Gross, K.C. (2003). Effect of packaging film on the quality of fresh-cut salad savoy. *Postharvest Biology and Technology*, 32: 99-107.
- Lai, H. M., Minimum-packaging technology for processed foods. APO Multi-Country Study Mission: Thailand. 2001.
- Lamikanra, O., Chen, J. C., Banks, D. and Hunter, P. A. (2000). Biochemical and microbial changes during the storage of minimally processed cantaloupe. *Journal of Agricultural and Food Chemistry*, 48: 5955-5961.
- Lamikanra, O. (2002). Preface. In O. Lamikanra (Ed.), *Fresh-cut fruits and vegetables. Science, technology and market*. Boca Raton, FL: CRC Press.

- Lamikanra, O., Juarez, B., Watson, M. A. and Richard, O. A. (2003). Effect of cutting and storage on sensory traits of cantaloupe melon cultivars with extended postharvest shelf life. *Journal of the Science of Food and Agriculture*, 83: 702-708.
- Lamikanra, O. and Watson, M. A. (2007). Mild heat and calcium treatment effects on fresh-cut cantaloupe melon during storage. *Food Chemistry*, 102: 1383-1388.
- Lange, D. (1998). Eastern cantaloupe varieties and their potential for fresh-cut melon products. *FreshCut*, (pp17-18).
- Larson, A. E. and Johnson, E. A. (1999). Evaluation of botulinal toxin production in packaged fresh-cut cantaloupe and honeydew melons. *Journal of Food Protection*, 62: 948-952.
- Latifah, M. N., Ab Aziz, I., Zaulia, O., Fauziah, O. and Talib, Y. (Eds.). (2011). Proceedings from ISHS 2011: *The 7th International Pineapple Symposium on Effect of Oxygen Scavenger Application on the Quality of Fresh-Cut Pineapple*. Mardi, Serdang: Selangor.
- Lee, L., Arul, J., Lencki, R. and Castaigne, F. (1995). A review on modified atmosphere packaging and preservation of fresh fruits and vegetables: physiological basis and practical aspects - part 1. *Journal of Packaging Technology Science*, 8:315-31.
- Lester, G. (1997). Melon (*Cucumis melo L*) fruit nutritional quality and health functionality. *Horticulture Technology*, 7: 222-7.
- Li, X., Li, W., Jiang, Y., Ding, Y., Yun, J., Tang, Y. and Zhang, P. (2011). Effect of nano-ZnO-coated active packaging on quality of fresh-cut 'Fuji' apple. *International Journal of Food Science and Technology*, 46: 1947-1955.
- Luna-Guzmán, I., Barrett, D. M. and Cantwell, M. *Effects of Calcium Chloride Dips and Heat Treatments on the Quality of Fresh-Cut Cantaloupe Melons*. Paper presented at the meeting of the Annual Meeting of Institute of Food Technologists. New Orleans, La. June 1996.
- Luna-Guzmán, I. (1997). Food safety and fresh-cut Cantaloupe. *Perishables Handling Quarterly*, 91: 13.
- Luna-Guzmán, I., Cantwell, M. and Barrett, D. M. (1999). Fresh-cut cantaloupe: effects of CaCl₂ dips and heat treatments on firmness and metabolic activity. *Postharvest Biology and Technology*, 17: 201-213.
- Luna-Guzmán, I. and Barrett, D. M. (2000). Comparison of calcium chloride and calcium lactate effectiveness in maintaining shelf stability and quality of fresh-cut cantaloupes. *Postharvest Biology and Technology*, 19(1):61-72.
- Machado, F. L. C., Alves, R. E. and Figueiredo, R. W. (2008). Application of 1-methylcyclopropene, calcium chloride and calcium amino acid chelate on fresh-cut cantaloupe muskmelon. *Pesquisa Agropecuária Brasileira*, 43(5): 569-574.

- Mangaraj, S., Goswami, T. K. and Mahajan, P. V. (2009). Applications of plastic films for modified atmosphere packaging of fruits and vegetables: a review. *Food Engineering Reviews*, 1: 133-158.
- Marsh, K. and Bugusu, B. (2007). Food packaging-roles, materials, and environmental issues. *Journal of Food Science*, 72(3): R39–R55.
- Marti´n-Diana, A. B., Rico, D., Fri´as, J. M., Barat, J. M., Henehan, G. T. M. and Barry-Ryan, C. (2007). Calcium for extending the shelf life of fresh whole and minimally processed fruits and vegetables: a review. *Food Science and Technology*, 18(4): 210-218.
- Martinez, M. V. and Whitaker, J. R. (1995). The biochemistry and control of enzymatic browning. *Trends in Food Science and Technology*, 6:195-200.
- Martinez-Romero, D. and Bail´en, G. (2007). Tools to maintain postharvest fruit and vegetables quality through the inhibition of ethylene action: a review. *Critical Reviews in Food Science and Nutrition*, 47:543-560.
- Mathooko, F. M. (1996): Regulation of respiration metabolism in fruits and vegetables by CO₂. *Postharvest Biology and Technology*. 9(3): 247-264.
- Mattheis, J.P. and Fellman, J.P. (2000). Impact of modified atmosphere packaging and controlled atmosphere on aroma, flavor and quality of horticultural produce. *Hort Technology*, 10:507-510.
- Mattos, L. M. Moretti, C. L. and Ferreira, M. D, Modified atmosphere packaging for perishable plant products, polypropylene. In F. Dogan (Ed.), 2012, <http://www.intechopen.com/books/polypropylene/modified-atmosphere-for-perishable-plant-products> (accessed 29 January 2014).
- Mayen, C. and Marshall, M. I, Opportunities in the Fresh-Cut Fruit Sector for Indiana Melon Growers by Purdue University Extension Publication EC-732, 2005, <https://www.extension.purdue.edu/extmedia/EC/EC-732.pdf> (accessed 2 September 2013)
- McGuire, R. G. (1992). Reporting of objective colour measurements. *Journal of HortScience*, 27 (12): 1254-1255.
- Mehyar, G. F. and Han, J. H. (2010). Active packaging for fresh-cut fruits and vegetables. In A.L. Brody, H. Zhuang and J. H. Han (Ed.), *Modified atmosphere packaging for fresh-cut fruits and vegetables* (pp. 267-283). Oxford: Blackwell Publishing Ltd.
- Mir, N. and Beaudry, R.M. (2004). Modified atmosphere packaging. In: K.C. Gross, C.Y. Wang, and M.E. Saltveit (Eds.), *The Commercial Storage of Fruits, Vegetables, and Florist and Nursery Stocks*, USDA Handbook 66., <http://www.ba.ars.usda.gov/hb66/ModAtmospherePack.pdf> (accessed 11 January 2013).

- Mohamed, S., Taufik, B. and Karim, M. N. A. (1995). Effect of modified atmosphere packaging on the physiochemical characteristics of ciku (*Achras sapota* L) at various storage temperature. *Journal of the Science of Food and Agriculture*, 70: 231-240.
- Montero-Calderón, M., Rojas-Graub, M. A. and Martín-Bellosob, O. (2008). Effect of packaging conditions on quality and shelf-life of fresh-cut pineapple (*Ananas comosus*). *Postharvest Biology and Technology*, 50:182–189.
- Moodley, R. S., Govinden, R. and Odhav, B. (2002). The effect of modified atmospheres and packaging on patulin production in apples. *Journal of Food Protection*, 65(5): 867-871.
- Ngcobo, M. E. K., Opara, U. L. and Thiart, G. D. (2012). Effects of packaging liners on cooling rate and quality attributes of table grape (cv. Regal Seedless). *Packaging Technology and Science*, 25:73–84.
- Nguyen-the, C. and Carlin, F. (1994). The microbiology of minimally processed fresh fruits and vegetables. *Critical Reviews in Food Science and Nutrition*, 34: 371-401.
- Nur Aida, M. P., Zaulia, O., Hairiyah, D., Che Omar, D. and Habsah, M. (Eds.). (2007). Proceedings from National Horticulture Conference on Effect of Washing Treatment on Microbial Changes and Sensory of Ready-to-Cook Mung Bean Sprout. Mardi, Serdang: Selangor.
- O’Beirne, D. and Francis, G. A. (2003). Reducing the Pathogen Risk in MAP-Prepared Produce. In: R. Ahvenainen (Ed.), *Novel food packaging techniques* (pp.231-286). Cambridge, UK: Woodhead Publishing Limited.
- O’Connor-Shaw, R. E., Roberts, R., Ford, A. L. and Nottingham, S. M. (1994). Shelf life of minimally processed honeydew melon, kiwifruit, papaya, pineapple and cantaloupe. *Journal of Food Science*, 59: 1202–1206.
- Olivas, G. I. and Barbosa-Canovas, G. V. (2005). Edible coatings for fresh-cut fruits. *Critical Reviews in Food Science and Nutrition*, 45: 657-663.
- Oms-Oliu, G., Soliva-Fortuny, R. and Mart’in-Belloso, O. (2007). Effect of ripeness on the shelf-life of fresh-cut melon preserved by modified atmosphere packaging. *European Food Research and Technology*, 225:301-311.
- Oms-Oliu, G., Soliva-Fortuny, R. and Mart’in-Belloso, O. (2008). Physiological and microbiological changes in fresh-cut pears stored in high oxygen active packaging compared with low oxygen active and passive modified atmosphere packaging. *Postharvest Biology and Technology*, 48:295-301.
- Orzolek, M. D., Bogash S. M., Kime, L. F., Harper, J. K, Cantaloupe Production. Agricultural Alternatives. Information and Communication Technologies in the College of Agricultural Sciences. The Pennsylvania State University by Penn State Cooperative Extension, 2006a, <http://pubs.cas.psu.edu/FreePubs/pdfs/ua436.pdf>(accessed 23 December 2013).

- Orzolek, M. D., Lamont, W. J., Kime, L. F., Bogash, S. M. and Harper, J. K, Watermelon Production. Agricultural Alternatives Series by Penn State Cooperative Extension, 2010, <http://extension.psu.edu/business/ag-alternatives/horticulture/melons-and-pumpkins/watermelon-production> (accessed 6 December 2013).
- Ozdemir, M. and Floros, J. D. (2004). Active food packaging technologies. *Critical Reviews in Food Science and Nutrition*, 44: 185-193.
- Park, C. M. and Beuchat, L. R. (1999). Evaluation of sanitizers for killing *Escherichia coli* O157:H7, *Salmonella* and naturally occurring microorganisms on cantaloupes, honeydew melons, and asparagus. *Dairy, Food and Environmental Sanitation*, 19: 842-847.
- Parker, S. P. (1997). McGraw-Hill Encyclopedia of Science and Technology. 8th edition. In McGraw-Hill (Ed.), (pp.194–197). New York, USA: McGraw-Hill Professional.
- Parnell, T. L., Suslow, T. and Harris, L. J, Cantaloupe: safe method to store, preserve and enjoy by University of California, Agriculture and Natural Resources Publication ANR 8095, 2003, <http://anrcatalog.ucdavis.edu>(accessed 12 June 2013).
- Pech, J. C., Bouzayen, M. and Latche, A. (2008). Climacteric fruit ripening: Ethylene-dependent and independent regulation of ripening pathways in melon fruit. *Plant Science*, 175: 114–120.
- Pocas, M. F, Innovations in Intelligent Packaging Technologies for Perishable Foods, 2001,<http://www.esb.ucp.pt/twt/embalagem/MyFiles/biblioteca/publicacoes/NATO3.pdf>(accessed 15 August 2014).
- Poovaiah, B.W. (1986). Role of calcium in prolonging the storage-life of fruits and vegetables. *Food Technology*, 40:86-89.
- Portela, S. I. and Cantwell, M. I. (2001). Cutting blade sharpness affects appearance and other quality attributes of fresh-cut Cantaloupe melon. *Journal of Food Science*, 66(9):1265-1270.
- Portela, S. I. and Cantwell, M. I. *Quality Changes During Storage of Intact and Fresh-Cut Cantaloupe Melon*. Paper presented at the 1999 Annual Meeting of American Society for Horticultural Science. Minneapolis, Minn. 27-31 July 1999.
- Powrie, W. D., Wu, C. H. and Skura, B. J. (1988). Preservation of cut and segmented fresh fruit pieces. *European Patent Application*.
- Powrie, W. D. and Skura, B. J. (1991). Modified atmosphere packaging of fruits and vegetables. In: B, Oraikul, M. E. Stiles and E, Horwood (Eds.), *Modified atmosphere packaging of food* (pp 169-245). New York, USA: Publisher unknown.

- Qi, L., Wu, T. and Watada, A. E. (1999). Quality changes of fresh-cut honeydew melons during controlled atmosphere storage. *Journal of Food Quality*, 22: 513-521.
- Ragaert, P., Jacxsens, L., Vandekinderen, I., Baert, L. and Devlieghere, F. (2011). Microbiological and Safety Aspects of Fresh-Cut Fruits and Vegetables. In O. Martin-Belloso and R. Soliva-Fortuny (Ed.), *Advances in Fresh Cut Fruits and Vegetables Processing* (pp.54-75). Boca Raton, Florida: CRC Press.
- Rathore, H. A., Masud, T., Sammi, S. and Soomro, A. H. (2007). Effect of storage on physico-chemical composition and sensory properties of mango (*Mangifera indica* L.) variety dosehari. *Pakistan Journal of Nutrition*, 6: 143-148.
- Rattanapanone, N., Lee, Y., Wu, T. and Watada, A. (2001). Quality and microbial changes of fresh-cut mango cubes held in controlled atmosphere. *HortScience*, 36: 1091-1095.
- Robert, C., Fortuny, S. and Belloso, O. M. (2003). New advances in extending the shelf life of fresh-cut fruits: a review. *Trends in Food Science and Technology*, 14: 341-353.
- Rodriguez-Aguilera, R. and Oliveira, J. C. (2009). Review of design engineering methods and applications of active and modified atmosphere packaging systems. *Food Engineering Reviews*, 1: 66-83.
- Rojas-Grau, M. A., Oms-Oliu, G., Soliva-Fortuny, R. and Martí'n-Belloso, O. (2009). The use of packaging techniques to maintain freshness in fresh-cut fruits and vegetables: a review. *International Journal of Food Science and Technology*, 44: 875-889.
- Saftner, R. A., Baj, J., Abbott, J. A. and Lee, Y. S. (2003). Sanitary dips with calcium propionate, calcium chloride, or calcium amino acid chelates maintain quality and shelf stability of fresh-cut honeydew chunks. *Postharvest Biology and Technology*, 29: 257-269.
- Saftner, R., Yaguang, L., James, M., Abbott, J. A. and Vinyard, B. (2006). Quality characteristics of fresh-cut watermelon slices from non-treated and 1-methylcyclopropene- and/or ethylene-treated whole fruit. *Postharvest Biology and Technology*.
- Sahelian, R., Cantaloupe Health Benefit, 2000, <http://www.raysahelian.com/cantaloupe.html> (accessed 21 August 2013).
- Saladie', M., Matas, A. J., Isaacson, T., Matthew, A., Jenks, S., Goodwin, M., Niklas, K. J., Xiaolin, R., Labavitch, J. M., Shackel, K. A., Fernie, A. R., Lytovchenko, A., O'Neill, M. A., Chris, B., Watkins. and Rose, J. K. C. (2007). A re-evaluation of the key factors that influence tomato fruit softening and integrity. *Plant Physiology*, 144: 1012-1028.
- Saltveit, M. E. (Eds.). (1993). Proceedings from *The Sixth National CA Conference on A summary of CA and MA requirements and recommendations for the storage of harvested vegetables*. Cornell University, Ithaca: New York.

- Saltveit, M. E. (1997). Physical and physiological changes in minimally processed fruits and vegetables. In F.A Tomás-Barberán and R.J Robins (Ed.), *Phytochemistry of fruits and vegetables* (pp. 205-220). Oxford, UK: Clarendon Press.
- Saltveit, M. E., Respiratory Metabolism, 2004, <http://www.ba.ars.usda.gov/hb66/respiratoryMetab.pdf> (accessed 8 August 2013).
- Sammi, S. and Masud T. (2007). Effect of different packaging systems on storage life and quality of tomato (*Lycopersicon esculentum* var. *rio grande*) during different ripening stages. *Journal of Food Safety*, 9: 37-44.
- Sammi, S. and Masud, T. (2008). Effect of different packaging systems on the quality of tomato (*Lycopersicon esculentum* var. Rio Grande) fruits during storage. *International Journal of Food Science and Technology*, 6: 1-9.
- Sandhya. (2010). Modified atmosphere packaging of fresh produce: Current status and future needs. *Food Science and Technology*, 43: 381-392.
- Sanjeev, K. and Ramesh, M. N. (2006). Low oxygen and inert gas processing of foods. *Critical Reviews in Food Science and Nutrition*, 46: 423-451.
- Sanz, C., Pérez, A. G., Olías, R. and Olías, J. M. (1999). Quality of strawberries packed with perforated polypropylene. *Journal of Food Science*, 64(4):748-752.
- Sapers, G. M., Miller, R. L., Pilizota, V. and Mattrazzo, A. M. (2006). Antimicrobial treatments for minimally processed cantaloupe melon. *Journal of Food Science*, 66(2): 345-349.
- Schlimme, D.V. and Rooney, M.L. (1994). Packaging of minimally processed fruits and vegetables. In: R.C. Wiley (Ed.), *Minimally processed refrigerated fruits and vegetables* (pp. 156-157). New York, USA: Chapman & Hall
- Senesi, E., Prinzevalli, C., Sala, M. and Gennari, M. (2000). Physicochemical and microbiological changes in fresh-cut green bell peppers as affected by packaging and storage. *Italian journal of food science*, 12:55-64.
- Shewfelt, R. (1994). Quality characteristics of fruits and vegetables. In R. P. Singh and F. A. R. Oliveira (Ed.), *Minimal Processing of Foods and Process Optimization: An Interface* (pp. 171-189). Boca Raton, FL: CRC Press.
- Silveira, A. C., Aguayo, E., Chisari, M. and Artés, F. (2011). Calcium salts and heat treatment for quality retention of fresh-cut 'Galia' melon. *Postharvest Biology and Technology*, 62(1): 77-84.
- Simmons, G. F., Rij, R., Smilanick, J. L. and John, S. *Microbial Population Reduction on Cantaloupe Destined for the Minimally Processed Market using Hydrogen Peroxide or SO₂*. Paper presented at the 1996 Annual Meeting of American Society for Horticultural Science. Lexington, Kentucky. October 1996.

- Simpson, R., Carevic, E. and Rojas, S. (2006). Modelling a modified atmosphere packaging system for fresh scallops (*Argopecten purpuratus*). *Packaging Technology and Science*, 20: 87-97.
- Smith, J. P., Ramaswamy, H. S. and Simpson, B. K. (1990). Developments in food packaging technology. Part II: Storage aspects. *Trends in Food Science and Technology*, 1:111-118.
- Smith, D. A. Selection of Food Containers: Glass Jars by Food Processing for Entrepreneurs Series, 2007, https://www.foodsafety.wisc.edu/assets/pdf_Files/Glassjars_Neb_Entre.pdf(accessed 11 September 2013).
- Soliva-Fortuny, R. C., Grigelmo-Miguel, N., Hernando, I., Lluch, M. A. and Martín-Belloso, O. (2002). Effect of minimal processing on the texture and structural properties of fresh-cut pears. *Journal of the Science of Food and Agriculture*, 82: 1682-1688.
- Soliva-Fortuny, R. C. and Martin-Belloso, O. (2003). New advances in extending the shelf life of fresh-cut fruits: a review. *Trends in Food Science and Technology*, 14: 341-353.
- Solomos, T. (1997). Principles Underlying Modified Atmosphere Packaging. In R.C Wiley (Ed.), *Minimally Processed Refrigerated Fruits and Vegetables* (pp.183–225). New York: Chapman and Hall.
- Song, J., W. and Deng, Fan, L. (1998). Aroma volatiles and quality changes in modified atmosphere packaging. In J.R. Gorny (Ed.), *Fresh-Cut Fruits and Vegetables and MAP* (pp. 89-95). University of California, Davis, USA
- Supapvanich, S. and Tucker, G. A. (2011). Physicochemical changes in fresh-cut Honeydew melon fruit during storage. *African Journal of Agricultural Research*, 6(12): 2737-2742.
- Tamplin, M. (1997). Salmonella and cantaloupes. *Dairy, Food and Environmental Sanitation*, 17: 284-286.
- Tarr, C.R. and Clingeffer, P.R. (2005). Use of an oxygen absorber for disinfestation of consumer packages of dried vine fruit and its effect on fruit colour. *Journal of Stored Product Research*, 41:77-89.
- Tice, P., Packaging materials 4. Polyethylene for food packaging applications. International Life Sciences Institute Report, 2003, [http:// orig. ils. org/ file/ PM4 _ Polyethylene.pdf](http://orig.ilsi.org/file/PM4_Polyethylene.pdf)(accessed 23 November 2013).
- Toivonen, P. M. A. and De-Ell, J. R. (2002). Physiology of fresh-cut fruits and vegetables. In O. Lamikanra (Ed.), *Fresh-Cut Fruits and Vegetables. Science, Technology and Market* (p.480) Boca Raton, FL: CRC Press.
- Toivonen, P. M. A. and Brummell, D. A. (2008). Biochemical bases of appearance and texture changes in fresh-cut fruit and vegetables. *Postharvest Biology and Technology*, 48: 1-14.

- Toivonen, P.M.A., Brandenburg, J.S. and Luo, Y. Modified Atmosphere Packaging for Fresh-Cut Produce, 2009, <http://naldc.nal.usda.gov/download/33518/PDF>(accessed 8 August 2015).
- Ukuku, D.O. (2004). Effect of hydrogen peroxide treatment on microbial quality and appearance of whole and fresh-cut melons contaminated with *Salmonella* spp. *International Journal of Food Microbiology*, 95: 137- 146.
- Ukuku, D. O. and Sapers, G. M. (2000). Effect of sanitizer treatments on *Salmonella* Stanley attached to the surface of Cantaloupe and cell transfers to the fresh-cut tissue during cutting practices. *Journal of Food Protection*, 64(9): 1286-91.
- Vermeiren, L., Devlieghere, F., Van Beest, M., de Kruijf, N. and Debevere, J. (1999). Developments in the active packaging of foods. *Trends in Food Science and Technology*, 10: 77-86.
- Watada, A. E., Ko, N. P. and Minott, D. A. (1996). Factors affecting quality of fresh-cut horticultural products. *Postharvest Biology and Technology*, 9: 115-125.
- Wei, C. I., Huang, T. S., Kim, J. M., Tamplin, M. L. and Bartz, J. A. (1995). Growth and survival of *Salmonella montevideo* on tomatoes and disinfection with chlorinated water. *Journal of Food Protection*, 58:829–836.
- Wright, K. P. and Kader, A. A. (1997). Effect of slicing and controlled atmosphere storage on the ascorbate content and quality of strawberries and persimmons. *Journal of Postharvest Biology and Technology*, 10:39–48.
- Yam, K. L., Takhistov, P. T. and Miltz, J. (2005). Intelligent packaging: concepts and applications. *Journal of Food Science*, 70: 1-10.
- Yamaguchi, M., Hughes, D. L., Yabumoto, K. and Jennings, W. G. (1977). Quality of cantaloupe muskmelons: variability and attributes. *Scientia Horticulturae*, 6: 59-70.
- Yang, H. H. and Lawsless, H. T. (2003). Descriptive analysis of divalent salts. *Journal of Sensory Studies*, 20: 97-113.
- Yeh, J. T., Cui, L., Chang, C. J., Jiang, T. and Chen, K. N. (2008). Investigation of the oxygen depletion properties of novel oxygen-scavenging plastics. *Journal of Applied Polymer Science*, 110: 1420-1434.
- Zagory, D. and Kader, A. A. (1988). Modified atmosphere packaging of fresh produce. *Food Technology*, 42:70-77.
- Zagory, D. (1995). Principles and practice of modified atmosphere packaging of horticultural commodities. In: J. M. Farber and K. L. Dodds (Ed.), *Principles of modified-atmosphere and sous-vide product packaging* (pp 175-204). Lancaster, PA: Technomic Publishing Co Inc
- Zainal Abidin, M., Shamsudin, R., Othman, Z. and Abdul Rahman, R. (2013). Effect of postharvest storage of whole fruit on physico-chemical and microbial changes

of fresh-cut Cantaloupe (*Cucumis melo L. Reticulatus cv. Glamour*).
International Food Research Journal, 20(1): 501-508.

Zaulia Othman. Effects of Postharvest Coatings and Heat Treatment on Quality of Stored Pineapple Fruits. PhD Thesis, Universiti Putra Malaysia.

Zhang, M. and Chen, S. Application of Modified Atmosphere Packaging of Fruit and Vegetables in China School of Food Science and Technology, Jiangnan University, Wuxi 214036, Jiangsu, China. 1996.

