



UNIVERSITI PUTRA MALAYSIA

***EFFECTS OF QUORUM SENSING AND ITS DEGRADER ON THE
VIRULENCE OF *Vibrio harveyi* TOWARDS TIGER GROUPER,
Epinephelus fuscoguttatus Forsskål LARVAE***

NOORASHIKIN BINTI MD. NOOR

FP 2015 17



**EFFECTS OF QUORUM SENSING AND ITS DEGRADER ON THE
VIRULENCE OF *Vibrio harveyi* TOWARDS TIGER GROUPER, *Epinephelus
fuscoguttatus* Forsskål LARVAE**

By

NOORASHIKIN BINTI MD. NOOR

**Thesis Submitted to the School of Graduate Studies, Universiti Putra Malaysia, in
Fullfilment of the Requirements for the Degree of Master of Science**

May 2015

All material contained within the thesis, including without limitation text, logos, icons, photographs and all other artwork, is copyright material of Universiti Putra Malaysia unless otherwise stated. Use may be made of any material contained within the thesis for non-commercial purposes from the copyright holder. Commercial use of material may only be made with the express, prior, written permission of Universiti Putra Malaysia.

Copyright © Universiti Putra Malaysia



© COPYRIGHT UPM

Abstract of thesis presented to the Senate of Universiti Putra Malaysia in fulfillment of the requirement for the degree of Master of Science

**EFFECTS OF QUORUM SENSING AND ITS DEGRADER ON THE
VIRULENCE OF *Vibrio harveyi* TOWARDS TIGER GROUPEL, *Epinephelus
fuscoguttatus* Forsskål LARVAE**

By

NOORASHIKIN BINTI MD. NOOR

May 2015

Chair: Natrah Fatin Mohd Ikhsan, PhD

Faculty: Agriculture

Quorum sensing (QS) is a bacterial cell-to-cell communication with small signal molecules such as acyl-homoserine lactones (AHL) that control a number of phenotypes including the regulation of virulence determinants in pathogenic bacteria. Therefore, quorum sensing degrader has been suggested as one of the biocontrol strategy to fight bacterial infections as an alternative to the use of antibiotics. In this study, the link between quorum sensing (QS) in *Vibrio harveyi* and its virulence towards tiger grouper (*Epinephelus fuscoguttatus*) were investigated. The virulence activity was studied using *V. harveyi* wild type and its QS mutants with constitutively maximal or minimal quorum sensing activity and different signal molecule (autoinducer) synthase gene mutants. The potential of live feed *Artemia* as the carrier of bacterial quorum sensing degrader for the tiger grouper larvae was also investigated. The results showed that the wild type *Vibrio harveyi* BB120 was pathogenic to grouper (*Epinephelus fuscoguttatus*) larvae causing more than 50% larval mortality after four days of challenge. Furthermore, the mortality of grouper larvae challenged with *V. harveyi* mutant JAF483 with maximally active QS (QS+) was significantly higher than the *V. harveyi* wild type BB120. Meanwhile, high survival was observed in the grouper larvae challenged to JAF548 with minimally active QS (QS-) compared to the JAF483 (QS+). High survival of larvae were also observed in the QS autoinducer mutant strains of JMH634 (QS triple autoinducer synthase mutant), BB152 (Harveyi autoinducer-1 (HAI-1) synthase mutant) and MM30 (Autoinducer-2 (AI-2) synthase mutant). In contrast, low survival was observed in the strain JMH603 (Cholerae autoinducer-1 (CAI-1) synthase mutant). This indicated that the HAI-1 and AI-2 QS signal molecules might play important roles for the virulence of *V. harveyi* towards grouper larvae but not CAI-1. To support this, addition of HAI-1 in the water restored the virulence of the mutant. The effects of the infection on the larvae could also be seen histologically. In addition, the *in vivo* expression data showed that the HAI-1 signal molecule upregulated the innate immune genes consisting of tripartite motif-containing protein 39 (TRIM39), Hecpudin-1 (Hep-1), peptidoglycan recognition protein SC2 (PGRP) and toll like receptor 5 (TLR5) at different hours compared to control. The expression of both control and the bacterial mutant BB152 without HAI-molecules remained low throughout the experimental period. Next, different bacterial QS degrader strains were isolated from *Artemia* and screened using *Chromobacterium violaceum* CV026

bioassay. The results showed six bacterial strains (four Gram-positive and two Gram-negative) isolated from *Artemia* are able to degrade AHL in two different *in vitro* assays. The highest bacterial AHL degrader, identified as *Bacillus litoralis* BP-ART/6 fully degraded 10 ppm AHL from 9 hours. Encapsulation of the *Bacillus* strain to *Artemia* also significantly improve the survival and total length of the live feed compared to control without addition of any bacteria. This study showed that bacterial strain isolated from *Artemia* can act as a quorum sensing degrader and can be encapsulated to *Artemia*. It was later observed that *B. litoralis* could also be used as probiotic bacteria administered through *Artemia* nauplii to protect grouper larvae against infection of *V. harveyi*.

Keywords: Tiger grouper; *Vibrio harveyi*; Quorum sensing; Quorum sensing degrader.



Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan untuk Ijazah Master Sains

KESAN PENDERIAAN KUORUM DAN PERENCATNYA TERHADAP KEVIRULENAN *Vibrio harveyi* KEPADA LARVA KERAPU HARIMAU, *Epinephelus fuscoguttatus* Forsskål

Oleh

NOORASHIKIN BINTI MD. NOOR

Mei 2015

Pengerusi: Natrah Fatin Mohd Ikhsan, PhD

Fakulti: Pertanian

Penderiaan kuorum (QS) adalah komunikasi bakteria antara bakteria sel ke sel dengan menggunakan molekul isyarat kecil dikenali sebagai 'acylated homoserine lactones' (AHL) yang mengawal beberapa fenotip termasuk pengaturan virulensi dalam bakteria patogenik. Oleh itu, perencatnya telah dicadangkan sebagai salah satu strategi untuk melawan jangkitan bakteria sebagai alternatif kepada antibiotik. Dalam kajian ini, hubungan antara penderiaan kuorum (QS) di antara *Vibrio harveyi* dan kevirulennanya terhadap kerapu harimau (*Epinephelus fuscoguttatus*) telah disiasat. Aktiviti ini dikaji menggunakan *V. harveyi* jenis liar dan mutan QS dengan aktiviti penderiaan kuorum maksimum atau minimum serta mutan gen bagi sintesis molekul isyarat yang berbeza. Potensi *Artemia* (makanan hidup ikan) sebagai pembawa bakteria perencat QS bagi larva kerapu harimau turut disiasat. Hasil kajian menunjukkan bahawa mutan liar *Vibrio harveyi* BB120 adalah patogenik kepada larva kerapu (*Epinephelus fuscoguttatus*) dengan lebih daripada 50 % kematian selepas empat hari. Tambahan pula, kadar kematian larva kerapu dicabar mutan JAF483 dengan aktiviti QS maksima (QS+) adalah jauh lebih tinggi daripada jenis liar BB120. Selain itu, hasil kajian menunjukkan kadar hidup larva kerapu tinggi apabila dicabar dengan JAF548 dengan aktiviti QS minimum (QS-) berbanding dengan JAF483 (QS+). Hasil kajian juga menunjukkan kadar hidup larva yang tinggi dalam JMH634 (mutan QS *triple autoinducer synthase*), BB152 (mutan *Harveyi autoinducer-1 (HAI-1) synthase*) dan MM30 (mutan *Autoinducer-2 (AI-2) synthase*). Sebaliknya, kadar hidup rendah diperhatikan dalam mutan JMH603 (mutan *Cholerae autoinducer-1 (CAI-1) synthase*). Ini menunjukkan bahawa molekul isyarat QS HAI-1 dan AI-2 mungkin memainkan peranan penting bagi kevirulenan *V. harveyi* terhadap larva kerapu tetapi tidak CAI-1. Untuk menyokong pendapat ini, penambahan HAI-1 di dalam air didapati memulihkan kevirulenan mutan. Kesan jangkitan juga boleh dilihat secara histologi. Molekul isyarat HAI-1 juga meningkatkan gen sistem imun *tripartite motif-containing protein 39 (TRIM39)*, *Hepcidin 1 (Hep-1)*, *peptidoglycan recognition protein SC2 (PGRP)* dan *toll like receptor 5 (TLR5)* pada waktu yang berbeza. Manakala, ekspresi gen imun adalah kekal rendah sepanjang tempoh eksperimen dalam rawatan terkawal dan mutan BB152 tanpa molekul HAI-1. Seterusnya, perencat QS telah diasingkan daripada *Artemia* menggunakan bioassai *Chromobacterium violaceum* CV026. Hasil kajian menunjukkan enam jenis bakteria (empat Gram-positif dan dua Gram-negatif) telah diasingkan daripada *Artemia* dan dapat merendahkan AHL dalam dua asai berbeza. Bakteria perencat QS yang terbaik dikenal pasti sebagai *Bacillus litoralis* (BP/Art-6), mampu merendahkan 10 ppm AHL sepenuhnya dalam 9 jam. Dalam

kajian ini, peningkatan yang ketara dalam hidup dan pertumbuhan dalam *Artemia* dicapai apabila dikultur bersama *B. litoralis*. Ia juga diperhatikan bahawa *B. litoralis* boleh digunakan sebagai bakteria probiotik melalui makanan hidup ikan *Artemia* yang mampu melindungi larva kerapu daripada jangkitan *V. harveyi*.

Kata Kunci: Kerapu harimau, *Vibrio harveyi*, penderiaan kuorum, perencat penderiaan kuorum



ACKNOWLEDGEMENTS

There are a number of people without whom this thesis might not have been written, and to whom I am greatly indebted.

To my parents, Norani and Md. Noor and members of my family who have been a source of encouragement and inspiration to me, a very special thank you for supported me in my determination to find and realize my potential.

I would like to express my deepest gratitude towards my main supervisor, Dr. Natrah Fatin Mohd Ikhsan who gives me guidance, encouragement and helpful advices in completing this research. Besides that, I would also like thank Assoc. Prof. Dr. Hassan Hj. Daud and Dr. Murni Marlina Karim for the meaningful consultation they gave as my co-supervisors.

I would also like to take this opportunity to thank the directors, lecturers, laboratory assistants and staffs of Department of Aquaculture, Faculty of Agriculture and Marine Biotechnology Laboratory, Institute of Bioscience especially Prof. Salleh, Prof. Fatimah, Mr. Farhan, Mrs. Norfarra, Mr. Zainan, Mrs. Norhasliza, Miss Zarirah, Mr. Zawawi, Mr. Jasni and Mrs. Shafika for their help during my study.

My sincere gratitude also goes to all of my postgraduate and undergraduate friends especially Nur Izzati, Wen Chen, Yan Li, Farha, Maria, Delphanie, Yenix, Chon, Aiman and my loved one Mohamad Khairol Amilin who were willing to spend their quality time to help me in completing my research. I would also like to thank Bonnie Bassler who has been provided me with the *Vibrio harveyi* quorum sensing strains. Lastly, I would like to thank to all who help me in completing this research as planned and hopefully this piece of work will be useful and beneficial to all.

I certify that a Thesis Examination Committee has met on 26th May 2015 to conduct the final examination of Noorashikin binti Md. Noor on her thesis entitled “Effects of quorum sensing and its degrader on the virulence of *Vibrio harveyi* towards tiger grouper, *Epinephelus fuscoguttatus* (Forsskål, 1775) larvae” in accordance with the Universities and University Colleges Act 1971 and the Constitution of the Universiti Putra Malaysia [P.U.(A) 106] 15 March 1988. The Committee recommends that the student be awarded the Master of Science.

Members of the Thesis Examination Committee were as follows:

Annie Christianus, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Ina Salwany Md Yassin, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Internal Examiner)

Yeong Yik Sung, PhD

Associate Professor
Institute of Marine Biotechnology
Universiti Malaysia Terengganu
(External Examiner)



(ZULKARNAIN ZAINAL, PhD)

Professor and Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 12 August 2015

This thesis was submitted to the Senate of Universiti Putra Malaysia and has been accepted as fulfilment of the requirement for the degree of (Master of Science). The members of the Supervisory Committee were as follows:

Natrah Fatin Mohd Ikhsan, PhD

Senior Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Chairman)

Hassan Mohd Daud, PhD

Associate Professor
Faculty of Veterinary Medicine
Universiti Putra Malaysia
(Member)

Murni Marlina Abdul Karim, PhD

Lecturer
Faculty of Agriculture
Universiti Putra Malaysia
(Member)

BUJANG BIN KIM HUAT, PhD

Professor and Dean
School of Graduate Studies
Universiti Putra Malaysia

Date:

Declaration by graduate student

I hereby confirm that:

- this thesis is my original work;
- quotations, illustrations and citations have been duly referenced;
- this thesis has not been submitted previously or concurrently for any other degree at any other institutions;
- intellectual property from the thesis and copyright of thesis are fully-owned by Universiti Putra Malaysia, as according to the Universiti Putra Malaysia (Research) Rules 2012;
- written permission must be obtained from supervisor and the office of Deputy Vice-Chancellor (Research and Innovation) before thesis is published (in the form of written, printed or in electronic form) including books, journals, modules, proceedings, popular writings, seminar papers, manuscripts, posters, reports, lecture notes, learning modules or any other materials as stated in the Universiti Putra Malaysia (Research) Rules 2012;
- there is no plagiarism or data falsification/fabrication in the thesis, and scholarly integrity is upheld as according to the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) and the Universiti Putra Malaysia (Research) Rules 2012. The thesis has undergone plagiarism detection software.

Signature: _____

Date:

Name and Matric No.: Noorashikin binti Md. Noor (Gs35651)

Declaration by Members of Supervisory Committee

This is to confirm that:

- the research conducted and the writing of this thesis was under our supervision;
- supervision responsibilities as stated in the Universiti Putra Malaysia (Graduate Studies) Rules 2003 (Revision 2012-2013) are adhered to.

Signature:

Name of
Chairman of
Supervisory

Committee: Natrah Fatin Mohd Ikhsan, PhD



DR. NATRAH IKHSAN
Lecturer
Department of Aquaculture
Faculty of Agriculture
Universiti Putra Malaysia
43400 UPM Serdang, Selangor.

Signature:

Name of
Member of
Supervisory
Committee:

Hassan Mohd. Daud, PhD



ASSOC PROF DR HASSAN HJ. MOHD. DAUD
DVM (UPM) MS (Stirling) PhD (Kingston, England)
Fish Health (Fish Virology)
Department of Veterinary Clinical Studies
Faculty of Veterinary Medicine
Universiti Putra Malaysia
43400 UPM Serdang, Selangor D. E.

Signature:

Name of
Member of
Supervisory
Committee:

Murni Marlina Abdul Karim, PhD



Dr. Murni Marlina Abd. Karim
Lecturer
Department of Aquaculture
Faculty of Agriculture
43400 UPM Serdang, Selangor

TABLE OF CONTENTS

		Page
ABSTRACT		i
ABSTRAK		iii
ACKNOWLEDGEMENTS		v
APPROVAL		vi
DECLARATION		viii
LIST OF TABLES		xiii
LIST OF FIGURES		xiii
LIST OF ABBREVIATIONS		xv
CHAPTER		
1	INTRODUCTION	1
2	LITERATURE REVIEW	4
	2.1 <i>Epinephelus fuscoguttatus</i>	4
	2.2 Larval rearing of grouper	5
	2.2.1 Live feed in grouper larviculture	5
	2.2.2 Problems related to grouper larviculture	6
	2.3 Bacterial management in overcome diseases in larviculture	7
	2.3.1 Matured seawater	7
	2.3.2 Prebiotics	8
	2.3.3 Probiotics	8
	2.3.4 Synbiotics	13
	2.4 Disruption of quorum sensing	13
	2.4.1 Quorum sensing in general	13
	2.4.2 Quorum sensing in <i>V. harveyi</i>	14
	2.4.3 Impacts of quorum sensing on aquaculture	15
	2.4.4 Disruption of quorum sensing	16
3	IMPACT OF QUORUM SENSING ON THE VIRULENCY OF <i>Vibrio harveyi</i> TOWARDS TIGER GROUPER, <i>Epinephelus fuscoguttatus</i> LARVAE	20
	3.1 Introduction	20
	3.2 Materials and methods	21
	3.2.1 Bacterial strains and growth condition	21
	3.2.2 Tiger grouper larvae experiment	21
	3.2.3 Determination of <i>Vibrio harveyi</i> in the grouper larvae	22
	3.2.4 Histopathological analysis	22
	3.2.5 Quantitative real time PCR	22
	3.2.5.1 RNA Extraction and quantification	22
	3.2.5.2 Primers and real time amplification	23

3.2.6	Statistical data analysis	24
3.3	Results and discussion	24
3.3.1	Impact of quorum sensing on tiger grouper	24
3.3.2	Impact of quorum sensing on immune related genes profile	29
3.4	Conclusion	33
4	SCREENING AND IDENTIFICATION OF QUORUM SENSING DEGRADERS FROM LIVE FEED <i>Artemia</i>	34
4.1	Introduction	34
4.2	Materials and method	35
4.2.1	QS molecules	35
4.2.2	Enrichment of AHL degraders from <i>Artemia</i>	35
4.2.3	QS degradation assay	35
4.2.4	Rifampicin resistant AHL degrader	36
4.2.5	Identification of AHL degrader strains	36
4.2.3.1	Gram staining	36
4.2.3.2	Biochemical test	37
4.2.3.3	Bacterial spore staining with malachite green	37
4.2.3.4	DNA extraction	37
4.2.3.5	Polymerase Chain Reaction (PCR)	37
4.2.6	Statistical analysis	39
4.3	Results and discussion	39
4.3.1	Isolation of QS degrader	39
4.3.2	QS degradation assay	39
4.3.3	Identification of QS degrader	44
4.4	Conclusion	46
5	EVALUATION ON THE EFFECTIVENESS OF BACTERIAL QUORUM SENSING DEGRADERS ENCAPSULATED TO <i>Artemia</i> ON <i>Epinephelus fuscoguttatus</i> LARVAE	48
5.1	Introduction	48
5.2	Materials and methods	49
5.2.1	Bacterial strain and growth	49
5.2.2	Axenic <i>Artemia</i> nauplii	49
5.2.3	Encapsulation of QS degrader to <i>Artemia</i> nauplii	50
5.2.4	Experimental challenge of QS degrader encapsulated to <i>Artemia</i> with <i>Vibrio harveyi</i>	50
5.2.5	Experimental challenge of grouper <i>E. fuscoguttatus</i> larvae with <i>V. harveyi</i> and evaluation of the probiotic <i>B. litoralis</i> effectiveness	50

5.2.6	Statistical analysis	51
5.3	Results and discussion	51
5.3.1	Encapsulation of QS degrader to <i>Artemia</i>	51
5.3.2	Experimental challenge of QS degrader encapsulated to <i>Artemia</i> with <i>V. harveyi</i>	54
5.3.3	<i>In vivo</i> grouper larvae	55
5.4	Conclusion	60
6	SUMMARY, GENERAL CONCLUSION AND RECOMMENDATIONS	61
	REFERENCES	64
	APPENDICES	87
	BIODATA OF STUDENT	91
	LIST OF PUBLICATIONS	92

LIST OF TABLES

Table	Page
2.1: Probiotics as biological control agents in larviculture	11
2.2: Disruption of quorum sensing for microbial control	18
3.1: List of <i>Vibrio harveyi</i> strains used in this study	21
3.2: List of primers used in qPCR gene expression analysis	23
3.3: Survival percentage of tiger grouper larvae after four days of challenge with <i>Vibrio harveyi</i> wildtype and quorum sensing mutants	24
3.4: <i>Vibrio harveyi</i> density in the grouper larvae after four days of challenge (log CFU/mL) on TCBS agar	25
4.1: PCR mastermix of 16S rRNA gene	38
4.2: AHL degradation of different bacterial samples at different hours	43
4.3: Characterization of QS degrader strains based on 16sr RNA gene	46
5.1: Bacterial count (Log 10 CFU/mL) in <i>Artemia</i> with QS degrader co-culture of different days	54
5.2: <i>Vibrio</i> density (Log 10 CFU/mL) in TCBS agar from grouper larvae during the first and last day of challenge	58
5.3: <i>Bacillus</i> density (Log 10 CFU/mL) in Marine agar supplemented with rifampicin from grouper larvae during the first and last day of challenge	58

LIST OF FIGURES

Figure	Page
2.1: <i>Epinephelus fuscoguttatus</i>	4
2.2: Multi-channel signalling in <i>Vibrio harveyi</i>	15
3.1: Histopathology changes of tiger grouper larvae gills with and without <i>V. harveyi</i> infection	27
3.2: Real time PCR expression result for immune genes responding to <i>Vibrio harveyi</i> strains infection in the tissue of tiger grouper (<i>Epinephelus fuscoguttatus</i>) larvae	31
4.1: Agarose gel (1.0% electrophoresis of Polymerase Chain Reaction (PCR) amplified DNA product (BP-ART/1,2,3&6) of 16S rRNA gene	38
4.2: Agarose gel (1.0% electrophoresis of Polymerase Chain Reaction (PCR) amplified DNA product (BP-ART/4 &5) of 16S rRNA gene.	38
4.3: Six degrader strains isolated from <i>Artemia</i> with AHL degradation properties on well plate.	41
4.4: Purple pigmentation for different AHLs concentration (ppm) for standard curve	41
4.5: Standard curve based on the diameter of the violacein at known AHL concentration	42
4.6: Degradation of AHL by BP-ART/6	42
4.7: <i>Bacillus</i> stained with malachite green under 100x magnification	47
5.1: An illustration of encapsulation	49
5.2: Experimental flow on larvae fed with encapsulated <i>Artemia</i> and challenged with <i>Vibrio</i>	51
5.3: Survival percentage (mean±standard deviation) of untreated and treated <i>Artemia</i> with QS degrader at different concentrations	52
5.4: Total length (mean±standard deviation) of untreated and treated <i>Artemia</i> with QS degrader at different concentration in fifteen days	53
5.5: Survival percentage (mean±standard deviation) of <i>Artemia</i> encapsulated with QS degrader following challenge with <i>V. harveyi</i>	55
5.6: Survival percentage (mean±standard deviation) of <i>E. fuscoguttatus</i> larvae three days post challenged with <i>V. harveyi</i> after fed with <i>B. litoralis</i> enriched <i>Artemia</i>	56
5.7: Histological changes (Magnification 400x, Haematoxylin & Eosin stain) of <i>E. fuscoguttatus</i> larvae gills in treatments	59

LIST OF ABBREVIATIONS

AHL	Acyl homoserine lactone
AI-2	Autoinducer-2
ATCC	American Type Culture Collection
CAI-1	Cholerae autoinducer-1
cDNA	complementary DNA
cfu	colony forming unit
DNA	Deoxyribonucleic acid
DO	dissolved oxygen
dpi	days post infection
FAO	Food and Agriculture Organization
H&E	Haematoxylin and Eosin
HAI-1	Harveyi autoinducer-1
HHL	Hexanoyl homoserine lactone
LB	Luria Bertani
MA	Marine agar
MB	Marine broth
MC	microbial communities
mRNA	Messenger RNA
NaCl	Sodium chloride
NCBI	National Centre for Biotechnology Information
OD	optical density
PCR	Polymerase chain reaction
ppm	parts per million
ppt	parts per thousand
QS	Quorum sensing
QSI	Quorum sensing inhibitor
RT-qPCR	Real time quantitative polymerase chain reaction
rpm	rotation per minute
rRNA	Ribosomal ribonucleic acid
SD	standard deviation
TCBS	Thiosulphate citrate bile salt sucrose
µg	microgram
µM	micromolar

CHAPTER 1

INTRODUCTION

Aquaculture is the farming of aquatic animals and plants (Costa-Pierce *et al.*, 2006). It involved the farming of freshwater and saltwater animal species including fishes, molluscs, crustaceans and aquatic plants. Unlike fishing where men remove aquatic organisms out of its habitat for consumption, aquaculture implies the cultivation of aquatic populations under controlled or semi-controlled conditions. Aquaculture has been growing steadily as a feedback to the declining of the world's fisheries due to overfishing. The aquaculture industry continues to expand, diversify and advance technologically dominating other animal-producing sectors in terms of growth (FAO, 2012).

Among the target species under the aquaculture sector are groupers. Currently groupers are cultured in large scales especially in Asian countries due to the advantages of this fish as efficient feed converters, fast growth and good flesh quality. The fish is much sought after by local and international markets, particularly in Asia such as Hong Kong, Singapore and Japan with high market price (FAO, 2012; Ottolenghi *et al.*, 2004). Compared to other grouper species, tiger grouper (*Epinephelus fuscoguttatus*) have high adaptability in captivity (Gunben *et al.*, 2014). However, one of the constraint in grouper production is that the supply of the juvenile fish is limited due to inadequate seed productions. In the hatchery, larval mortality of *Epinephelus fuscoguttatus* at the first feeding was common (Liao *et al.*, 2005). High densities of larvae and organic matter (live feed, faeces or from dead larvae debris) lead to the development of pathogens e.g., viruses, parasites and pathogenic bacteria causing poor growth or mass mortality. The control of pathogens in the live feed production and rearing systems are among the determinant aspects for the larvae endurance (Dhert *et al.*, 2001; Shirakashi *et al.*, 2013).

Of all pathogens, vibriosis among the common bacterial disease which affect various grouper culture stages (Manin and Ransangan, 2011). *Vibrio harveyi* are among the important bacterial pathogen in finfishes (Pietrak *et al.*, 2012), molluscs (Romalde *et al.*, 2013) and shrimps (Manilal *et al.*, 2010) culture which result in crucial financial losses in the farms and hatcheries (Lafferty *et al.*, 2015). Mainly, the ways for the control of bacterial infection is either to inhibit the growth of the microorganism or to attenuate its virulence leading to the failure of the organism. In aquaculture, antibiotics have been used to treat vibriosis. However, several pathogenic strains have shown resistance towards antibiotics. In juvenile fish, vaccination is given to treat diseases and it has lowered the use of chemotherapeutics in aquaculture and increased the survival of cultured fish. However, Vadstein (1997) stated vaccines are not suitable for larvae since the organisms are small and does not have a developed immune system and mainly depend on the non specific immune response and maternal antibodies. Thus, several researches continue to find new approach to a more environmentally friendly and sustainable control of vibriosis in the larvae culture systems as an alternative to antibiotics and vaccination.

The discoveries that bacteria use quorum sensing molecules, to regulate the production of virulence determinants and secondary metabolites could be a novel target to overcome bacterial diseases (Hamza *et al.*, 2015). In quorum sensing, bacteria communicate with one another by releasing, detecting and responding to the small signal molecules, also known as autoinducers. Detection of autoinducers allow bacteria to distinguish between low and high cell population density and regulate gene expression in response to changes in cell numbers. Many bacterial behaviours such as symbiosis, virulence, antibiotic production and biofilm are regulated by quorum sensing (Schauder and Bassler, 2001). It is shown that highly specific as well as universal quorum sensing languages exist, which enabled bacteria to communicate within and between species.

Interference of quorum sensing has been suggested as a new anti-infective strategy in bacteria. Several methods has been proposed to interfere the quorum sensing. This strategy involved any organism that is able in degrading quorum sensing signal molecules without inhibiting the pathogens growth. Since quorum sensing degraders do not disrupt the bacterial growth, it could reduce the risk of resistance due to the low selective pressure (Rutherford and Bassler, 2012). Furthermore, it allows the host's immune system to naturally recognize and destroy the pathogens.

In larviculture, live feed are necessary for first feeding because of their small mouth. The gut of the larvae is also not fully developed at the time of first feeding. Thus, the use of live feed may contribute to some exo-enzymes e.g., protease, lipase, amylase increment of production that would aid in the feed digestion of the fish (Metges, 2000). Furthermore, larvae is attracted to moving prey than inert particles or artificial feed. One of the common live feed used in aquaculture is the brine shrimp, *Artemia*. In aquaculture, the nauplii of *Artemia* are generally the most commonly used live feed organisms. *Artemia* cyst is easy to obtain and providing live *Artemia* nauplii to grouper larvae (after the rotifer-feeding stage) still remains essential in commercial hatchery. Annually, thousands of metric tons of dry *Artemia* cysts are marketed globally for use in marine aquaculture. However, *Artemia* shows low biochemical composition, therefore, enrichment is necessary for first feeding of larvae (Sorgeloos *et al.*, 2001).

Interestingly, the live nauplii of brine shrimp (*Artemia* spp.) have also been used as mediums for carrying compounds of diverse nutritional and therapeutic value to larval stages of fish through a process known as bioencapsulation. *Artemia* bioencapsulated with lactic acid bacteria have been successfully introduced into orange-spotted grouper larvae with significant improvements in survival (Sun *et al.*, 2013).

A survey made by Tanjong Demong Marine Fish Production and Research Centre (PPIL), Terengganu revealed that *Vibrio alginolyticus* and *Vibrio harveyi* are the most virulent *Vibrio* isolated from infected grouper during disease outbreak (Ransangan *et al.*, 2012; Ali *et al.*, 2008). Although antibiotics have been used as one of the prevention method, the use of antibiotics have had restricted success since studies have shown that the pathogen has developed resistance against antibiotics (Hamza *et al.*, 2015). This condition showed that there is a need to discover another option against *Vibrios*. Given that there are researches that showed pathogenicity of *V. harveyi* is regulated by quorum

sensing in other organisms as being demonstrated by Yang and Defoirdt (2014), Natrah *et al.*(2011a) and Ruwandeepika *et al.*(2011) thus, in this study we focused on the impacts of quorum sensing on the virulence of *V. harveyi* towards tiger grouper larvae.

To our knowledge there is no study of bacterial QS on grouper larvae. Furthermore, the potential of live feed *Artemia* as the carrier of bacterial quorum sensing degrader for tiger grouper larvae was also investigated. The specific objectives of the study are:

1. To investigate the virulence of *Vibrio harveyi* and the impact of *V. harveyi* QS towards tiger grouper *E. fuscoguttatus* larvae.
2. To isolate and screen QS degraders from live feed *Artemia*.
3. To evaluate the effectiveness of the QS degraders encapsulated to *Artemia* towards tiger grouper *E. fuscoguttatus* larvae.



REFERENCES

- Abbas, A. K., and Lichtman, A. H. (2006). *Basic Immunology*. USA:Saunders Elsevier.
- Abowei, J. F. N., and Briyai, O. F. (2011). A review of some bacteria diseases in Africa culture fisheries. *Asian Journal of Medical Sciences*, 3(5): 206-217.
- Ai Q., Xu H., Mai K., Xu W., and Wang J. (2011) Effects of dietary supplementation of *Bacillus subtilis* and fructooligosaccharide on growth performance, survival, non-specific immune response and disease resistance of juvenile large yellow croaker, *Larimichthys crocea*. *Aquaculture*, 317:155-161.
- Ali, H.M., Yusoff, N.H.N., Om, A.D., Musa, C.U.C. and Othman, M.F. (2008). Status and prospectus of grouper aquaculture in Malaysia. In: I.C. Liao and E.M. Leño, (eds). *The Aquaculture of Grouper*. (pp155-175). Taiwan: Asian Fisheries Society, World Aquaculture Society, the Fisheries Society of Taiwan and National Taiwan Ocean University.
- Arıg, N., Süzer, C., Gökvardar, A., Başaran, F., Çoban, D., Yıldırım, Ş., and Saka, Ş. (2013). Effects of probiotic (*Bacillus* sp.) supplementation during larval development of Gilthead sea bream (*Sparus aurata*, L.). *Turkish Journal of Fisheries and Aquatic Sciences*, 13(3).
- Attramadal, K. J., Truong, T. M. H., Bakke, I., Skjermo, J., Olsen, Y., and Vadstein, O. (2014). RAS and microbial maturation as tools for K-selection of microbial communities improve survival in cod larvae. *Aquaculture*, 432:483-490.
- Austin, B. and Zhang, X.H. (2006), *Vibrio harveyi*: a significant pathogen of marine vertebrates and invertebrates. *Letters in Applied Microbiology*, 43:119-124.
- Austin, B., and Brunt, J. W. (2009). The use of probiotics in aquaculture. *Aquaculture Microbiology and Biotechnology*, 1:185-207.
- Avella, M. A., Gioacchini, G., Decamp, O., Makridis, P., Bracciatelli, C., and Carnevali, O. (2010). Application of multi-species of *Bacillus* in sea bream larviculture. *Aquaculture*, 305(1): 12-19.
- Bai, F., Han, Y., Chen, J., and Zhang, X. H. (2008). Disruption of quorum sensing in *Vibrio harveyi* by the AiiA protein of *Bacillus thuringiensis*. *Aquaculture*, 274(1):36-40.
- Bairagi, A., Ghosh, K. S., Sen, S. K., and Ray, A. K. (2002). Enzyme producing bacterial flora isolated from fish digestive tracts. *Aquaculture International*, 10(2): 109-121.
- Bairagi, A., Sarkar Ghosh, K., Sen, S. K., and Ray, A. K. (2004). Evaluation of the nutritive value of *Leucaena leucocephala* leaf meal, inoculated with fish intestinal bacteria *Bacillus subtilis* and *Bacillus circulans* in formulated diets for rohu, *Labeo rohita* (Hamilton) fingerlings. *Aquaculture Research*, 35(5): 436-446.

- Bassler, B. L., Wright, M., and Silverman, M. R. (1994). Multiple signalling systems controlling expression of luminescence in *Vibrio harveyi*: sequence and function of genes encoding a second sensory pathway. *Molecular microbiology*, 13(2): 273-286.
- Bassler B. L., Greenberg E.P. and Stevens A. M. (1997). Cross-species induction of luminescence in the quorum-sensing bacterium *Vibrio harveyi*. *Journal Bacteriology*, 179:4043-4045.
- Basu, M., Swain, B., Maiti, N. K., Routray, P., and Samanta, M. (2012). Inductive expression of toll-like receptor 5 (TLR5) and associated downstream signaling molecules following ligand exposure and bacterial infection in the Indian major carp, mrigal (*Cirrhinus mrigala*). *Fish & shellfish immunology*, 32(1): 121-131.
- Beveridge, T. J. (2001). Use of the Gram stain in microbiology. *Biotechnic & Histochemistry*, 76(3): 111-118.
- Birkeland, C., and Dayton, P. K. (2005). The importance in fishery management of leaving the big ones. *Trends in Ecology and Evolution*, 20(7):356-358.
- Boonyaratpalin M. (1997) Nutrient requirements of marine food fish cultured in Southeast Asia. *Aquaculture*, 151:283-313.
- Boyer, M., and Wisniewski-Dyé, F. (2009). Cell–cell signalling in bacteria: not simply a matter of quorum. *FEMS microbiology ecology*, 70(1): 1-19.
- Brackman, G., Defoirdt, T., Miyamoto, C., Bossier, P., Van Calenbergh, S., Nelis, H., and Coenye, T. (2008). Cinnamaldehyde and cinnamaldehyde derivatives reduce virulence in *Vibrio* spp. by decreasing the DNA-binding activity of the quorum sensing response regulator LuxR. *BMC microbiology*, 8(1): 149-152.
- Brito, R., Rosas, C., Chimal, M. E., and Gaxiola, G. (2001). Effect of different diets on growth and digestive enzyme activity in *Litopenaeus vannamei* (Boone, 1931) early post larvae. *Aquaculture Research*, 32(4):257-266.
- Cam, D. T. V., Nhan, D. T., Ceuppens, S., Hao, N. V., Dierckens, K., Wille, M., and Bossier, P. (2009). Effect of N-acyl homoserine lactone-degrading enrichment cultures on *Macrobrachium rosenbergii* larviculture. *Aquaculture*, 294(1): 5-13.
- Cao, J. G., and Meighen, E. A. (1989). Purification and structural identification of an autoinducer for the luminescence system of *Vibrio harveyi*. *Journal of Biological Chemistry*, 264(36):21670-21676.
- Cao, M., Feng, Y., Wang, C., Zheng, F., Li, M., Liao, H., and Tang, J. (2011). Functional definition of LuxS, an autoinducer-2 (AI-2) synthase and its role in full virulence of *Streptococcus suis* serotype 2. *The Journal of Microbiology*, 49(6): 1000-1011.
- Carnevali, O., Zamponi, M. C., Sulpizio, R., Rollo, A., Nardi, M., Orpianesi, C., and Cresci, A. (2004). Administration of probiotic strain to improve sea bream wellness during development. *Aquaculture International*, 12(4): 377-386.

- Cerezuela R., Mesguer J. and Esteban M.A. (2011) Current Knowledge in Synbiotic Use for Fish Aquaculture: A Review. *Journal of Aquaculture Research and Development*, 10:1-8.
- Cerezuela, R., M. Fumanal, S.T. Tapia-Paniagua, J. Mesguer, M.A. Morinigo, and M.A. Esteban. (2012). Histological alterations and microbial ecology of the intestine in gilthead seabream (*Sparus aurata* L.) dietary probiotics and microalgae. *Cell and tissue research*, 350:477-489.
- Chakrabarti, R., Rathore, R. M., Mittal, P., and Kumar, S. (2006). Functional changes in digestive enzymes and characterization of proteases of silver carp (♂) and bighead carp (♀) hybrid, during early ontogeny. *Aquaculture*, 253(1): 694-702.
- Chang C.I., and Liu W.Y. (2002). An evaluation of two probiotic bacterial strains, *Enterococcus faecium* SF68 and *Bacillus toyoi*, for reducing edwardsiellosis in cultured European eel, *Anguilla anguilla*L. *Journal of Fish Diseases*, 25: 311–315.
- Chang, I. A., Kim, I. H., Kang, S. C., Hou, C. T., and Kim, H. R. (2007). Production of 7, 10-dihydroxy-8 (E)-octadecenoic acid from triolein via lipase induction by *Pseudomonas aeruginosa* PR3. *Applied microbiology and biotechnology*, 74(2):301-306.
- Chelladurai, G., Felicitta, J., and Nafarajan, R. (2013). Protective effects of probiotic diets on haematobiochemical and histopathology changes of *Mystus montanus* against *Aeromonas hydrophila*. *Journal of Coastal Life Medicine*, 1(4): 259-264.
- Chen, X., Schauder, S., Potier, N., Van Dorselaer, A., Pelczer, I., Bassler, B. L., and Hughson, F. M. (2002). Structural identification of a bacterial quorum-sensing signal containing boron. *Nature*, 415(6871): 545-549.
- Chen, F., Wang, M., Zheng, Y., Luo, J., Yang, X., and Wang, X. (2010). Quantitative changes of plant defense enzymes and phytohormone in biocontrol of cucumber *Fusarium wilt* by *Bacillus subtilis* B579. *World Journal of Microbiology and Biotechnology*, 26(4):675-684.
- Chen, Y., Yan, F., Chai, Y., Liu, H., Kolter, R., Losick, R., and Guo, J. H. (2013). Biocontrol of tomato wilt disease by *Bacillus subtilis* isolates from natural environments depends on conserved genes mediating biofilm formation. *Environmental microbiology*, 15(3): 848-864.
- Ching, B., Jamieson, S., Heath, J. W., Heath, D. D., & Hubberstey, A. (2010). Transcriptional differences between triploid and diploid Chinook salmon (*Oncorhynchus tshawytscha*) during live *Vibrio anguillarum* challenge. *Heredity*, 104(2): 224-234.
- Chiu, C. H., Cheng, C. H., Gua, W. R., Guu, Y. K., and Cheng, W. (2010). Dietary administration of the probiotic, *Saccharomyces cerevisiae* P13, enhanced the growth, innate immune responses, and disease resistance of the grouper, *Epinephelus coioides*. *Fish and shellfish immunology*, 29(6): 1053-1059.

- Christiaen, S. E., Matthijs, N., Zhang, X. H., Nelis, H. J., Bossier, P., and Coenye, T. (2014). Bacteria that inhibit quorum sensing decrease biofilm formation and virulence in *Pseudomonas aeruginosa* PAO1. *Pathogens and disease*, 70(3): 271-279.
- Chu, W., Lu, F., Zhu, W., and Kang, C. (2011). Isolation and characterization of new potential probiotic bacteria based on quorum sensing system. *Journal of applied microbiology*, 110(1): 202-208.
- Cordero, H., Esteban, M. Á., and Cuesta, A. (2014). Use of Probiotic Bacteria against Bacterial and Viral Infections in Shellfish and Fish Aquaculture.
- Costa-Pierce B., Desbonnet A., Edwards P. and Baker D. (2006). *Urban Aquaculture*. London, England: CABI Publishing.
- Cuesta, A., Meseguer, J., & Esteban, M. Á. (2008). The antimicrobial peptide hepcidin exerts an important role in the innate immunity against bacteria in the bony fish gilthead seabream. *Molecular immunology*, 45(8): 2333-2342.
- Czajkowski, R., and Jafra, S. (2009). Quenching of acyl-homoserine lactone-dependent quorum sensing by enzymatic disruption of signal molecules. *Acta Biochemical Pollution*, 56(1): 1-16.
- Das, B., Manna, S. K., Sarkar, P., and Batabyal, K. (2009). Occurrence of *Vibrio parahaemolyticus* in different finfish and shellfish species. *Journal of food safety*, 29(1):118-125.
- Dash, H. R., Mangwani, N., and Das, S. (2014). Characterization and potential application in mercury bioremediation of highly mercury-resistant marine bacterium *Bacillus thuringiensis* PW-05. *Environmental Science and Pollution Research*, 21(4):2642-2653.
- de Nys, R., Givskov, M., Kumar, N., Kjelleberg, S., and Steinberg, P. D. (2006). Furanones. In *Antifouling compounds* (pp.55-86). Berlin Heidelberg: Springer.
- Defoirdt T., Boon N., Bossier P., and Verstraete W. (2004). Disruption of bacterial quorum sensing: an unexplored strategy to fight infections in aquaculture. *Aquaculture*, 240:69-88.
- Defoirdt T., Bossier P., Sorgeloos P., and Verstraete W. (2005). The impact of mutations in the quorum sensing systems of *Aeromonas hydrophila*, *Vibrio anguillarum* and *Vibrio harveyi* on their virulence towards gnotobiotically cultured *Artemia franciscana*. *Environmental Microbiology*, 7:1239-1247.
- Defoirdt, T., Crab, R., Wood, T. K., Sorgeloos, P., Verstraete, W., and Bossier, P. (2006). Quorum sensing-disrupting brominated furanones protect the gnotobiotic brine shrimp *Artemia franciscana* from pathogenic *Vibrio harveyi*, *Vibrio campbellii*, and *Vibrio parahaemolyticus* isolates. *Applied and environmental microbiology*, 72(9): 6419-6423.

- Defoirdt, T., Boon, N., Sorgeloos, P., Verstraete, W., and Bossier, P. (2007). Quorum sensing and quorum quenching in *Vibrio harveyi*: lessons learned from *in vivo* work. *The ISME journal*, 2(1): 19-26.
- Defoirdt T., Verstraete W. and Bossier P. (2008). Luminescence, virulence and quorum sensing signal production by pathogenic *Vibrio campbellii* and *Vibrio harveyi* isolates. *Journal of Applied Microbiology*, 104:1480–1487.
- Defoirdt, T., Boon, N., and Bossier, P. (2010). Can bacteria evolve resistance to quorum sensing disruption?. *PLoS pathogens*, 6(7):98-105.
- Defoirdt, T., Sorgeloos, P., and Bossier, P. (2011). Alternatives to antibiotics for the control of bacterial disease in aquaculture. *Current opinion in microbiology*, 14(3): 251-258.
- Defoirdt, T. (2013). Virulence mechanisms of bacterial aquaculture pathogens and antivirulence therapy for aquaculture. *Reviews in Aquaculture*, 10:142-150.
- Dhert, P., Rombaut, G., Suantika, G., and Sorgeloos, P. (2001). Advancement of rotifer culture and manipulation techniques in Europe. *Aquaculture*, 200(1): 129-146.
- Dhert, P., King, N., & O'Brien, E. (2014). Stand-alone live food diets, an alternative to culture and enrichment diets for rotifers. *Aquaculture*, 431: 59-64.
- Diaz-Rosales, P., Salinas, I., Rodríguez, A., Cuesta, A., Chabrillón, M., Balebona, M. C., and Meseguer, J. (2006). Gilthead seabream (*Sparus aurata* L.) innate immune response after dietary administration of heat-inactivated potential probiotics. *Fish and shellfish immunology*, 20(4):482-492.
- Dimitroglou, A., Merrifield, D. L., Moate, R., Davies, S. J., Spring, P., Sweetman, J., and Bradley, G. (2009). Dietary mannan oligosaccharide supplementation modulates intestinal microbial ecology and improves gut morphology of rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Animal Science*, 87(10): 3226-3234
- Dimitroglou, A., Merrifield, D. L., Carnevali, O., Picchiatti, S., Avella, M., Daniels, C., and Davies, S. J. (2011). Microbial manipulations to improve fish health and production—a Mediterranean perspective. *Fish and Shellfish Immunology*, 30(1):1-16.
- Dobretsov, S., Teplitski, M., Alagely, A., Gunasekera, S. P., and Paul, V. J. (2010). Malynogolide from the cyanobacterium *Lyngbya majuscula* interferes with quorum sensing circuitry. *Environmental microbiology reports*, 2(6): 739-744.
- Dong, Y. H., Xu, J. L., Li, X. Z., and Zhang, L. H. (2000). AiiA, an enzyme that inactivates the acylhomoserine lactone quorum-sensing signal and attenuates the virulence of *Erwinia carotovora*. *Proceedings of the National Academy of Sciences*, 97(7): 3526-3531.

- Dong, Y. H., and Zhang, L. H. (2005). Quorum sensing and quorum-quenching enzymes. *Journal of Microbiology*, 43(5):101-109.
- Drancourt, M., Bollet, C., Carlioz, A., Martelin, R., Gayral, J. P., and Raoult, D. (2000). 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *Journal of clinical microbiology*, 38(10):3623-3630.
- Elward, K., and Gasque, P. (2003). “Eat me” and “don’t eat me” signals govern the innate immune response and tissue repair in the CNS: emphasis on the critical role of the complement system. *Molecular immunology*, 40(2): 85-94.
- Fang, A., and Demain, A. L. (2000). The natural functions of secondary metabolites. *Advanced Biochemical Engineering Biotechnology*, 69:1–39.
- Farahi, A., Kasiri, M., and Alamshahi, F. (2011). The effects on growth, survival and tolerance against environmental stressor (high temperature) of different concentrations probiotic *Bacillus* sp., fed to angelfish (*Pterophyllum scalare* Schultze, 1823) larvae. *J Anim Vet Adv*, 10(17): 2305-2311.
- Fisheries, F. A. O. (2012). Aquaculture Department. *Global Aquaculture Production Statistics for the year*.
- Feinerman, O., Jentsch, G., Tkach, K. E., Coward, J. W., Hathorn, M. M., Sneddon, M. W., & Altan-Bonnet, G. (2010). Single-cell quantification of IL-2 response by effector and regulatory T cells reveals critical plasticity in immune response. *Molecular systems biology*, 6(1): 201-204.
- Fernandez, R. G. (2001). *Artemia* bioencapsulation I. Effect of particle sizes on the filtering behavior of *Artemia franciscana*. *Journal of Crustacean Biology*, 21(2):435-442.
- Flagan, S., Ching, W. K., and Leadbetter, J. R. (2003). Arthrobacter strain VAI-A utilizes acyl-homoserine lactone inactivation products and stimulates quorum signal biodegradation by *Variovorax paradoxus*. *Applied and environmental microbiology*, 69(2): 909-916.
- Forberg, T., Arukwe, A., and Vadstein, O. (2011). A protocol and cultivation system for gnotobiotic Atlantic cod larvae (*Gadus morhua* L.) as a tool to study host microbe interactions. *Aquaculture*, 315(3): 222-227.
- Freeman, J. A., and Bassler, B. L. (1999). A genetic analysis of the function of LuxO, a two-component response regulator involved in quorum sensing in *Vibrio harveyi*. *Molecular microbiology*, 31(2): 665-677.
- Fuller, R. (1989). Probiotics in man and animals. *The Journal of applied bacteriology*, 66(5): 365-378.
- Gatesoupe F.J. (1999). The use of probiotic in Aquaculture. *Aquaculture*, 180:147-165.

- Gatesoupe, F. J. (2002). Probiotic and formaldehyde treatments of *Artemianauplii* as food for larval pollack, *Pollachius pollachius*. *Aquaculture*, 212(1): 347-360.
- Gatlin III, D. M., Li, P., Wang, X., Burr, G. S., Castille, F., & Lawrence, A. L. (2006). Potential application of prebiotics in aquaculture. *Avances en nutrición acuícola VIII. VIII Simposium I de Nutrición Acuícola*, 371-376.
- Geng X., Dong X.H., Tan B.P., Yang Q.H., and Chi S.Y. (2011) Effects of dietary chitosan and *Bacillus subtilis* on the growth performance, non-specific immunity and disease resistance of cobia, *Rachycentron canadum*. *Fish and Shellfish Immunology*, 31: 400-406.
- Gerwick, L., Corley-Smith, G., and Bayne, C. J. (2007). Gene transcript changes in individual rainbow trout livers following an inflammatory stimulus. *Fish & shellfish immunology*, 22(3): 157-171.
- Gomez-Gil, B., Tron-Mayen, L., Roque, A., Turnbull, J. F., Inglis, V., and Guerra-Flores, A. L. (1998). Species of *Vibrio* isolated from hepatopancreas, haemolymph and digestive tract of a population of healthy juvenile *Penaeus vannamei*. *Aquaculture*, 163(1): 1-9.
- Gomez-Gil, B., Roque, A., and Turnbull, J. F. (2000). The use and selection of probiotic bacteria for use in the culture of larval aquatic organisms. *Aquaculture*, 191(1): 259-270.
- Gomez-Gil B., Rogue A., and Velasco-Blanco G. (2002). Culture of *Vibrio alginolyticus* C7b, a potential probiotic bacterium, with the microalga *Chaetoceros muelleri*. *Aquaculture*, 211: 43-48.
- Gomez-Gil B., Soto-Rodriguez S., Garcia-Gasca A., Roque A., Vazquez-Juarez R., Thompson F.L., and Swings J. (2004). Molecular identification of *Vibrio harveyi*-related isolates associated with diseased aquatic organisms. *Microbiology*, 150:1769-1777.
- Gowda K.L., Marie M.A., John J., Pradeep C.S., Dabwan K.H., and Sangeetha G. (2014). Cytopathic effects of toxogenic strains on *Helicobacter pylori* on different cell lines. *Indian Journal of Pathology Microbiology*, 57(2): 187-190.
- Gram L., Løvold T., Nielsen J., Melchiorson J. and Spanggaard B. (2001). *In vitro* antagonism of the probiont *Pseudomonas fluorescens* strain AH2 against *Aeromonas salmonicida* does not confer protection of salmon against furunculosis. *Aquaculture*, 199: 1-11.
- Gunben, E. M., Senoo, S., Yong, A., and Shapawi, R. (2014). High Potential of Poultry by-product Meal as a Main Protein Source in the Formulated Feeds for a Commonly Cultured Grouper in Malaysia (*Epinephelus fuscoguttatus*). *Sains Malaysiana*, 43(3): 399-405.
- Halver J. E. (2002). *Fish Nutrition*. 3rd Ed. San Diego, USA:Academic Press Inc.

- Hamouda, T., Shih, A. Y., and Baker, J. R. (2002). A rapid staining technique for the detection of the initiation of germination of bacterial spores. *Letters in applied microbiology*, 34(2): 86-90.
- Hamza, F., Kumar, A. R., and Zinjarde, S. (2015). Biotechnological Applications of Quorum-Sensing Inhibitors in Aquacultures. In *Quorum Sensing vs Quorum Quenching: A Battle with No End in Sight* (pp. 371-383). India: Springer.
- Hansen, G. H., and Olafsen, J. A. (1999). Bacterial interactions in early life stages of marine cold water fish. *Microbial Ecology*, 38(1): 1-26.
- Harikrishnan, R., Balasundaram, C., and Heo, M. S. (2011). Fish health aspects in grouper aquaculture. *Aquaculture*, 320(1): 1-21.
- He, F., Morita, H., Kubota, A., Ouwehand, A. C., Hosoda, M., Hiramatsu, M., and Kurisaki, J. I. (2005). Effect of Orally Administered Non-Viable *Lactobacillus* Cells on Murine Humoral Immune Responses. *Microbiology and immunology*, 49(11): 993-997.
- Henke, J. M., and Bassler, B. L. (2004). Three parallel quorum-sensing systems regulate gene expression in *Vibrio harveyi*. *Journal of bacteriology*, 186(20):6902-6914.
- Hidalgo, M. C., Urea, E., and Sanz, A. (1999). Comparative study of digestive enzymes in fish with different nutritional habits. Proteolytic and amylase activities. *Aquaculture*, 170(3): 267-283.
- H.M. Ali, N.H.N. Yusoff, A.D. Daud, C.U.C. Musa and M.F. Othman. (2008). Status and Prospects of Grouper Aquaculture in Malaysia. *World Aquaculture Society*, 155-175.
- Hoj, L., Bourne, D.G., and Hall, M.R. (2009). Localization, abundance and community structure of bacteria associated with *Artemia*: Effects of nauplii enrichment and antimicrobial treatment. *Aquaculture*, 293(3): 278-285.
- Hoseinifar, S. H., Mirvaghefi, A., Amoozegar, M. A., Sharifian, M., & Esteban, M. Á. (2015). Modulation of innate immune response, mucosal parameters and disease resistance in rainbow trout (*Oncorhynchus mykiss*) upon synbiotic feeding. *Fish & shellfish immunology*, 45(1): 27-32.
- Hsieh, J. C., Pan, C. Y., and Chen, J. Y. (2010). Tilapia hepcidin (TH) 2-3 as a transgene in transgenic fish enhances resistance to *Vibrio vulnificus* infection and causes variations in immune-related genes after infection by different bacterial species. *Fish & shellfish immunology*, 29(3): 430-439.
- Hu, G. B., Lou, H. M., Dong, X. Z., Liu, Q. M., and Zhang, S. C. (2012). Characteristics of the interferon regulatory factor 5 (IRF5) and its expression in response to LCDV and poly I: C challenges in Japanese flounder, *Paralichthys olivaceus*. *Developmental and Comparative Immunology*, 38(2): 377-382.

- Hueck, C. J. (1998). Type III protein secretion systems in bacterial pathogens of animals and plants. *Microbiology and molecular biology reviews*, 62(2):379-433.
- Hynes, N. A., Furnes, C., Fredriksen, B. N., Winther, T., Bøggwald, J., Larsen, A. N., and Dalmo, R. A. (2011). Immune response of Atlantic salmon to recombinant flagellin. *Vaccine*, 29(44): 7678-7687.
- Interaminense, J. A., Ferreira Calazans, N., do Valle, B. C., Lyra Vogeley, J., Peixoto, S., Soares, R., and Lima Filho, J. V. (2014). *Vibrio* spp. Control at Brine Shrimp, *Artemia*, Hatching and Enrichment. *Journal of the World Aquaculture Society*, 45(1): 65-74.
- Irianto A., and Austin B. (2002). Use of probiotics to control furunculosis in rainbow trout, *Oncorhynchus mykiss* (Walbaum). *Journal of Fish Diseases*, 25: 333–342.
- Itoh, N., and Takahashi, K. G. (2008). Distribution of multiple peptidoglycan recognition proteins in the tissues of Pacific oyster, *Crassostrea gigas*. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 150(4): 409-417.
- Jafaryan, H., Taati, M. M., and Jafarzadeh, M. (2011). The enhancement of growth parameters in common carp (*Cyprinus carpio*) larvae using probiotic in rearing tanks and feeding by various *Artemia* nauplii. *AACL Bioflux*, 4(4): 511-518.
- Jamali H, Tafi AA, Jafaryan H., and Patimar R. (2014). Effects of Enriched *Artemia* with Probiotic (*Bacillus* spp.) on Growth , Survival, Fecal Production and Nitrogenous Excretion in Rainbow Trout (*Oncorhynchus mykiss*) Larvae. *Journal Fisheries Livestock Production*, 2:5-11.
- Kalia, D. (2015). Synthetic Quorum Sensing Inhibitors: Signal Analogues. In *Quorum Sensing vs Quorum Quenching: A Battle with No End in Sight* (pp. 285-302). India: Springer.
- Kanemitsu, K., Nishino, T., Kunishima, H., Okamura, N., Takemura, H., Yamamoto, H., and Kaku, M. (2001). Quantitative determination of gelatinase activity among enterococci. *Journal of microbiological methods*, 47(1): 11-16.
- Kawai, T., and Akira, S. (2005). Pathogen recognition with Toll-like receptors. *Current opinion in immunology*, 17(4): 338-344.
- Kim C., Kim S., and Shin S. (2012). Iron-mediated regulation of metalloprotease VvpE Production in *Vibrio vulnificus*, 20:481–486.
- Kolkovski, S. (2001). Digestive enzymes in fish larvae and juveniles—implications and applications to formulated diets. *Aquaculture*, 200(1):181-201.
- Korsnes K., Nicolaisen O., Skar C. K., Nerland A. H. and Bergh Ø. (2006). Bacteria in the gut of juvenile cod *Gadus morhua* fed live feed enriched with four different commercial diets. *ICES Journal of Marine Science*, 63: 296-301.

- Lafferty, K. D., Harvell, C. D., Conrad, J. M., Friedman, C. S., Kent, M. L., Kuris, A. M., & Saksida, S. M. (2015). Infectious diseases affect marine fisheries and aquaculture economics. *Annual review of marine science*, 7:471-496.
- Lamari, F., Sadok, K., Bakhrouf, A., and Gatesoupe, F. J. (2014). Selection of lactic acid bacteria as candidate probiotics and *in vivo* test on *Artemia* nauplii. *Aquaculture International*, 22(2): 699-709.
- Lara-Flores, M. (2011). The use of probiotic in aquaculture: an overview. *Int Res Journal Microbiology* 2(12): 471-478.
- Lauth, X., Babon, J. J., Stannard, J. A., Singh, S., Nizet, V., Carlberg, J. M., and Westerman, M. E. (2005). Bass hepcidin synthesis, solution structure, antimicrobial activities and synergism, and *in vivo* hepatic response to bacterial infections. *Journal of Biological Chemistry*, 280(10): 9272-9282.
- Lazzarini, A., Cavaletti, L., Toppo, G., and Marinelli, F. (2000). Rare genera of actinomycetes as potential producers of new antibiotics. *Antonie van Leeuwenhoek*, 78(3-4): 399-405.
- Lee, S. J., Park, S. Y., Lee, J. J., Yum, D. Y., Koo, B. T., and Lee, J. K. (2002). Genes encoding the N-acyl homoserine lactone-degrading enzyme are widespread in many subspecies of *Bacillus thuringiensis*. *Applied and Environmental Microbiology*, 68(8):3919-3924.
- Lee, S. M., Lee, S. B., Park, C. H., and Choi, J. (2006). Expression of heat shock protein and hemoglobin genes in *Chironomus tentans* (Diptera, chironomidae) larvae exposed to various environmental pollutants: A potential biomarker of freshwater monitoring. *Chemosphere*, 65(6): 1074-1081.
- Leung, H. M., Leung, S. K. S., Au, C. K., Cheung, K. C., Wong, Y. K., Leung, A. O. W., & Yung, K. K. L. (2015). Comparative assessment of water quality parameters of mariculture for fish production in Hong Kong Waters. *Marine pollution bulletin*, 94(1):318-322.
- Li, X., Wang, S., Qi, J., Echtenkamp, S. F., Chatterjee, R., Wang, M., and Gupta, D. (2007). Zebrafish peptidoglycan recognition proteins are bactericidal amidases essential for defense against bacterial infections. *Immunity*, 27(3): 518-529.
- Li, X., Yeung, D. F., Fiegen, A. M., and Sodroski, J. (2011). Determinants of the higher order association of the restriction factor TRIM5 α and other tripartite motif (TRIM) proteins. *Journal of Biological Chemistry*, 286(32): 27959-27970.
- Liao J.J, Chen Y-M, and Chen Tzong-Yueh.2005. Cloning and Characterization of the Grouper (*Epinephelus coioides*) SPARC Gene. 2005 Annual Conference of the Fisheries Society of Taiwan, Taipei, Taiwan. June 2005.
- Lilly, B. N., & Bassler, B. L. (2000). Regulation of quorum sensing in *Vibrio harveyi* by LuxO and sigma-54. *Molecular microbiology*, 36(4): 940-954.

- Liu, P. C., Lee, K. K., Yii, K. C., Kou, G. H., and Chen, S. N. (1996). News & Notes: Isolation of *Vibrio harveyi* from Diseased Kuruma Prawns *Penaeus japonicus*. *Current microbiology*, 33(2):129-132.
- Liu, K.F., Chiu, C.H., Shiu, Y.L., Cheng, W., and Liu, C.H. (2010). Effects of the probiotic *Bacillus subtilis* E20 on the survival, development, stress tolerance and immune status of white shrimp, *Litopenaeus vannamei* larvae. *Fish and shellfish immunology*, 28(5): 837-844.
- Liu, C. H., Chiu, C. H., Wang, S. W., and Cheng, W. (2012). Dietary administration of the probiotic, *Bacillus subtilis* E20, enhances the growth, innate immune responses, and disease resistance of the grouper, *Epinephelus coioides*. *Fish and shellfish immunology*, 33(4): 699-706.
- Livak, K. J., and Schmittgen, T. D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the $2\Delta\Delta CT$ Method. *Methods*, 25(4): 402-408.
- Looi, C.L. and Senoo S. (2008) Egg development and larval development of the grouper hybrid fish *Epinephelus fuscoguttatus* × Giant grouper *E. lanceolatus*. *Aquaculture*, 56(4):505-512.
- Magnadóttir B. Innate immunity of fish (overview) (2006). *Fish Shellfish Immunol*, 20:137–151.
- Mahdhi, A., Esteban, M. Á., Hmila, Z., Bekir, K., Kamoun, F., Bakhrouf, A., and Krifi, B. (2012). Survival and retention of the probiotic properties of *Bacillus* sp. strains under marine stress starvation conditions and their potential use as a probiotic in *Artemia* culture. *Research in veterinary science*, 93(3): 1151-1159.
- Mahious, A. S., Gatesoupe, F. J., Hervi, M., Metailler, R., and Ollevier, F. (2006). Effect of dietary inulin and oligosaccharides as prebiotics for weaning turbot, *Psetta maxima* (Linnaeus, C. 1758). *Aquaculture International*, 14(3): 219-229.
- Makridis, P., Fjellheim, A. J., Skjermo, J., and Vadstein, O. (2000). Control of the bacterial flora of *Brachionus plicatilis* and *Artemia franciscana* by incubation in bacterial suspensions. *Aquaculture*, 185(3): 207-218.
- Manefield, M., de Nys, R., Naresh, K., Roger, R., Givskov, M., Peter, S., and Kjelleberg, S. (1999). Evidence that halogenated furanones from *Delisea pulchra* inhibit acylated homoserine lactone (AHL)-mediated gene expression by displacing the AHL signal from its receptor protein. *Microbiology*, 145(2):283-291.
- Manefield M., Harris L., Rice S.A., Nys R. D., and Kjelleberg S. (2000). Inhibition of luminescence and virulence in the black tiger prawn (*Penaeus monodon*) pathogen *Vibrio harveyi* by intercellular signal antagonists. *Applied Environmental Microbiology*, 66:2079-2084.

- Manilal, A., Sujith, S., Selvin, J., Shakir, C., Gandhimathi, R., and Kiran, G. S. (2010). Virulence of vibrios isolated from diseased black tiger shrimp, *Penaeus monodon*, Fabricius. *Journal of the World Aquaculture Society*, 41(3): 332-343.
- Manin B.O., and Ransangan J. (2011). Experimental evidence of horizontal transmission of Betanodavirus in hatchery-produced Asian seabass, *Lates calcarifer* and brown-marbled grouper, *Epinephelus fuscoguttatus* fingerling. *Aquaculture*, 321:157-165.
- Mc Clean, K. H., Winson, M. K., Fish, L., Taylor, A., Chhabra, S. R., Camara, M., and Williams, P. (1997). Quorum sensing and *Chromobacterium violaceum*: exploitation of violacein production and inhibition for the detection of N-acylhomoserine lactones. *Microbiology*, 143(12): 3703-3711.
- Metges, C. C. (2000). Contribution of microbial amino acids to amino acid homeostasis of the host. *The Journal of nutrition*, 130(7):1857-1864.
- Michaud, L., Blancheton, J. P., Bruni, V., and Piedrahita, R. (2006). Effect of particulate organic carbon on heterotrophic bacterial populations and nitrification efficiency in biological filters. *Aquacultural Engineering*, 34(3):224-233.
- Miller, B. and Bassler B.L. (2001). "Quorum sensing in bacteria". *Annual Revised Microbiology*, 55: 165–199.
- Miyamoto, C. M., Smith, E. E., Swartzman, E., Cao, J. G., Graham, A. F., and Meighen, E. A. (1994). Proximal and distal sites bind LuxR independently and activate expression of the *Vibrio harveyi* lux operon. *Molecular microbiology*, 14(2): 255-262.
- Mohapatra, S., Chakraborty, T., Prusty, A. K., Das, P., Paniprasad, K., and Mohanta, K. N. (2012). Use of different microbial probiotics in the diet of rohu, *Labeo rohita* fingerlings: effects on growth, nutrient digestibility and retention, digestive enzyme activities and intestinal microflora. *Aquaculture Nutrition*, 18(1): 1-11.
- Mohapatra, S., Chakraborty, T., Prusty, A. K., Das, P., Paniprasad, K., and Mohanta, K. N. (2014). Dietary multispecies probiotics supplementation enhances the immunohematological responses and reduces mortality by *Aeromonas hydrophilla* in *Labeo rohita* fingerlings. *Journal of the World Aquaculture Society*, 45(5): 532-544.
- Mok K.C., Wingreen N.S., and Bassler B.L. (2003). *Vibrio harveyi* quorum sensing: A coincidence detector for two autoinducers controls gene expression. *EMBO Journal*, 22:870–281.
- Morohoshi, T., Kato, M., Fukamachi, K., Kato, N., and Ikeda, T. (2008). N-Acyl homoserine lactone regulates production of *Chromobacterium violaceum* type strain ATCC 12472. *FEMS microbiology letters*, 279(1):124-130.
- Mongkoltharuk, W., and Dharmstithi, S. (2002). Biodegradation of lipid-rich wastewater by a mixed bacterial consortium. *International biodeterioration and biodegradation*, 50(2):101-105.

- Moraes, F. R., and Martins M. L. (2004): Predisposing conditions and principal diseases of intensive fish farming teleosts. In: *Especial Topics in Intensive Freshwater Fish Culture in the Tropics*. (pp343-382). São Paulo, Brazil: TecArt.
- Motlagh, H. R. A., Farhangi, M., Rafiee, G., and Noori, F. (2012). Modulating gut microbiota and digestive enzyme activities of *Artemia urmiana* by administration of different levels of *Bacillus subtilis* and *Bacillus licheniformis*. *Aquaculture International*, 20(4):693-705.
- Mukai, T., T. Asasaka, E. Sato, K. Mori, M. Matsumoto and Ohori H. (2002). Inhibition of binding of *Helicobacter pylori* to the glycolipid receptors by probiotic *Lactobacillus reuteri*. *Fems Immunology Medicine Microbiology*, 32: 105-110.
- Muñoz, I., Sepulcre, M. P., Meseguer, J., and Mulero, V. (2013). Molecular cloning, phylogenetic analysis and functional characterization of soluble Toll-like receptor 5 in gilthead seabream, *Sparus aurata*. *Fish and shellfish immunology*, 35(1): 36-45.
- Musthafa, K. S., Saroja, V., Pandian, S. K., and Ravi, A. V. (2011). Antipathogenic potential of marine *Bacillus* sp. SS4 on N-acyl-homoserine-lactone-mediated virulence factors production in *Pseudomonas aeruginosa* (PAO1). *Journal of biosciences*, 36(1): 55-67.
- Natrah F.M.I., Darshanee Ruwandepika HA, Pawar S, Karunasagar I, Sorgeloos P, Bossier P, and Defoirdt T. (2011). Regulation of virulence factors by quorum sensing in *Vibrio harveyi*. *Veterinary Microbiology*, 154:124-129.
- Natrah, F. M. I., Defoirdt, T., Sorgeloos, P., and Bossier, P. (2011). Disruption of bacterial cell-to-cell communication by marine organisms and its relevance to aquaculture. *Marine Biotechnology*, 13(2): 109-126.
- Natrah, F. M. I., Alam, M. I., Pawar, S., Harzevili, A. S., Nevejan, N., Boon, N., and Defoirdt, T. (2012). The impact of quorum sensing on the virulence of *Aeromonas hydrophila* and *Aeromonas salmonicida* towards burbot (*Lota lota* L.) larvae. *Veterinary microbiology*, 159(1): 77-82.
- Nayak, S. K. (2010). Probiotics and immunity: a fish perspective. *Fish and shellfish immunology*, 29(1): 2-14.
- Neto, J. A., Martins, A. C. P., Silva, L. D. M. D., Tiraboschi, R. B., Domingos, A. L. A., Cologna, A. J., and Tucci Jr, S. (2003). Community acquired urinary tract infection: etiology and bacterial susceptibility. *Acta Cirurgica Brasileira*, 18: 33-36.
- Ng, W. L., Perez, L. J., Wei, Y., Kraml, C., Semmelhack, M. F., & Bassler, B. L. (2011). Signal production and detection specificity in *Vibrio* CqsA/CqsS quorum-sensing systems. *Molecular microbiology*, 79(6): 1407-1417.
- Nhan, D. T., Wille, M., De Schryver, P., Defoirdt, T., Bossier, P., and Sorgeloos, P. (2010). The effect of poly β -hydroxybutyrate on larviculture of the giant freshwater prawn *Macrobrachium rosenbergii*. *Aquaculture*, 302(1): 76-81.

- Nicholson, W. L. (2002). Roles of *Bacillus* endospores in the environment. *Cellular and Molecular Life Sciences CMLS*, 59(3): 410-416.
- Nielsen, A., Månsson, M., Bojer, M. S., Gram, L., Larsen, T. O., Novick, R. P., & Ingmer, H. (2014). *Solonamide B* inhibits quorum sensing and reduces *Staphylococcus aureus* mediated killing of human neutrophils. *PloS one*, 9(1): 84-92.
- Nikoskelainen, S., Ouwehand, A., Salminen, S., and Bylund, G. (2001). Protection of rainbow trout (*Oncorhynchus mykiss*) from furunculosis by *Lactobacillus rhamnosus*. *Aquaculture*, 198: 229–236.
- Nithya, C., Aravindraja, C., & Pandian, S. K. (2010). *Bacillus pumilus* of Palk Bay origin inhibits quorum-sensing-mediated virulence factors in Gram-negative bacteria. *Research in microbiology*, 161(4): 293-304.
- Niu, Y., Defoirdt, T., Rekecki, A., De Schryver, P., Van den Broeck, W., Dong, S., and Bossier, P. (2012). A method for the specific detection of resident bacteria in brine shrimp larvae. *Journal of microbiological methods*, 89(1): 33-37.
- Novick, R. P., and Muir, T. W. (1999). Virulence gene regulation by peptides in staphylococci and other Gram-positive bacteria. *Current opinion in microbiology*, 2(1): 40-45.
- Nylund, G. M., Persson, F., Lindegarth, M., Cervin, G., Hermansson, M., and Pavia, H. (2010). The red alga *Bonnemaisonia asparagoides* regulates epiphytic bacterial abundance and community composition by chemical defence. *FEMS microbiology ecology*, 71(1): 84-9.
- Oelschlaeger, T. A. (2010). Mechanisms of probiotic actions—a review. *International Journal of Medical Microbiology*, 300(1): 57-62.
- Ottolenghi, F., Silvestri, C., Giordano, P., Lovatelli, A., and New, M. B. (2004). *Capture-based aquaculture: the fattening of eels, groupers, tunas and yellowtails*. FAO.
- Ozato, K., Shin, D. M., Chang, T. H., and Morse, H. C. (2008). TRIM family proteins and their emerging roles in innate immunity. *Nature Reviews Immunology*, 8(11): 849-860.
- Pande, G. S. J., Scheie, A. A., Benneche, T., Wille, M., Sorgeloos, P., Bossier, P., and Defoirdt, T. (2013). Quorum sensing-disrupting compounds protect larvae of the giant freshwater prawn *Macrobrachium rosenbergii* from *Vibrio harveyi* infection. *Aquaculture*, 406: 121-124.
- Panigrahi, A., Kiron, V., Puangkaew, J., Kobayashi, T., Satoh, S., and Sugita, H. (2005). The viability of probiotic bacteria as a factor influencing the immune response in rainbow trout *Oncorhynchus mykiss*. *Aquaculture*, 243(1): 241-254.
- Park, S. C., Shimamura, I., Fukunaga, M., Mori, K. I., and Nakai, T. (2000). Isolation of bacteriophages specific to a fish pathogen, *Pseudomonas plecoglossicida*, as a

candidate for disease control. *Applied and environmental microbiology*, 66(4):1416-1422

- Park, S.Y., Hwang, B. J., Shin, M. H., Kim, J., Kim, H. K., and Lee, J. K. (2006). N-Acylhomoserine lactonase producing *Rhodococcus* spp. with different AHL-degrading activities. *FEMS microbiology letters*, 261(1):102-108.
- Pfaffl, M. W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic acids research*, 29(9), 40-45.
- Picchietti, S., Mazzini, M., Taddei, A. R., Renna, R., Fausto, A. M., Mulero, V., and Abelli, L. (2007). Effects of administration of probiotic strains on larval gilthead seabream: immunohistochemical and ultrastructural studies. *Fish and shellfish immunology*, 22(1): 57-67.
- Picchietti, S., Fausto, A. M., Randelli, E., Carnevali, O., Taddei, A. R., Buonocore, F., and Abelli, L. (2009). Early treatment with *Lactobacillus delbrueckii* strain induces an increase in intestinal T-cells and granulocytes and modulates immune-related genes of larval *Dicentrarchus labrax* (L.). *Fish and shellfish immunology*, 26(3): 368-376.
- Pietrak, M. R., Molloy, S. D., Bouchard, D. A., Singer, J. T., and Bricknell, I. (2012). Potential role of *Mytilus edulis* in modulating the infectious pressure of *Vibrio anguillarum* 02 β on an integrated multi-trophic aquaculture farm. *Aquaculture*, 326: 36-39.
- Pietretti, D., and Wiegertjes, G. F. (2014). Ligand specificities of Toll-like receptors in fish: indications from infection studies. *Developmental and Comparative Immunology*, 43(2): 205-222.
- Porsby, C. H., Nielsen, K. F., and Gram, L. (2008). *Phaeobacter* and *Ruegeria* species of the *Roseobacter* clade colonize separate niches in a Danish turbot (*Scophthalmus maximus*)-rearing farm and antagonize *Vibrio anguillarum* under different growth conditions. *Applied and environmental microbiology*, 74(23):7356-7364.
- Quigley E.M. (2010) Prebiotics and probiotics; modifying and mining the microbiota. *Pharmacology Resource* 61: 213-218.
- Rajkumar, M., and Kumaraguru K.P. (2006) Suitability of the copepod, *Acartia clausi* as a live feed for sea bass larvae (*Lates calcarifer*): Compared to traditional live food organisms with special emphasis on the nutritional value. *Aquaculture*, 261(2): 649-658.
- Ransangan, J., and Mustafa S. (2009). Identification of *Vibrio harveyi* isolated from Asian seabass (*Lates calcarifer*) by use of 16S ribosomal DNA sequencing. *Journal of Aquatic Animal Health*, 21: 150-155.
- Ransangan, J., Lal, T. M., and Al-Harbi, A. H. (2012). Characterization and experimental infection of *Vibrio harveyi* isolated from diseased Asian seabass (*Lates calcarifer*). *Malaysian Journal of Microbiology*, 8(2): 104-115.

- Rantakari, A., Virtaharju, O., Vähämiko, S., Taira, S., Palva, E. T., Saarilahti, H. T., and Romantschuk, M. (2001). Type III secretion contributes to the pathogenesis of the soft-rot pathogen *Erwinia carotovora*: partial characterization of the hrp gene cluster. *Molecular plant-microbe interactions*, 14(8): 962-968.
- Rasch M., Buch C., Austin B., Slierendrecht W.J., Ekmann K. S., Larsen J.C., Johansen C., Riedel K., Eberl L., Givskov M., and Gram L. (2004). An inhibitor of bacterial quorum sensing reduces mortalities caused by vibriosis in rainbow trout (*Oncorhynchus mykiss*, Walbaum). *Applied Microbiology*, 27:350-359.
- Ravi, A.V., Musthafa, K.S., Jegathammbal, G., Kathiresan K., and Pandian, S.K. (2007). Screening and evaluation of probiotics as a biocontrol agent against pathogenic *Vibrios* in marine aquaculture. *Letters in Applied Microbiology*. 45 (2): 219-223.
- Rekecki, A., Gunasekara, R., Dierckens, K., Laureau, S., Boon, N., Favoreel, H., and Van Den Broeck, W. (2012). Bacterial host interaction of GFP-labelled *Vibrio anguillarum* HI-610 with gnotobiotic sea bass, *Dicentrarchus labrax* (L.), larvae. *Journal of fish diseases*, 35(4): 265-273.
- Ringo, E., Olsen, R. E., Gifstad, T. Ø., Dalmo, R. A., Amlund, H., Hemre, G. I., and Bakke, A. M. (2010). Prebiotics in aquaculture: a review. *Aquaculture Nutrition*, 16(2):117-136.
- Rodriguez-Estrada U., Satoh S., Haga Y., Fushimi H., and Sweetman J. (2009) Effects of single and combined supplementation of *Enterococcus faecalis*, mannanoligosaccharide and polyhydrobutyric acid on growth performance and immune response of rainbow trout *Oncorhynchus mykiss*. *Aquaculture Science*, 57: 609-617.
- Romalde, J. L., Diéguez, A. L., Lasa, A., and Balboa, S. (2013). New *Vibrio* species associated to molluscan microbiota: a review. *Frontiers in microbiology*, 4:122-128.
- Rombaut, G., Dhert, P., Vandenberghe, J., Verschuere, L., Sorgeloos, P., and Verstraete, W. (1999). Selection of bacteria enhancing the growth rate of axenically hatched rotifers (*Brachionus plicatilis*). *Aquaculture*, 176(3): 195-207.
- Romero, M., Martin-Cuadrado, A.B., Roca-Rivalda, A., Cabello, A.M., and Otero, A. (2011). Quorum quenching in cultivable bacteria from dense marine coastal microbial communities. *FEMS microbiology ecology*, 75(2):205-217.
- Rønnestad, I., Conceição, L. E., Aragão, C., & Dinis, M. T. (2000). Free amino acids are absorbed faster and assimilated more efficiently than protein in postlarval Senegal sole (*Solea senegalensis*). *The Journal of nutrition*, 130(11): 2809-2812.
- Rønnestad, I., Yúfera, M., Ueberschär, B., Ribeiro, L., Sæle, Ø., and Boglione, C. (2013). Feeding behaviour and digestive physiology in larval fish: current knowledge, and gaps and bottlenecks in research. *Reviews in Aquaculture*, 5(1):59-68.

- Rurangwa, E., Laranja, J. L., Van Houdt, R., Delaedt, Y., Geraylou, Z., Van de Wiele, T., and Ollevier, F. (2009). Selected nondigestible carbohydrates and prebiotics support the group of probiotics *in vitro*. *Journal of applied microbiology*, 106(3): 932-940.
- Rutherford, S. T., and Bassler, B. L. (2012). Bacterial quorum sensing: its role in virulence and possibilities for its control. *Cold Spring Harbor perspectives in medicine*, 2(11):124-127.
- Ruwandepika, H. A., Defoirdt, T., Bhowmick, P. P., Karunasagar, I., Karunasagar, I., and Bossier, P. (2011). *In vitro* and *in vivo* expression of virulence genes in *Vibrio* isolates belonging to the *Harveyi* clade in relation to their virulence towards gnotobiotic brine shrimp (*Artemia franciscana*). *Environmental microbiology*, 13(2): 506-517.
- Ruwandepika, D., Arachchige, H., Sanjeewa Prasad Jayaweera, T., Paban Bhowmick, P., Karunasagar, I., Bossier, P., and Defoirdt, T. (2012). Pathogenesis, virulence factors and virulence regulation of vibrios belonging to the *Harveyi* clade. *Reviews in Aquaculture*, 4(2): 59-74.
- Salinas, I., Díaz-Rosales, P., Cuesta, A., Meseguer, J., Chabrilón, M., Morinigo, M. A., and Esteban, M. A. (2006). Effect of heat-inactivated fish and non-fish derived probiotics on the innate immune parameters of a teleost fish (*Sparus aurata* L.). *Veterinary immunology and immunopathology*, 111(3): 279-286.
- Salvesen I., Skjermo J., and Vadstein O. (1999). Growth of turbot (*Scophthalmus maximus* L.) during first feeding in relation to the fraction of r/K-strategists in the bacterial community of the rearing water. *Aquaculture*, 175:337-350.
- Schauder, S., and Bassler, B. L. (2001). The languages of bacteria. *Genes and Development*, 15(12): 1468-1480.
- Secombes, C. J., Wang, T., Hong, S., Peddie, S., Crampe, M., Laing, K. J., and Zou, J. (2001). Cytokines and innate immunity of fish. *Developmental & Comparative Immunology*, 25(8):713-723.
- Seenivasan, C., Bhavan, P. S., Radhakrishnan, S., & Shanthi, R. (2012). Enrichment of *Artemia* nauplii with *Lactobacillus sporogenes* for enhancing the survival, growth and levels of biochemical constituents in the post-larvae of the freshwater prawn *Macrobrachium rosenbergii*. *Turkish Journal of Fisheries and Aquatic Sciences*, 12(1): 202-208.
- Setlow, P., Johnson, E. A., Doyle, M. P., and Beuchat, L. R. (2007). Spores and their significance. *Food microbiology: fundamentals and frontiers*, 3:35-67.
- Sfanos, K., Harmody, D., Dang, P., Ledger, A., Pomponi, S., McCarthy, P., and Lopez, J. (2005). A molecular systematic survey of cultured microbial associates of deep-water marine invertebrates. *Systematic and applied microbiology*, 28(3): 242-264.

- Shirakashi, S., Hirano, C., Asmara, A. B., Noorashikin M.N., Ishimaru, K., and Miyashita, S. (2013). Shading Reduces *Neobenedenia girellae* Infection on Cultured Greater Amberjack *Seriola dumerili*. *Fish Pathology*, 48(1): 25-28.
- Showalter, R. E., Martin, M. O., & Silverman, M. R. (1990). Cloning and nucleotide sequence of luxR, a regulatory gene controlling bioluminescence in *Vibrio harveyi*. *Journal of bacteriology*, 172(6): 2946-2954.
- Skjermo J., Salvesen, I., Oie, G., Olsen, Y., and Vadstein, O. (1997). Microbially matured water: a technique for selection of a non-opportunistic bacterial flora in water that may improve performance of marine larvae. *Aquaculture International*, 5:13–28.
- Sletta, H., Degnes, K. F., Herfindal, L., Klinkenberg, G., Fjærvik, E., Zahlsen, K., and Zotchev, S. B. (2014). Anti-microbial and cytotoxic 1, 6-dihydroxyphenazine-5, 10-dioxide (iodinin) produced by *Streptosporangium* sp. DSM 45942 isolated from the fjord sediment. *Applied microbiology and biotechnology*, 98(2): 603-610.
- Smelt, J. P. P. M., Bos, A. P., Kort, R., and Brul, S. (2008). Modelling the effect of sub (lethal) heat treatment of *Bacillus subtilis* spores on germination rate and outgrowth to exponentially growing vegetative cells. *International journal of food microbiology*, 128(1): 34-40.
- Smith J.N. and Ahmer B.M. (2003). "Detection of other microbial species by *Salmonella*: expression of the SdiA regulon". *Jornal of Bacteriology*, 185 (4): 1357–1366.
- Son, V. M., Chang, C. C., Wu, M. C., Guu, Y. K., Chiu, C. H., and Cheng, W. (2009). Dietary administration of the probiotic, *Lactobacillus plantarum*, enhanced the growth, innate immune responses, and disease resistance of the grouper *Epinephelus coioides*. *Fish and Shellfish Immunology*, 26(5): 691-698.
- Sorgeloos, P., Dhert, P., and Candreva, P. (2001). Use of the brine shrimp *Artemia* spp., in marine fish larviculture. *Aquaculture*, 200(1): 147-159.
- Sorroza, L., Padilla, D., Acosta, F., Román, L., Grasso, V., Vega, J., and Real, F. (2012). Characterization of the probiotic strain *Vagococcus fluvialis* in the protection of European sea bass (*Dicentrarchus labrax*) against vibriosis by *Vibrio anguillarum*. *Veterinary microbiology*, 155(2): 369-373.
- Spanggaard B., Huber I., Nielsen J., Sick E.B., Pipper C.B., Martinussen, T., Slierendrecht, W.J., and Gram, L., (2001). The probiotic potential against vibriosis of the indigenous microflora of rainbow trout. *Environmental Microbiology*, 3(12): 755–765.
- Sreedevi, P. R., and Ramasubramanian, V. (2011). Biocontrol of ammonia pollution in the rearing water of fish by inducing a heterotrophic bacterial-based food chain in the medium. *Aquaculture International*, 19(4): 789-796.
- Stottrup, J. G. (2003). Production and nutritional value of copepods. *Live feeds in marine aquaculture*:145-205.

- Sun, Y. Z., Yang, H. L., Huang, K. P., Ye, J. D., and Zhang, C. X. (2013). Application of autochthonous *Bacillus* bioencapsulated in copepod to grouper *Epinephelus coioides* larvae. *Aquaculture*, 392: 44-50.
- Surette M.G, and Bassler B.L. (1999). Regulation of autoinducer production in *Salmonella typhimurium*. *Molecular Microbiology*, 31:585-95.
- Suzer, C., Çoban, D., Kamaci, H. O., Saka, Ş., Firat, K., Otgucuoğlu, Ö., and Küçüksari, H. (2008). *Lactobacillus* spp. bacteria as probiotics in gilthead sea bream (*Sparus aurata* L) larvae: Effects on growth performance and digestive enzyme activities. *Aquaculture*, 280(1):140-145.
- Swartzman, E., Silverman, M., & Meighen, E. A. (1992). The luxR gene product of *Vibrio harveyi* is a transcriptional activator of the lux promoter. *Journal of bacteriology*, 174(22): 7490-7493.
- Teasdale, M. E., Liu, J., Wallace, J., Akhlaghi, F., & Rowley, D. C. (2009). Secondary metabolites produced by the marine bacterium *Halobacillus salinus* that inhibit quorum sensing-controlled phenotypes in gram-negative bacteria. *Applied and environmental microbiology*, 75(3): 567-572.
- Tendencia, E. A., dela Peña, M. R., Fermin, A. C., Lio-Po, G., Choresca, C. H., and Inui, Y. (2004). Antibacterial activity of tilapia *Tilapia hornorum* against *Vibrio harveyi*. *Aquaculture*, 232(1):145-152.
- Tinh, N. T. N., Linh, N. D., Wood, T. K., Dierckens, K., Sorgeloos, P., & Bossier, P. (2007). Interference with the quorum sensing systems in a *Vibrio harveyi* strain alters the growth rate of gnotobiotically cultured rotifer *Brachionus plicatilis*. *Journal of applied microbiology*, 103(1):194-203.
- Tinh, N. T. N., Yen, V. H. N., Dierckens, K., Sorgeloos, P., & Bossier, P. (2008). An acyl homoserine lactone-degrading microbial community improves the survival of first-feeding turbot larvae (*Scophthalmus maximus* L.). *Aquaculture*, 285(1): 56-62.
- Torres, M., Romero, M., Prado, S., Dubert, J., Tahrioui, A., Otero, A., and Llamas, I. (2013). N-acyl homoserine lactone-degrading bacteria isolated from hatchery bivalve larval cultures. *Microbiological research*, 168(9):547-554.
- Touraki, M., Karamanlidou, G., Karavida, P., and Chrysi, K. (2012). Evaluation of the probiotics *Bacillus subtilis* and *Lactobacillus plantarum* bioencapsulated in *Artemia* nauplii against vibriosis in European seabass larvae. *World Journal of Microbiology and Biotechnology*, 28(6): 2425-2433.
- Vadstein, O., Øie, G., Olsen, Y., Salvesen, I., Skjermo, J., & Skjåk-Bræk, G. (1993). A strategy to obtain microbial control during larval development of marine fish. *Fish farming technology*: 69-75.
- Vadstein, O. (1997). The use of immunostimulation in marine larviculture: possibilities and challenges. *Aquaculture*, 155(1): 401-417.

- Vamsee-Krishna, C., Mohan, Y., and Phale, P. S. (2006). Biodegradation of phthalate isomers by *Pseudomonas aeruginosa* PP4, *Pseudomonas* sp. PPD and *Acinetobacter lwoffii* ISP4. *Applied microbiology and biotechnology*, 72(6): 1263-1269.
- van der Geize, R., and Dijkhuizen, L. (2004). Harnessing the catabolic diversity of rhodococci for environmental and biotechnological applications. *Current opinion in microbiology*, 7(3): 255-261.
- Vaseeharan, B. A. R. P., and Ramasamy, P. (2003). Control of pathogenic *Vibrio* spp. by *Bacillus subtilis* BT23, a possible probiotic treatment for black tiger shrimp *Penaeus monodon*. *Letters in applied microbiology*, 36(2): 83-87.
- Verschuere, L., Rombaut, G., Sorgeloos, P., & Verstraete, W. (2000). Probiotic bacteria as biological control agents in aquaculture. *Microbiology and molecular biology reviews*, 64(4): 655-671.
- Villamil, L., Figueras, A., Planas, M., & Novoa, B. (2003). Control of *Vibrio alginolyticus* in *Artemia* culture by treatment with bacterial probiotics. *Aquaculture*, 219(1): 43-56.
- Vine, N. G., Leukes, W. D., Kaiser, H., Daya, S., Baxter, J., & Hecht, T. (2004). Competition for attachment of aquaculture candidate probiotic and pathogenic bacteria on fish intestinal mucus. *Journal of fish diseases*, 27(6): 319-326.
- Vine, N. G., Leukes, W. D., & Kaiser, H. (2006). Probiotics in marine larviculture. *FEMS microbiology reviews*, 30(3): 404-427.
- Vinoj, G., Vaseeharan, B., Thomas, S., Spiers, A. J., and Shanthi, S. (2014). Quorum-quenching activity of the AHL-Lactonase from *Bacillus licheniformis* DAHB1 inhibits *Vibrio* biofilm formation in vitro and reduces shrimp intestinal colonisation and mortality. *Marine Biotechnology*, 16(6): 707-715.
- Vinoj, G., Jayakumar, R., Chen, J. C., Withyachumnarnkul, B., Shanthi, S., and Vaseeharan, B. (2015). N-hexanoyl-L-homoserine lactone-degrading *Pseudomonas aeruginosa* PsDAHP1 protects zebrafish against *Vibrio parahaemolyticus* infection. *Fish & shellfish immunology*, 42(1): 204-212.
- Wagner, V. E., Gillis, R. J., and Iglewski, B. H. (2004). Transcriptome analysis of quorum-sensing regulation and virulence factor expression in *Pseudomonas aeruginosa*. *Vaccine*, 22:S15-S20.
- Wang, L. T., Lee, F. L., Tai, C. J., and Kasai, H. (2007). Comparison of gyrB gene sequences, 16S rRNA gene sequences and DNA-DNA hybridization in the *Bacillus subtilis* group. *International Journal of Systematic and Evolutionary Microbiology*, 57(8): 1846-1850.
- Waters, C. M., and Bassler, B. L. (2005). Quorum sensing: cell-to-cell communication in bacteria. *Annual Review of Cell Division in Biology*, 21: 319-346.

- White, C. E., and Finan, T. M. (2009). Quorum quenching in *Agrobacterium tumefaciens*: chance or necessity?. *Journal of bacteriology*, 191(4): 1123-1125.
- Winkaler, E. U., Silva, A. D. G., Galindo, H. C., and Martine, C. R. (2001). Histological and physiological biomarkers to assess fish health in Londrina streams, Parana State. *Acta scientiarum. Maringa*, 23(2): 507-514.
- Wu L., Estrada O., Zaborina O., Bains M., Shen L., Kohler J.E. (2005). Recognition of host immune activation by *Pseudomonas aeruginosa*. *Science* , 309:774–7777.
- Xiang, L. X., He, D., Dong, W. R., Zhang, Y. W., & Shao, J. Z. (2010). Deep sequencing-based transcriptome profiling analysis of bacteria-challenged *Lateolabrax japonicus* reveals insight into the immune-relevant genes in marine fish. *BMC genomics*, 11(1): 472.
- Yamaoka, K., Nanbu, T., Miyagawa, M., Isshiki, T., and Kusaka, A. (2000). Water surface tension-related deaths in prelarval red-spotted grouper. *Aquaculture*, 189(1): 165-176.
- Yang, Q., and Defoirdt, T. (2014). Quorum sensing positively regulates flagellar motility in pathogenic *Vibrio harveyi*. *Environmental microbiology*, 12:24-30.
- Yashiro, R. (2008). Overview of grouper aquaculture in Thailand. *The Aquaculture of Grouper*. High Point Press, Taiwan, 143-154.
- Yates, E. A., Philipp, B., Buckley, C., Atkinson, S., Chhabra, S. R., Sockett, R. E., and Williams, P. (2002). N-acylhomoserine lactones undergo lactonolysis in a pH-, temperature-, and acyl chain length-dependent manner during growth of *Yersinia pseudotuberculosis* and *Pseudomonas aeruginosa*. *Infection and immunity*, 70(10): 5635-5646.
- Ye, J., Ma, Y., Liu, Q., Zhao, D. L., Wang, Q. Y., and Zhang, Y. X. (2008). Regulation of *Vibrio alginolyticus* virulence by the LuxS quorum-sensing system. *Journal of fish diseases*, 31(3):161-169.
- Yeh, R. Y., Shiu, Y. L., Shei, S. C., Cheng, S. C., Huang, S. Y., Lin, J. C., & Liu, C. H. (2009). Evaluation of the antibacterial activity of leaf and twig extracts of stout camphor tree, *Cinnamomum kanehirae*, and the effects on immunity and disease resistance of white shrimp, *Litopenaeus vannamei*. *Fish & shellfish immunology*, 27(1):26-32.
- Yii, K. C., Yang, T. I., & Lee, K. K. (1997). Isolation and characterization of *Vibrio carchariae*, a causative agent of gastroenteritis in the groupers, *Epinephelus coioides*. *Current Microbiology*, 35(2): 109-115.
- Zambonino-Infante, J. L., Koumoundouros, G., and Tandler, A. (2009). The influence on nutrition at the larval stages in marine fish. *Control of Malformations in Fish Aquaculture: Science and Practice*, 85-92.

- Zhang, P., Setlow, P., and Li, Y. (2009). Characterization of single heat-activated *Bacillus* spores using laser tweezers Raman spectroscopy. *Optics express*, 17(19):16480-16491.
- Zhao, J., Jiang, H., Cheng, W., Wu, J., Zhao, J., Wang, J., and Dong, L. (2014). The role of quorum sensing system in antimicrobial induced ampC expression in *Pseudomonas aeruginosa* biofilm. *Journal of basic microbiology*, 5:800-807.
- Zhonghua, C., Chunpin, G., Yong, Z., Kezhi, X., & Yaou, Z. (2008). Cloning and bioactivity analysis of a CXC ligand in black seabream *Acanthopagrus schlegeli*: the evolutionary clues of ELR+ CXC chemokines. *BMC immunology*, 9(1): 66.
- Zhou, Q. C., Wu, Z. H., Chi, S. Y., & Yang, Q. H. (2007). Dietary lysine requirement of juvenile cobia (*Rachycentron canadum*). *Aquaculture*, 273(4): 634-640.
- Zhou, Q., Li, K., Jun, X., & Bo, L. (2009). Role and functions of beneficial microorganisms in sustainable aquaculture. *Bioresource technology*, 100(16): 3780-3786.
- Ziaei-Nejad, S., Rezaei, M. H., Takami, G. A., Lovett, D. L., Mirvaghefi, A. R., and Shakouri, M. (2006). The effect of *Bacillus* spp. bacteria used as probiotics on digestive enzyme activity, survival and growth in the Indian white shrimp *Fenneropenaeus indicus*. *Aquaculture*, 252(2):516-524.
- Ziemer, C. J., & Gibson, G. R. (1998). An overview of probiotics, prebiotics and synbiotics in the functional food concept: perspectives and future strategies. *International Dairy Journal*, 8(5): 473-479.