

# ARC '18

مؤتمر مؤسسة قطر  
السنوي للبحوث

QATAR FOUNDATION  
ANNUAL RESEARCH CONFERENCE

البحث والتطوير: التركيز على  
الأولويات، وإحداث الأثر

R&D: FOCUSING ON PRIORITIES,  
DELIVERING IMPACT

20-19 مارس  
19-20 MARCH



مؤسسة قطر  
Qatar Foundation

إطلاق قدرات الإنسان.  
Unlocking human potential.

## Health and Biomedical - Poster Display

<http://doi.org/10.5339/qfarc.2018.HBPD390>

# DOES WEIGHT LOSS THROUGH MEANS OF BARIATRIC SURGERY REDUCE THE RISK OF TYPE 2 DIABETES IN OBESE QATARI PATIENT A RETROSPECTIVE ANALYSIS

Fahad Hanna\*, Montaha Ghallab Mohammed, Rahma Ahmad Saad,  
Anwar Abdulkarem Al-hwsali, Manar Mamdoh Dalal, Mohamed Aly El-Sherif

Qatar University  
\* fhanna@qu.edu.qa


**Background:** The use of bariatric surgeries such as Gastric Bypass and Sleeve Gastrectomy in managing obesity and associated diseases such as type 2 diabetes mellitus (T2DM) has been induced in clinical practice. Weight reduction through means of bariatric surgery has metabolic benefits and may improve the management of T2DM. **Objectives:** The aim of this study was to investigate if weight loss through bariatric surgery can reduce the risk of T2DM in patients without the onset of T2DM. **Study design:** A retrospective analysis was conducted on post-bariatric patients at the department of bariatric and metabolic surgery at Hamad General Hospital. **Methods:** Two hundred and two eligible pre-diabetic Qatari patients who have undergone bariatric surgery in 2016 and satisfied the inclusion and exclusion criteria of the study were analyzed. Data on Glucose, Insulin and C-peptides levels at baseline and follow-up were extracted in order to compare the change of these variables at baseline, 6 and 10 months follow up before and after 10 months from the date of surgery. **Results:** Seventy one males with mean age of  $32.73 \pm 10.37$  and one hundred and thirty one females with mean age of  $33.90 \pm 9.88$  were included in the analysis. Change in weight was strongly and positively associated with change in insulin level (0.701, 95% CI: 0.027, 1.347,  $p=0.042$ ) also, as weight changes fasting glucose changes (1.993, 95% CI: 0.359, 3.627,  $p=0.017$ ). Follow-up period greater than 6 months was not found to be significantly associated with weight loss (2.049, 95% CI: -2.249, 6.349,  $p=0.313$ ). **Conclusion:** Our study confirms results from international studies that weight

© 2018 The Author(s), licensee HBKU Press. This is an open access article distributed under the terms of the Creative Commons Attribution license CC BY 4.0, which permits unrestricted use, distribution and reproduction in any medium, provided the original work is properly cited.

دار جامعة حمد بن خليفة للنشر  
HAMD BIN KHALIFA UNIVERSITY PRESS 

Cite this article as: Hanna F et al. (2018). DOES WEIGHT LOSS THROUGH MEANS OF BARIATRIC SURGERY REDUCE THE RISK OF TYPE 2 DIABETES IN OBESE QATARI PATIENT A RETROSPECTIVE ANALYSIS. Qatar Foundation Annual Research Conference Proceedings 2018: HBPD390  
<http://doi.org/10.5339/qfarc.2018.HBPD390>.

The statements and opinions contained in the publication are solely those of the individual authors and do not necessarily reflect those of the publisher. HBKU Press does not accept any legal responsibility or liability for any errors or omissions that may be made.



loss through bariatric surgery can reduce the risk of developing Type 2 Diabetes Qatari obese patients. The results of the study suggest that post-surgery periods can be detrimental to the fate of fasting glucose and insulin levels and therefore compliance maybe of great importance to ensure success and sustainability of weight loss and diabetes prevention. Larger samples size and longer follow-up period is required to confirm these findings.