

Contents lists available at ScienceDirect

Chemosphere

journal homepage: www.elsevier.com/locate/chemosphere

Trace elements in loggerhead turtles (*Caretta caretta*) stranded in mainland Portugal: Bioaccumulation and tissue distribution



Lídia Nicolau ^{a, b, *}, Sílvia S. Monteiro ^{a, b}, Andreia T. Pereira ^b, Ana Marçalo ^{a, b},
Marisa Ferreira ^{b, c}, Jordi Torres ^{d, e}, José Vingada ^{b, c, f}, Catarina Eira ^{a, b}

^a Departamento de Biologia & Centro de Estudos do Ambiente e do Mar (CESAM), Universidade de Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

^b Sociedade Portuguesa de Vida Selvagem, Universidade do Minho, Departamento de Biologia, Campus de Gualtar, 4710-057, Braga, Portugal

^c Departamento de Biologia & Centro Biologia Molecular e Ambiental (CBMA), Universidade de Minho, Campus de Gualtar, 4710-047 Braga, Portugal

^d Departament de Biologia, Sanitat i Medi ambient, Facultat de Farmàcia i Ciències de l'Alimentació, Universitat de Barcelona, Av. Joan XXIII, sn, 08028 Barcelona, Spain

^e Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona, Av. Joan XXIII, sn, 08028 Barcelona, Spain

^f Centro de Estudos do Ambiente e do Mar (CESAM), Campus Universitário de Santiago, 3810-193 Aveiro, Portugal

HIGHLIGHTS

- High levels of cadmium in loggerheads stranded in mainland Portugal.
- Evidence of bioaccumulation of renal cadmium.
- Cadmium-Zinc correlations were observed in liver and kidney tissues.
- The 3 largest loggerheads showed lower Cd concentrations than smaller turtles.

ARTICLE INFO

Article history:

Received 5 January 2017

Received in revised form

13 March 2017

Accepted 26 March 2017

Available online 27 March 2017

Handling Editor: Martine Leermakers

Keywords:

Bioaccumulation

Cadmium

Marine turtle

Northeast Atlantic

ABSTRACT

Pollution is among the most significant threats that endanger sea turtles worldwide. Waters off the Portuguese mainland are acknowledged as important feeding grounds for juvenile loggerheads. However, there is no data on trace element concentrations in marine turtles occurring in these waters. We present the first assessment of trace element concentrations in loggerhead turtles (*Caretta caretta*) occurring off the coast of mainland Portugal. Also, we compare our results with those from other areas and discuss parameters that may affect element concentrations. Trace element concentrations (As, Cd, Cu, Pb, Mn, Hg, Ni, Se, Zn) were determined in kidney, liver and muscle samples from 38 loggerheads stranded between 2011 and 2013. As was the only element with higher concentrations in muscle ($14.78 \mu\text{g g}^{-1}$ ww) than in liver or kidney. Considering non-essential elements, Cd presented the highest concentrations in kidney ($34.67 \mu\text{g g}^{-1}$) and liver ($5.03 \mu\text{g g}^{-1}$). Only a weak positive link was found between renal Cd and turtle size. Inter-elemental correlations were observed in both liver and kidney tissues. Hepatic Hg values ($0.30 \pm 0.03 \mu\text{g g}^{-1}$) were higher than values reported in loggerheads in the Canary Islands but lower than in Mediterranean loggerheads. Cd concentrations in the present study were only exceeded by values found in turtles from the Pacific. Although many endogenous and exogenous parameters related with complex life cycle changes and wide geographic range may influence trace element accumulation, the concentrations of Cd are probably related to the importance of crustaceans in loggerhead diet in the Portuguese coast.

© 2017 Elsevier Ltd. All rights reserved.

* Corresponding author. Departamento de Biologia & Centro de Estudos do Ambiente e do Mar (CESAM), Universidade de Aveiro, Campus Universitário de Santiago, 3810-193 Aveiro, Portugal.

E-mail address: lvnicolau@ua.pt (L. Nicolau).

1. Introduction

Along with other large marine vertebrates, marine turtles are known as key animal groups acting as sentinels of environmental disturbances, reflecting natural and anthropogenic threats on a

wider portion of the marine ecosystem (McAlouse and Newton, 2009; Fossi et al., 2012). According to the IUCN Marine Turtle Specialist Group, the most significant threats that endanger sea turtles include direct take, coastal development, global warming, fisheries impacts, pathogens and pollution (Mast et al., 2005).

Evaluating the effects of anthropogenic factors is presently a top global research priority for marine turtle conservation, including the impacts of pollution on marine turtles (Hamann et al., 2010). In fact, pollution has already been associated to the decline of some sea turtle populations (Gordon et al., 1998; Godley et al., 1999; Sakai et al., 2000). Sublethal effects caused by several pollutants, including trace elements, include impaired reproductive success (Perrault et al., 2011) and also deterioration of the immune system and other physiological functions in marine turtles (Aguirre et al., 1994; Day et al., 2007; Komoroske et al., 2011; Camacho et al., 2013).

Several studies reported the concentrations of trace elements in loggerhead turtles (*Caretta caretta*) in different areas of its wide distribution range (e.g. Gordon et al., 1998; Caurant et al., 1999; Godley et al., 1999; Saeki et al., 2000; Sakai et al., 2000; Storelli et al., 1998, 2005; Storelli and Marcotrigiano, 2003; Franzellitti et al., 2004; Torrent et al., 2004; Day et al., 2005; Maffucci et al., 2005; Andreani et al., 2008; García-Fernández et al., 2009; Jerez et al., 2010; D'Ilio et al., 2011; Camacho et al., 2013; López-Castro et al., 2013). These studies revealed how several ecological and biological variables (age, sex, diet or migration areas) influence trace element concentrations in loggerhead turtles (Caurant et al., 1999; Jerez et al., 2010; López-Castro et al., 2013).

In mainland Portugal, the analysis of a relatively high number of stranded marine turtles revealed that loggerhead turtles (242 loggerheads stranded in the 2009–2013 period) clearly include the Portuguese mainland coast in their oceanic pathways (Nicolau et al., 2016). The loggerhead turtle is currently categorized as “Vulnerable” by the IUCN (Casale and Tucker, 2015). Considering the loggerhead turtles Regional Management Units (RMUs, Wallace et al., 2010), the Atlantic Northwest and the Mediterranean RMU overlap along the Portuguese mainland coast. These sub-populations have recently been re-categorized by the IUCN as Least Concern (Ceriani and Meylan, 2015; Casale, 2015) although they represent Low Risk–High Threat units (Wallace et al., 2011), indicating the need to mitigate threats that could lead to abundance declining in the future.

Despite the extensive literature regarding the concentrations of contaminants in loggerhead turtles, there is currently no information on trace element concentrations in marine turtles occurring off the coast of mainland Portugal. This is particularly relevant considering that the southern Portuguese coastal region represents an important hotspot for the neritic loggerhead turtle juveniles (Nicolau et al., 2016), where they may be exposed to contaminant-enriched coastal waters (van Geen et al., 1991; Mil-Homens et al., 2014).

Considering the importance of pollution as a threat to sea turtles, the present study aims at providing the first assessment of trace element concentrations in loggerhead turtles stranded in the Portuguese mainland coast, and relate them to concentrations reported in loggerheads elsewhere in the world.

2. Materials and methods

2.1. Sample collection

Loggerhead turtles stranded in the Portuguese mainland coast are routinely investigated (for example, following an alert given by the Maritime Authority) by experienced personnel belonging to the Portuguese stranding network, coordinated by the Institute for

Nature Conservation and Forests (ICNF) and the Portuguese Wildlife Society (SPVS). Detailed necropsies were performed, where biometric data and samples were collected according to standard protocols (Wyneken, 2001). Kidney, liver and muscle samples were collected from 38 loggerhead turtles stranded between 2011 and 2013 (Fig. 1). Samples of loggerhead turtles were stored in glass vials and frozen (-20°C) for posterior trace element analysis.

Curved carapace length (CCL) was registered ($n = 38$; CCL mean = 50.1 cm; CCL SD = 9.2 cm; CCL range = 35.5–75.5) and used as a proxy for age. Considering the minimum size of nesting females for the western North Atlantic stocks (CCL = 87.2 cm; TEWG, 2009), all specimens were identified as juveniles or sub-adults. It is also generally accepted that loggerheads from the North Atlantic presenting a CCL between 46 and 64 cm transit from oceanic waters to neritic habitats (Bjorndal et al., 2000).

2.2. Analytical procedure

Approximately 100–150 mg (wet weight, ww) of kidney, liver and muscle of loggerhead turtles were digested in teflon vessels with 2 mL of HNO_3 and 1 mL of H_2O_2 (Merck, Suprapure). Acid digestion of the samples was performed in a drying oven at 90°C , overnight (14 h). All materials used in the digestion process were




Fig. 1. Stranding locations of loggerhead sea turtles used in the present study.

Table 1
Trace element concentrations (mean \pm SE, $\mu\text{g g}^{-1}$ ww) in different tissues of loggerhead turtles (*Caretta caretta*) stranded in mainland Portugal. Concentrations are provided for the total number of animals and separately for oceanic juveniles (CCL < 46 cm), individuals transitioning from oceanic to neritic areas (CCL = 46–64 cm) and neritic subadults (CCL > 64 cm), according to Bjorndal et al. (2000).

	Liver				Kidney				Muscle			
	Total	<46	46–64	>64	Total	<46	46–64	>64	Total	<46	46–64	>64
Zn	24.01 \pm 0.94	23.85 \pm 1.36	23.84 \pm 1.28	25.88 \pm 5.18	30.50 \pm 1.49	30.35 \pm 2.42	31.23 \pm 2.10	25.49 \pm 2.66	19.79 \pm 0.82	18.62 \pm 0.92	19.77 \pm 1.19	24.64 \pm 1.73
Mn	1.78 \pm 0.09	1.99 \pm 0.18	1.71 \pm 0.12	1.39 \pm 0.13	2.09 \pm 0.14	2.26 \pm 0.28	2.05 \pm 0.18	1.70 \pm 0.50	0.14 \pm 0.01	0.14 \pm 0.02	0.14 \pm 0.01	0.12 \pm 0.02
Pb	0.10 \pm 0.01	0.08 \pm 0.01	0.12 \pm 0.02	0.08 \pm 0.03	0.23 \pm 0.03	0.22 \pm 0.06	0.26 \pm 0.04	0.11 \pm 0.03	0.01 \pm 0.00	0.01 \pm 0.00	0.01 \pm 0.00	0.03 \pm 0.02
Cd	5.03 \pm 0.54	3.97 \pm 0.50	5.99 \pm 0.78	1.86 \pm 0.37	34.67 \pm 3.21	28.78 \pm 4.37	41.06 \pm 4.13	9.22 \pm 0.72	0.16 \pm 0.01	0.17 \pm 0.02	0.18 \pm 0.01	0.03 \pm 0.00
As	4.49 \pm 0.35	4.87 \pm 0.69	4.13 \pm 0.45	5.71 \pm 0.49	6.28 \pm 0.72	6.69 \pm 1.52	5.61 \pm 0.82	9.83 \pm 2.52	14.78 \pm 1.47	13.99 \pm 2.46	14.68 \pm 2.03	18.66 \pm 4.15
Cu	5.99 \pm 0.48	4.94 \pm 0.71	6.70 \pm 0.67	4.72 \pm 0.24	1.73 \pm 0.09	1.81 \pm 0.13	1.75 \pm 0.12	1.25 \pm 0.24	0.55 \pm 0.04	0.49 \pm 0.06	0.58 \pm 0.06	0.59 \pm 0.03
Hg	0.30 \pm 0.03	0.30 \pm 0.03	0.30 \pm 0.03	0.34 \pm 0.22	0.21 \pm 0.02	0.21 \pm 0.03	0.21 \pm 0.03	0.14 \pm 0.09	0.05 \pm 0.01	0.03 \pm 0.01	0.06 \pm 0.00	0.05 \pm 0.04
Ni	0.14 \pm 0.02	0.14 \pm 0.03	0.14 \pm 0.03	0.06 \pm 0.01	0.44 \pm 0.04	0.42 \pm 0.04	0.48 \pm 0.06	0.19 \pm 0.08	0.08 \pm 0.03	0.04 \pm 0.01	0.10 \pm 0.05	0.05 \pm 0.03
Se	5.23 \pm 0.20	5.07 \pm 0.30	5.28 \pm 0.26	5.46 \pm 1.23	4.92 \pm 0.29	4.87 \pm 0.56	4.90 \pm 0.37	5.35 \pm 1.18	2.34 \pm 0.09	2.24 \pm 0.13	2.19 \pm 0.11	3.06 \pm 0.53

thoroughly acid-rinsed. After digestion, samples were diluted with ultrapure water and analyzed for nine trace elements [arsenic (As), cadmium (Cd), copper (Cu), lead (Pb), manganese (Mn), mercury (Hg), nickel (Ni), selenium (Se), zinc (Zn)], by ICP-MS (Perkin Elmer Elan 6000). To determine analytical accuracy, several blanks and standard reference material (*Squalus acanthias* - Dogfish liver (DOLT-3) and muscle (DORM-2)) (National Research Council, Canada) were prepared and analyzed along with samples. All of the analyzed trace elements exhibited concentrations above the detection limits of the analytical instruments, except for Hg in muscle samples of some individuals, which were considered missing values. Trace element concentrations are reported in $\mu\text{g g}^{-1}$, based on wet weight values (ww).

2.3. Statistical procedure

All data series were explored for outliers, collinearity, heterogeneity of variance and for visualization of potential relationships between response and explanatory variables, following Zuur et al. (2010).

Linear Models were used to determine the effect of curved carapace length (CCL) on the concentrations of trace elements in loggerhead turtles. Since the concentration values of each trace element were continuous, a Gaussian distribution was applied. Validation of the final model involved checking the assumptions of normality, homogeneity and independence of residuals, together with the lack of highly influential data points (“hat” values), through the use of plots involving residuals against fitted values and Q–Q plots, among others (Zuur et al., 2007). Assumptions of normality and homogeneity were also tested using the Shapiro–Wilk (Royston and Remark, 1995) and Breusch–Pagan (Breusch and Pagan, 1979) tests, respectively. Model validation decisions were based on both approaches. Turtles with CCL >64 cm ($n = 3$) were excluded from this analysis, due to the low number of samples in this size category.

In order to investigate the potential inter-elemental correlations, the Pearson correlation coefficient was used and the assumptions of normality and homogeneity of the residuals were checked, as described above for validation of the linear models.

Separate analyses were performed for each tissue type used in




Fig. 2. Significant Pearson correlation coefficients (r) between several trace elements in kidney (a) and liver (b) of loggerhead turtles (*Caretta caretta*) stranded in mainland Portugal. Only correlations yielding an $r > 0.50$ are shown.




Fig. 3. Non-essential trace element concentrations in liver (a), kidney (b) and muscle (c) of loggerhead turtles stranded in mainland Portugal, plotted against curved carapace length (CCL, in cm). Fitted regression line is included. The vertical dash and solid lines refer to the CCL limits for oceanic juveniles (CCL < 46 cm), individuals transitioning from oceanic to neritic areas (CCL = 46–64 cm) and neritic sub-adults (CCL > 64 cm), respectively, described by Bjorndal et al. (2000).

the present study. Statistical tests were performed in R v.3.2.3 (R Core Team, 2015).

3. Results

Trace element concentrations detected in the different loggerhead turtle tissues are presented in Table 1. Arsenic is the only element that shows higher concentrations in muscle ($14.78 \mu\text{g g}^{-1}$, $\text{SE} = 1.47$) compared to liver ($4.49 \mu\text{g g}^{-1}$, $\text{SE} = 0.35$) or kidney ($6.28 \mu\text{g g}^{-1}$, $\text{SE} = 0.72$) (Table 1). In turn, Cu, Hg and Se presented higher mean concentrations in hepatic samples, while Cd, Mn, Pb, Ni and Zn showed higher values in renal samples. Considering non-essential elements, Cd presented the highest concentrations in liver and kidney tissues (Table 1).

Several inter-elemental correlations were observed in both liver and kidney tissues of loggerhead turtles, as described in Fig. 2. In particular, a correlation between Cd and Zn was observed in both tissues. In renal samples, Zn concentrations also correlated with Pb and Cu, whereas Cu concentrations correlated with Mn concentrations. No strong correlations were observed in muscle samples.

Linear models revealed no relationship between any of the trace elements analyzed in the present study and turtle CCL, with the exception of a weak positive link between CCL and renal Cd (t -test = 2.109, p -value = 0.043, $r^2 = 0.119$). Fig. 3 shows turtle CCL plotted against the non-essential elements As, Cd and Hg and, in fact, no specific trends are detected except for renal Cd. Also, turtles with a CCL superior to 64 cm (excluded from linear models, see section describing the Statistical procedure) apparently show lower toxic element concentrations in comparison to smaller turtles, except for As.

4. Discussion

In general, the loggerhead turtles stranded in the Portuguese mainland presented trace element concentrations within the range of values reported in the Atlantic and Pacific oceans (Table 2). As for one of the most concerning toxic elements in the marine environment, Hg values in hepatic tissues obtained in the present study were lower than concentrations reported in loggerhead turtles from the acknowledged Hg-elevated Mediterranean basin (Storelli et al., 2005). However, Hg values in the present study were higher than values reported in loggerhead turtles in the Canary Islands (Torrent et al., 2004) (Table 2). Even though Hg levels in the analyzed tissues do not seem to be a cause of concern, according to

Day et al. (2007) relatively low levels of Hg may affect health parameters in loggerhead sea turtles, including their immune function.

Relatively higher concentrations of renal Cd were also found in the present study ($34.67 \pm 3.21 \mu\text{g g}^{-1}$ ww) compared to animals from the Atlantic coast of France and Spain (Caurant et al., 1999; Torrent et al., 2004), as well as the Adriatic sea and the Alboran sea (Storelli et al., 2005; García-Fernández et al., 2009). In fact, Cd concentrations in the present study were only exceeded by values found in turtles from the Pacific (Sakai et al., 1995; Gordon et al., 1998) (Table 2).

Knowledge about the actual toxic effects of many contaminants in higher marine vertebrates remains scarce and mostly based on values from laboratory experiments or surrogate species. Nonetheless, Cd is a non-essential element known to correlate with several health markers (Komoroske et al., 2011). Cd induces negative effects on essential elements metabolism and on endocrine and renal functions, being teratogenic and carcinogenic (Hopkins et al., 1999; Noel et al., 2004; Kitana and Callard, 2008; Ikonmopoulou et al., 2009; Simoniello et al., 2011). No toxicological data have been published specifically for marine turtles describing threshold concentrations of Cd above which detrimental effects would likely occur. Nevertheless, the renal concentrations of Cd found in the present study exceeded the toxic threshold previously suggested for evidence of Cd effects on vertebrates ($10 \mu\text{g g}^{-1}$ fresh weight, Eisler, 1985). It is noteworthy, however, that Torrent et al. (2004) reported seven turtles with Cd values between 20 and $60 \mu\text{g g}^{-1}$ (ww) with no evidence of renal lesions.

Arsenic is the only element presenting higher concentrations in muscular tissue than in hepatic or renal tissues (Table 1). The highest As accumulation in muscle is in agreement with previous studies on loggerhead turtles (Saeki et al., 2000; Storelli et al., 1998; Storelli and Marcotrigiano, 2003). On the other hand, Torrent et al. (2004) revealed a higher As concentration in liver, where its toxicity could cause liver damage, and proposed that across their life cycle, changes in marine turtle feeding habitats and type of available prey in these habitats, may be responsible for the variation of As accumulation patterns.

Several marine vertebrates, including marine turtles, have developed detoxification strategies to mitigate the toxic effects of non-essential elements. For example, metallothioneins (MTs) are metal-binding proteins implicated in the detoxification of toxic trace elements such as Cd, Hg, Ag and Pb (Roesijadi, 1992; Das et al., 2000; Anan et al., 2002) and in the homeostasis of essential

Table 2
Mean trace element concentrations ($\mu\text{g g}^{-1}$ w.w.) of loggerhead turtles (*Caretta caretta*) available in literature.

Ref.	Location	Liver					Kidney					Muscle				
		Pb	Cd	As	Hg	Se	Pb	Cd	As	Hg	Se	Pb	Cd	As	Hg	Se
Atlantic																
(1)	Portugal mainland	0.10	5.03	4.49	0.30	5.23	0.23	34.67	6.28	0.21	4.92	0.01	0.16	14.78	0.05	2.34
(2)	Canary Islands	2.94	2.53	17.07	0.04		2.44	5.01	13.80	0.04		2.26	1.14	7.35		
(3)	Atlantic – France		2.58					13.3					0.08			
(4)	South Carolina				0.594					0.214					0.155	
(5)	Brasil south				0.74									0.23		
Mediterranean																
(6)	Andalusia	0.69	5.85				0.17	10.49				0.05	0.04			
(7)	Adriatic + Ionian sea	0.16	3.36		0.43	3.54	0.12	8.35		0.16	2.20	0.04	0.07		0.18	1.65
(8)	Adriatic (Italy)		2.84										0.36			
Pacific																
(9)	Japan		9.29		1.51			39.40		0.25				0.06		0.11
(10)	Queensland (Australia)		16.4	0.46	0.015	2.21		28.30	0.71	0.045	1.52					

(1). Present study; (2). [Torrent et al., 2004](#); (3). [Caurant et al., 1999](#); (4). [Day et al., 2005](#); (5). [Soto et al., 2005](#); (6). [García-Fernández et al., 2009](#); (7). [Storelli et al., 2005](#); (8). [Franzellitti et al., 2004](#); (9). [Sakai et al., 1995](#); (10). [Gordon et al., 1998](#).

elements (e.g. Cu, Zn, [Roesijadi, 1992](#); [Anan et al., 2002](#)). Zinc is important in the regulation of MTs gene expression and, at least in mammals, MTs seem to occur more predominantly as ZnMT (reviewed in [Sakulsak, 2012](#)). A similar process in turtles might explain the strong correlations found in the present study between Zn-Cd, Zn-Pb and Zn-Cu. In fact, elements such as Cd, Pb and Cu show a higher binding affinity to MTs than Zn, being more capable of displacing Zn through a metal-metal exchange reaction, forming more stable MT complexes than other elements ([Waalkes et al., 1984](#); [Hamer, 1986](#); [Sabolic et al., 2010](#); [Sakulsak, 2012](#)). Similar correlations have already been reported in other marine turtles ([Sakai et al., 2000](#); [Maffucci et al., 2005](#); [Camacho et al., 2013](#)), marine mammals ([Méndez-Fernández et al., 2014](#)) and seabirds ([Mendes et al., 2008](#); [Ribeiro et al., 2009](#)).

The concentrations of pollutants in loggerhead turtles may result from endogenous (e.g. sex, age) or exogenous factors (diet, prevailing environmental conditions), or from a combination of both ([Aguilar et al., 1999](#); [Caurant et al., 1999](#)). Diet of loggerhead turtles stranded in mainland Portugal is mostly constituted by crustaceans (55.8% by number; 36.5% by weight) ([Nicolau, 2016](#)). Similar results were reported in other locations in the distribution range of the loggerhead Atlantic Northwest RMU, such as the West Atlantic nesting areas ([Seney and Musick, 2007](#); [Wallace et al., 2009](#)). Crustaceans are known to be an important source of essential elements (e.g. Se and Zn), but also non-essential elements such as Cd ([Turoczy et al., 2001](#); [Nuñez-Nogueira and Rainbow, 2005](#); [Karouna-Renier et al., 2007](#); [Barrento et al., 2008](#); [Reed et al., 2010](#); [Maulvault et al., 2011](#)). Therefore, the concentrations of Cd found in the present study may be associated to the dietary preferences of turtles in the Portuguese mainland coast.

A reduced importance of fish (recognized Hg vectors, e.g. [Chouvelon et al., 2012](#)) was reported in loggerhead turtle diet in Portuguese waters ([Nicolau, 2016](#)). The low fish intake may explain the lower Hg concentrations in the present study in comparison to other large vertebrate species in the same region ([Mendes et al., 2008](#); [Ribeiro et al., 2009](#); [Ferreira et al., 2016](#); [Monteiro et al., 2016a,b](#)). Also, no Hg-Se correlation was detected in the present study. The non-critical concentrations of Hg were probably insufficient to trigger the formation of mercury and selenium complexes (HgSe) in hepatic tissues described as a detoxification Hg strategy in marine turtles and mammals ([Storelli et al., 1998](#); [Jerez et al., 2010](#); [Frouin et al., 2012](#); [Lailson-Brito et al., 2012](#)).

The chronic exposure to non-essential trace elements in the marine environment associated to a limited capacity of excretion may lead to their bioaccumulation in long lived predators such as

cetaceans ([Ferreira et al., 2016](#); [Monteiro et al., 2016a,b](#); [2017](#)), seabirds ([Mendes et al., 2008](#); [Ribeiro et al., 2009](#)) and turtles ([Jerez et al., 2010](#); [Komoroske et al., 2011](#)). A positive, yet weak, link between renal Cd concentrations and turtle CCL was detected in the present study (excluding the three largest individuals from the sample, due to low sample size in that category). Considering the animals' CCL and previous stomach content analysis, which revealed the presence of both oceanic and neritic prey species ([Nicolau, 2016](#)), all animals in the present study were immature individuals inhabiting either oceanic waters (CCL < 46 cm) or a mix between oceanic and neritic animals (46 cm < CCL < 64 cm). Even the 3 largest loggerheads in our sample (CCL > 64 cm) were not yet mature individuals (considering the minimum size of nesting females for the western North Atlantic stocks; [TEWG, 2009](#)). Also, according to [Mansfield and Putman \(2013\)](#) both sub-adult and adult stage individuals probably undergo reversible oceanic-neritic transitions. As such, apart from the importance of crustaceans in the loggerhead turtle diet in the present study, the enhanced Cd concentrations may result from: 1) natural sources, due to the Cd enrichment in deeper waters due to biogeochemical processes ([Bruland et al., 1978](#); [Wu and Roshan, 2015](#)), in the case of oceanic immature turtles, or to a lesser extent 2) anthropogenic sources, due to non-essential elements enrichment (e.g. Cd, Hg) in coastal waters owing to the proximity to contaminant sources ([Delgado et al., 2011](#); [Mil-Homens et al., 2014](#)) in the case of transitioning turtles from oceanic to neritic habitats.

It is noteworthy that strong negative correlations were reported in previous studies between turtle CCL and non-essential elements such as Cd or Hg, which have been attributed to changes in physiological processes or in habitat use across life stages ([Storelli et al., 1998](#); [Sakai et al., 2000](#); [Komoroske et al., 2011](#)). In the present study, the 3 individuals with CCL > 64 cm generally showed lower concentrations of most trace elements than the remaining animals. Some authors argued that the negative correlations between some trace elements and turtle CCL may be due to the ontogenetic shifts in dietary preferences associated with the transition from oceanic to neritic habitats ([Sakai et al., 2000](#)). If that was the case, higher concentrations of Hg and Cd would have been expected in larger turtles in the present study since those turtles ingest relatively more fish and cephalopods ([Nicolau, 2016](#)), which are known vectors of Hg and Cd, respectively ([Bustamante et al., 1998](#); [Chouvelon et al., 2012](#)). Hence, the observed patterns in larger turtles may instead relate with trace elements biotransformation or elimination processes, including physiological aptitudes dependent on turtle maturity or exposure over time, such as up-regulation of

metallothionein or dose-dependent assimilation changes (Davis and Cousins, 2000; Guirlet and Das, 2012; Sharma and Ebadi, 2014).

Apart from a baseline assessment of trace element concentrations in loggerheads in the study area, in particular, the present study revealed high levels of cadmium in loggerheads stranded in the Portuguese mainland. However, the effects of high cadmium levels on turtle physiological processes and their implications on turtle survival and reproduction remain unknown. Recently, Finlayson et al. (2016) stressed the need for further research on marine turtle ecotoxicology, particularly using non-invasive *in vitro* methods. In addition, considering the transoceanic range of loggerhead turtles, ecotoxicology studies should be accompanied by the assessment of diet variation and health markers across life cycle stages, from pelagic juveniles to neritic adults.

Acknowledgments

The authors thank the members of Sociedade Portuguesa de Vida Selvagem for their assistance with data and sample collection and the staff at the Centres Científics i Tècnics of the Universitat de Barcelona. The authors also thank the Oceanário de Lisboa for their support. LN, MF and AM were supported by the Portuguese Foundation for Science and Technology (FCT) grants (SFRH/BD/51416/2011, SFRH/BD/30240/2006, SFRH/BPD/64889/2009, respectively). Also, LN, SM and CE were supported by CESAM UID/AMB/50017/2013 (FCT) co-funded by FCT/MEC and FEDER, within PT2020 and Compete 2020. Sample collection was partially supported by Project LIFE MarPro (Life09 NAT/PT/000038) co-funded by the European Commission and Project CetSenti RECI/AAG-GLO/0470/2012 (FCOMP-01-0124-FEDER- 027472), FCT/MCTES (PIDDAC) and FEDER - COMPETE (POFC).

References

- Aguilar, A., Borrel, A., Pastor, T., 1999. Biological factors affecting variability of persistent pollutant levels in cetaceans. *J. Cetacean Res. Manage. Spec. Issue* 1, 83–116.
- Aguirre, A.A., Balaz, G.H., Zimmerman, B., Galey, F.D., 1994. Organic contaminants and trace metals in the tissues of green turtles (*Chelonia mydas*) afflicted with fibropapillomas in the Hawaiian Islands. *Mar. Pollut. Bull.* 28, 109–114.
- Anan, Y., Kunito, T., Sakai, H., Tanabe, S., 2002. Subcellular distribution of trace elements in the liver of sea turtles. *Mar. Pollut. Bull.* 45, 224–229.
- Andreani, G., Santoro, M., Cottignoli, S., Fabbri, M., Carpenè, E., Isani, G., 2008. Metal distribution and metallothionein in loggerhead (*Caretta caretta*) and green (*Chelonia mydas*) sea turtles. *Sci. Total Environ.* 390, 287–294.
- Barreto, S., Marques, A., Teixeira, B., Vaz-Pires, P., Carvalho, M.L., Nunes, M.L., 2008. Essential elements and contaminants in edible tissues of European and American lobsters. *Food Chem.* 111, 862–867.
- Bjørndal, K.A., Bolten, A.B., Martins, H.R., 2000. Somatic growth model of juvenile loggerhead sea turtles *Caretta caretta*: duration of pelagic stage. *Mar. Ecol. Prog. Ser.* 202, 265–272.
- Breusch, T.S., Pagan, A.R., 1979. A simple test for heteroscedasticity and random coefficient variation. *Econometrica* 47, 1287–1294.
- Bruland, K.W., Knauer, G.A., Martin, J.N., 1978. Cadmium in northeast Pacific waters. *Limnol. Oceanogr.* 23, 618–625.
- Bustamante, P., Caurant, F., Fowler, S.W., Miramand, P., 1998. Cephalopods as a vector for the transfer of cadmium to top marine predators in the north-east Atlantic Ocean. *Sci. Total Environ.* 220, 71–80.
- Camacho, M., Orós, J., Boada, L.D., Zaccaroni, A., Silvi, M., Formigaro, C., López, P., Zumbado, M., Luzardo, O.P., 2013. Potential adverse effects of inorganic pollutants on clinical parameters of loggerhead sea turtles (*Caretta caretta*): results from a nesting colony from Cape Verde, West Africa. *Mar. Environ. Res.* 92, 15–22.
- Casale, P., 2015. *Caretta caretta* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2015. <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T83644804A83646294.en> eT83644804A83646294.
- Casale, P., Tucker, A.D., 2015. *Caretta caretta*. The IUCN Red List of Threatened Species 2015. eT3897A83157651. <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T3897A83157651.en>.
- Caurant, F., Bustamante, P., Bordes, M., Miramand, P., 1999. Bioaccumulation of cadmium, copper and zinc in some tissues of three species of marine turtles stranded along the French Atlantic coasts. *Mar. Pollut. Bull.* 38, 1085–1091.
- Ceriani, S.A., Meylan, A.B., 2015. *Caretta caretta* (North West Atlantic Subpopulation). The IUCN Red List of Threatened Species 2015. eT84131194A84131608.en. <http://dx.doi.org/10.2305/IUCN.UK.2015-4.RLTS.T84131194A84131608.en>.
- Chouvelon, T., Spitz, J., Caurant, F., Mendez-Fernandez, P., Autier, J., Lassus-Débat, A., Chappuis, A., Bustamante, P., 2012. Enhanced bioaccumulation of mercury in deep sea fauna from the Bay of Biscay (North-East Atlantic) revealed by stable isotope analysis. *Deep-Sea Res. Part I Oceanogr. Res. Pap.* 65, 113–124.
- D'Illo, S., Mattei, D., Blasi, M.F., Alimonti, A., Bogianni, S., 2011. The occurrence of chemical elements and POPs in loggerhead turtles (*Caretta caretta*): an overview. *Mar. Pollut. Bull.* 62, 1606–1615.
- Das, K., Debacker, V., Bouquegneau, J., 2000. Metallothioneins in marine mammals. *Cell. Mol. Biol.* 46, 283–294.
- Davis, S.R., Cousins, R.J., 2000. Metallothionein expression in animals: a physiological perspective on function. *J. Nutr.* 130, 1085–1088.
- Day, R.D., Christopher, S.J., Becker, P.R., Whitaker, D.W., 2005. Monitoring mercury in the loggerhead sea turtle, *Caretta caretta*. *Environ. Sci. Technol.* 39, 437–446.
- Day, R.D., Segars, L., Arendt, M.D., Lee, A.M., Peden-Adams, M.M., 2007. Relationship of blood mercury levels to health parameters in the loggerhead sea turtle (*Caretta caretta*). *Environ. Health Perspect.* 115, 1421–1428.
- Delgado, J., Nieto, J.M., Boski, T., 2011. Analysis of the spatial variation of heavy metals in the Guadiana Estuary sediments (SW Iberian Peninsula) based on GIS-mapping techniques. *Estuar. Coast. Shelf. Sci.* 88, 71–83.
- Eisler, R., 1985. Cadmium Hazards to Fish, Wildlife and Invertebrates: a Synoptic Review. Contaminant Hazard Reviews Report No 2, p. 30.
- Ferreira, M., Monteiro, S.S., Torres, J., Oliveira, I., Sequeira, M., López, A., Vingada, J., Eira, C., 2016. Biological variables and health status affecting inorganic element concentrations in harbour porpoises (*Phocoena phocoena*) from Portugal (western Iberian Peninsula). *Environ. Pollut.* 210, 293–302.
- Finlayson, K.A., Leusch, F.D.L., van der Merwe, J.P., 2016. The current state and future directions of marine turtle toxicology research. *Environ. Int.* 94, 113–123.
- Fossi, M.C., Panti, C., Guerranti, C., Coppola, D., Giannetti, M., Marsili, L., Minutoli, R., 2012. Are baleen whales exposed to the threat of microplastics? A case study of the Mediterranean fin whale (*Balaenoptera physalus*). *Mar. Pollut. Bull.* 64, 2374–2379.
- Franzellitti, S., Locatelli, C., Gerosa, G., Vallini, C., Fabbri, E., 2004. Heavy metals in tissues of loggerhead turtles (*Caretta caretta*) from the northwestern Adriatic Sea. *Comp. Biochem. Physiol. C* 138, 187–194.
- Frouin, H., Loseto, L.L., Stern, G.A., Haulena, M., Ross, P.S., 2012. Mercury toxicity in beluga whale lymphocytes: limited effects of selenium protection. *Aquat. Toxicol.* 109, 185–193.
- García-Fernández, A.J., Gómez-Ramírez, P., Martínez-López, E., Hernández-García, A., María-Mojica, P., Romero, D., Jiménez, P., Castillo, J.J., Bellido, J.J., 2009. Heavy metals in tissues from loggerhead turtles (*Caretta caretta*) from the southwestern Mediterranean (Spain). *Ecotoxicol. Environ. Saf.* 72, 557–563.
- Godley, B.J., Thompson, D.R., Furness, R.W., 1999. Do heavy metal concentrations pose a threat to marine turtles from the Mediterranean Sea? *Mar. Pollut. Bull.* 38, 497–502.
- Gordon, A.N., Pople, A.R., Ng, J., 1998. Trace metal concentrations in livers and kidneys of sea turtles from south-eastern Queensland, Australia. *Mar. Freshw. Res.* 49, 409–414.
- Guirlet, E., Das, K., 2012. Cadmium toxicokinetics and bioaccumulation in turtles: trophic exposure of *Trachemys scripta elegans*. *Ecotoxicology* 21, 18–26.
- Hamann, M., Godfrey, M.H., Seminoff, J.A., Arthur, K., Barata, P.C.R., Bjørndal, K.A., Bolten, A.B., Broderick, A.C., Campbell, L.M., Carreras, C., Casale, P., Chaloupka, M., Chan, S.K.F., Coyne, M.S., Crowder, L.B., Diez, C.E., Dutton, P.H., Epperly, S.P., FitzSimmons, N.N., Formia, A., Girondot, M., Hays, G.C., Cheng, I.J., Kaska, Y., Lewison, R., Mortimer, J.A., Nichols, W.J., Reina, R.D., Shanker, K., Spotila, J.R., Tomás, J., Wallace, B.P., Work, T.M., Zbinden, J., Godley, B.J., 2010. Global research priorities for sea turtles: informing management and conservation in the 21st century. *Endanger. Species Res.* 11, 245–269.
- Hamer, D.H., 1986. Metallothionein. *Ann. Rev. Biochem.* 55, 913–951.
- Hopkins, W.A., Rowe, C.L., Congdon, J.D., 1999. Elevated trace element concentrations and standard metabolic rate in banded water snakes (*Nerodia fasciata*) exposed to coal combustion wastes. *Environ. Toxicol. Chem.* 18, 1258–1263.
- Ikonopoulou, M.P., Olszowy, H., Hodge, M., Bradley, A.J., 2009. The effect of organochlorines and heavy metals on sex steroid-binding proteins *in vitro* in the plasma of nesting green turtles, *Chelonia mydas*. *J. Comp. Physiol. B* 179, 653–662.
- Jerez, S., Motas, M., Cánovas, R.Á., Talavera, J., Almela, R.M., Del Río, A.B., 2010. Accumulation and tissue distribution of heavy metals and essential elements in loggerhead turtles (*Caretta caretta*) from Spanish Mediterranean coastline of Murcia. *Chemosphere* 78, 256–264.
- Karouna-Renier, N.K., Snyder, R.A., Allison, J.G., Wagner, M.G., Rao, K.R., 2007. Accumulation of organic and inorganic contaminants in shellfish collected in estuarine waters near Pensacola, Florida: contamination profiles and risks to human consumers. *Environ. Pollut.* 145, 474–488.
- Kitana, N., Callard, I.P., 2008. Effect of cadmium on gonadal development in freshwater turtle (*Trachemys scripta*, *Chrysemys picta*) embryos. *J. Environ. Sci. Health A* 43, 262–271.
- Komoroske, L.M., Lewison, R.L., Seminoff, J.A., Deheyn, D.D., Dutton, P.H., 2011. Pollutants and the health of green sea turtles resident to an urbanized estuary in San Diego, CA. *Chemosphere* 84, 544–552.
- Lailson-Brito, J., Cruz, R., Dorneles, P.R., Andrade, L., Azevedo, A.F., Fragoso, A.B., Vidal, L.G., Costa, M.B., Bisi, T.L., Almeida, R., Carvalho, D.P., Bastos, W.R., Malm, O., 2012. Mercury-selenium relationships in liver of Guiana dolphin: the possible role of Kupffer cells in the detoxification process by tiemannite formation. *PLoS ONE* 7, e42162.

- López-Castro, M.C., Bjørndal, K.A., Kamenov, G.D., Zenil-Ferguson, R., Bolten, A.B., 2013. Sea turtle population structure and connections between oceanic and neritic foraging areas in the Atlantic revealed through trace elements. *Mar. Ecol. Prog. Ser.* 490, 233–246.
- Maffucci, F., Caurant, F., Bustamante, P., Bentivegna, F., 2005. Trace element (Cd, Cu, Hg, Se, Zn) accumulation and tissue distribution in loggerhead turtles (*Caretta caretta*) from the Western Mediterranean Sea (southern Italy). *Chemosphere* 58, 535–542.
- Mansfield, K.L., Putman, N.F., 2013. Oceanic habits and habitats *Caretta caretta*. In: Wyneken, J., Lohmann, K.J., Musick, J.A. (Eds.), *The Biology of Sea Turtles*, vol. III. CRC Press, Boca Raton, FL, USA.
- Mast, R.B., Hutchinson, B.J., Howgate, E., Pilcher, N.J., 2005. MTSG update: IUCN/SSC marine turtle specialist group host the 2nd burning issues assessment workshop. *Mar. Turt. Newsl.* 110, 13–15.
- Maulvault, A.L., Machado, R., Afonso, C., Lourenço, H.M., Nunes, M.L., Coelho, I., Langerholc, T., Marques, A., 2011. Bioaccessibility of Hg, Cd and as in cooked black scabbard fish and edible crab. *Food Chem. Toxicol.* 49, 2808–2815.
- McAloose, D., Newton, A.L., 2009. Wildlife cancer: a conservation perspective. *Nat. Rev. Cancer* 9, 517–526.
- Mendes, P., Eira, C., Torres, J., Soares, A.M.V.M., Melo, P., Vingada, J., 2008. Toxic element concentration in the Atlantic Gannet *Morus bassanus* (Pelecaniformes, sulidae) in Portugal. *Arch. Environ. Contam. Toxicol.* 55, 503–509.
- Méndez-Fernandez, P., Chouvelon, T., Bustamante, P., Ferreira, M., González, A.F., López, A., Moffat, C.F., Pierce, G.J., Read, F.L., Russell, M., Santos, M.B., Spitz, J., Vingada, J.V., Caurant, F., 2014. An assessment of contaminant concentrations in toothed whale species of the NW Iberian Peninsula: Part II. Trace element concentrations. *Sci. Total Environ.* 484, 206–217.
- Mil-Homens, M., Vale, C., Raimundo, J., Pereira, P., Brito, P., Caetano, M., 2014. Major factors influencing the elemental composition of surface estuarine sediments: the case of 15 estuaries in Portugal. *Mar. Pollut. Bull.* 84, 135–146.
- Monteiro, S.S., Torres, J., Ferreira, M., Marçalo, A., Nicolau, L., Vingada, J.V., Eira, C., 2016a. Ecological variables influencing trace element concentrations in Bottlenose dolphins (*Tursiops truncatus*, Montagu 1821) stranded in continental Portugal. *Sci. Total Environ.* 544, 837–844.
- Monteiro, S.S., Pereira, A.T., Costa, E., Torres, J., Oliveira, I., Bastos-Santos, J., Araujo, H., Ferreira, M., Vingada, J., Eira, C., 2016b. Bioaccumulation of trace element concentrations in common dolphins (*Delphinus delphis*) from Portugal. *Mar. Pollut. Bull.* 113, 400–407.
- Monteiro, S.S., Caurant, F., López, A., Cedeira, J., Ferreira, M., Vingada, J.V., Eira, C., Méndez-Fernandez, P., 2017. Sympatric *Globicephala* species: feeding ecology and contamination status based on stable isotopes and trace elements. *Mar. Ecol. Prog. Ser.* 563, 233–247.
- Nicolau, L., 2016. Influência de fatores antropogénicos na comunidade de tartarugas marinhas em águas continentais Portuguesas. PhD Thesis. Department of Biology, University of Aveiro, Portugal.
- Nicolau, L., Ferreira, M., Santos, J., Araújo, H., Sequeira, M., Vingada, J., Eira, C., Marçalo, A., 2016. Sea turtle strandings along the Portuguese mainland coast: spatio-temporal occurrence and main threats. *Mar. Biol.* 163, 1–13.
- Noel, L., Guérin, T., Kolf-Clauw, M., 2004. Subchronic dietary exposure of rats to cadmium alters the metabolism of metals essential to bone health. *Food. Chem. Toxicol.* 42, 1203–1210.
- Nuñez-Nogueira, G., Rainbow, P.S., 2005. Cadmium uptake and accumulation by the decapod crustacean *Penaeus indicus*. *Mar. Environ. Res.* 60, 339–354.
- Perrault, J., Wyneken, J., Thompson, L.J., Johnson, C., Miller, D.L., 2011. Why are hatching and emergence success low? Mercury and selenium concentrations in nesting leatherback sea turtles (*Dermochelys coriacea*) and their young in Florida. *Mar. Pollut. Bull.* 62, 1671–1682.
- R Core Team, 2015. R: a Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna. www.R-project.org.
- Reed, L.A., Pennington, P.L., Wirth, E., 2010. A survey of trace element distribution in tissues of stone crabs (*Menippe mercenaria*) from South Carolina Coastal Waters. *Mar. Pollut. Bull.* 60, 2297–2302.
- Ribeiro, A.R., Eira, C., Torres, J., Mendes, P., Miquel, J., Soares, A.M.V.M., Vingada, J., 2009. Toxic element concentrations in the Razorbill *Alca torda* (Charadriiformes, Alcidae) in Portugal. *Arch. Environ. Contam. Toxicol.* 56, 588–595.
- Roesijadi, G., 1992. Metallothioneins in metal regulation and toxicity in aquatic animals. *Aquat. Toxicol.* 22, 81–114.
- Royston, P., Remark, A.S., 1995. R94: a remark on algorithm AS 181: the W-test for normality. *J. R. Stat. Soc. Ser. C Appl. Stat.* 44, 547–551.
- Sabolic, I., Breljak, D., Skarica, M., Herak-Kramberger, C.M., 2010. Role of metallothionein in cadmium traffic and toxicity in kidneys and other mammalian organs. *Biometals* 23, 897–926.
- Saeki, K., Sakakibara, H., Sakai, H., Kunito, T., Tanabe, S., 2000. Arsenic accumulation in three species of sea turtles. *Biometals* 13, 241–250.
- Sakai, H., Ichihashi, H., Suganuma, H., Tatsukawa, R., 1995. Heavy metal monitoring in sea turtles using eggs. *Mar. Pollut. Bull.* 30, 347–353.
- Sakai, H., Saeki, K., Ichihashi, H., Suganuma, H., Tanabe, S., Tatsukawa, R., 2000. Species-specific distribution of heavy metals in tissues and organs of loggerhead turtle (*Caretta caretta*) and green turtle (*Chelonia mydas*) from Japanese coastal waters. *Mar. Pollut. Bull.* 40, 701–709.
- Sakulsak, N., 2012. Metallothionein: an overview on its metal homeostatic regulation in mammals. *Int. J. Morphol.* 30, 1007–1012.
- Seney, E.E., Musick, J.A., 2007. Historical diet analysis of Loggerhead sea turtles (*Caretta caretta*) in Virginia. *Copeia* 2, 478–489.
- Sharma, S., Ebadi, M., 2014. Significance of metallothioneins in aging brain. *Neurochem. Int.* 65, 40–48.
- Simoniello, P., Motta, C.M., Scudiero, R., Trinchella, F., Filosa, S., 2011. Cadmium-induced teratogenicity in lizard embryos: correlation with metallothionein gene expression. *Comp. Biochem. Physiol. C* 153, 119–127.
- Soto, J.M., Soares, J.T., Celini, A.A.O.S., Santos, R.C., 2005. Concentração de mercúrio total em tecidos de *Caretta caretta* (Linnaeus, 1758) (Reptilia, Cheloniidae) encahadas na costa sul do Rio Grande do Sul, Brasil. In: 2ª Jornada de Conservação e Pesquisa de Tartarugas Marinhas no Atlântico Sul Ocidental, Rio Grande, pp. 25–27.
- Storelli, M.M., Ceci, E., Marcotrigiano, G.O., 1998. Distribution of heavy metal residues in some tissues of *Caretta caretta* (Linnaeus) specimen beached along the Adriatic Sea (Italy). *Bull. Environ. Contam. Toxicol.* 60, 546–552.
- Storelli, M.M., Marcotrigiano, G.O., 2003. Heavy metal residues in tissues of marine turtles. *Mar. Pollut. Bull.* 46, 397–400.
- Storelli, M.M., Storelli, A., D'Addabbo, R., Marano, C., Bruno, R., Marcotrigiano, G.O., 2005. Trace elements in loggerhead turtles (*Caretta caretta*) from the eastern Mediterranean Sea: overview and evaluation. *Environ. Pollut.* 135, 163–170.
- Torrent, A., González-Díaz, O.M., Monagas, P., Orós, J., 2004. Tissue distribution of metals in loggerhead turtles (*Caretta caretta*) stranded in the Canary Islands. Spain. *Mar. Pollut. Bull.* 49, 854–874.
- Turoczy, N.J., Mitchell, B.D., Levings, A.H., Rajendram, V.S., 2001. Cadmium, copper, mercury, and zinc concentrations in tissues of the King Crab (*Pseudocarcinus gigas*) from southeast Australian waters. *Environ. Int.* 27, 327–334.
- TEWG, 2009. An Assessment of the Loggerhead Turtle Population in the Western North Atlantic Ocean. NOAA Technical Memorandum NMFS-SEFSC-575.
- van Geen, A., Boyle, E.A., Moore, W.S., 1991. Trace metal enrichments in waters of the Gulf of Cadiz, Spain. *Geochim. Cosmochim. Acta* 55, 2113–2191.
- Waalkes, M.P., Harvey, M.J., Klaassen, C.D., 1984. Relative in vitro affinity of hepatic metallothionein for metals. *Toxicol. Lett.* 20, 33–39.
- Wallace, B.P., Avens, L., Braun-McNeill, J., McClellan, C.M., 2009. The diet composition of immature loggerheads: insights on trophic niche, growth rates, and fisheries interactions. *J. Exp. Mar. Biol. Ecol.* 373, 50–57.
- Wallace, B.P., DiMatteo, A.D., Bolten, A.B., Chaloupka, M.Y., Hutchinson, B.J., Abreu-Grobois, F.A., Mortimer, J.A., Seminoff, J.A., Amorcho, D., Bjørndal, K.A., Bourjea, J., Bowen, B.W., Dueñas, R.B., Casale, P., Choudhury, B.C., Costa, A., Dutton, P.H., Fallabrino, A., Finkbeiner, E.M., Girard, A., Girondot, M., Hamann, M., Hurley, B.J., López-Mendilaharsu, M., Marcovaldi, M.A., Musick, J.A., Nel, R., Pilcher, N.J., Troeng, S., Witherington, B., Mast, R.B., 2011. Global conservation priorities for marine turtles. *PLoS ONE* 6, e24510.
- Wallace, B.P., DiMatteo, A.D., Hurley, J.B., Finkbeiner, E.M., Bolten, A.B., Chaloupka, M.Y., Hutchinson, B.J., Abreu-Grobois, A.A., Amorcho, D., Bjørndal, K.A., Bourjea, J., Bowen, B.W., Dueñas, R.B., Casale, P., Choudhury, B.C., Costa, A., Dutton, P.H., Fallabrino, A., Girard, A., Girondot, M., Godfrey, M.H., Hamann, M., López-Mendilaharsu, M., Marcovaldi, M.A., Mortimer, J.A., Musick, J.A., Nel, R., Pilcher, N.J., Seminoff, J.A., Troeng, S., Witherington, B., Mast, R.B., 2010. Regional Management Units for Marine Turtles: a novel framework for prioritizing conservation and research across multiple scales. *PLoS ONE* 5, e15465.
- Wu, J., Roshan, S., 2015. Cadmium in the North Atlantic: implication for global cadmium–phosphorus relationship. *Deep Sea Res. Part II* 116, 226–239.
- Wyneken, J., 2001. *The Anatomy of Sea Turtles*. US Department of Commerce NOAA Technical Memorandum NMFS-SEFSC-470.
- Zuur, A.F., Ieno, E.N., Elphick, S., 2010. A protocol for data exploration to avoid common statistical problems. *Methods Ecol. Evol.* 1, 3–14.
- Zuur, A.F., Ieno, E.N., Smith, G.M., 2007. *Analysing Ecological Data*. Springer, New York.