


## Mild *N*-Alkylation of Amines with Alcohols Catalyzed by Acetate Ruthenium Complexes

Walter Baratta,<sup>a</sup> Rosario Figliolia,<sup>a</sup> Salvatore Baldino,<sup>a</sup> Hans Günter Nedden,<sup>b</sup> Antonio Zanotti-Gerosa<sup>b</sup>

<sup>a</sup> Dipartimento DI4A, Università di Udine, Via Cotonificio 108, 33100 Udine, Italy; <sup>b</sup> Johnson Matthey Fine Chemicals Division 28 Cambridge Science Park, Milton Road, Cambridge, CB4 0FP, United Kingdom; E-mail: [walter.baratta@uniud.it](mailto:walter.baratta@uniud.it)

The formation of C-N bonds for the preparation of amines compounds is a reaction of high relevance for the synthesis of bulk and fine chemicals (1). The preparation of several drug molecules involves *N*-substitution transformations that are usually performed by reaction of amines with alkylating agents or via reductive amination. In this context, the catalytic *N*-alkylation of amines using environmentally friendly alcohols as alkylating reagents and affording water as only byproduct, is an attractive atom-economic way for the C-N bond formation (2,3).

We report here the straightforward synthesis of the carboxylate ruthenium complexes of formula Ru(OAc)<sub>2</sub>(diphosphane)(CO)<sub>n</sub> (n = 0, 1). These compounds are efficient catalysts for the *N*-alkylation of amines using primary alcohols under mild reaction conditions, with an alcohol / amine molar ratio of 10-100. Evidence has been provided that in catalysis a monohydride species is formed through an equilibrium reaction.



### References:

1. J. S. Carey, D. Laffan, C. Thomson, M. T. Williams, *Org. Biomol. Chem.* **2006**, *4*, 2337.
2. X. Ma, C. Su, Q. Xu, *Top. Curr. Chem. (Z)* **2016**, *374*, 27.
3. J. Leonard, A. J. Blacker, S. P. Marsden, M. F. Jones, K. R. Mulholland, R. Newton, *Org. Process Res. Dev.* **2015**, *19*, 1400.