

Contents lists available at [ScienceDirect](http://ScienceDirect.com)

Journal of Pediatric Surgery Case Reports

journal homepage: www.jpascalereports.com

Segmental intestinal dilatation associated with omphalocele



F. Molinaro ^a, R. Angotti ^{a,*}, E. Bindi ^a, C. Pellegrino ^a, M. Messina ^b, M. Aglianò ^b,
M. Messina ^a

^a Unit of Pediatric Surgery, Department of Medical Sciences, Surgical Sciences and Neurosciences, University of Siena, Italy

^b Section of Human Anatomy, Department of Medical Sciences, Surgical Sciences and Neurosciences, University of Siena, Italy

ARTICLE INFO

Article history:

Received 28 April 2017

Received in revised form

8 August 2017

Accepted 9 August 2017

Available online 12 August 2017

1. Introduction

Segmental intestinal dilatation (SID) is a rare congenital malformation, defined by the presence of a limited bowel dilatation with an abrupt transition between normal and dilated bowel, in absence of an extrinsic or intrinsic stop and with a normal pattern of neuronal plexus [1]. Concomitant SID and omphalocele is extremely rare, and sometimes associated with Meckel's diverticulum. We describe a patient with SID associated with perforated Meckel's diverticulum and omphalocele. After a literature review (PubMed database) we found 16 cases with the association between SID and omphalocele.

2. Case report

A 1-day-old boy (weight 3.1 kg), born through vaginal delivery in our hospital at 37 weeks of gestation, was evaluated for a leakage of meconium from the umbilical cord. An ultrasound performed at 33 weeks referred a cyst of the umbilical cord. An ultrasound revealed a minor omphalocele (2.5 cm of diameter). Based on clinical and radiological evaluation we put a suspicion on perforated omphalocele and we planned a surgery. At surgery a

perforated Meckel diverticulum within the abdominal defect (Fig. 1) and a dilated bowel loop at 20 cm from the ileocecal valve (Fig. 2) were found. Our surgical procedure included: (i) resection of Meckel diverticulum and end-to-end anastomosis, (ii) resection of segmental bowel dilatation and end-to-end anastomosis, (iii) umbilicoplasty (Figs. Figs 3–5). Postoperative course was uneventful. Patient started orally on 8th postoperative day. Stool pass was on 6th post-operative day. At last follow up, after 9 months, patient is healthy.

3. Discussion

Segmental intestinal dilatation (SID) is a rare congenital malformation. It usually involves the distal ileum, and any cases of colon and jejunum are described [2,3]. It is usually located at 12–25 cm from the ileocecal valve [4]. Our child, the first in our experience, had a SID about 20 cm from the ileocecal valve as it is described. Resection and anastomosis is the surgical approach described in literature as we performed in our child.

Swenson and Rathauer described firstly the intestinal dilatation segmental (SID) and established the diagnostic criteria that are used to date: (i) limited bowel dilation with three or four times higher AI: dimensions, (ii) sharp transition between normal and dilated intestinal (iii) any extrinsic or intrinsic causes of obstruction (iv) normal neuronal plexus (v) clinical presentation intestinal obstruction [5].

It can be associated with other malformations such as intestinal malrotation, herniated Bochdaleck, intestinal atresia, spina bifida and bladder exstrophy. However, the omphalocele is the most frequent association (85–90%).

Based on literature review, we found 16 cases of SID-omphalocele association, as shown in Table 1 [6–13].

In most of cases, as in our patient, the SID was found during surgery at birth, but in a few cases it was undiscovered until a

* Corresponding author. Department of Medical Sciences, Surgery and Neuroscience, Section of Pediatric Surgery, University of Siena, Policlinico Le Scotte, Viale Bracci 14, 53100 Siena, Italy.

E-mail address: rossellaangotti@me.com (R. Angotti).



Fig. 1. Minor omphalocele.



Fig. 2. Perforated Meckel's diverticulum within the minor omphalocele.



Fig. 3. Segmental intestinal dilatation (SID) at nearly 20 cm from the ileocecal valve.

symptomatic intestinal occlusion/sub-occlusion [14]. In two patients, SID was discovered due to the appearance of intestinal occlusion symptoms, after surgical closure of the omphalocele. In one patient, after the closing of the omphalocele and the execution of a biopsy, symptoms of intestinal occlusion appeared and so he was subjected to intestinal resection.

While the association SID/omphalocele is common and known, the co-presence of SID and Meckel's diverticulum is rare [15–17].




Fig. 4. Resection of segmental bowel dilatation.



Fig. 5. Umbilicoplasty.

Table 1 shows 5 patients with Meckel's diverticulum, of which 2 had a perforation.

These data show the importance of an exploration of the abdomen and bowel in the case of omphalocele.

The treatment of congenital segmental dilatation of intestine is resection of the dilated segment and end-to-end anastomosis [18]. Temporary stomas can be made in critically sick patients [19]. Our review shows only one patient treated with temporary ileostomy [14].

Histology of the resected segment is usually normal in most of the cases. We found 4 cases where histological examination showed mucous hypertrophy, 2 with mucous thinning, 1 with thickening, 2 with ectopic gastric mucosa (one in the context of the Meckel's diverticulum). In all cases, the ganglion cells were normally represented.

In conclusion, nevertheless SID is considered a less frequent malformations, it should be suspected in all patients with omphalocele.

The presence of a SID in association with omphalocele justifies the execution of a single surgical intervention with resection and intestinal anastomosis, in complete safety.

Table 1

This table summarizes the largest series of SID [11–18].

Series	N° of cases	Age of presentation	Symptoms	Site of dilatation	Associated malformations	Treatment	Histological findings
Molinaro et al.	1	Neonatal	Perforated Meckel's diverticulum in omphalocele (intestinal perforation)	Ileum	Meckel's diverticulum Omphalocele minor	Surgical resection and end-to-end anastomosis	Normal
Sam CJ, 2011	1	8 y.o.	Abdominal pain, bilious vomiting failure to pass stools	Ileum	Omphalocele, Meckel's diverticulum	Surgical resection and end-to-end anastomosis	Normal bowel musculature and normal ganglion cells with ectopic gastric mucosa in the region of Meckel's diverticulum
Wei CH et al., 2010	1	Neonatal	Omphalocele, vomiting	Ileum	Omphalocele	First operation, closure of omphalocele and intestinal biopsy, second operation surgical resection and end-to-end anastomosis	Normal
Thambidorai CR et al., 2009	1	Neonatal	Perforated Meckel's diverticulum in omphalocele (intestinal perforation)	Ileum	Meckel's diverticulum, Omphalocele minor	Surgical resection and end-to-end anastomosis	Normal
Brahim MB et al., 2006	8 (6 with omphalocele and SID)	7 Neonatal 1 y.o.	6 Omphalocele, 1 Neonatal obstruction, 1 Constipation	ileum	6 Omphalocele, 1 Esophageal Atresia, 1 Hirschsprung disease	Surgical resection and end-to-end anastomosis	3 hypertrophic muscularis, 4 normal, 1 heterotopic gastric mucosa 1 Thin muscularis
Basaran UN et al., 2005	1	Neonatal	Omphalocele	ileum	omphalocele	Ileostomy, Surgical resection and end-to-end anastomosis	Normal ganglion cells, Hypertrophic and thin muscularis layer
Balik E et al., 1993	1	4 y.o.	Abdominal distension, bilious vomiting, failure to pass stools	ileum	Small omphalocele	Surgical resection and end-to-end anastomosis	Normal ganglion cells, thickening of muscular layer
Bell MJ et al., 1982	7 (2 with omphalocele and SID)	2 Neonatal 5 Children	2 Omphalocele, 5 pallor, fatigue, and anemia, intestinal obstruction and recurrent abdominal pain	ileum	2 Omphalocele 2 Meckel's diverticulum	Surgical resection and end-to-end anastomosis	No luminal stenosis
Aboulola M et al., 1979	2	2 Neonatal	2 Omphalocele	ileum	2 Omphalocele, 2 intestinal malrotation	Surgical resection and end-to-end anastomosis	–

Conflict of interest

None declared.

Source of support

Nil.

References

- Ben Brahim M, Belghith M, Mekki M, Jouini R, Sahnoun L, Maazoun K, et al. Segmental dilatation of the intestine. *J Pediatr Surg* 2006;41(6):1130–3.
- Fang SB, Lee HC, Sheu JC, Chen BF. Segmental dilatation of the ileum as an unusual cause of gastrointestinal bleeding: report of one case. *Acta Paediatr Sin* 2000;41(2):93–7.
- Brandigi E, Molinaro F, Meucci D, Sica M, Bindi E, Messina M. Ileal duplication cyst in new born complicated with volvulus: 2 cases report and literature review. *Chirurgia (Turin)* 2014;27(5):297–9.
- Ratan S, Rattan K, Kalra R, Maheshwari J, Parihar D, Ratan J. Omphalomesenteric duct cyst as a content of omphalocele. *Indian J Pediatr* 2007;74(5):500–2.
- Swenson O, Rathauser F. Segmental dilatation of the colon. *Am J Surg* 1959;97(6):734–8.
- Sam CJ. Segmental ileal dilatation in a child. *Trop Gastroenterol* 2011;32(3):221–3.
- Wei CH, Sheu JC. Concomitant segmental intestinal dilatation and Omphalocele. *Formos J Surg* 2011;44:168–70.
- Thambidorai CR, Arief H, Noor Afidah MS. Ileal perforation in segmental intestinal dilatation associated with omphalocele. *Singap Med J* 2009;50(12):e412.
- Brahim Mohamed Ben, Belghith Mohsen, Mekki Mongi, Jouini Riadh, Sahnoun Lassaad, Maazoun Kaies, et al. Segmental dilatation of the intestine. *J Pediatr Surg* 2006;41:1130–3.
- Basaran UN, Sayin C, Oner N, Celtik C. Segmental intestinal dilatation associated with omphalocele. *Pediatr Int* 2005;47:227–9.
- Balik E, Taneli C, Yazici M, Demircan M, Herek O. Segmental dilatation of intestine: a case report and review of the literature. *Eur J Pediatr Surg* 1993 Apr;3(2):118–20.
- Bell MJ, Ternberg JL, Bower RJ. Ileal dysgenesis in infants and children. *J Pediatr Surg* 1982 Aug;17(4):395–9.
- Aboulola M, Boukheloua B, Asselah F. Segmental dilatation of the ileum associated with an omphalocele and a common mesentery. Discussion around 2 cases : chance or morphogenically "related"? *Chir Pediatr* 1979;20(3):197–202.
- Basaran U, Sayin C, Oner N, Celtik C. Segmental intestinal dilatation associated with omphalocele. *Pediatr Int* 2005;47(2):227–9.
- Bertozzi M, Melissa B, Radicioni M, Magrini E, Appignani A. Symptomatic Meckel's diverticulum in newborn: two interesting additional cases and review of literature. *Pediatr Emerg Care* 2013 Sep;29(9):1002–5.
- Sakaguchi T, Hamada Y, Masumoto K, Taguchi T. Segmental dilatation of the intestine: results of a nationwide survey in Japan. *Pediatr Surg Int* 2015;31(11):1073–6.
- Messina M, Ferrucci E, Meucci D, Di Maggio G, Molinaro F, Buonocore G. Lit-tre's hernia in newborn infants: report of two cases. *Pediatr Surg Int* 2005 Jun;21(6):485–7. Epub 2005 Apr 12.
- Rai BK, Mirza B, Hashim I, Saleem M. Varied presentation of congenital segmental dilatation of the intestine in neonates: report of three cases. *J Neonatal Surg* 2016 Oct 10;5(4):55.
- Ferrara F, Angotti R, Burgio A, Di Maggio G, Molinaro F, Messina M. Ileostomy in extremely low birth weight and premature neonates. *Minerva Pediatr* 2013 Aug;65(4):411–5.