



Emerging Contaminants Such as Perfluorinated Compounds: Toxicity and Oxidative Stress

Paola Irato*, Sara Trabucco and Laura Tallandini

Department of Biology, University of Padua, Italy

Corresponding author: Irato P, Department of Biology, University of Padua, Italy, Tel: +39 049 8276315; E-mail: paola.irato@unipd.it

Rec date: Feb 13, 2017; Acc Date: Mar 1, 2017; Pub date: Mar 08, 2017

Copyright: © 2017 Irato P, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Letter to Editor

Perfluoroalkyl and polyfluoroalkyl substances (PFASs) are widely applied in a variety of industrial and consumer products since 1949 and comprise a diverse group of chemicals with significant economic value. Due to their unique properties, mainly given by the strong C-F bond, they have been used in a wide variety of technological and industrial application, such as surfactants in fluoropolymer production, metal plating, aqueous film-forming foams, paper, textile, and household products [1]. PFASs are highly stable and difficult to degrade in the environment [2]. They also have the capability for long-range transport through the atmosphere and water [1]. Their presence in soils raised concerns over their potential ecological risks in terrestrial food chain [2].

Several episodes of pollution took place in the phreatic zone of medium-lower Agno Valley (Vicenza Province), between Lessini Mountains and the contiguous plain in Southern direction until the area of Colli Berici and the Municipality of Vicenza, due to the high density of production sites and industries. Nowadays the total catchment area involves three Provinces (Vicenza, Padova and Verona).

According to the length of the carbon chain, PFAS can be divided into: long chain PFAS (>C8), short-chain PFAS (C4-C7) and ultra-short chain PFAS (C2-C4). Between the whole PFAS, two sub-categories are of environmental interest: perfluoroalkyl sulfonate sub-category that includes also acid salts and precursors (PFOS is the most important) and perfluoroalkyl carboxylate including perfluorooctanoic acid (PFOA) and homologues and their acid salts and precursors. Other perfluoroalkyl chemicals (PFCs) include perfluorobutane sulfonate (PFBS).

Perfluorooctanoic acid (PFOA; C7 fluorocarbon) has a low bioaccumulation potential and is relatively similar among species from different trophic levels. This low bioaccumulation potential may be driven by the short perfluorocarbon chain length and particular functional group [1]. Unlike most other persistent and bioaccumulative organic toxicants, PFOA is water-soluble and does not bind well to soil, allowing for easy transportation through and contamination of human drinking water [3].

PFOA and other PFAS with shorter and ultra-shorter carbon chain are considered emerging contaminants and are included in the national environmental legislations. A high number of studies report high levels of PFAS in the environment, as well as in human samples such as blood, tissues, and breast milk [4]. PFAS are not stored in body fat. Several studies on laboratory animals indicate that PFOA exposure can impact male fertility. Altered polyunsaturated fatty acid compositions, concomitant with excessive oxidative stress, as demonstrated by increased malonaldehyde and decreased glutathione peroxidase, were observed [5-20].

References

- Adams JM, Gallagher JM, Donnison IS (2009). Fermentation study on Saccharina latissima for bioethanol production considering variable pretreatments. *J Appl Phycol* 21: 569-574.
- AOAC. (1990). Official methods of analysis. 15th edn. Assoc. of official analytical chemists. Washington DC, USA.
- Balat M (2007). Global bio-fuel processing and production trends. *Energy Explore Exploit* 25, pp. 195-218.
- Balat M, Balat H, Oz C, (2008). Progress in bioethanol processing. *Progress in Energy and Combustion Sci* 34: 551-573.
- Bothast RJ, Schlicher MA (2005) Biotechnological processes for conversion of corn into ethanol. *Appl Microbiol Biotechnol* 67: 19-25.
- Bouthilet RJ, Caputi A, Ueda M, (1961). Analysis of ethanol in wine by gas liquid partition chromatography. *J Assoc Off Anal Chem* 44: 410-414.
- Caputi A, Ueda M, Brown T (1968). Spectrophotometric determination of ethanol in wine. *Am J Enol Vitic*. 19: 160-165.
- Chinese National Standard (1983). Method of test for beverage determination of alcohol content. Chinese National Standard, Taiwan. Code No. 10292 (N6181).
- Demirbas MF, Balat M, (2006). Recent advances on the production and utilization trends of bio-fuels: A global perspective. *Energy Convers Manage* 47: 2371-2381.
- Flora G, Maria Victorial Rani S, (2013). GC-MS analysis of *Acanthophora spicifera*. *Int J Pharm. Bio. Sci* 4: 649-653.
- Hahn-Hagerdal B, Karhumaa K, Fonseca C, Spencer-Martins I, Gorwa-Grauslund MF (2007). Towards industrial pentose-fermenting yeast strains. *Appl Microbiol. Biotechnol* 74: 937-953.
- Hall DO, House JI, Scrase JI (2000). Introduction: Overview of Biomass Energy, In: *Industrial uses of Biomass Energy*, Rosillo-Calle, F. and Bajay, R. H. (Eds.), Taylor and Francis, London, United Kingdom.
- Horn SJ (2000). Bioenergy from brown seaweeds. Thesis Ph.D. Department of Biotechnology, Norwegian University of Science and Technology, Trondheim, Norway.
- Karakashev D, Thomsen AB, Angelidaki I (2007) Anaerobic biotechnological approaches for production of liquid energy carriers from biomass. *Biotechnol. Lett* 29:1005-1012.
- Koshy BE, Pandey FK, Bhatnagar T (2014). Quantitative estimation of bioethanol produced from lignocellulosic and household wastes. *International Journal of Life Sciences Research*, 2: 130-145.
- McHugh DJ (2003) A guide to the seaweed industry, Food and Agricultural Organization: 106.
- Richards OW (1928). The growth of the yeast *Saccharomyces cerevisiae*: the growth curve, its mathematical analysis and the effects of temperature on the yeast growth, *Annals of Botany* 42: 271-283.
- Santhanam RN, Remanathan G, Jagathusan (1990). *Coastal Aquaculture in India*. C.B.S. Publishers and Distributors. pp: 159-162.
- Stackler R, Christensen EN (1974). Quantitative determination of ethanol in wine by gas chromatography. *Am J Enol. Vitic* 25: 202-207.
- Zhao XQ, Bai FW (2009) Yeast flocculation: New story in fuel ethanol production. *Biotechnol Adv* 27: 849-856.

