

Technical University of Denmark



An emerging European Doppler lidar network for meteorological applications

O'Connor, Ewan; Hirsikko, Anne ; Halios, Christos; Gryning, Sven-Erik; Leinweber, Ronny ; Manninen, Antti; Marke, Tobias; Petersen, Nina; Preissler, Jana ; Päsche, Eileen; Saeed, Umar; schween, jan; Shu, Yang; Suomi, Irene; Tuononen, Minttu; Vakkari, Ville; Thobois, Ludovic; Pearson, Guy; Dabas, Alain; Buehl, Johannes

Published in:
EMS Annual Meeting Abstracts

Publication date:
2017

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
O'Connor, E., Hirsikko, A., Halios, C., Gryning, S-E., Leinweber, R., Manninen, A., ... Buehl, J. (2017). An emerging European Doppler lidar network for meteorological applications. In EMS Annual Meeting Abstracts (Vol. 14). [745]

DTU Library
Technical Information Center of Denmark

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



An emerging European Doppler lidar network for meteorological applications

Ewan O'Connor (1,2), Anne Hirsikko (1), Christos Halios (2), Sven-Erik Gryning (3), Ronny Leinweber (4), Antti Manninen (5), Tobias Marke (6), Nína Petersen (7), Jana Preissler (8), Eileen Päschrke (4), Umar Saeed (9), Jan Schween (6), Yang Shu (10), Irene Suomi (1), Minttu Tuononen (1), Ville Vakkari (1), Ludovic Thobois (11), Guy Pearson (12), Alain Dabas (13), and Johannes Buehl (14)

(1) FMI, Finland, (2) University of Reading, UK, (3) DTU, Denmark, (4) DWD, Germany, (5) University of Helsinki, Finland, (6) University of Köln, Germany, (7) IMO, Iceland, (8) NUIG, Ireland, (9) Universitat Politècnica de Catalunya, Spain, (10) Reykjavik University, Iceland, (11) Leosphere, France, (12) Halo Photonics, UK, (13) Meteo France, France, (14) TROPOS, Germany

Doppler lidars are capable of providing winds and turbulent parameters, such as dissipation rate at high spatial and temporal resolution. The implications for boundary-layer retrieval are very exciting as this will allow the diagnosis of various aspects of the dynamical boundary layer, such as low-level jets and gusts, together with boundary layer mixing and height.

Within the EU Cost Action TOPROF (Towards operational ground based profiling with ceilometers, Doppler lidars and microwave radiometers for improving weather forecasts), the Working Group on Doppler lidar has advanced the objective of creating a European Doppler lidar network for meteorological applications, through coordinating the Doppler lidars performing meteorological research in Europe. This group is tasked with assessing the performance of the various instruments in use, and establishing operational procedures for the provision of quality-controlled products for a wide variety of end-users, including operational evaluation and assimilation of winds and BL classification in NWP models.

Here, we provide an overview of the results produced during TOPROF. These include: in-depth characterisation of instrument performance, together with post-processing routines to improve sensitivity and uncertainty analysis; designing suitable scanning strategies to optimise retrievals of winds and turbulence for multiple purposes; flexibility for different scan selections in different locations. The harmonisation of data from different instrument types and scanning sequences then permits unified retrievals such as boundary layer type classification and low-level jet detection to be applied across the network.